Intragenomic conflict over soldier allocation in polyembryonic parasitoid wasps

Petri Rautiala^{1,*} & Andy Gardner²

1. University of Jyväskylä, Department of Biological and Environmental Science, PO Box 35, FI-40014 University of Jyväskylä, Finland & Centre of Excellence in Biological Interactions. Email: petri.rautiala@jyu.fi

2. School of Biology, University of St Andrews, St Andrews KY16 9TH, United Kingdom. Email: andy.gardner@st-andrews.ac.uk

* Corresponding author

Short Title: Intragenomic conflict over soldiering

Article Type: E-Note.

Keywords: Genetic conflict, genomic imprinting, kin selection, parent-of-origin effects, sex allocation, spiteful behaviour.

Colour Figures: Figure 1, Figure 2.

Online Enhancements: None.

The authors wish to be identified to the reviewers.

26 **Abstract** | Understanding the selection pressures that have driven the evolution of sterile insect castes
27 has been the focus of decades of intense scientific debate. An amenable empirical testbed for theory on
28 this topic is provided by the sterile soldier caste of polyembryonic parasitoid wasps. The function of
29 these soldiers has been a source of controversy, with two basic hypotheses emerging: the "brood-
30 benefit" hypothesis that they provide an overall benefit for their siblings; and the "sex-ratio-conflict"
31 hypothesis that the soldiers mediate a conflict between brothers and sisters, by killing their opposite-sex
32 siblings. Here, we investigate the divergent sex-ratio optima of a female embryo's maternal-origin and
33 paternal-origin genes, to determine the potential for, and direction of, intragenomic conflict over
34 soldiering. We then derive contrasting empirically-testable predictions, concerning the patterns of
35 genomic imprinting that are expected to arise out of this intragenomic conflict, for the brood-benefit
36 versus sex-ratio-conflict hypotheses of soldier function.

Introduction

Understanding the selection pressures that have driven the evolution of sterile insect castes has been the focus of decades of intense scientific debate (Hamilton 1964, 1972; Wilson 2005; Foster et al 2006; Boomsma 2007, 2009, 2013; Nowak et al 2010; Abbott et al 2011; Gardner et al 2012; Liao et al 2015). An amenable empirical testbed for theory on this topic is provided by the sterile soldier caste of polyembryonic parasitoid wasps of the genus *Copidosoma* (Cruz 1986; Strand 2009). These are wasps that inject their eggs into the bodies of other insects, and whose young devour their hosts from the inside before emerging as adults to mate and find new hosts to parasitize. A curious aspect of their biology is that each egg proliferates clonally to give rise to a very large number of genetically-identical embryos, which then compete for resources within the host; such polyembryony has arisen independently in four families of the parasitoid Hymenoptera: the Braconidae, Platygasteridae, Dryinidae and Encyrtidae, with *Copidosoma* belonging to the latter (Ivanova-Kasas 1972). An even curiouser aspect of their biology is that some of these embryos - mostly, but not solely, females - develop precociously as soldier larvae that patrol the interior of the host and do not emerge as reproductive adults.

The function of these soldiers has been a source of controversy, with two basic hypotheses emerging (Gardner et al 2007a). First, the "brood-benefit" hypothesis suggests that their primary function is to provide an overall benefit for their siblings, either by macerating host tissues to facilitate release of nutrients (Silvestri 1906) or, more likely, by attacking the young of other parasitoids that may also be present in the host (Cruz 1981; Strand et al 1990; Harvey et al 2000; Giron et al 2004). Second, the "sex-ratio-conflict" hypothesis suggests that the soldiers' primary function is to mediate a conflict between brothers and sisters, over the sex ratio of the reproductive adults that will emerge from the host, by killing their opposite-sex siblings (Godfray 1992; Hardy 1994; Ode & Hunter 2002; Giron et al 2004). Gardner et al (2007a) provided mathematical analyses of both putative functions and showed that, if individuals of either sex are equally capable of developing and acting as soldiers, then male-

62 biased soldiering is expected under the brood-benefit hypothesis and female-biased soldiering is
63 expected under the sex-ratio-conflict hypothesis, because females value their brothers relatively less
64 than males value their sisters. Accordingly, since female-biased soldiering is observed (Doutt 1947;
65 Grbic et al 1992; Ode & Strand 1995; Giron et al 2004; Keasar et al 2006), Gardner et al's (2007a)
66 analysis lends support to the idea that the soldiers have a sex-ratio conflict function.

67 However, an alternative explanation for the observed sex bias in soldiering is that the sexes may
68 differ in their intrinsic ability to develop and behave as soldiers (Doutt 1947; Gardner et al 2007a). This
69 view mirrors the more general understanding of why sterile workers among the social Hymenoptera are
70 always female: although this sex-bias was traditionally attributed to relatedness asymmetries arising
71 from haplodiploid inheritance (Hamilton 1964, 1972), more recent empirical analysis instead supports
72 the idea that females are simply better workers, being already equipped with adaptations for nursing
73 young owing to the presence of maternal but not paternal care among the ancestors of this insect group
74 (Ross et al 2013). Accordingly, the empirically-observed female-biased soldiering of polyembryonic
75 parasitoid wasps need not rule out a primarily brood-benefit function for soldiers.

76 Here, we develop a further set of empirically-testable predictions, that may be used to
77 discriminate between the brood-benefit and sex-ratio-conflict hypotheses for soldier function, and that
78 do not depend upon the relative preadaptation of females and males to soldiering. In particular, we
79 follow up on West's (2009, p287; see also Wild & West 2009) suggestion that there may be an
80 intragenomic conflict of interests, between a female's maternal-origin and paternal-origin genes, over
81 the decision to develop as a soldier, and that this may drive the evolution of parent-of-origin-specific
82 patterns of gene expression, i.e. "genomic imprinting" (Moore & Haig 1991; Haig 1997). We first adapt
83 the mathematical model of Gardner et al (2007a) to investigate the sex-ratio optima of a female's
84 maternal-origin and paternal-origin genes, to ascertain the potential for, and direction of, intragenomic
85 conflict over soldiering. We then derive contrasting predictions, as to the patterns of genomic imprinting

86 that are expected to arise out of this intragenomic conflict, for the brood-benefit versus sex-ratio-conflict
87 hypotheses with regards to soldier function.

88

89 **Model and analysis**

90

91 *Basic model* - Following Gardner et al (2007a), whose model focuses mostly upon the biology of
92 *Copidosoma floridanum*, we consider that a single foundress wasp injects two eggs - one fertilised (i.e.
93 female) and one unfertilised (i.e. male) - into a parasitised host, with each egg proliferating clonally to
94 give a large number of embryos, such that each embryo is genetically identical to its same-sex brood
95 mates and is related to its opposite-sex brood mates according to the usual brother-sister relationship.
96 Some proportion of female and male embryos develop as soldiers, which modulates the number and sex
97 ratio of the embryos that will successfully emerge from the host as adults. After emerging, a proportion
98 $1-d_f$ of females and a proportion $1-d_m$ of males remain close to their host, where they form a mating
99 group, whereas a proportion d_f of females and a proportion d_m of males disperse to other mating groups.
100 Mating then occurs at random within each mating group, after which all males perish and the mated
101 females parasitize the next generation of hosts.

102

103 *Inclusive fitness* - We take an inclusive-fitness approach to capture the evolutionary interests of each
104 member of the family unit. In particular, we express the inclusive fitness of any actor A as:

$$105 \quad H_A = N_m M p_{mA} + 2N_f p_{fA}, \quad (1)$$

106 where: N_m is the number of males emerging from a focal host; M is the average number of successful
107 matings enjoyed by each of these males; p_{mA} is the consanguinity of each of these males to the actor (i.e.
108 the probability that a gene drawn at random from a male is identical by descent with one drawn at
109 random from the actor, from the same locus; Bulmer 1994); N_f is the number of females emerging from
110 the focal host; p_{fA} is the consanguinity of each of these females to the actor; and the factor 2 reflects that

each female has twice the reproductive value of the male with whom she mates, under haplodiploidy (Hamilton 1972). The average number of successful matings per local male may itself be expressed as $M = d_m(\bar{N}_f/\bar{N}_m) + (1 - d_m) [(1 - d_f)N_f + d_f\bar{N}_f]/[(1 - d_m)N_m + d_m\bar{N}_m]$, where $\bar{N}_m$ and $\bar{N}_f$ are the average number of males and females, respectively, emerging from each host in the population.

Intragenomic conflict over sex ratio - Different actors may have different preferences with respect to the sex ratio of the emerging adults, and this disagreement may be investigated by consideration of the inclusive fitness function. Specifically, defining $N = N_f + N_m$ and $z = N_m/N$, and hypothetically assigning the actor full control over the sex ratio z , their inclusive fitness may be written as:

$$H_A = Nz \left[d_m \frac{1 - \bar{z}}{\bar{z}} + (1 - d_m) \frac{(1 - d_f)(1 - z) + d_f(1 - \bar{z})}{(1 - d_m)z + d_m\bar{z}} \right] p_{mA} + 2N(1 - z)p_{fA}, \quad (2)$$

where $\bar{z}$ is the population average sex ratio. The actor prefers a higher than population average sex ratio whenever their marginal inclusive fitness is positive at that population average, i.e. when $\partial H_A / \partial z|_{z=\bar{z}} = [(2 - d_m)d_m/\bar{z} + (d_f - d_m)(1 - d_m) - 1]Np_{mA} - 2Np_{fA} > 0$. Accordingly, setting marginal inclusive fitness equal to zero, and solving for $z = \bar{z} = z_A$, yields the sex ratio optimum for actor A:

$$z_A = \frac{(2 - d_m)d_m}{1 + (1 - d_m)(d_m - d_f) + 2p_{fA}/p_{mA}}. \quad (3)$$

Different actors may have different sex ratio optima because they may have different consanguinities to the female and male brood (p_{fA} & p_{mA}). The coefficients of consanguinity for the different family members are listed in table 1. Equations (3)-(5) of Gardner et al (2007a; see also Gardner et al 2007b) provide the sex ratio optima from the perspective of a female embryo, a male embryo and their mother, respectively: these are recovered by substituting the appropriate coefficients of consanguinity into equation (3), above, and are illustrated in figure 1.

If all males disperse prior to mating ($d_m = 1$), then there is full outbreeding and no local mate competition. In this case, mothers prefer an equal sex allocation ($z = 1/2$): although daughters have twice the reproductive value of sons under haplodiploid inheritance, when the sex ratio is unbiased, sons are

twice as consanguinous to their mothers than are daughters, owing to all of the son's genes deriving from his mother, and these two effects exactly cancel to recover the usual equal sex allocation optimum (Fisher 1930; Hamilton 1967; Gardner 2014). In contrast: female embryos, being clonally related to their sisters, prefer a female-biased sex allocation ($z < \frac{1}{2}$); male embryos, being clonally related to their brothers, prefer a male-biased sex allocation ($z > \frac{1}{2}$); and fathers, being entirely unrelated to the sons of their mating partners, prefer all offspring to be female ($z = 0$). If there is incomplete dispersal of males prior to mating ($0 < d_m < 1$) then this results in local mate competition (i.e. related males competing for mating opportunities), which results in mothers, daughters and sons all preferring a sex-allocation that is relatively female biased (lower z). Moreover, if there is also incomplete dispersal of females prior to mating ($d_f < 1$) then, owing to inbreeding, fathers are related to the sons of their mating partners and accordingly prefer nonzero investment into males ($z > 0$). Finally, in the limit of the complete absence of male dispersal prior to mating ($d_m \rightarrow 0$), local mate competition is complete and all parties prefer vanishingly small investment into males ($z \rightarrow 0$; Hamilton 1967).

An intragenomic conflict over sex ratio arises when a female embryo's maternal-origin and paternal-origin genes have different sex-ratio optima. Substituting the appropriate parent-of-origin-specific coefficients of consanguinity (table 1) into equation (3) yields the optima:

$$z_{f|M} = \frac{(2 - d_m)d_m}{3 + (1 - d_m)(d_m - d_f)} \quad (4)$$

for the female embryo's maternal-origin genes; and

$$z_{f|P} = \frac{(1 - d_f)(1 - d_m)(2 - d_m)d_m}{3 + d_f^2(1 - d_m)^2 + (1 - d_m)(2d_m + d_f d_m^2) + d_m^3} \quad (5)$$

for her paternal-origin genes. These distinct sex ratio optima are illustrated in figure 1.

We find that the female embryo's maternal-origin genes prefer a greater proportion of males among the emerging adults than do the female embryo's paternal-origin genes ($z_{f|M} > z_{f|P}$ for all $d_i, d_m > 0$). This is because the female embryo is more related to her brothers through her mother than through her father, on account of the entire brood having the same mother but only the female embryos having a

159 father. Note that, although the female embryo's father makes no direct genetic contribution to her
 160 brothers, her paternal-origin genes are nevertheless consanguinous with her brother's genes to the extent
 161 that the female's mother and father were relatives (i.e. insofar as there is inbreeding). Accordingly, the
 162 female's paternal-origin genes need not always favour an entirely female-biased sex ratio. Also note
 163 that, whilst the sex-ratio optimum for the female embryo's maternal-origin genes is distinct from that of
 164 her mother's genes, the sex-ratio optimum for the female embryo's paternal-origin genes is exactly the
 165 same as that of her father's genes. This is because, whilst the female embryo's maternal genome is
 166 genetically distinct from her mother's genome (the former is a random haploid subset of of the latter's
 167 diploid set of genes), the female embryo's paternal genome is genetically identical to her father's
 168 genome (*de novo* mutation aside; her father has only a single haploid genome to contribute to each of his
 169 daughters). More generally, this point clarifies that conflicts between maternal-origin versus paternal-
 170 origin genes are conceptually distinct from conflicts between an individual's parents.

171

172 *Intragenomic conflict over soldiering* - We now investigate the evolutionary interests of a female's
 173 maternal-origin versus paternal-origin genes with respect to soldiering. We consider that the proportion
 174 x of female embryos developing as soldiers modulates both the sex ratio z and the number N of adults
 175 emerging from the host. The male mating success M is modulated by the number of males and females
 176 emerging from the host, and therefore by x (and, to be precise, also by the population average $\bar{x}$).
 177 Rewriting equation (1) as $H_A(x) = N_m(x)M(x)p_{mA} + 2N_f(x)p_{fA}$, to make this dependency explicit,
 178 any actor favours a greater than the population average female soldiering when $\partial H_A / \partial x|_{x=\bar{x}} =$
 179 $p_{mA} \partial(N_m M) / \partial x|_{x=\bar{x}} + 2p_{fA} \partial N_f / \partial x|_{x=\bar{x}} > 0$. And so, upon the assumption that the focal individual's
 180 and population-average probability of developing as a soldier are both at the female's own optimum (i.e.
 181 $x = \bar{x} = x_f$ which is hereafter abbreviated “*” for ease of presentation), and that this takes an
 182 intermediate value (i.e. $0 < x_f < 1$), we may write

$$\left. \frac{\partial(N_m M)}{\partial x} \right|_* = -2 \frac{p_{ff}}{p_{mf}} \left. \frac{\partial N_f}{\partial x} \right|_* \quad (6)$$

It follows that, from the perspective of the female embryo's maternal-origin genes, the marginal inclusive fitness is $\partial H_{f|M}/\partial x|_* = -2p_{mf|M}(p_{ff}/p_{mf})\partial N_f/\partial x|_* + 2p_{ff|M}\partial N_f/\partial x|_*$
 $= -2(p_{mf|M}/p_{mf} - 1)p_{ff}\partial N_f/\partial x|_*$ and, from the perspective of the female embryo's paternal-origin genes, the marginal inclusive fitness is $\partial H_{f|P}/\partial x|_* = -2p_{mf|P}(p_{ff}/p_{mf})\partial N_f/\partial x|_* + 2p_{ff|P}\partial N_f/\partial x|_* =$
 $-2(p_{mf|P}/p_{mf} - 1)p_{ff}\partial N_f/\partial x|_*$, where we have made use of the fact that $p_{ff|M} = p_{ff|P} = p_{ff}$, i.e. the consanguinity of a female to herself or to her clonal sister is the same for her maternal-origin and her paternal-origin genes. In the appendix, found online, we show that $\partial N_f/\partial x|_* < 0$ under the brood-benefit hypothesis and that $\partial N_f/\partial x|_* > 0$ under the sex-ratio-conflict hypothesis. Since $p_{mf|P} < p_{mf} < p_{mf|M}$, it follows that $\partial H_{f|P}/\partial x|_* < 0 < \partial H_{f|M}/\partial x|_*$ under the brood-benefit hypothesis, and $\partial H_{f|M}/\partial x|_* < 0 < \partial H_{f|P}/\partial x|_*$ under the sex-ratio-conflict hypothesis.

We have found that the female embryo's maternal-origin genes are favoured to increase her probability of developing as a soldier under the brood-benefit hypothesis and they are favoured to decrease her probability of developing as a soldier under the sex-ratio-conflict hypothesis, whereas her paternal-origin genes are favoured to decrease her probability of developing as a soldier under the brood-benefit hypothesis and they are favoured to increase her probability of developing as a soldier under the sex-ratio-conflict hypothesis. That is, we predict an intragenomic conflict of interests with respect to female soldier development, with a direction that depends on the function of the soldier caste.

Genomic imprinting - Having ascertained the existence and direction of the conflict of interest between the female embryo's maternal-origin genes and her paternal-origin genes with respect to her probability of developing as a sterile soldier, we now elaborate predictions for patterns of genomic imprinting (figure 2). For loci whose gene products modulate a female embryo's probability of developing as a soldier we expect there to be a disagreement between her maternal-origin genes and her paternal-origin

207 genes over the optimal level of gene expression and, where parent-of-origin-specific gene expression is
208 feasible, we expect such genomic imprinting to evolve as a consequence of this disagreement.
209 According to the "loudest voice prevails" principle, the gene that prefers a lower level of expression
210 ultimately silences itself whilst the gene that prefers a greater level of expression ultimately wins the
211 conflict and expresses at a level corresponding to its optimum (Haig 1996; Úbeda & Haig 2003).

212 Under the brood-benefit hypothesis, we expect the female embryo's maternal-origin genes will
213 prefer a greater allocation to soldiering than will her paternal-origin genes. Accordingly, considering
214 loci whose gene products promote soldier development ("soldier promoters") we expect her maternal-
215 origin genes will prefer a greater level of gene expression and her paternal-origin genes will prefer a
216 lower level of gene expression, and so we predict soldier promoters will be maternally-expressed and
217 paternally-silenced (figure 2A). And considering loci whose gene products inhibit soldier development
218 ("soldier inhibitors") we expect her maternal-origin genes will prefer a lower level of gene expression
219 and her paternal-origin genes will prefer a greater level of gene expression, and so we predict soldier
220 inhibitors will be maternally-silenced and paternally-expressed (figure 2A).

221 Conversely, under the sex-ratio-conflict hypothesis, we expect the female embryo's maternal-
222 origin genes will prefer a lower allocation to soldiering than will her paternal-origin genes. Accordingly,
223 considering loci whose gene products promote soldier development, we expect her maternal-origin
224 genes will prefer a lower level of gene expression and her paternal-origin genes will prefer a greater
225 level of gene expression, and so we predict soldier promoters will be maternally-silenced and paternally-
226 expressed (figure 2B). And considering loci whose gene products inhibit soldier development, we
227 expect her maternal-origin genes will prefer a greater level of gene expression and her paternal-origin
228 genes will prefer a lower level of gene expression, and so we predict soldier inhibitors will be
229 maternally-expressed and paternally-silenced (figure 2B).

230 Such genomic imprinting is expected to modulate the phenotypic consequences of gene
231 knockouts. A loss-of-function mutation that prevents the affected gene from expressing a functional

gene product is expected to have no impact on the phenotype if that gene is predicted to be silenced anyway. Accordingly, under the brood-benefit hypothesis: a knockout mutation is expected to have no impact on the soldiering phenotype when the gene is a paternal-origin soldier promoter or a maternal-origin soldier inhibitor (figure 2A); but the knockout mutation is expected to reduce soldier development when the gene is a maternal-origin soldier promoter, and to enhance soldier development when it is a paternal-origin soldier inhibitor (figure 2A). Conversely, under the sex-ratio-conflict hypothesis: a knockout mutation is expected to have no impact on the soldiering phenotype when the gene is a maternal-origin soldier promoter or a paternal-origin soldier inhibitor (figure 2B); but the knockout mutation is expected to reduce soldier development when it is a paternal-origin soldier promoter, and to enhance soldier development when it is a maternal-origin soldier inhibitor (figure 2B).

Discussion

Our analysis concerns the function of the sterile soldier caste of polyembryonic parasitoid wasps. We have demonstrated that an intragenomic conflict of interest may arise between a female embryo's maternal-origin genes and her paternal-origin genes, ultimately with respect to the sex ratio of the reproductive adults emerging from the parasitised host, and more proximately with respect to her own propensity for developing as a sterile soldier. In particular, we have found that, owing to the female embryo being relatively more related to her brothers through her mother than through her father, her maternal-origin genes prefer the sex ratio to be less female biased than do her paternal-origin genes. Consequently: if the primary function of soldiers is to altruistically benefit the brood overall, such that female soldiers tend to reduce the reproductive success of the female brood and increase the reproductive success of the male brood, then her maternal-origin genes prefer a greater probability of developing as a soldier than do her paternal-origin genes; whereas, if the primary function of the soldiers is to spitefully distort the sex ratio in favour of their own sex, at a cost to the overall

257 reproductive success of the brood, such that female soldiers tend to increase the reproductive success of
258 the female brood and decrease the reproductive success of the male brood, then her maternal-origin
259 genes prefer a lower probability of developing as a soldier than do her paternal-origin genes.

260 Moreover, we have related this intragenomic conflict of interests to patterns of genomic
261 imprinting of loci underpinning soldier development, deriving contrasting predictions that may be used
262 to discriminate between the brood-benefit versus sex-ratio-conflict hypotheses for soldier function. In
263 particular, we predict that: under the brood-benefit hypothesis, soldier-promoter genes will tend to be
264 maternally-expressed and soldier-inhibitor genes will tend to be paternally-expressed; whereas under the
265 sex-ratio-conflict hypothesis, soldier-promoter genes will tend to be paternally-expressed and soldier-
266 inhibitor genes will tend to be maternally-expressed. Furthermore, we have related these patterns of
267 genomic imprinting to predictions for when loss-of-function mutations will have an impact on the
268 phenotype, and in which direction, which will further aid empirical discrimination between the brood-
269 benefit and sex-ratio-conflict hypotheses for soldier function. Importantly, the “loudest voice prevails”
270 (Haig 1996; Úbeda & Haig 2003) logic underpinning our predictions of genomic imprinting depend
271 only upon the existence and direction – and not the intensity – of intragenomic conflict. Accordingly,
272 our predictions are robust to variation in demographic assumptions concerning, for example, patterns of
273 dispersal that modulate the intensity but not the existence or direction of conflict (cf Farrell et al 2015).

274 Sterile (or reduced-reproductive) soldier castes are known from a number of taxa and – the
275 copidosomatine encyrtids excepted – their primary function is generally understood to be one of brood-
276 benefit, e.g. nest defence. As the patterns of genomic imprinting predicted by the present analysis owe to
277 the basic asymmetry of haplodiploid inheritance (and not the bizarre biology of polyembryony per se),
278 we expect that these predictions will apply widely to female soldiers in many haplodiploid taxa (e.g.
279 eusocial thrips; Crespi 1992). Other asymmetries – such as multiple mating and sex-biases in dispersal,
280 mortality and variance in reproductive success (e.g. Úbeda & Gardner 2012) – have been suggested to
281 drive the evolution of genomic imprinting under diploid inheritance, but here the predicted patterns of

282 imprint are less clear cut. Accordingly, the scope for genomic imprinting in relation to soldiering in
283 diploids (e.g. eusocial trematodes; Hechinger et al 2011) represents an avenue for future study.

284 Parent-of-origin-specific gene expression is well documented in mammals and flowering plants:
285 here, modification of DNA by means of the addition of a methyl group provides a mechanism for
286 regulating gene expression and associated differentiation of cellular tissues, and the differential
287 transmission of methyl modifications via female and male gametes provides the molecular paradigm for
288 parent-of-origin gene effects (Ferguson-Smith 2011). In contrast, the scope for such effects among
289 insects is highly controversial. Previously, the main reason for suspecting that they are unimportant has
290 been the lack of key DNA methylation enzymes in fruit flies (Yan et al 2014). However, there is now
291 strong evidence of methylation-mediated caste differentiation in the social Hymenoptera (Wang et al
292 2006; Kucharski et al 2008; Herb et al 2012; Amarasignhe et al 2014; Yan et al 2014, 2015), where
293 DNA methylation is widespread (Kronforst et al 2008), and disruption of DNA methylation has recently
294 been shown to affect sex allocation in the parasitoid wasp *Nasonia vitripennis* (Cook et al 2015). It is
295 also conceivable that insects could also employ other molecular mechanisms to achieve parent-of-
296 origin-specific gene expression. Moreover, parent-of-origin-specific phenotypic effects have recently
297 been described in relation to social traits of honeybees (Oldroyd et al 2014), and some retention of
298 parent-of-origin information presumably occurs in those insect taxa in which males routinely eliminate
299 their entire paternal genome during spermatogenesis (Ferguson-Smith 2011; Gardner & Ross 2014).

300 The present study of soldiering in polyembryonic parasitoid wasps has demonstrated that the
301 kinship theory of genomic imprinting may provide a powerful tool for exploring social evolution, not
302 only in terms of understanding the adaptations of genes engaged in intragenomic conflicts but also in
303 terms of elucidating the adaptations of individual organisms. Here, we have highlighted the problem of
304 confounding in comparative analyses, i.e. that different populations and different individuals may differ
305 in many respects, and rarely for a single explanatory variable. Specifically: Gardner et al (2007a)
306 interpreted an observed sex difference in soldier allocation as a reflection of sex difference in selection

pressures and, accordingly, inferred that the function of soldiers lies in sex-ratio conflict as opposed to brood benefit; but it is feasible that a sex difference in inherent soldiering ability is instead responsible for this pattern, and that the sex difference in soldiering does not provide any clues as to the soldiers' function. By reframing our comparative analysis at the within-individual level, i.e. between a single individual's maternal-origin and paternal-origin genes, we have eliminated the confounding effect of sex differences in inherent soldiering ability (and many other confounds) and have derived a new set of empirically testable predictions for discriminating the function of soldiers.

Moreover, the relative lack of existing data on parent-of-origin-specific patterns of gene expression provides exciting avenues for truly-independent tests of social evolution theory (Queller & Strassmann 2002; Queller 2003; Wild & West 2009). Often, new theoretical developments on the topic of social evolution are put to empirical test using much the same sources of data that have served as inspiration for the theory in the first place: such circularity is inevitable considering how intensely biological research is focused into a small number of study species. So the possibility for deriving clear-cut predictions about parent-of-origin-specific patterns of gene expression, and resulting phenotypic effects of gene knockouts, in a taxon for which there is no a priori information about such patterns, represents a rare opportunity for subjecting social evolution theory to proper empirical evaluation.

323

324 **Acknowledgments**

325

326 This work was supported by an ESEB Godfrey Hewitt Mobility Award (PR), the Finnish Centre
327 of Excellence in Biological Interactions (PR) and a NERC Independent Research Fellowship (AG; grant
328 number NE/K009524/1). The authors would like to thank J. Boomsma, H. Helanterä, A. Micheletti, M.
329 Puurtinen, L. Ross, T. S. Salminen and J. Wilkins for helpful comments and discussion.

330

331

Literature Cited

- Abbot, P., et al. 2011. Inclusive fitness theory and eusociality. *Nature* 471:E1-E4.
- Amarasinghe, H. E., C. I. Clayton, and E. B. Mallon. 2014. Methylation and worker reproduction in the
bumble-bee (*Bombus terrestris*). *Proceedings of the Royal Society B: Biological Sciences*
281:20132502.
- Boomsma, J. J. 2007. Kin selection versus sexual selection: why the ends do not meet. *Current Biology*
17:R673-R683.
- Boomsma, J. J. 2009. Lifetime monogamy and the evolution of eusociality. *Philosophical Transactions
of the Royal Society of London. Series B, Biological Sciences* 364:3191-3207.
- Boomsma, J. J. 2013. Beyond promiscuity: mate-choice commitments in social breeding. *Philosophical
Transactions of the Royal Society of London. Series B, Biological Sciences* 368:20120050.
- Bulmer, M. 1994. *Theoretical Evolutionary Ecology*. Sinauer Associates, Sunderland, MA.
- Cook, N., B. A. Pannebakker, E. Tauber, and D. M. Shuker. 2015. DNA methylation and sex allocation
in the parasitoid wasp *Nasonia vitripennis*. *The American Naturalist* 186. doi: 10.1086/682950
- Crespi, B. J. 1992. Eusociality in Australian gall thrips. *Nature* 359: 724-726.

356 Cruz, Y. P. 1981. A sterile defender morph in a polyembryonic hymenopterous parasite. Nature
357 294:446-447.

358

359 Cruz, Y. P. 1986. The defender role of the precocious larvae of *Copidosomopsis tanytmemus*
360 Caltagirone (Encyrtidae, Hymenoptera). Journal of Experimental Zoology 237:309-318.

361

362 Doult, R. L. 1947. Polyembryony in *Copidosoma koehleri* Blanchard. The American Naturalist 81:435-
363 453.

364

365 Farrell, E. J., F. Úbeda, and A. Gardner. 2015. Intragenomic conflict over dispersal. The American
366 Naturalist 186. doi: 10.1086/682275

367

368 Ferguson-Smith, A. C. 2011. Genomic imprinting: the emergence of an epigenetic paradigm. Nature
369 Reviews Genetics 12:565-575.

370

371 Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.

372

373 Foster, K. R., T. Wenseleers, and F. L. W. Ratnieks. 2006. Kin selection is the key to altruism. Trends in
374 Ecology and Evolution 21:57-60.

375

376 Gardner, A. 2014. Total reproductive value of juvenile females is twice that of juvenile males under X-
377 linkage and haplodiploidy. Journal of Theoretical Biology 359: 236-237.

378

379 Gardner, A., J. Alpedrinha, and S. A. West. 2012. Haplodiploidy and the evolution of eusociality: split
380 sex ratios. The American Naturalist 179:240-256.

381

382 Gardner, A., I. C. W. Hardy, P. D. Taylor, and S. A. West. 2007a. Spiteful soldiers and sex ratio conflict

383 in polyembryonic parasitoid wasps. *The American Naturalist* 169:519-533.

384

385 Gardner, A., I. C. W. Hardy, P. D. Taylor, and S. A. West. 2007b. Correction. *The American Naturalist*

386 169:837-837.

387

388 Gardner, A., and L. Ross. 2014. Mating ecology explains patterns of genome elimination. *Ecology*

389 *Letters* 17:1602-1612.

390

391 Giron, D., D. W. Dunn, I. C. W. Hardy, and M. R. Strand. 2004. Aggression by polyembryonic wasp

392 soldiers correlates with kinship but not resource competition. *Nature* 430:676-679.

393

394 Godfray, H. C. J. 1992. Strife among siblings. *Nature* 360:213-214.

395

396 Grbić, M., P. J. Ode, and M. R. Strand. 1992. Sibling rivalry and brood sex ratios in polyembryonic

397 wasps. *Nature* 360:254-256.

398

399 Haig, D. 1996. Placental hormones, genomic imprinting, and maternal-fetal communication. *Journal of*

400 *Evolutionary Biology* 9:357-380.

401

402 Haig, D. 1997. Parental antagonism, relatedness asymmetries, and genomic imprinting. *Proceedings of*

403 *the Royal Society B: Biological Sciences* 264:1657-1662.

404

405 Hamilton, W. D. 1964. The genetical evolution of social behavior I & II. *Journal of Theoretical Biology*
 406 7:1-52.
 407
 408 Hamilton, W. D. 1967. Extraordinary sex ratios. *Science* 156: 477-488.
 409
 410 Hamilton, W. D. 1972. Altruism and related phenomena, mainly in social insects. *Annual Review of*
 411 *Ecology and Systematics* 3:193-232.
 412
 413 Hardy, I. C. W. 1994. Sex ratio and mating structure in the parasitoid Hymenoptera. *Oikos* 69:3-20.
 414
 415 Harvey, J. A., L. S. Corley, and M. R. Strand. 2000. Competition induces adaptive shifts in caste ratios
 416 of a polyembryonic wasp. *Nature* 406:183-186.
 417
 418 Hechinger, R. F., A. C. Wood, and A. M. Kuris. 2011. Social organization in a flatworm: trematode
 419 parasites form soldier and reproductive castes. *Proceedings of the Royal Society B* 278: 656-665.
 420
 421 Herb, B. R., F. Wolschin, K. D. Hansen, M. J. Aryee, B. Langmead, R. Irizarry, G. V Amdam, and A. P.
 422 Feinberg. 2012. Reversible switching between epigenetic states in honeybee behavioral subcastes.
 423 *Nature Neuroscience* 15:1371-1373.
 424
 425 Ivanova-Kasas, O. M. 1972. Polyembryony in insects. In: *Developmental Systems: Insects*, vol. 1 (S. J.
 426 Counce & C. H. Waddington, eds). Academic Press, New York.
 427

428 Keasar, T., M. Segoli, R. Barak, S. Steinberg, D. Giron, M. R. Strand, A. Bouskila, et al. 2006. Costs
 429 and consequences of superparasitism in the polyembryonic parasitoid *Copidosoma koehleri*
 430 (Hymenoptera: Encyrtidae). *Ecological Entomology* 31:277-283.

431

432 Kronforst, M. R., D. C. Gilley, J. E. Strassmann, and D. C. Queller. 2008. DNA methylation is
 433 widespread across social Hymenoptera. *Current Biology* 18:287-288.

434

435 Kucharski, R., J. Maleszka, S. Foret, and R. Maleszka. 2008. Nutritional control of reproductive status
 436 in honeybees via DNA methylation. *Science* 319:1827-1830.

437

438 Liao, X., S. Rong, and D. C. Queller. 2015. Relatedness, conflict, and the evolution of eusociality. *PLoS*
 439 *Biology* 13:e1002098.

440

441 Moore, T., and D. Haig. 1991. Genomic imprinting in mammalian development: a parental tug-of-war.
 442 *Trends in Genetics* 7:45-49.

443

444 Nowak, M. A., C. E. Tarnita, and E. O. Wilson. 2010. The evolution of eusociality. *Nature* 466:1057-
 445 1062.

446

447 Ode, P. J., and M. S. Hunter. 2002. Sex ratios of parasitic Hymenoptera with unusual life histories. In I.
 448 C. W. Hardy, ed., *Sex Ratios: concepts and research methods* (pp. 218-234). Cambridge University
 449 Press, Cambridge.

450

451 Ode, P. J., and M. R. Strand. 1995. Progeny and sex allocation decisions of the polyembryonic wasp
 452 *Copidosoma floridanum*. *Journal of Animal Ecology* 64:213-224.

453

454 Oldroyd, B. P., M. H. Allsopp, K. M. Roth, E. J. Remnant, R. A. Drewell, and M. Beekman. 2014. A
455 parent-of-origin effect on honeybee worker ovary size. *Proceedings of the Royal Society B: Biological*
456 *Sciences* 281:20132388.

457

458 Queller, D. C., and J. E. Strassmann. 2002. The many selves of social insects. *Science* 296:311–313.

459

460 Queller, D. C. 2003. Theory of genomic imprinting conflict in social insects. *BMC Evolutionary*
461 *Biology* 3:15.

462

463 Ross, L., A. Gardner, N. Hardy, and S. A. West. 2013. Ecology, not the genetics of sex determination,
464 determines who helps in eusocial populations. *Current Biology* 23:2383-2387.

465

466 Silvestri, F. 1906. Contribuzioni alla conoscenza biologica degli imenotteri parassiti. *Biologia del*
467 *Litomastix truncatellus* (Dalm.) (2° nota preliminare). *Annali della Regia Scuola Superiore*
468 *d'Agricoltura di Portici* 6:3?51.

469

470 Strand, M. R. 2009. Polyembryony. *Encyclopedia of Insects* (pp. 821-825).

471

472 Strand, M. R., J. A. Johnson, and J. D. Culin. 1990. Intrinsic interspecific competition between the
473 polyembryonic parasitoid *Copidosoma floridanum* and solitary endoparasitoid *Microplitis demolitor* in
474 *Pseudoplusia includens*. *Entomologia Experimentalis et Applicata* 55:275-284.

475

476 Úbeda, F., and A. Gardner. 2012. A model for genomic imprinting in the social brain: elders. *Evolution*
477 66: 1567-1581.

478

479 Úbeda, F., and D. Haig. 2003. Dividing the child. *Genetica* 117:103-110.

480

481 Wang, Y., M. Jorda, P. L. Jones, R. Maleszka, X. Ling, H. M. Robertson, C. A. Mizzen, et al. 2006.

482 Functional CpG methylation system in a social insect. *Science* 314:645-647.

483

484 West, S. A. 2009. *Sex Allocation*. Princeton University Press, Princeton, NJ.

485

486 Wild, G., and S. A. West. 2009. Genomic imprinting and sex allocation. *The American Naturalist*

487 173:E1–E14.

488

489 Wilson, E. O. 2005. Kin selection as the key to altruism: its rise and fall. *Social Research* 72:159-166.

490

491 Yan, H., R. Bonasio, D. F. Simola, J. Liebig, S. L. Berger, and D. Reinberg. 2015. DNA methylation in

492 social insects: how epigenetics can control behavior and longevity. *Annual Review of Entomology*

493 60:435-452.

494

495 Yan, H., D. F. Simola, R. Bonasio, J. Liebig, S. L. Berger, and D. Reinberg. 2014. Eusocial insects as

496 emerging models for behavioural epigenetics. *Nature Reviews Genetics* 15:677-688.

497

Genealogical relationship	Symbol	Value	In terms of model parameters
Mating partners	p_{MP}	$(1 - d_f)(1 - d_m)p_{mf}$	$\frac{(1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Male to himself/ brother to brother	p_{mm}	1	1
Sister to brother	p_{fm}	$\frac{p_{ff} + p_{MP}}{2}$	$\frac{1}{4 - 3(1 - d_f)(1 - d_m)}$
Female to herself/ sister to sister	p_{ff}	$\frac{1 + p_{MP}}{2}$	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Sister to sister maternal	$p_{ff M}$	p_{ff}	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Sister to sister paternal	$p_{ff P}$	p_{ff}	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Brother to sister	p_{mf}	p_{fm}	$\frac{1}{4 - 3(1 - d_f)(1 - d_m)}$
Brother to sister maternal	$p_{mf M}$	p_{ff}	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$

Brother to sister paternal	$p_{mf P}$	p_{MP}	$\frac{(1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Daughter to father	p_{fP}	p_{ff}	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Son to father	p_{mP}	p_{MP}	$\frac{(1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$
Daughter to mother	p_{fM}	$\frac{p_{ff} + p_{MP}}{2}$	$\frac{1}{4 - 3(1 - d_f)(1 - d_m)}$
Son to mother	p_{mM}	p_{ff}	$\frac{2 - (1 - d_f)(1 - d_m)}{4 - 3(1 - d_f)(1 - d_m)}$

500

501 **Table 1:** *Summary of the consanguinities used in the analysis* - Because of the possible inbreeding via
502 failure to disperse, the consanguinity coefficients depend on each other. By using those dependences we
503 can solve and express the consanguinity coefficients with the model parameters. The focal dispersing
504 tendencies and the number of emerging males and females are assumed to follow the population
505 average. Therefore the probability that mating partners are siblings can be simplified to a probability
506 that they both failed to disperse $((1-d_f)(1-d_m))$. The dependancies are presented in the 'Value' column,
507 from which the exact values are solved and represented in the 'In terms of model parameters' column. In
508 the indices f represents female embryo, m male embryo, M mother and maternal genome, and P father
509 and paternal genome.

510

Figure Legends

Figure 1: *Illustration of the sex ratio optima z_A* - The graphs represent the sex ratio optima with zero female dispersal ($d_f = 0$) in respect to the proportion of males dispersing (d_m). They are calculated from equation (3) by substituting in the appropriate consanguinity coefficients given in table 1. The actors whose optima are presented in order from the top: the male embryo (dashed gray); the mother (dotted gray); the maternal-origin genome of the female embryo (red); the female embryo as an individual (solid gray); the paternal-origin genome of the female embryo (blue); and the father of the brood (also blue). The sex ratio optima for the male and female embryos, and the mother (gray graphs) were previously presented by Gardner et al (2007a, b) and they are recovered from our equation (3).

Figure 2: *Predictions for patterns of genomic imprinting* - Under the brood-benefit hypothesis (column A) we predict that soldier promoters will be maternally-expressed (green) and paternally-silenced (red), and that soldier inhibitors will be maternally-silenced and paternally-expressed. Under the sex-ratio-conflict hypothesis (column B) the predictions are reversed: the soldier promoters will be maternally-silenced and paternally-expressed, and the soldier inhibitors maternally-expressed and paternally-silenced. Under both hypotheses, a knockout mutation is expected to reduce soldier development (arrow down) when the gene is a soldier promoter, enhance soldier development (arrow up) when it is a soldier inhibitor, and have no effect when the gene is silenced.

Online appendix

Here we show how the number of emerging females behaves at the female's soldiering optimum under both hypotheses, i.e. when $\partial N_f / \partial x|_*$ is positive and when it is negative. The consequences of increasing female soldiering can be divided into three effects: (i) increasing female soldiering leads to a partial decrease in the number of emerging females; (ii) the action of these extra soldiers increases the number of emerging males under the brood-benefit hypothesis ($\partial N_m / \partial x|_* > 0$), and decreases the number of emerging males under the sex-ratio-conflict hypothesis ($\partial N_m / \partial x|_* < 0$); (iii) more resources are freed up (i.e. from: soldier-killed competitors or soldier-macerated host tissue, under the brood-benefit hypothesis; soldier-killed brothers, under the sex-ratio-conflict hypothesis; and from soldiers potentially requiring fewer resources to develop than do reproductive-destined larvae, under both hypotheses) and this leads to a partial increase in the number of emerging females under both hypotheses. The total change in the number of emerging females $\partial N_f / \partial x$ is the combined effect of (i) and (iii). If the effect (ii) were null, then this would mean that increasing the female soldiering does not free up any resources, and so the effect (iii) would also be null, i.e. the only consequence of increasing the allocation to female soldiering would be effect (i), and hence fewer emerging females. Therefore the change in the number of emerging males ($\partial N_m / \partial x$) and females ($\partial N_f / \partial x$) cannot both be zero at the same time, especially at the female's soldiering optimum x_f .

Multiplying the male mating success M with the number of emerging males N_m gives the total mating success for the progenitor male egg. This total number can be divided into two components: matings achieved by non-dispersing males, and matings achieved by dispersing males. Both of these components increase as the number of emerging males increases: the first component, because with more emerging males the local mating pool has a larger frequency of focal males competing for an unchanged amount of available matings; the second component, because then there are more males pursuing matings outside the focal host, which has an insignificant effect on the male mating success.

556 Therefore, the total mating success of the male egg increases with the number of emerging males
 557 $(\partial(N_m M)/\partial N_m > 0)$. Increasing the number of emerging females can only increase the total mating
 558 success of the male egg, by increasing the number of mating opportunities in the local mating pool
 559 $(\partial(N_m M)/\partial N_f \geq 0)$.

560 Rewriting equation (6), using the two dimensional chain rule, we have

561
$$\left. \frac{\partial(N_m M)}{\partial N_m} \right|_* \left. \frac{\partial N_m}{\partial x} \right|_* = - \left. \frac{\partial N_f}{\partial x} \right|_* \left(2 \frac{p_{ff}}{p_{mf}} + \left. \frac{\partial(N_m M)}{\partial N_f} \right|_* \right). \quad (A1)$$

562 From the above line of argument – especially the effect (ii) of increasing female soldiering – we see
 563 from equation (A1) that $\partial N_f / \partial x|_* < 0$ under the brood-benefit hypothesis and $\partial N_f / \partial x|_* > 0$ under the
 564 sex-ratio-conflict hypothesis.

Figure 2