

1 **Implications of ^{36}Cl exposure ages from Skye, northwest Scotland for**
2 **the timing of ice stream deglaciation and deglacial ice dynamics.**

3 David Small ^{a*,1}, Vincent Rinterknecht ^{a,b}, William E. N. Austin ^{a,c}, Richard Bates ^a,
4 Douglas I. Benn ^a, James D. Scourse ^d, Didier L. Bourlès ^e, ASTER Team ^{e,±}, Fiona D.
5 Hibbert ^f

6

7 *^a School of Geography and Geosciences, University of St Andrews, St Andrews, KY16*
8 *9AL, UK.*

9 *^b Université Paris 1 Panthéon-Sorbonne, CNRS Laboratoire de Géographie Physique, F-*
10 *92195 Meudon, France.*

11 *^c Scottish Marine Institute, Scottish Association for Marine Sciences, Oban, PA37 1QA,*
12 *UK.*

13 *^d School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB,*
14 *UK.*

15 *^e Aix-Marseille Université, CNRS-IRD-Collège de France, UM 34 CEREGE,*
16 *Technopôle de l'Environnement Arbois-Méditerranée, BP80, 13545 Aix-en-Provence,*
17 *France.*

18 *^e Research School of^f Ocean and Earth Sciences, The Australian National University,*
19 *Canberra, ACT 2601, Australia*

20 ¹ To whom correspondence should be addressed: David.Small@glasgow.ac.uk

21 ^{*} Now at Department of Geographical and Earth Sciences, University of Glasgow,
22 Glasgow, G12 8QQ, UK. +44 141 330 5442.

23 [±] Maurice Arnold, Georges Aumaître, Karim Keddadouche.

24

25 **Abstract**

26 Constraining the past response of marine terminating ice streams during episodes of
27 deglaciation provides important insights into potential future changes due to climate
28 change. This paper presents new ^{36}Cl cosmic ray exposure dating from boulders located
29 on two moraines (Glen Brittle and Loch Scavaig) in southern Skye, northwest Scotland.
30 Ages from the Glen Brittle moraines constrain deglaciation of a major marine
31 terminating ice stream, the Barra-Donnegal Ice Stream that drained the former British-
32 Irish Ice Sheet, depending on choice of production method and scaling model this
33 occurred $19.9 \pm 1.5 - 17.6 \pm 1.3$ ka. We compare this timing of deglaciation to existing
34 geochronological data and changes in a variety of potential forcing factors constrained
35 through proxy records and numerical models to determine what deglaciation age is most
36 consistent with existing evidence. Another small section of moraine, the Scavaig
37 moraine, is traced offshore through multibeam swath-bathymetry and interpreted as
38 delimiting a later stillstand/readvance stage following ice stream deglaciation.
39 Additional cosmic ray exposure dating from the onshore portion of this moraine indicate
40 that it was deposited $16.3 \pm 1.3 - 15.2 \pm 0.9$ ka ago. When calculated using the most up-
41 to-date scaling scheme this time of deposition is, within uncertainty, the same as the
42 timing of a widely identified readvance, the Wester Ross Readvance, observed
43 elsewhere in northwest Scotland. This extends the area over which this readvance is
44 potentially occurred, reinforcing the view that it was climatically forced.

45

46 **1. Introduction**

47 Concerns over the stability of the remaining ice sheets have been raised by suggestions
48 that irreversible collapse of some marine based sectors is possible or has already begun,
49 with attendant effects on associated terrestrial glaciers (Joughin et al., 2014; Wouters et

50 al., 2015). Marine terminating ice streams are important components of the
51 interconnected ocean-cryosphere system because they discharge large volumes of ice
52 directly into the ocean through calving (Alley and MacAyeal, 1994; Bradwell and
53 Stoker, 2015; Deschamps et al., 2012). While modern observations provide useful
54 information, the temporal coverage is not sufficient to capture the complete response of
55 a marine terminating ice stream to rapid climate change. Researchers are therefore
56 increasingly drawn to analogous palaeo-settings where the complete deglaciation record
57 can be observed (Serjup et al., 2000; Dowdeswell et al., 2014; Svendsen et al., 2015).

58 Of the Pleistocene ice sheets, the British-Irish Ice Sheet (BIIS) provides a useful
59 analogue. Its western margin was marine terminating while its position next to a major
60 surficial artery of the Atlantic Meridional Overturning Circulation (AMOC) rendered it
61 potentially sensitive to small climatic perturbations (Knutz et al., 2007). This sensitivity
62 is captured in proxy data (Scourse et al., 2009; Hibbert et al., 2010) and numerical
63 modelling experiments (Hubbard et al., 2009). Past reconstructions of the BIIS relied
64 heavily on onshore mapping of landforms that can be inferred to represent former ice
65 limits including terminal, lateral and recessional moraines (Sissons et al., 1973;
66 Ballantyne, 1989, Bennett and Boulton, 1993; Clark et al., 2004). Recent advances in
67 offshore geomorphological mapping, particularly the use of bathymetric and seismic
68 data, have allowed workers to identify sediments and landforms associated with ice
69 extending onto the continental shelf (Bradwell et al., 2008a; Dunlop et al., 2010;
70 Ó'Cofaigh et al., 2012). This has allowed delimitation of fast flowing ice streams that
71 drained much of the western sector of the former BIIS (Scourse et al., 2000; Stoker and
72 Bradwell, 2005; Howe et al., 2012; Bradwell and Stoker, 2015; Dove et al., 2015).
73 Further identification of subsequent landforms associated with confined ice flow casts
74 light on post-ice streaming behaviour inshore of the onset zone of the BDIS (Howe et

75 al., 2012; Dove et al., 2015).

76 The Barra-Donegal Ice Stream (BDIS) drained a large portion of the western
77 BIIS and, at the Last Glacial Maximum (LGM), reached the shelf edge (Knutz et al.,
78 2001) where it deposited glaciogenic sediments in the Barra-Donegal Fan (BDF), the
79 southernmost glaciogenic fan on the Eurasian continental margin (Figure 1). Recent
80 observations using swath bathymetry have revealed a suite of glaciogenic landforms at
81 the bed of the former BDIS, stretching from Skye in the north to Islay in the south
82 (Howe et al., 2012; Dove et al., 2015). The BDIS flowed southwest from the Inner
83 Hebrides before turning west around the Outer Hebrides towards the outer shelf (Howe
84 et al., 2012). Large scale erosional features such as glacially over-deepened basins and
85 streamlined bedrock are observed across large areas of the BDIS and provide important
86 information on past ice flow directions. In comparison, large moraines are confined to
87 the mid-outer shelf with smaller recessional moraines being more abundant in the
88 nearshore (Dunlop et al., 2010; Dove et al., 2015).

89 Offshore evidence from ice rafted detritus (IRD) demonstrates that ice sourced in
90 Scotland reached the shelf edge by 29 ka with a significant reduction in IRD delivery
91 after 23 ka (Knutz et al., 2001; Scourse et al., 2009; Hibbert et al., 2010). To the north,
92 basal marine radiocarbon ages show deglaciation of mid-shelf (Figure 1; Table 1) prior
93 to 16.7 ± 0.3 ka (Peacock et al., 1992; Austin and Kroon, 1996; Small et al., 2013) while
94 cosmogenic exposure and radiocarbon ages (Figure 1) show initial deglaciation of the
95 southern sector of the BDIS before ~ 20.0 ka (McCabe et al., 2003; Clark et al., 2009).
96 Complete deglaciation of the southern sector occurred before 16.8 ka (Figure 1)
97 (Ballantyne et al., 2014), an inference supported by IRD evidence that the BIIS
98 maintained calving margins throughout the period 23.0-16.0 ka (Scourse et al., 2009,
99 Small et al., 2013). All available geochronological data related to the BIIS was

100 synthesised to produce 1 ka time-slices of the pattern of deglaciation (Clark et al., 2012),
101 this was subsequently refined to include maximum and minimum ice-extents at the same
102 temporal resolution (Hughes et al., 2016). In the BDIS sector both reconstructions depict
103 initial deglaciation from the shelf edge at c.25 ka with ice persisting on the mid-shelf
104 until 19-17 ka. Rapid deglaciation occurs 17-16 ka by which time is located near the
105 present day coastline (Figure 1).

106 While the submarine geomorphology and retreat pattern of the BDIS is relatively well
107 established (Howe et al., 2012; Dove et al., 2015), post-ice streaming behaviour and
108 geochronological data relating to deglaciation of the northern sector of the BDIS is still
109 comparatively limited. In northwest Scotland a regional scale readvance, the Wester
110 Ross readvance has been delimited from a suite of onshore moraines and dated with ^{10}Be
111 exposure ages to ~ 16 ka (Robinson and Ballantyne, 1979; Bradwell et al., 2008;
112 Ballantyne et al., 2009). However to date, this readvance has not been identified south of
113 Skye. In this contribution we present bathymetric data from inshore waters near Skye,
114 which highlights ice dynamics following ice stream retreat. Cosmogenic ^{36}Cl cosmic ray
115 exposure (CRE) ages from moraine boulders provide geochronological constraints on
116 the timing of this deglaciation.

117

118 **2. Study Site**

119 Skye is located off the west coast of Scotland, >200 km upstream from the maximum
120 extent of the BDIS at the shelf break (Figure 1). During the LGM the mountains of
121 central Skye (the Cuillin) nourished an independent ice dome, the Skye Ice Dome (SID),
122 which deflected ice moving from the mainland to the west and acted as an ice divide
123 between the BDIS and the Minch Ice Stream (MIS). Together, these ice streams drained
124 the majority of the northern sector of the BIIS (Bradwell et al., 2008a). To the north of

125 Skye, the zone of confluence between mainland ice and the SID is inferred to follow the
126 narrow straits between Skye and the islands of Scalpay and Raasay (Harker, 1901)
127 (Figure 2). To the south, mainland erratics occur on the island of Soay and the
128 orientation of striae on the southern margin of the Cuillin suggest that locally nourished
129 ice was strongly deflected westwards by mainland ice. This implies that the zone of ice
130 confluence lay between Skye and the neighbouring island of Soay (Ballantyne et al.,
131 1991). The southern branch of mainland ice, along with ice flowing south from the
132 Cuillin, fed the embryonic BDIS with ice stream onset beyond Rum (Howe et al., 2012;
133 Dove et al., 2015). The northern branch fed the MIS (Stoker and Bradwell, 2005). Given
134 its central position within the BDIS, Skye is an important location for constraining
135 deglaciation of the BDIS and comparing the deglacial history of neighbouring ice
136 streams that drained a dynamic, marine-based ice sheet.

137 Deglaciation of the MIS is constrained by several CRE ages. Two ^{36}Cl CRE ages
138 from ice smoothed bedrock on a col in Trotternish (Figure 2) show deglaciation at
139 altitude in Northern Skye before ~16 ka (Stone et al., 1998). Further constraint on final
140 deglaciation of the MIS is provided by five ^{10}Be CRE ages with a mean age of $15.9 \pm$
141 1.0 ka from a boulder moraine at Strollamus (Small et al., 2012), above the strait that
142 separates Skye from Scalpay (Figure 2). In contrast, the only CRE ages from southern
143 Skye are from a moraine related to the later Loch Lomond Readvance (LLR) (Small et
144 al., 2012).

145 Our study focuses on two locations in Southern Skye where there are moraines
146 outside the well mapped LLR limits., Glen Brittle to the west of the Cuillin, and Loch
147 Scavaig, to the south (Figures 2 and 5). At both sites the moraines represent the
148 innermost pre-LLR limit yet identified but without geochronological control it is not
149 possible to determine if they were deposited contemporaneously. In lower Glen Brittle

150 the up-valley termination of raised shorelines coincides with a series of low moraine
151 ridges littered with basalt boulders which have been interpreted as terminal moraines
152 (Walker *et al.*, 1988). These moraines occur well outside the mapped limits of the LLR
153 (Ballantyne, 1989) and thus clearly pre-date them (Figure 3). In Glen Brittle there are
154 two main parallel moraine ridges up to 100 m long and 2-3m high (Figure 3). The ridges
155 are separated by ~50 m.

156 On Soay which forms the western margin of Loch Scavaig, a small section of
157 moraine comes onshore at the northeastern corner of the island (Clough and Harker,
158 1904). This moraine section is ~200-300 m in length and 4-5 m high in places. Large
159 erratic gabbro boulders are found on its crest indicating that at some time following
160 deglaciation of the BDIS ice sourced from the Cuillin extended into Loch Scavaig and
161 reached Soay which itself is composed entirely of Torridonian sandstone with some
162 Tertiary basalt dykes.

163

164 **3. Methods**

165 *3.1 Bathymetry*

166 To constrain deglaciation of the BDIS we confirmed the presence of ice margin
167 positions in southern Skye from onshore fieldwork in Glen Brittle and a bathymetric
168 survey of Loch Scavaig. This study used a SEA SwathPlus High Frequency System with
169 a central frequency of 468 kHz and a ping rate of up to 30 pings per second giving a
170 potential footprint of less than 5 cm at standard survey speed. Data were acquired with a
171 TSSDMS205 motion reference unit and positioning provided by a Topcon Hiper RTK
172 dGPS. The RTK dGPS base system was established on the loch shore and tied to the
173 BNG datum using Rinex corrections from the OS. An Applied Microsystems MicroSV

174 sound velocity probe was mounted at the sonar head in order to record changes in
175 velocity due to mixing of different waters (and thus potential salinity changes) in the
176 relatively enclosed waters of the loch. Final data were recorded to a position accuracy of
177 better than +/-5 cm, however the final data set was processed to a bin resolution of 2 m
178 with vertical heights given to ± 20 cm. The data was processed using SwathPlus and
179 GridProcessor (SEA Ltd) with further editing using IVS Fledermaus. Bathymetric data
180 points were converted from WGS84 to OSGP using the OSGB36 datum (origin 49°N
181 and 2°W). Final data processing was accomplished within ArcGIS (v10).

182

183 *3.2 Surface exposure dating using ^{36}Cl .*

184 *3.2.1 Sampling*

185 Moraines with suitable material for CRE dating using *in situ*-produced cosmogenic
186 ^{36}Cl were identified in Glen Brittle and on the island of Soay where the onshore
187 continuation of an offshore moraine is located. Eleven samples, four from Glen Brittle
188 and seven from Soay, were collected from basic igneous boulders (basalt and gabbro)
189 for CRE dating. In Glen Brittle two samples were collected from the outer moraine ridge
190 (BRI01 and BRI04) and two samples from the inner moraine ridge (BRI02-03). On Soay
191 7 samples were collected from the onshore moraine section (Figure 4).

192 We selected boulders from moraine crests with the largest *b*-axis to minimise the
193 potential for disturbance and snow cover. Where possible we sampled sub-rounded
194 boulders considered indicative of sub-glacial transport (Ballantyne and Stone, 2009) to
195 minimise the potential for inheritance. Similarly we sampled boulders with intact top
196 surfaces as they are least likely to have suffered significant chemical weathering and to
197 minimise the potential influence of spallation of material. Samples were collected from

198 the top surfaces of boulders using hammer and chisel. When possible, we sampled flat
199 surfaces but, where necessary, strike and dip were recorded using a compass-clinometer.
200 Detailed site descriptions (e.g. geomorphological context, boulder dimensions,
201 weathering) were made for each sample. Sample locations and elevations were recorded
202 using a hand-held GPS with elevations checked against 1:25000 maps. Skyline
203 measurements were taken using a compass-clinometer at all sites with the topographic
204 shielding factors calculated using the skyline calculator within the CRONUS online
205 calculator (Balco et al., 2008;
206 http://hess.ess.washington.edu/math/general/skyline_input.php; accessed on 14th
207 September 2015). Sample information is shown in Table 2. Sample photos are shown in
208 Figures 5 (Glen Brittle) and 6 (Soay).

209

210 *3.2.2 Processing*

211 The thickness and dry bulk density of samples from each site was measured before
212 samples for ³⁶Cl analysis were crushed and sieved to 250-500 µm at the University of St
213 Andrews. About 2 g of material was retained for elemental analysis with the remainder
214 sent to University of New Hampshire for further preparation and isotopic extraction.
215 Chlorine was extracted and purified from whole-rock samples to produce AgCl for
216 accelerator mass spectrometry (AMS) analysis, following a modified version of
217 procedures developed by Stone et al. (1996). Crushed samples were sonicated first in
218 distilled water and then in 2% HNO₃ to remove any secondary material attached to
219 grains. 13-20 g of pretreated rock was prepared from each sample for subsequent
220 chemical procedures. Samples were spiked with ~0.48 g of isotopically enriched carrier
221 (³⁵Cl/³⁷Cl = 999 ± 4, total Cl concentration = 3.65 mg g⁻¹) before dissolution in an HF –
222 HNO₃ solution. Following complete dissolution, aqueous samples were separated from

223 solid fluoride residue by centrifuging, and ~1 ml of 5% AgNO₃ solution was added to
224 precipitate AgCl (and Ag₂SO₄ if sulfates were present). The precipitate was collected by
225 centrifuging and dissolved in NH₄OH solution. To remove sulfates, ~1 ml of saturated
226 (BaNO₃)₂ was added to precipitate BaSO₄. Final precipitation of AgCl from the aqueous
227 solution was accomplished by addition of 2 M HNO₃ and 5% AgNO₃. The final AgCl
228 precipitate was collected by centrifuging, washed repeatedly with 18.2 MΩ-cm
229 deionized water, and dried. Approximately 1.5 – 1.75 mg of purified AgCl target
230 material was produced from each sample for AMS measurement.

231

232 *3.2.2 Analysis and age calculations*

233 ³⁶Cl measurements were carried out at the 5 MV French accelerator mass
234 spectrometry national facility ASTER at CEREGE (Arnold et al., 2013). Use of an
235 isotopically enriched carrier allows simultaneous measurement of ³⁵Cl/³⁷Cl and
236 determination of the natural Cl content of the dissolved samples. For normalization of
237 ³⁶Cl/³⁵Cl ratios, calibration material ‘KN1600’ prepared by K. Nishiizumi, was used.
238 This has a given ³⁶Cl/³⁵Cl value of $2.11 \pm 0.06 \times 10^{-12}$ (Fifield et al., 1990). Typical
239 uncertainties for raw AMS data are 0.3 – 1.2% for ³⁵Cl/³⁷Cl and 4.8 – 8.0% for ³⁶Cl/³⁵Cl.
240 All samples have ³⁶Cl/³⁵Cl ratios in the range of $3.8 - 6.9 \times 10^{-14}$ compared to two
241 process blanks (CLBLK7 & 8) with ³⁶Cl/³⁵Cl ratios of 7.83 ± 1.0 and $4.15 \pm 0.75 \times 10^{-15}$,
242 respectively. Resulting blank corrections therefore range between 3.4 and 18.1%.
243 Measurement results and calculated concentrations with uncertainties are shown in
244 Table 3.

245 ³⁶Cl CRE ages were calculated using the CRONUScale online calculator
246 (<http://web1.ittc.ku.edu:8888>; accessed 09/02/2016; Marrero et al., 2016a) and a freely

247 available spreadsheet (Schimmelpfennig et al., 2009). ^{36}Cl production rates for
248 spallation (Ca, K) have recently been updated by Marrero et al. (2016b). Consequently,
249 we calculated our exposure ages using sea level-high latitude ^{36}Cl production rates of
250 56.0 ± 4.1 , 155 ± 11 , 13 ± 3 and 1.9 ± 0.2 atoms $^{36}\text{Cl g}^{-1} \text{a}^{-1}$, for Ca, K, Ti and Fe,
251 respectively (Marrero et al., 2016b; Schimmelpfennig et al., 2009). In comparison,
252 previous production rates for Ca and K were 42.2 ± 4.8 , 145.5 ± 7.7 atoms $^{36}\text{Cl g}^{-1} \text{a}^{-1}$
253 (Schimmelpfennig et al., 2011, 2014; also see Braucher et al., 2011). We report CRE
254 ages calculated using both Ca and K production rates and scaled for latitude and altitude
255 according to Stone (2000), as adapted by Balco et al. (2008), and Lifton et al. (2014) for
256 comparison. CRE ages were calculated assuming no erosion. Correcting for 1 mm ka^{-1}
257 erosion would vary exposure ages by 1-2%. The chemical composition of representative
258 bulk material was determined for each individual sample at the Facility for Earth and
259 Environmental Analysis at the University of St Andrews using X-ray fluorescence
260 (XRF) for major elements and inductively coupled plasma mass spectrometry (ICP-MS)
261 for minor and trace elements. The composition of individual samples is shown in Table
262 4.

263

264 *3.3 Comparison to proximal marine cores:*

265 We compare our surface exposure dating of the marine terminating Barra-Donegal
266 Ice Stream with two proximal marine records, MD02-2822 (Hibbert, 2011; Hibbert et
267 al., 2010) and MD01-2461 (Peck et al., 2006, 2008). Giant piston core MD04-2822 was
268 recovered by the RV *Marion Dufresne* from the deep-water margins of the BDF in the
269 Rockall Trough (Figure 1; $56^{\circ} 50.54' \text{ N}$, $11^{\circ} 22.96' \text{ W}$; 2344 m water depth, recovered
270 in 2004). MD01-2461 was collected from the north-western flank of the Porcupine

271 Seabight approximately 550 km to the southwest (51°45'N, 12°55'W; 1153 m water
272 depth, recovered in 2001). This region lies within the zone of meridional oscillation of
273 the North Atlantic Polar Front during the last glacial (Knutz et al., 2007; Scourse et al.,
274 2009; Hibbert et al., 2010) and as a result is ideally positioned to record both the
275 prevailing hydrographic conditions and the dynamics of the proximal BIIS.

276 Each core is plotted on their own age model based on tuning to the Greenland $\delta^{18}\text{O}$
277 ice core records (using NGRIP on the GICC05 timescale for MD04-2822 and GISP2 for
278 MD01-2461) and calibrated ^{14}C dates (Figure 10). We have updated the age model for
279 MD04-2822 using: the most recent calibration dataset (IntCal13; Reimer et al., 2013);
280 age uncertainty estimates for each tie-point (a mean squared estimate incorporating
281 uncertainties from both the ice core chronology and tuning procedure) and; a Bayesian
282 deposition model (OxCal 'Poisson' function; Bronk Ramsey and Lee, 2013)
283 (Supplementary Table 1).

284

285 **4. Results**

286 *4.1 Multibeam bathymetry*

287 The multibeam bathymetric survey of Loch Scavaig reveals numerous features –
288 both glaciogenic and post glacial – of interest. The most conspicuous of these is a large
289 arcuate ridge that spans Loch Scavaig and connects with the observed onshore moraine
290 section found on Soay. The ridge is ~4.5 km long and up to 10 m high in places (Figures
291 7 and 8). A further small extension (~1 km) of this ridge crosses the Sound of Soay to
292 come onshore on the southern margin of the Cuillin. This ridge is interpreted as a
293 terminal moraine, the Scavaig moraine, that clearly delimits the extent of a glacier that
294 flowed from the central rock basin of the Cuillin and into Loch Scavaig.

295 The glacial land-system preserved in Loch Scavaig is very different, both in

296 morphology and scale, from that associated with surging tidewater glaciers in Svalbard
297 (Ottesen et al., 2008) with a lack of megascale glacial lineations, crevasse fills and
298 eskers. In addition, the scale and shape of the Scavaig moraine is strikingly different
299 from thrust moraines in Svalbard, which are up to 1 km across with large debris flow
300 lobes on their distal slopes (Ottesen et al., 2008; Kristensen et al., 2009). The Scavaig
301 moraine is a much smaller feature with a well-defined crest, it is generally arcuate in
302 planform, with an asymmetric profile. These features are consistent with a push moraine
303 formed at the margin of the former glacier, indicating that the Scavaig moraine was not
304 formed by a surging glacier but instead marks a readvance of ice from the Cuillin or a
305 still-stand during overall retreat. The Scavaig moraine is traceable across the floor of
306 Loch Scavaig and onto the island of Soay (Figure 4 and 8). The onshore section aligns
307 exactly with the offshore moraine, is composed of material from the Cuillin where the
308 glacier that deposited the Scavaig moraine must have been sourced. It is therefore
309 clearly part of the same feature.

310 Within the limits of the large moraine is a suite of shorter but conspicuous linear
311 ridges, most prominent in the east of the survey area and immediately inboard of the
312 large moraine (Figure 8). These are up to 2 km long and 5 m high and are interpreted as
313 recessional moraines formed during deglaciation from the outer limit demarked by the
314 Scavaig moraine.

315 In the east of the survey area, an area of the sea floor is covered with chaotic,
316 hummocky topography (Figure 8). This bears resemblance to features identified as
317 submarine slope failures in bathymetric studies carried out elsewhere in Scotland
318 (Stoker et al., 2010). In addition, the features occur immediately below a conspicuous
319 failure scarp that occurs on Ben Cleat which forms the eastern shore of Loch Scavaig.
320 This feature is interpreted as a post-glacial rock slope failure. Similar terrestrial features

321 in Scotland have been linked to glacial debuitressing and seismic activity associated
322 with post-glacial isostatic rebound (Ballantyne and Stone, 2013).

323

324 *4.2 Surface exposure dating using ^{36}Cl .*

325 The exposure ages calculated following Schimmelfennig et al. (2009) and Marrero et
326 al., (2016a, b) are shown in Table 5. Due to the differing ways in which each calculator
327 deals with the numerous production pathways of ^{36}Cl and the varying compositions of
328 our samples the difference in calculated CRE age is not consistent between samples
329 although the ages calculated using the Lm scaling show general agreement between the
330 Schimmelfennig calculator (Schimmelfennig et al., 2009) and CRONUScalc (Marrero
331 et al., 2016a). Notably, the choice of scaling is important when using the new
332 CRONUScalc online calculator with CRE ages calculated using the Lm scaling (Stone et
333 al., 2000; Balco et al., 2008) being up to 14% older than when calculated with the SA
334 scaling (Lifton et al., 2014). The cause of this discrepancy is currently enigmatic. The
335 dependency of the CRE ages on choice scaling scheme makes interpretation difficult as
336 there is the danger of selecting CRE ages to fit pre-existing or favoured hypotheses.
337 However, given the range of production rate calibrations included in the CRONUScalc
338 programme, the improved agreement with observed atmospheric cosmic-ray fluxes
339 obtained using the SA scaling scheme and for simplicity, we focus discussion on CRE
340 ages calculated using CRONUScalc and the SA scaling. We present the alternative CRE
341 age calculations for completeness.

342 The ^{36}Cl CRE ages range from 19.4 ± 1.7 to 12.9 ± 1.2 ka. The Glen Brittle
343 samples (BRI-01-04) yield CRE ages between 19.4 ± 1.7 and 15.5 ± 1.7 ka while the
344 Soay samples (SOAY-1-7) yield CRE ages between 16.4 ± 1.5 and 12.9 ± 1.2 ka. A plot
345 of all ^{36}Cl CRE ages reveals significant overlap in ages from both locations (Figure 9,

346 Table 5). The 11 samples combined have a reduced Chi-square ($\chi^2_{\text{R}} = 4.51$) indicating
347 that they are not a single population and are influenced by geological uncertainty.
348 Additionally, a Student's t test ($p < 0.01$) suggests that the CRE ages from the two
349 valleys are significantly different. Given this, and the absence of direct
350 geomorphological correlation between the sampled moraines in Glen Brittle and Soay
351 we consider each sample site individually. The Glen Brittle samples have $\chi^2_{\text{R}} = 1.59$
352 which is an acceptable value for a population with three degrees of freedom (Bevington
353 and Robinson, 2003).

354 The Soay samples have $\chi^2_{\text{R}} = 2.06$ indicating the CRE ages are not a single
355 population. Figure 9 shows two CRE age clusters at ~ 13 ka and ~ 15 ka ($\chi^2_{\text{R}} = 0.02$ and
356 0.05 , respectively). There are two potential interpretations of these CRE ages. The first
357 is that the younger CRE age population reflects the age of deposition of the Scavaig
358 moraine and that the older CRE ages reflect nuclide inheritance from a previous
359 exposure. An alternative interpretation is that the older CRE ages are representative of
360 the true moraine age and the young CRE ages are the result of some post-depositional
361 adjustment and/or exhumation.

362

363 **5. Discussion**

364 *5.2 Time of moraine deposition*

365 A compilation of exposure ages from boulders suggests that they are more likely to
366 underestimate the true CRE age (Heyman et al., 2011). However, this compilation was
367 solely comprised of ^{10}Be CRE ages. The greater importance of muons in ^{36}Cl production
368 (e.g. Stone et al., 1998; Braucher et al., 2013) means ^{36}Cl CRE ages have a greater
369 propensity for inheritance and thus overestimation of ages. Similarly the more

370 complicated evolution of production rate with depth (cf. Gosses and Philips, 2001)
371 means that erosion and or spalling of boulder surfaces can make CRE ages appear older
372 than the true boulder age. Despite careful sample selection (Section 3.2.1) the spread in
373 our ages demonstrates that some of our samples were influenced by geological
374 uncertainty. We therefore outline what ages we believe best represent the true moraine
375 age and use these ages as the basis for our interpretation with a general note of caution
376 that our ages may overestimate the true moraine age. We outline some reasons why we
377 consider this less likely however acknowledge it as a possibility.

378 Given the agreement between the CRE ages from Glen Brittle we consider an
379 arithmetic mean to best represent the timing of moraine deposition. Thus we infer that
380 the Glen Brittle moraines were most likely deposited at 17.6 ± 1.3 ka, the mean of our
381 ages. At this time relative sea level (RSL) around the south coast of Skye was high
382 (Figure 11) and the termination of high shorelines is associated with the dated moraines
383 in Glen Brittle. This led Walker et al. (1988) to speculate that at the time of high RSL
384 ice occupied Glen Brittle, a view supported by our CRE ages. We note that there is
385 considerable spread in the ages from Glen Brittle and that the mean age may over- or
386 underestimate the true moraine age.

387 As stated in section 4.2 there are two possible interpretations of the exposure
388 ages from Soay. We consider it unlikely that nuclide inheritance would affect the other
389 boulders to the same degree such that they yielded internally consistent CRE ages that
390 give an acceptable χ^2_R value. Additionally, the young CRE ages suggest moraine
391 deposition prior to the LLR (\approx Younger Dryas - 12.9–11.7 ka b2k; Lowe et al., 2008).
392 This would imply ice survival throughout the warm Bølling-Allerød interstadial, a
393 scenario that is considered unlikely in Scotland (Ballantyne and Stone, 2012). If the
394 older CRE age cluster is to be inferred as best representing the true moraine age it does

395 however raise the question of how the three other boulders were exhumed at the same
396 time. We note that these three boulders are located in very close proximity (Figure 4)
397 and that in comparison to the other sampled boulders they are relatively low lying.
398 Boulder height has been shown to influence the clustering of CRE ages with taller
399 boulders being favoured over shorter boulders (Heyman et al., 2016). Thus while we
400 cannot speculate on the specific mechanism of exhumation the boulder-height
401 relationship identified by Heyman et al. (2016) and the close spatial proximity of the
402 three young Soay samples suggests that contemporaneous exhumation is possible. Given
403 all of these considerations, we favour the second scenario and infer that the Scavaig
404 moraine was most likely deposited 15.2 ± 0.9 ka.

405 The mean ages from the moraines do not agree within their analytical
406 uncertainties which, given the proximity of the sample locations, suggests that they may
407 represent separate glacial events. However, we note that there is considerable overlap
408 between the ages from Glen Brittle and Soay thus we can not definitively make this
409 conclusion. We therefore propose, as a hypothesis, that two separate readvances
410 occurred on the southern margin of the SID during deglaciation. This hypothesis
411 requires further testing with geochronological data.

412

413 *5.1 Implications for local ice dynamics*

414 Evidence for readvance of locally nourished ice on Skye has been documented from
415 several localities on the low ground that surrounds the Cuillin (Benn, 1997). Glacio-
416 tectonised sediments, patterns of erratic dispersal and changes in the marine limit, all
417 suggest that locally nourished ice remained dynamically active after its separation from
418 mainland ice. Benn (1997) delimited potential readvance limits of the SID, but whether
419 these were contemporaneous has, thus far, remained untested.

420 It has previously been suggested that readvance of the SID may have resulted from
421 the removal of constraints imposed by confluent ice allowing the ice to drain radially
422 away from the high ground (Benn, 1997). To the north of the SID a readvance/stillstand
423 is inferred from ice-thrust subaqueous outwash at Suisnish in southern Raasay (Benn,
424 1997). This site is likely to have been proximal to an ice margin when the Strollamus
425 moraine was deposited at 15.9 ± 1.0 ka (Small et al., 2012). This similarity in age to the
426 older CRE exposure ages from Soay suggests that readvance of the northern and
427 southern sectors of the SID may have been synchronous within dating uncertainties.
428 Additionally, the CRE ages of the Scavaig moraine from Soay and the ^{10}Be CRE ages
429 from the Strollamus moraine are the same as a suite of ^{10}Be CRE ages from moraines
430 delimiting the Wester Ross Readvance (Figure 2), ~60 km to the northwest (Robinson
431 and Ballantyne, 1979; Bradwell et al., 2008b, Ballantyne et al., 2009). While the
432 Strollamus moraine has been interpreted as a medial moraine and thus does not record a
433 readvance, it does indicate the existence of a significant ice mass at the time of the
434 WRR. If the Scavaig moraine represents a later readvance, or our CRE ages
435 overestimate the age the Glen Brittle moraine, then, in combination with the evidence
436 for readvance at Suisnish, it is possible that the Wester Ross Readvance may have been
437 more widespread than previously recognized, and involved readvance of local ice on
438 Skye. If this is the case then it implies a common, and likely climatic trigger such as an
439 increase in precipitation associated with climatic warming (c.f. Ballantyne and Stone,
440 2012). We note however that the uncertainties associated with our ages prevent
441 definitive correlation of the Scavaig moraine to moraines dated elsewhere in Scotland.

442

443 *5.2 Deglaciation of the BDIS*

444 The deposition age of the Glen Brittle moraine provides a constraint on final

445 deglaciation of the BDIS as its morphology and lithology demonstrates deposition by
446 valley glaciers fed from the locally nourished SID. As such, it would not be possible to
447 form moraines in Glen Brittle until BDIS deglaciation was complete. Taken at face
448 value, the ^{36}Cl CRE ages from Glen Brittle presented here suggest deglaciation of the
449 northern sector of the BDIS had occurred by 17.6 ± 1.3 ka (SA scaling). Use of the Lm
450 scaling makes deglaciation considerably earlier (19.9 ± 1.1 ka) although the ages do
451 overlap at 1σ . Considered alongside existing geochronological control from the north
452 coast of Ireland and Jura (McCabe and Clark, 2003; Clark et al., 2009; Ballantyne et al.,
453 2014) (Figure 1), our data suggest that the entire marine portion of the former BDIS was
454 deglaciated by 17.6 ± 1.3 ka. Notably, this timing of deglaciation compares well to a
455 reduction in delivery of IRD to the adjacent deep-sea core MD04-2822 (Hibbert et al.,
456 2010) (Figure 10G). Previous reconstructions of the BIIS (Clark et al., 2012; Hughes et
457 al., 2016) depict ice persisting on the mid-inner shelf until ~ 17 ka with ice reaching the
458 coastline at 16 ka. Our data from Glen Brittle suggest that deglaciation occurred earlier
459 and that ice may have reached the coastline several ka earlier than previously inferred.
460 Notably use of the Lm scaling to calculate the CRE age would exacerbate this
461 difference.

462 Numerous oceanic forcing mechanisms have been linked to observations of marine
463 deglaciation within the palaeoenvironment. Eustatically forced changes in sea-level
464 (ESL) rise has been cited as a potentially important factor in deglaciation of other
465 palaeo-ice streams that drained the BIIS (Scourse and Furze, 2001; Haapaniemi et al.,
466 2010; Chiverrell et al., 2013) and an initial eustatic sea level rise occurs at 19 ka (e.g.,
467 DeDeckker and Yokoyama, 2009; Lambeck et al., 2014), prior to BDIS deglaciation at
468 17.6 ± 1.3 ka, as constrained by our data (Figure B).

469 Additionally, it has been shown that tidal mechanical forcing can impact on

470 grounded ice streams (Murray et al., 2007; Arbic et al., 2008; Rosier et al., 2015). The
471 palaeotidal regime influencing the western ice streams draining the BIIS was enhanced
472 compared to the present day because the open glacial North Atlantic was characterized
473 by megatidal amplitudes (tidal ranges > 10 m) in many sectors south of the Iceland-
474 Faroe-Scotland ridge (Uehara et al., 2006; Scourse et al., submitted). Hitherto it has been
475 difficult to disentangle the relative influence of tidal amplitudes *vis-à-vis* relative sea
476 level (RSL) changes but recent modelling efforts have addressed this issue for the BIIS
477 (Scourse et al., submitted) and generated simulations of the potential influence of
478 palaeotides on the BDIS (Figure 11). These show an enhanced tidal regime in the period
479 immediately prior to deglaciation as constrained by the CRE ages from Glen Brittle in
480 the inner BDIS sector (Figure 11). This raises the possibility that this mechanism is a
481 potentially important driver of deglaciation. However, these large tidal amplitudes are
482 associated, in this area, with falling RSL driven by rapid glacio-isostatic uplift which
483 will have mitigated the impact of large tidal range on, for instance, calving rates and ice
484 stream velocities. Similarly, the deposition of the Scavaig moraine occurred during a
485 period of enhanced palaeotidal amplitude but falling RSL (Figure 11). The continuity of
486 these RSL and palaeotidal trends throughout deglaciation imply that other factors; e.g.
487 climate, topography, ice sheet internal dynamics; were controlling the higher frequency
488 BDIS advance/readvance phases documented by the new data.

489 Finally, changes in ocean circulation that allow warmer water to access the calving
490 front (e.g. Holland et al., 2008) have been cited as a major factor in past deglaciations
491 (Marcott et al., 2011, Rinterknecht et al., 2014). Records of Nps% and $\delta^{18}\text{O}_{\text{Nps}}$ in MD04-
492 2822 and a Mg/Ca sea surface temperature estimate from MD01-2461 (Peck et al.,
493 2008) (Figure 10C, D, E) show a consistent trend indicating northerly migration of the
494 polar front during Greenland Interstadial 2 (GI-2). Scourse et al. (2009) cite this oceanic

495 warming as a driver of a major phase of BIIS deglaciation represented by high IRD
496 fluxes to the deep sea record from ~23 ka. That the BDIS was likely involved in this is
497 indicated by the IRD records from the proximal cores MD95-2006 (Knutz et al., 2001)
498 and MD04-2822 (Hibbert et al., 2010). The rate at which the BDIS deglaciated in
499 response to GI-2 remains unclear. The IRD record from MD04-2822 retains high IRD
500 fluxes 22-18 ka (Hibbert et al., 2010) indicating that the BIIS, and most likely the BDIS,
501 retained calving margins throughout this period. This implies deglaciation may have
502 been a continuous process with punctuated retreat across the shelf although additional
503 geochronological data from the mid-outer shelf is needed to provide further constraints
504 on the nature of BDIS deglaciation in response to GI-2.

505

506 **6. Conclusions**

507 The data presented here provide insights into the timing of deglaciation of a major
508 palaeo-ice stream that drained a large portion of the former BIIS as well as indicating
509 post-ice stream dynamics of the remnant ice mass. Following de-coupling of ice sourced
510 from mainland Scotland and ice sourced in Skye, our data lead us to hypothesise that
511 there were possibly two local readvances/stillstands at ~17.6 and ~15.2 ka demarked by
512 moraines in Glen Brittle and Loch Scavaig, respectively. Evidence for local readvance
513 of ice sourced in Skye occurs around the periphery of Cuillin and our data suggests that
514 the latter readvance, north and south of the Cuillin, was contemporaneous with the
515 Wester Ross Readvance recorded elsewhere in northern Scotland, strengthening the
516 conclusion that it was climatically forced.

517 The ³⁶Cl CRE ages from Glen Brittle provide constraints on the timing of final
518 deglaciation of a major ice stream that drained the former BIIS. They indicate that
519 deglaciation of the BDIS was complete by 17.6 ± 1.3 ka, in general agreement with

520 offshore IRD evidence. The complex production pathways associated with *in situ*-
521 produced ^{36}Cl lead to large inherent uncertainties on our data that prevent us from
522 definitively linking deglaciation of the BDIS and subsequent readvance to any one
523 forcing factor. Ultimately, disentangling the relative contribution of the various forcing
524 factors requires further data constraining ice margin retreat on the shelf combined with
525 new and more precise geochronological data that constrains final deglaciation.

526

527 **Acknowledgements**

528 We thank Joe Licciardi for laboratory access at the University of New Hampshire,
529 USA and preparation of ^{36}Cl targets. The French national AMS facility ASTER
530 (CEREGE, Aix en Provence) is supported by the INSU/CNRS, the ANR through the
531 "Projets thématiques d'excellence" program for the "Equipements d'excellence"
532 ASTER-CEREGE action, IRD and CEA. We would like to thank Shasta Marrero for
533 helpful and informative discussion on the CRONUScale online calculator. DS was
534 supported by a SAGES studentship and fieldwork by funds from the QRA and BSG.
535 Detailed comments from two anonymous reviewers have improved the quality and
536 clarity of this manuscript.

537 **References**

- 538 Alley, R.B. and MacAyeal, D.R., 1994. Ice-rafted debris associated with binge/purge
539 oscillations of the Laurentide Ice Sheet. *Paleoceanography*, 9, 503-511.
- 540 Arbic, B.K., Mitrovica, J.X., MacAyeal, D.R. and Milne, G.A., 2008. On the factors
541 behind large Labrador Sea tides during the last glacial cycle and the potential
542 implications for Heinrich events. *Paleoceanography*, 23, PA3211.
- 543 Austin, W.E.N. and Kroon, D., 1996. Late glacial sedimentology, foraminifera and
544 stable isotope stratigraphy of the Hebridean Continental Shelf, northwest
545 Scotland. *Geological Society, London, Special Publications*, 111, 187-213.
- 546 Arnold, M., Aumaître, G., Bourlès, D.L., Keddadouche, K., Braucher, R., Finkel, R.C.,
547 Nottoli, E., Benedetti, L. and Merchel, S., 2013. The French accelerator mass
548 spectrometry facility ASTER after 4years: Status and recent developments on
549 ^{36}Cl and ^{129}I . *Nuclear Instruments and Methods in Physics Research Section B:*
550 *Beam Interactions with Materials and Atoms*, 294, 24-28.
- 551 Ballantyne, C.K., 1989, The Loch Lomond Readvance on the Isle of Skye, Scotland:
552 glacier reconstruction and palaeoclimatic implications: *Journal of Quaternary*
553 *Science*, 4, 95-108.
- 554 Ballantyne, C.K., and Stone, J.O., 2012, Did large ice caps persist on low ground in
555 north-west Scotland during the Lateglacial Interstade?: *Journal of Quaternary*
556 *Science*, 27, 297-306.
- 557 Ballantyne, C.K. and Stone, J.O., 2013. Timing and periodicity of paraglacial rock-slope
558 failures in the Scottish Highlands. *Geomorphology*, 186, 150-161.
- 559 Ballantyne C.K., Benn, D.I., Lowe, J.J., Walker, M.J.C., (Eds.), 1991, The Quaternary
560 of the Isle of Skye: Field Guide, Quaternary Research Association, Cambridge.

561 Ballantyne, C.K., Schnabel, C. and Xu, S., 2009. Readvance of the last British–Irish ice
562 sheet during Greenland interstade 1 (GI-1): the Wester Ross readvance, NW
563 Scotland. *Quaternary Science Reviews*, 28, 783-789.

564 Ballantyne, C.K., Wilson, P., Gheorghiu, D., and Rodés, À., 2014, Enhanced rock-slope
565 failure following ice-sheet deglaciation: timing and causes: *Earth Surface
566 Processes and Landforms*, 39, 900-913.

567 Benn, D.I., 1997. Glacier fluctuations in western Scotland. *Quaternary International*,
568 38, 137-147.

569 Bennett, M.R., 2003. Ice streams as the arteries of an ice sheet: their mechanics, stability
570 and significance. *Earth-Science Reviews*, 61, 309-339.

571 Bennett, M.R. and Boulton, G.S., 1993. Deglaciation of the Younger Dryas or Loch
572 Lomond Stadial ice-field in the northern Highlands, Scotland. *Journal of
573 Quaternary Science*, 8, 133-145.

574 Baltzer, A., Bates, R., Mokeddem, Z., Clet-Pellerin, M., Walter-Simonnet, A.V.,
575 Bonnot-Courtois, C. and Austin, W.E., 2010. Using seismic facies and pollen
576 analyses to evaluate climatically driven change in a Scottish sea loch (fjord) over
577 the last 20 ka. *Geological Society, London, Special Publications*, 344, 355-369.

578 Berger, A., and Loutre, M.F., 1991, Insolation values for the climate of the last 10
579 million years. *Quaternary Science Reviews*, 10, 297-317.

580 Bevington, PR and Robinson, D.K., 2003, Data reduction and error analysis: McGraw–
581 Hill, New York, p. 320.

582 Bradwell, T. and Stoker, M.S., 2015. Submarine sediment and landform record of a
583 palaeo-ice stream within the British– Irish Ice Sheet. *Boreas*, 44, 255-276.

584 Bradwell, T., Stoker, M.S., Golledge, N.R., Wilson, C.K., Merritt, J.W., Long, D.,
585 Everest, J.D., Hestvik, O.B., Stevenson, A.G., Hubbard, A.L. and Finlayson,

586 A.G., 2008a. The northern sector of the last British Ice Sheet: maximum extent
587 and demise. *Earth-Science Reviews*, 88, 207-226.

588 Bradwell, T., Fabel, D., Stoker, M., Mathers, H., McHargue, L. and Howe, J., 2008b. Ice
589 caps existed throughout the Lateglacial Interstadial in northern Scotland. *Journal*
590 *of Quaternary Science*, 23, 401-407.

591 Braucher, R., Merchel, S., Borgomano, J. and Bourlès, D.L., 2011. Production of
592 cosmogenic radionuclides at great depth: A multi element approach. *Earth and*
593 *Planetary Science Letters*, 309, 1-9.

594 Clark, C.D., Evans, D.J., Khatwa, A., Bradwell, T., Jordan, C.J., Marsh, S.H., Mitchell,
595 W.A. and Bateman, M.D., 2004. Map and GIS database of glacial landforms and
596 features related to the last British Ice Sheet. *Boreas*, 33, 359-375.

597 Clark, C.D., Hughes, A.L., Greenwood, S.L., Jordan, C., and Sejrup, H.P., 2012, Pattern
598 and timing of retreat of the last British-Irish Ice Sheet: *Quaternary Science*
599 *Reviews*, 44, 112-146.

600 Clark, J., McCabe, A.M., Schnabel, C., Clark, P.U., McCarron, S., Freeman, S.P.,
601 Maden, C. and Xu, S., 2009. Cosmogenic ¹⁰Be chronology of the last
602 deglaciation of western Ireland, and implications for sensitivity of the Irish Ice
603 Sheet to climate change. *Geological Society of America Bulletin*, 121, 3-16.

604 Clough, C.T. and Harker, A., 1904. *The Geology of West-Central Skye, with Soay:*
605 *Explanation of* (Vol. 70). HM Stationery Office.

606 Chiverrell, R.C., Thrasher, I.M., Thomas, G.S., Lang, A., Scourse, J.D., van
607 Landeghem, K.J., Mccarroll, D., Clark, C.D., Cofaigh, C.Ó., Evans, D.J. and
608 Ballantyne, C.K., 2013. Bayesian modelling the retreat of the Irish Sea Ice
609 Stream. *Journal of Quaternary Science*, 28(2), pp.200-209.

610 De Deckker, P. and Yokoyama, Y., 2009. Micropalaeontological evidence for Late
611 Quaternary sea-level changes in Bonaparte Gulf, Australia. *Global and*
612 *Planetary Change*, 66, 85-92.

613 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L.,
614 Henderson, G.M., Okuno, J.I. and Yokoyama, Y., 2012. Ice-sheet collapse and
615 sea-level rise at the Bolling warming 14,600 years ago. *Nature*, 483, 559-564.

616 Dove, D., Arosio, R., Finlayson, A., Bradwell, T. and Howe, J.A., 2015. Submarine
617 glacial landforms record Late Pleistocene ice-sheet dynamics, Inner Hebrides,
618 Scotland. *Quaternary Science Reviews*, 123, 76-90.

619 Dowdeswell, J.A., Hogan, K.A., Cofaigh, C.Ó., Fugelli, E.M.G., Evans, J. and
620 Noormets, R., 2014. Late Quaternary ice flow in a West Greenland fjord and
621 cross-shelf trough system: submarine landforms from Rink Isbrae to
622 Uummannaq shelf and slope. *Quaternary Science Reviews*, 92, 292-309.

623 Dunlop, P., Shannon, R., McCabe, M., Quinn, R., and Doyle, E., 2010. Marine
624 geophysical evidence for ice sheet extension and recession on the Malin Shelf:
625 New evidence for the western limits of the British Irish Ice Sheet: *Marine*
626 *Geology*, 276, 86-99.

627 Fabel, D., Ballantyne, C.K. and Xu, S., 2012. Trimlines, blockfields, mountain-top
628 erratics and the vertical dimensions of the last British–Irish Ice Sheet in NW
629 Scotland. *Quaternary Science Reviews*, 55, 91-102.

630 Fifield, L.K., Ophel, T.R., Allan, G.L., Bird, J.R. and Davie, R.F., 1990. Accelerator
631 mass spectrometry at the Australian National University's 14UD accelerator:
632 experience and developments. *Nuclear Instruments and Methods in Physics*
633 *Research Section B: Beam Interactions with Materials and Atoms*, 52, 233-237.

634 Haapanieni, A.I., Scourse, J.D., Peck, V.L., Kennedy, D.P., Kennedy, H., Hemming,
635 S.R., Furze, M.F.A., Pieńkowski-Furze, A.J., Walden, J., Wadsworth, E. and
636 Hall, I.R. 2010. Source, timing, frequency and flux of ice-rafted detritus to the
637 Northeast Atlantic margin, 30-12 ka: testing the Heinrich precursor hypothesis.
638 *Boreas*, 39, 576-591.

639 Harker, A., 1901. Ice-erosion in the Cuillin Hills, Skye: *Transactions of the Royal*
640 *Society of Edinburgh*, 40, 221–252.

641 Heyman, J., Stroeven, A.P., Harbor, J.M., and Caffee, M.W., 2011, Too young or too
642 old: evaluating cosmogenic exposure dating based on an analysis of compiled
643 boulder exposure ages: *Earth and Planetary Science Letters*, 302, 71-80.

644 Hibbert FD. 2011. Dynamics of the British Ice Sheet and Prevailing Hydrographic
645 Conditions for the Last 175,000 years: An investigation of marine sediment core
646 MD04-2822 from the Rockall Trough. *Unpublished PhD thesis, University of St*
647 *Andrews*.

648 Hibbert, F.D., Austin, W.E., Leng, M.J., and Gatliff, R.W., 2010, British Ice Sheet
649 dynamics inferred from North Atlantic ice-rafted debris records spanning the last
650 175 000 years: *Journal of Quaternary Science*, 25, 461-482.

651 Holland, D.M., Thomas, R.H., De Young, B., Ribergaard, M.H., and Lyberth, B., 2008,
652 Acceleration of Jakobshavn Isbrae triggered by warm subsurface ocean waters:
653 *Nature Geoscience*, 1, 659-664.

654 Howe, J.A., Dove, D., Bradwell, T. and Gafeira, J., 2012. Submarine geomorphology
655 and glacial history of the Sea of the Hebrides, UK. *Marine Geology*, 315, 64-76.

656 Hubbard, A., Bradwell, T., Golledge, N., Hall, A., Patton, H., Sugden, D., Cooper, R.
657 and Stoker, M., 2009. Dynamic cycles, ice streams and their impact on the

658 extent, chronology and deglaciation of the British–Irish ice sheet. *Quaternary*
659 *Science Reviews*, 28, 758-776.

660 Joughin, I., Smith, B.E., and Medley, B., 2014, Marine ice sheet collapse potentially
661 under way for the Thwaites Glacier Basin, West Antarctica: *Science*, 344, 735-
662 738.

663 Knutz, P.C., Austin, W.E., and Jones, E.J.W., 2001, Millennial-scale depositional cycles
664 related to British Ice Sheet variability and North Atlantic paleocirculation since
665 45 kyr BP, Barra Fan, UK margin: *Paleoceanography*, 16, 53-64.

666 Knutz, P.C., Zahn, R., and Hall, I.R., 2007, Centennial-scale variability of the British Ice
667 Sheet: Implications for climate forcing and Atlantic meridional overturning
668 circulation during the last deglaciation: *Paleoceanography* 22.

669 Kristensen, L., Benn, D.I., Hormes, A. and Ottesen, D. 2009. Mud aprons in front of
670 Svalbard surge moraines: evidence of subglacial deforming layers or proglacial
671 tectonics? *Geomorphology* 111, 206-221.

672 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M., 2014, Sea level and
673 global ice volumes from the Last Glacial Maximum to the Holocene:
674 *Proceedings of the National Academy of Science of the United States of America*,
675 111, 15296-15303.

676 Lifton, N., Sato, T. and Dunai, T.J., 2014. Scaling in situ cosmogenic nuclide production
677 rates using analytical approximations to atmospheric cosmic-ray fluxes. *Earth*
678 *and Planetary Science Letters*, 386, pp.149-160.

679 Livingstone, S.J., Cofaigh, C.Ó., Stokes, C.R., Hillenbrand, C.D., Vieli, A. and
680 Jamieson, S.S., 2013. Glacial geomorphology of Marguerite Bay palaeo-ice
681 stream, western Antarctic Peninsula. *Journal of Maps*, 9, 558-572.

682 Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J. and
683 Yu, Z.C., 2008. Synchronisation of palaeoenvironmental events in the North
684 Atlantic region during the Last Termination: a revised protocol recommended by
685 the INTIMATE group. *Quaternary Science Reviews*, 27, 6-17.

686 Marcott, S.A., Clark, P.U., Padman, L., Klinkhammer, G.P., Springer, S.R., Liu, Z.,
687 Otto-Bliesner, B.L., Carlson, A.E., Ungerer, A., Padman, J. and He, F., 2011.
688 Ice-shelf collapse from subsurface warming as a trigger for Heinrich events.
689 *Proceedings of the National Academy of Sciences*, 108, 13415-13419.

690 Marrero, S.M., Phillips, F.M., Borchers, B., Lifton, N., Aumer, R. and Balco, G., 2016a.
691 Cosmogenic nuclide systematics and the CRONUScale program. *Quaternary*
692 *Geochronology*, 31, 160-187.

693 Marrero, S.M., Phillips, F.M., Caffee, M.W. and Gosse, J.C., 2016b. CRONUS-Earth
694 cosmogenic ³⁶Cl calibration. *Quaternary Geochronology*, 31, 199-219.

695 McCabe, A.M., and Clark, P.U., 2003, Deglacial chronology from County Donegal,
696 Ireland: implications for deglaciation of the British–Irish ice sheet: *Journal of the*
697 *Geological Society of London*, 160, 847-855.

698 Murray, T., Smith, A.M., King, M.A. and Weedon, G.P., 2007. Ice flow modulated by
699 tides at up to annual periods at Rutford Ice Stream, West Antarctica. *Geophysical*
700 *Research Letters*, 34, L18503.

701 Ó' Cofaigh, C., Dunlop, P. and Benetti, S., 2012. Marine geophysical evidence for Late
702 Pleistocene ice sheet extent and recession off northwest Ireland. *Quaternary*
703 *Science Reviews*, 44, 147-159.

704 Ottesen, D., Dowdeswell, J.A., Benn, D.I., Kristensen, L., Christiansen, H.H.,
705 Christensen, O., Hansen, L., Lebesbye, E., Forwick, M. and Vorren, T.O., 2008.

706 Submarine landforms characteristic of glacier surges in two Spitsbergen fjords.
707 *Quaternary Science Reviews*, 27, 1583-1599.

708 Peacock, J.D., 2008, Late Devensian palaeoenvironmental changes in the sea area
709 adjacent to Islay, SW Scotland: implications for the deglacial history of the
710 island: *Scottish Journal of Geology*, 44, 183-190.

711 Peacock, J.D., Austin, W.E.N., Selby, I., Graham, D.K., Harland, R. and Wilkinson, I.P.,
712 1992. Late Devensian and Flandrian palaeoenvironmental changes on the
713 Scottish continental shelf west of the Outer Hebrides. *Journal of Quaternary*
714 *Science*, 7, 145-161.

715 Peck, V.L., Hall, I.R., Zahn, R., and Elderfield, H., 2008, Millennial-scale surface and
716 subsurface paleothermometry from the northeast Atlantic, 55–8 ka BP:
717 *Paleoceanography*, 23, PA3221.

718 Peck, V.L., Hall, I.R., Zahn, R., Grousset, F., Hemming, S.R. and Scourse, J.D., 2007.
719 The relationship of Heinrich events and their European precursors over the past
720 60ka BP: a multi-proxy ice-rafted debris provenance study in the North East
721 Atlantic. *Quaternary Science Reviews*, 26(7), 862-875.

722 Ramsey, C.B., 2008. Deposition models for chronological records. *Quaternary Science*
723 *Reviews*, 27, 42-60.

724 Ramsey, C.B., 2013. OxCal 4.2. *Manual [online] available at: [https://c14.arch.ox.ac.](https://c14.arch.ox.ac.uk/oxcalhelp/hlp_contents.html)*
725 *uk/oxcalhelp/hlp_contents.html*.

726 Ramsey, C.B. and Lee, S., 2013. Recent and planned developments of the program
727 OxCal. *Radiocarbon*, 55, 720-730.

728 Rasmussen, S.O., Seierstad, I.K., Andersen, K.K., Bigler, M., Dahl-Jensen, D., and
729 Johnsen, S.J., 2008, Synchronization of the NGRIP, GRIP, and GISP2 ice cores

730 across MIS 2 and palaeoclimatic implications: *Quaternary Science Reviews*, 27,
731 18-28.

732 Rinterknecht, V., Jomelli, V., Brunstein, D., Favier, V., Masson-Delmotte, V., Bourlès,
733 D., Leanni, L. and Schläppy, R., 2014. Unstable ice stream in Greenland during
734 the Younger Dryas cold event. *Geology*, 42, 759-762.

735 Robinson, M. and Ballantyne, C.K., 1979. Evidence for a glacial readvance pre-dating
736 the Loch Lomond Advance in Wester Ross. *Scottish Journal of Geology*, 15,
737 271-277.

738 Rosier, S.H.R., Gudmundsson, G.H. and Green, J.A.M. 2015. Temporal variations in
739 the flow of a large Antarctic ice-stream controlled by tidally induced changes in
740 the subglacial water system. *Cryosphere*, 9, 2397-2429.

741 Schimmelpfennig, I., Benedetti, L., Finkel, R., Pik, R., Blard, P. H., Bourlès, D.,
742 Burnard, P., and Williams, A., 2009, Sources of in-situ ³⁶Cl in basaltic rocks.
743 Implications for calibration of production rates: *Quaternary Geochronology*, 4,
744 441-461.

745 Schimmelpfennig, I., Benedetti, L., Garreta, V., Pik, R., Blard, P.H., Burnard, P.,
746 Bourlès, D., Finkel, R., Ammon, K., and Dunai, T., 2011, Calibration of
747 cosmogenic ³⁶Cl production rates from Ca and K spallation in lava flows from
748 Mt. Etna (38° N, Italy) and Payun Matru (36° S, Argentina): *Geochimica et*
749 *Cosmochimic Acta*, 75, 2611-2632.

750 Schimmelpfennig, I., Schaefer, J.M., Putnam, A.E., Koffman, T., Benedetti, L., Ivy-
751 Ochs, S., ASTER team, and Schlüchter, C., 2014, ³⁶Cl production rate from K-
752 spallation in the European Alps (Chironico landslide, Switzerland): *Journal of*
753 *Quaternary Science*, 29, 407-413.

754 Scourse, J.D. and Furze, M.F.A., 2001. A critical review of the glaciomarine model for
755 Irish Sea deglaciation: evidence from southern Britain, the Celtic shelf and
756 adjacent continental slope. *Journal of Quaternary Science*, 16, 419-434.

757 Scourse, J.D., Hall, I.R., McCave, I.N., Young, J.R. and Sugdon, C., 2000. The origin of
758 Heinrich layers: evidence from H2 for European precursor events. *Earth and*
759 *Planetary Science Letters*, 182(2), pp.187-195.

760 Scourse, J.D., Haapaniemi, A.I., Colmenero-Hidalgo, E., Peck, V.L., Hall, I.R., Austin,
761 W.E., Knutz, P.C. and Zahn, R., 2009. Growth, dynamics and deglaciation of the
762 last British–Irish ice sheet: the deep-sea ice-rafted detritus record. *Quaternary*
763 *Science Reviews*, 28(27), 3066-3084.

764 Scourse, J.D., Ward, S.L., Wainwright, A., Bradley, S.L. and Uehara, K. Submitted. The
765 role of megatides and relative sea level in controlling the deglaciation of the
766 British-Irish and Fennoscandian Ice Sheets. *Journal of Quaternary Science*.

767 Seierstad, I.K., Abbott, P.M., Bigler, M., Blunier, T., Bourne, A.J., Brook, E., Buchardt,
768 S.L., Buizert, C., Clausen, H.B., Cook, E. and Dahl-Jensen, D., 2014.
769 Consistently dated records from the Greenland GRIP, GISP2 and NGRIP ice
770 cores for the past 104 ka reveal regional millennial-scale $\delta^{18}\text{O}$ gradients with
771 possible Heinrich event imprint. *Quaternary Science Reviews*, 106, pp.29-46.

772 Sejrup, H.P., Larsen, E., Landvik, J., King, E.L., Haflidason, H. and Nesje, A., 2000.
773 Quaternary glaciations in southern Fennoscandia: evidence from southwestern
774 Norway and the northern North Sea region. *Quaternary Science Reviews*, 19,
775 667-685.

776 Sissons, J.B., Lowe, J.J., Thompson, K.S. and Walker, M.J.C., 1973. Loch Lomond
777 readvance in Grampian Highlands of Scotland: *Nature*, 244, 75-77.

778 Small, D., 2012 The Deglaciation of the northwest sector of the last British-Irish Ice
779 Sheet: Integrating onshore and offshore data relating to chronology and
780 behaviour. *Unpublished PhD Thesis, University of St Andrews*.

781 Small, D., Rinterknecht, V., Austin, W., Fabel, D. and Miguens-Rodriguez, M., 2012,
782 In situ cosmogenic exposure ages from the Isle of Skye, northwest Scotland:
783 implications for the timing of deglaciation and readvance from 15 to 11 ka.
784 *Journal of Quaternary Science*, 27, 150-158.

785 Small D., Austin W., and Rinterknecht V., 2013. Freshwater influx, hydrographic
786 reorganization and the dispersal of ice-rafted detritus in the sub-polar North
787 Atlantic Ocean during the last deglaciation: *Journal of Quaternary Science*, 28,
788 527-535.

789 Stoker, M. and Bradwell, T., 2005. The Minch palaeo-ice stream, NW sector of the
790 British–Irish Ice Sheet. *Journal of the Geological Society*, 162, 425-428.

791 Stoker, M.S., Wilson, C.R., Howe, J.A., Bradwell, T. and Long, D., 2010. Paraglacial
792 slope instability in Scottish fjords: examples from Little Loch Broom, NW
793 Scotland. *Geological Society, London, Special Publications*, 344, 225-242.

794 Stokes, C.R. and Clark, C.D., 2001. Palaeo-ice streams. *Quaternary Science Reviews*,
795 20, 1437-1457.

796 Stokes, C.R., Tarasov, L., Blomdin, R., Cronin, T.M., Fisher, T.G., Gyllencreutz, R.,
797 Hättestrand, C., Heyman, J., Hindmarsh, R.C., Hughes, A.L. and Jakobsson, M.,
798 2015. On the reconstruction of palaeo-ice sheets: Recent advances and future
799 challenges. *Quaternary Science Reviews*, 125, 15-49.

800 Stone, J.O., 2000, Air pressure and cosmogenic isotope production: *Journal of*
801 *Geophysical Research: Solid Earth*, 105, 23753-23759.

802 Stone, J.O., Allan, G.L., Fifield, L.K. and Cresswell, R.G., 1996. Cosmogenic chlorine-
803 36 from calcium spallation. *Geochimica et Cosmochimica Acta*, 60, 679-692.

804 Stone, J.O., Ballantyne, C.K. and Fifield, L.K., 1998. Exposure dating and validation of
805 periglacial weathering limits, northwest Scotland. *Geology*, 26, 587-590.

806 Svendsen, J.I., Briner, J.P., Mangerud, J. and Young, N.E., 2015. Early break-up of the
807 Norwegian channel ice stream during the last glacial maximum. *Quaternary*
808 *Science Reviews*, 107, 231-242.

809 Uehara, K., Scourse, J.D., Horsburgh, K.J., Lambeck, K. and Purcell, A.P., 2006. Tidal
810 evolution of the northwest European shelf seas from the Last Glacial Maximum
811 to the present. *Journal of Geophysical Research: Oceans*, 111, C09025.

812 Walker, M.J., Ballantyne, C.K., Lowe, J.J. and Sutherland, D.G., 1988. A
813 reinterpretation of the Lateglacial environmental history of the Isle of Skye,
814 Inner Hebrides, Scotland. *Journal of Quaternary Science*, 3, 135-146.

815 Wouters, B., Martin-Español, A., Helm, V., Flament, T., van Wessem, J.M., Ligtenberg,
816 S.R.M., van den Broeke, M.R., and Bamber, J. L., 2015, Dynamic thinning of
817 glaciers on the Southern Antarctic Peninsula: *Science*, 348, 899-903.

818

819 **Figure Captions**

820 Figure 1. Google Earth Image with extent of the BDIS and related glaciological features.
821 Existing geochronological dates are shown (Table 1) along with location of marine core
822 MD04-2822. Flowlines adjusted from Bradwell et al. (2008). Dashed box shows the
823 location of Figure 2, solid box shows location of Figure 7. Isochrones depicting the most
824 likely ice extent at 24 ka, 17 ka, and 16 ka (shaded for clarity) are taken from Hughes et
825 al., 2016. BDF = Barra/Donegal Fan, BDIS = Barra/Donegal Ice Stream, MIS = Minch

826 Ice Stream. All ^{10}Be CRE dates have been re-calculated using a local production rate
827 (Loch Lomond Production Rate) of 3.92 ± 0.18 atoms $\text{g}^{-1} \text{a}^{-1}$ (Fabel et al., 2012).

828

829 Figure 2. Location map of Skye and northwest Scotland showing locations mentioned in
830 text. Dashed lines demark inferred zones of confluence between mainland ice and the
831 Skye Ice Dome. Red stars show locations of existing exposure ages from (1) Trotternish
832 (Stone et al., 1998) and (2) the Strollamus moraine (Small et al., 2012). GB = Glen
833 Brittle, LS = Loch Scavaig. Arrows show generalized ice flow directions, MIS = Minch
834 Ice Stream, BDIS = Barra-Donegal Ice Stream. Also shown are inferred limits of Wester
835 Ross Readvance (WRR). Letters A and B denote the locations of the palaeotidal and
836 RSL simulations (Section 5.2; Figure 11). DEM derived from NASNA SRTM 90 m
837 data, available at <http://www.sharegeo.ac.uk/handle/10672/5>.

838

839 Figure 3. Map of Glen Brittle area showing sampled moraines, raised shorelines and
840 locations of sampled boulders. The limits of the Loch Lomond Readvance and
841 associated landforms are shown as adapted from Ballantyne (1989). Contours at 100 m
842 intervals. See Figure 5 for location.

843

844 Figure 4. Map of the northeast corner of Soay showing sampled moraine and locations
845 of sampled boulders. Dashed line shows crest of offshore moraine (Figure 8).

846

847 Figure 5. Site and sample photographs from Glen Brittle. (A) Glen Brittle looking North.
848 Showing two parallel moraine ridges. Southern (outer) moraine with person, northern
849 (inner) moraine with boulders on near horizon. Ice flow is towards the camera. (B)
850 BRI01 boulder. (C) BRI-02 boulder. (D) BRI-03 boulder. (E) BRI-04 boulder.

851

852 Figure 6. Site and sample photographs from Soay. (A) SOAY-01 boulder. (B) SOAY-02
853 boulder. (C) SOAY-03 boulder. (D) SOAY-04 boulder. (E) SOAY-05 boulder. (F)
854 SOAY-06 boulder. (G) SOAY-07 boulder. (H) Soay moraine onshore. The dashed
855 white line marks the crest. The offshore continuation stretches across Loch Scavaig to
856 the far shore (see Figures 5 and 6). Samples were located off-shot in the wooded area to
857 the right. Ice flow was from left to right.

858

859 Figure 7. Location of ^{36}Cl samples presented and onshore moraines in Glen Brittle and
860 on Soay. The multibeam bathymetry of Loch Scavaig is shown alongside mapped YD
861 ice limits modified from Ballantyne (1989). Note the distinctive offshore moraine that
862 impinges on Soay. Failure scarp and extent of inferred slope failure (SF) is also shown.
863 The red star in the upper right is the location of the dated Strollamus medial moraine
864 (Benn, 1990; Small et al., 2012). NEXTmap hillshade DEM by Intermap Technologies.

865

866 Figure 8. Interpreted bathymetric map of Loch Scavaig showing the distinctive arcuate
867 terminal moraine. Suites of recessional moraines are also highlighted. There is a
868 distinctive glacially over-deepened basin in the western portion of the survey area. The
869 trench in the northeastern sector is the offshore continuation of the Camasunary Fault.
870 The red star shows the location of the vibrocore VC57/-07/844 which yielded a basal
871 radiocarbon age of 12.8 ± 0.1 ka (Small, 2012). Also show is the failure scarp on Ben
872 Cleat and the associated landslide deposits. NEXTmap hillshade DEM by Intermap
873 Technologies.

874

875 Figure 9. Summary CRE age plot of ^{36}Cl samples presented here shown alongside the
876 NGRIP oxygen isotope record ($\delta^{18}\text{O}$, ‰) (Rasmussen et al., 2008). Grey boxes show
877 arithmetic means and uncertainties of Brittle and Soay samples respectively. The Soay
878 samples not included in calculating moraine ages shown with hollow circles.
879 Uncertainties are 1σ analytical uncertainties. The Younger Dryas stadial (YD) and
880 Bølling-Allerød interstadial are also shown (B-A).

881

882 Figure 10. Proxy records of deglacial forcing for the time period of BDIS deglaciation
883 indicated by the shaded column. (A) Greenland oxygen isotope records ($\delta^{18}\text{O}$, ‰) from
884 NGRIP, GRIP and GISP2 on the GICC05 timescale (Rasmussen et al., 2008; Seierstad
885 et al., 2014) [50 yr moving averages shown by black line] (B) Reconstructed ESL
886 (Lambeck et al., 2014). Proxies relating to oceanic forcing: (C) Mg/Ca (*G.bulloides*)
887 SST estimates from MD01-2461 (Porcupine Seabight, Peck et al., 2008); and MD04-
888 2822 (Rockall Trough, Hibbert, 2011; Hibbert et al., 2010) (D) $\delta^{18}\text{O}$ *N.pachyderma*
889 sinistral (‰ VPDB), (E) % *N.pachyderma* (sinistral), (F) XRF core scanning (ITRAX)
890 TiCa (proxy for terrigenous input) and, (G) total IRD flux ($> 150 \mu\text{m cm}^{-2} \text{ka}^{-1}$).

891

892 Figure 11. Relative sea level (RSL) and palaeotidal (PTM) simulations for two locations
893 in the inner part of the BDIS adjacent to Skye. A) 57.04°N , 6.88°W and, B) 57.12°N ,
894 6.13°W (see Figure 2 for locations). RSL simulations are based on the modified glacio-
895 isostatic adjustment model of Lambeck and PTM simulations on a modified version of
896 the Princeton Ocean Model forced with dynamic open ocean tide (Uehara et al., 2006).
897 These show mean *M2* tidal ranges $> 6 \text{ m}$ throughout the deglacial phase from the Last
898 Glacial Maximum to around 11 ka BP (spring tidal ranges would have been significantly

899 larger). The shaded boxes in A and B show the mean exposure ages from Glen Brittle

900 and Soay, respectively.

901

902 Table 1. Published ages referred to in the text and shown on Figure 1. Outliers are
 903 shown in italics. Clusters of CRE ages that yield acceptable χ_R^2 values are shown in
 904 bold, the mean of these is shown in Figure 1. Underlined radiocarbon ages are the oldest
 905 from a site and these are used in Figure 1. CRE ages calculated using CRONUS online
 906 calculator (<http://hess.ess.washington.edu>; accessed April 20th 2016), Lm scaling and,
 907 Loch Lomond Production Rate of 3.92 ± 0.18 atoms $\text{g}^{-1} \text{yr}^{-1}$ (Fabel et al. 2012). ^{14}C ages
 908 calibrated using OxCal 4.2 (Bronk-Ramsey 2013) and Marine14 (Reimer et al.,2013),
 909 $\Delta R=300$ yr.

Reference	Location (site no. Fig. 1)	Sample name	Technique	Age (yr)	Uncert. (yr)
Clark et al. (2009)	N Donegal coast (1)	BF-04-01	CRE	17607	1772
Clark et al. (2009)	N Donegal coast (1)	BF-04-03	CRE	33035	2940
Clark et al. (2009)	N Donegal coast (1)	BF-04-04	CRE	21463	1754
Clark et al. (2009)	N Donegal coast (1)	BF-04-05	CRE	20924	1863
Clark et al. (2009)	N Donegal coast (1)	BF-04-06	CRE	20949	2060
Clark et al. (2009)	N Donegal coast (1)	BF-04-08	CRE	23251	2135
Clark et al. (2009)	N Donegal coast (1)	BF-04-09	CRE	21428	2196
Clark et al. (2009)	N Donegal coast (1)	BF-04-10	CRE	21799	2190
McCabe & Clark (2003)	N Donegal coast (2)	AA32315	^{14}C	16602	178
McCabe & Clark (2003)	N Donegal coast (2)	AA45968	^{14}C	18676	168
McCabe & Clark (2003)	N Donegal coast (2)	AA45967	^{14}C	17997	188
McCabe & Clark (2003)	N Donegal coast (2)	AA45966	^{14}C	19093	496
McCabe & Clark (2003)	N Donegal coast (2)	AA33831	^{14}C	17913	130
<u>McCabe & Clark (2003)</u>	<u>N Donegal coast (2)</u>	<u>AA33832</u>	<u>^{14}C</u>	<u>20308</u>	<u>148</u>
Peacock (2008)	Islay (3)	SUERC-13122	^{14}C	14457	163
Peacock (2008)	Islay (3)	SUERC-13123	^{14}C	14337	149
<u>Peacock (2008)</u>	<u>Islay (3)</u>	<u>SUERC-13124</u>	<u>^{14}C</u>	<u>14498</u>	<u>166</u>
Ballantyne et al. (2014)	Jura (4)	SNC-02	CRE	14006	1690
Ballantyne et al. (2014)	Jura (4)	SNC-03	CRE	12352	1414
Ballantyne et al. (2014)	Jura (4)	SNC-06	CRE	16875	1102
Ballantyne et al. (2014)	Jura (4)	SNC-07	CRE	16819	1025
Baltzer et al. (2010)	W coast of Scotland (5)	UL2853	^{14}C	16587	311
Small et al., (2013)	Mid Shelf (5)	AAR-2606	^{14}C	16664	279

910

911

912 Table 2. Sample information for all ³⁶Cl samples from Glen Brittle and Soay.

Sample Name	Lat.	Long.	Elevation (m)	Shielding correction	Sample thickness (cm)	Lithology	Density (g/cm)
<u>Glen Brittle</u>							
BRI01	57.21595	-6.29651	10	0.9891	2.3	Basalt	2.6
BRI02	57.21652	-6.29641	11	0.9891	3.2	Basalt	2.6
BRI03	57.21667	-6.29678	10	0.9891	1.5	Basalt	2.6
BRI04	57.21602	-6.29554	11	0.9891	2.2	Basalt	2.6
<u>Isle of Soay</u>							
SOAY01	57.16073	-6.18362	13	0.9993	2.5	Gabbro	2.6
SOAY02	57.16079	-6.18352	14	0.9993	1.4	Gabbro	2.6
SOAY03	57.16118	-6.18385	15	0.9993	1.5	Gabbro	2.6
SOAY04	57.16125	-6.18392	15	0.9993	1.7	Gabbro	2.6
SOAY05	57.16120	-6.18389	9	0.9993	1.5	Gabbro	2.6
SOAY06	57.16067	-6.18340	10	0.9993	1.4	Gabbro	2.6
SOAY07	57.16076	-6.18362	15	0.9993	1.6	Gabbro	2.6

913

914

915 Table 3. Chemical and analytical data for all ^{36}Cl samples. Ratios are rounded to two
 916 significant figures. Calculated concentrations reflect precision of AMS measurements.

Sample Name	Sample mass (g)	Carrier added (g)	$^{35}\text{Cl}/^{37}\text{Cl}$	Uncert. (%)	$^{36}\text{Cl}/^{35}\text{Cl}$	Uncert. (%)	$^{36}\text{Cl}/^{37}\text{Cl}$	Uncert. (%)	^{36}Cl conc. (at g^{-1})	Uncert. (abs)
<u>Glen</u>										
<u>Brittle</u>										
BRI01	15.1294	0.4844	9.55E+01	0.931	5.73E-14	6.576	5.46E-12	6.548	110167	7943
BRI02	14.9876	0.4824	1.08E+02	1.216	3.81E-14	6.223	4.11E-12	6.180	70422	5140
BRI03	15.0566	0.4818	1.05E+02	0.646	5.87E-14	5.557	6.16E-12	5.509	112557	6893
BRI04	12.9649	0.4824	1.30E+02	0.692	4.55E-14	8.045	5.91E-12	8.012	98678	8902
<u>Soay</u>										
SOAY01	20.0777	0.4853	5.59E+01	0.571	6.92E-14	4.806	3.86E-12	4.751	98972	5545
SOAY02	20.0711	0.4853	1.73E+01	0.253	6.81E-14	5.188	1.18E-12	5.135	113848	6786
SOAY03	20.0162	0.4787	2.40E+01	0.345	5.70E-14	5.297	1.37E-12	5.247	86176	5447
SOAY04	20.0341	0.478	2.26E+01	0.535	3.79E-14	6.449	8.56E-13	6.408	53701	4515
SOAY05	16.8693	0.4781	2.33E+01	0.883	5.68E-14	5.147	1.32E-12	5.096	108742	6183
SOAY06	20.1048	0.4816	6.39E+00	0.276	6.34E-14	5.954	4.04E-13	5.909	179705	12009
SOAY07	19.9611	0.4818	7.73E+00	0.379	6.19E-14	5.31	4.78E-13	5.262	150704	8986

917

918

Table 4. Whole rock geochemistry of samples from Glen Brittle and Soay.

Sample Name	SiO ₂ (wt-%)	Na ₂ O (wt-%)	MgO (wt-%)	Al ₂ O ₃ (wt-%)	MnO (wt-%)	H ₂ O (wt-%)	Sm (ppm)	Gd (ppm)	K ₂ O (wt-%)	CaO (wt-%)	Cl (ppm)	TiO ₂ (wt-%)	Fe ₂ O ₃ (wt-%)	P ₂ O ₅ (wt-%)	U (ppm)	Th (ppm)
<u>Glen Brittle</u>																
BRI01	46.19	1.96	8.62	13.17	0.17	2.42	2.61	3.25	0.301	12.16	2.76	1.85	13.06	0.01	0.04	0.16
BRI02	41.99	1.60	13.1	12.04	0.2	2.18	3.9	4.39	0.18	7.90	2.13	2.78	17.85	0.06	0.08	0.32
BRI03	47.64	1.99	8.52	13.68	0.16	2.01	7.26	3.09	0.18	11.97	2.25	1.77	11.98	0.02	0.04	0.15
BRI04	46.71	1.75	8.92	12.05	0.18	1.83	2.66	3.3	0.26	13.08	1.53	2.10	13.01	0.02	0.04	0.12
<u>Soay</u>																
SOAY01	45.77	0.96	11.07	20.13	0.09	0.44	0.5	0.82	0.03	14.08	4.69	0.27	7.11	0.02	0.02	0.11
SOAY02	44.55	1.03	12.49	20.96	0.09	0.41	1.30	0.69	< 0.005	12.99	23.69	0.25	7.17	0.02	< 0.01	0.03
SOAY03	44.92	1.01	13.01	22.26	0.06	0.48	0.15	0.33	< 0.005	13.16	15.14	0.1	4.97	0.01	< 0.01	0.02
SOAY04	42.94	0.07	28.02	10.14	0.13	0.62	0.28	0.5	< 0.005	7.28	16.32	0.18	10.34	0.01	0.01	0.06
SOAY05	47.12	1.58	1.88	29.28	0.04	0.56	0.56	0.87	0.02	16.46	19.06	0.34	2.74	0.02	0.02	0.10
SOAY06	47.08	1.25	5.76	23.44	0.09	1.30	0.40	0.71	0.06	15.87	109.82	0.19	4.90	0.02	< 0.01	0.01
SOAY07	48.15	0.51	12.79	8.85	0.18	1.36	1.57	2.66	0.04	15.74	77.84	0.63	11.63	0.02	< 0.01	0.02

920 Table 5. Comparison of CRE ages from Skye calculated using alternative calculation
 921 methods and scaling schemes. Full uncertainties (analytical uncertainties). CRE ages
 922 used in interpretation highlighted in bold text.

Calc. method	<i>Schimmelpfenig et al. (2009)</i>		<i>Marrero et al. (2016a)</i>		<i>Marrero et al. (2016a)</i>	
	Age	Uncert.	Age	Uncert.	Age	Uncert.
Prod. rates	<i>Marrero et al. (2016b)</i>		<i>Marrero et al. (2016b)</i>		<i>Marrero et al. (2016b)</i>	
Scaling	<i>Lm</i>		<i>Lm</i>		<i>SA</i>	
SOAY1	17.2	2.1 (1.5)	17.0	1.8 (1.5)	15.0	1.3 (0.9)
SOAY2	19.0	2.3 (1.5)	19.0	2.0 (1.5)	16.4	1.5 (1.0)
SOAY3	14.9	1.8 (1.4)	14.6	1.6 (1.4)	12.9	1.2 (0.8)
SOAY4	15.0	1.9 (1.6)	14.7	1.8 (1.6)	13.0	1.4 (1.1)
SOAY5	15.2	1.8 (1.0)	14.9	1.5 (1.0)	13.1	1.1 (0.8)
SOAY6	17.6	2.5 (1.1)	16.9	2.4 (1.1)	14.8	1.8 (1.0)
SOAY7	17.2	2.2 (0.9)	17.0	2.0 (0.9)	14.6	1.5 (0.9)
BRI01	19.0	2.4 (1.4)	20.6	2.2 (1.5)	18.2	1.7 (1.3)
BRI02	18.9	2.2 (1.4)	19.4	2.1 (1.5)	17.3	1.6 (1.3)
BRI03	21.9	2.6(1.4)	22.0	2.3 (1.4)	19.4	1.7 (1.2)
BRI04	17.3	2.1 (1.6)	17.5	2.1 (1.6)	15.5	1.7 (1.4)

923