

Landscape-scale effects of single- and multiple small wind turbines on bat activity

Journal:	<i>Animal Conservation</i>
Manuscript ID	ACV-02-16-OM-038.R3
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Minderman, Jeroen; University of St Andrews, School of Biology Gillis, Mairi; University of Stirling, Biological and Environmental Sciences Daly, Helen; University of Stirling, Biological and Environmental Sciences Park, Kirsty; University of Stirling,
Keywords:	conservation, domestic turbines, microturbines, planning, renewable energy, cumulative effects

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Landscape-scale effects of single- and multiple small wind turbines on bat activity**

2 *Jeroen Minderman*^{1*}, *Mairi H. Gillis*^{1,2}, *Helen F. Daly*^{1,3} & *Kirsty J. Park*¹

3 1. Biological & Environmental Sciences, University of Stirling, Stirling, FK9 4LA, United
4 Kingdom.

5 2. 228 Galashiels Road, Stow, Galashiels, Selkirkshire, TD1 2RA, mgillis4a7i@gmail.com

6 3. 20/11 Duff Street, Dalry, Edinburgh, Midlothian, EH11 2HG, hfdaly@outlook.com

7 **Corresponding author*. Present address: School of Biology, University of St Andrews, St
8 Andrews, Fife, KY16 9ST, UK. Email: jm340@st-andrews.ac.uk. +44 1334 463362.

9
10 **Short title:** Landscape-scale effects of multiple small wind turbines

11
12 **Word count:** 4808 (excluding references/tables/figures)

13

1
2
3 14 **ABSTRACT**
4
5

6 15 While the effects of wind farms on bats are widely studied, effects of small wind turbines
7
8 16 (SWTs, here <25m hub height) remain understudied. SWTs are installed in a wider range of
9
10 17 habitats compared to wind farms and their effect on wildlife can therefore be different. While
11
12 18 single SWTs can adversely affect bat activity in their immediate vicinity, they are often installed
13
14 19 in clusters, and to date no data are available on whether installation of more than one turbine
15
16 20 increases the likelihood of adverse effects on bats.
17

18
19 21 Here, we test whether (1) SWT installations affect the activity of two species of bat (the common
20
21 22 pipistrelle *Pipistrellus pipistrellus* and soprano pipistrelle *P. pygmaeus*) on a landscape scale
22
23 23 (here defined as up to 500m from SWTs) and (2) such an effect is stronger when multiple (2-4)
24
25 24 SWTs are installed. We show that, after accounting for potentially confounding effects (e.g.
26
27 25 variation in habitat and weather), (1) mean *P. pipistrellus* activity is lower at 0-100m compared
28
29 26 to 200-500m from SWTs, and (2) the effect on *P. pygmaeus* activity tends to be similar and
30
31 27 stronger in multiple SWT sites, although evidence for the latter is limited.
32

33
34 28 We conclude that in some cases, adverse effects of SWTs on bat activity may be measurable
35
36 29 over longer spatial scales (within 100m) than previously thought. However, combined with
37
38 30 earlier findings, it is likely that the bulk of such effects operate within relative close proximity of
39
40 31 SWTs (<25m). Moreover, although these effects may be species-specific, with e.g. *P. pygmaeus*
41
42 32 potentially more strongly affected by multiple SWT sites, this requires further data. These
43
44 33 findings are highly relevant to decision-making aimed at minimizing any adverse effects of wind
45
46 34 turbines, specifically single- vs. multiple SWT developments, on wildlife.
47
48
49
50
51
52
53
54

55 35
56
57
58
59
60

1
2
3 36 **KEY WORDS:** cumulative effects, microturbines, domestic turbines, renewables, renewable
4
5 37 energy, conservation, planning
6
7

8
9 38 **INTRODUCTION**

10
11 39 Understanding and predicting how wildlife is affected by disturbance or habitat loss is becoming
12
13 40 increasingly important as anthropogenic pressures on the natural world continue to grow
14
15 41 (Tuomainen & Candolin 2011). This is particularly pertinent where incremental changes or
16
17 42 disturbances could combine to produce disproportionately greater (adverse) effects. For example,
18
19 43 whilst the rapid growth of the wind energy sector worldwide provides a vital contribution to
20
21 44 sustainable energy production, a growing number of studies show that wind farms can have
22
23 45 adverse effects on wildlife (Kuvlesky *et al.* 2007, Zwart *et al.* 2015), both directly through
24
25 46 collision mortality (e.g. Rydell *et al.* 2010) and indirectly through disturbance or displacement
26
27 47 effects (e.g. Pearce-Higgins *et al.* 2009). Moreover, wind farms with larger numbers of turbines
28
29 48 installed can have greater effects on wildlife compared to smaller sites, even if e.g. collision risk
30
31 49 per turbine is relatively low (Langston & Pullan 2003, Zwart *et al.* 2015).
32
33
34
35
36
37

38 50 By contrast, similar effects of small wind turbines (SWTs) on wildlife remain relatively
39
40 51 unknown. These turbines are not only much smaller in size (in the UK, typical units are between
41
42 52 6 and 25m hub height) than those in wind farms, they are also commonly installed in a much
43
44 53 wider range of habitats. Thus, their impacts on wildlife are likely to be different to that of large
45
46 54 turbines, but up until recently this had not been quantified systematically. In addition to direct
47
48 55 mortality (Minderman *et al.* 2014), our previous work found evidence for adverse effects of
49
50 56 SWTs on bat activity (e.g. avoidance behaviour, Minderman *et al.* 2012). Whilst this is likely to
51
52 57 lower the risk of collision, such avoidance behaviour could have important population-level
53
54 58 consequences in itself, as it can amount to effective habitat loss (Minderman *et al.* 2012).
55
56
57
58
59
60

1
2
3 59 However, these previous studies focused on a relatively limited spatial scale (e.g. within 25m of
4
5
6 60 installed SWTs) and on single SWTs only. Crucially, it is therefore unclear whether any such
7
8 61 adverse effects of SWTs on wildlife could be greater at sites with more than one turbine
9
10 62 installed, i.e. whether the installation of multiple SWTs causes a disproportionately greater effect.

11
12
13 63 This lack of understanding of the potential disproportionate effects of sites with multiple SWTs
14
15
16 64 is particularly important for two reasons. First, the number of SWTs installed is growing rapidly
17
18 65 with over 900,000 units installed worldwide by 2014 (compared to 656,000 in 2010) and this
19
20 66 growth is set to continue (Pitteloud and Gsänger 2016). Secondly, SWTs are increasingly being
21
22
23 67 installed in groups, for example between two and four turbines in close proximity. Indeed, some
24
25 68 companies specifically promote the installation of multiple SWTs to maximize yield and
26
27
28 69 eligibility for government grants or payback schemes. As a result, where SWT installation
29
30 70 requires planning permission (Park, Turner & Minderman 2013), decision makers are
31
32 71 increasingly asked to decide whether multiple-turbine installations would have greater effects on
33
34
35 72 wildlife, or if limiting the number of SWTs to be installed would be an effective mitigation
36
37 73 option. Currently, the evidence base for such decisions for SWTs specifically is entirely lacking.

38
39
40 74 Here, we aim to (1) quantify the effect of SWTs on bat activity (and thus concomitant habitat
41
42 75 loss) on a wider landscape scale, up to 500m from installed turbines (as opposed to 25m in
43
44
45 76 Minderman *et al.* 2014), and (2) to test whether this effect is different between single- and
46
47 77 multiple (2-4) turbine installations. Using data collected at 34 SWT sites throughout the UK we
48
49
50 78 test the following predictions: (1) bat activity is systematically lower in closer proximity (e.g. 0-
51
52 79 100m from SWTs compared to 100-200m, 200-300m, etc.) of operating SWTs, controlling for
53
54
55 80 the effects of habitat and environmental conditions; and (2) the effect of SWT proximity on bat
56
57 81 activity is stronger in sites with multiple (2-4) SWTs installed compared to single SWT sites.
58
59
60

1
2
3 82
4
5
6 83 **METHODS**
7
8
9 84 **Sites**
10
11
12 85 Data were collected at 34 SWT sites in central and eastern Scotland. Sites were selected from an
13 86 existing database of owners (Minderman *et al.* 2012; Park, Turner & Minderman 2013) based on
14 87 how representative they were of common UK SWT installation settings in terms of habitat,
15 88 turbine models and size. All turbines studied here were free-standing and between 6 and 25m
16 89 (mean 16m, 75% of sites had turbines between 15 and 18m; including hub height in the analyses
17 90 presented here does not affect results or conclusions – see Supplementary Material) in hub
18 91 height, and all were in rural settings but specific habitat in the surrounding area varied (see
19 92 [Habitat data and variable selection](#)). The number of SWTs installed in each site varied between
20 93 one and four (median 1.6) and the distance between individual turbines in a site was between 12
21 94 and 90m (with only one site >62m and a mean of 46.4m). SWTs installed in individual sites were
22 95 the same size and specification with the exception of one site. More than two SWTs were
23 96 installed in only five sites in our sample. We therefore restricted the analyses presented here to a
24 97 comparison of single and multiple turbine sites; hence we here refer to sites with 2-4 SWTs
25 98 installed as “multiple turbine” sites throughout the remainder of this manuscript. Although this
26 99 limits possible inference from our findings somewhat, we are confident that this reflects the
27 100 current distribution of numbers of turbines per site in the UK; i.e. the majority of sites have one
28 101 or two SWTs installed (*pers. obs.*).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53 102 **Bat data and transects**
54
55
56
57
58
59
60

1
2
3 103 Bat activity data were collected between 28 May and 1 September 2013 and 7 July and 4
4
5 104 September 2014. This period reflects the seasonal activity of the species studied here (Swift
6
7
8 105 2009). The time of data collection varied but started 30 minutes after sunset at the earliest and
9
10 106 finished at least 60 minutes before sunrise in all cases.

11
12
13 107 Bat activity was measured along transects by 1-2 observers walking the length of each transect at
14
15 108 a slow (approximately 2.5 km h⁻¹) and constant pace, using EchoMeter EM3+ bat detectors
16
17
18 109 (WildLife Acoustics, Mass., USA). A target of four transects was planned for all sites, running
19
20 110 out from the turbine (or the central point between turbines in the case of multiple turbine sites) in
21
22 111 four directions. However, because of physical constraints (e.g. walls, impassable fences or
23
24 112 ditches, houses or buildings) the actual number of transects per site as well as their length varied
25
26
27 113 (3-4 transects, mean 3.7 per site, length: 300-500m). All transects were placed using a
28
29 114 combination of GIS-based planning and on-site assessment of landscape variation, so that (1) the
30
31 115 combination of all transects within each site covered all major habitats present, and (2) the
32
33 116 overall distance separating each transect was maximized. Transects were divided into 100m
34
35 117 sections running out from the turbine centre point, giving up to five distance bands running away
36
37 118 from the turbine. See Figure A1 for an example transect and transect section layout.

38
39
40 119 One measure of ground level wind speed was taken at the end of each section (i.e. in each
41
42 120 distance band) on each survey visit using handheld anemometers. Minimum daily temperature
43
44 121 measures for each survey visit were obtained from the UK MIDAS weather station data at
45
46 122 Grangemouth (N 56° 1' 5.15, W 3° 43' 5.88, between 5.5 and 92.6 km, average 52.4km, from the
47
48 123 sites) (UK Meteorological Office 2006).

49
50
51
52
53
54
55 124 **Habitat data and variable selection**
56
57
58
59
60

1
2
3 125 To account for expected confounding effects of habitat variation on bat activity we collated
4
5 126 measurements of habitat availability per transect section. To do this, we placed 50m buffers
6
7
8 127 around digital maps of each transect route, resulting in approximately 100m x 100m transect
9
10 128 sections. The exact area of each section varied because of non-linear transect sections, but this
11
12
13 129 was accounted for in the analysis, see [Statistical analysis](#) and Figure A1. All habitat variables
14
15 130 were quantified in each transect section using 1:1250 UK Ordnance Survey MasterMap
16
17 131 Topography digital maps, using QGIS v. 1.8 (<http://www.qgis.org/en/site/>). We focused on two
18
19
20 132 key measures of habitat variability in each transect section. These two variables were selected on
21
22 133 the basis of a preliminary analysis (See Supplementary Material) of the effect of a full set of 10
23
24 134 habitat variables on bat activity. This was done to avoid both overparameterisation of the models,
25
26
27 135 as well as the inclusion of highly collinear habitat metrics. First, we used **edge density** (m m^{-2})
28
29 136 as a measure of the density of "edge" habitat (defined as woodland and water edges, hedgerows,
30
31 137 roads and tracks, roadsides, field boundaries and building edges), calculated as the total length of
32
33
34 138 all such features divided by the area of the transect section. Although the specific effect of these
35
36 139 pooled habitat features on bat activity may vary, all tend to be associated with high levels of bat
37
38 140 activity relative to open and/or homogenous habitat (e.g. through their use as commuting routes)
39
40
41 141 (Walsh & Harris 1996). Second, we used the **proportion of woodland** calculated as the sum of
42
43 142 all tree coverage (m^2) (coniferous, non-coniferous and unclassified trees) divided by the transect
44
45
46 143 section area.

49 144 **Data analysis and statistics**

52 145 ***Bat activity: probability of a pass per hectare surveyed***

54 146 Bat activity was initially quantified as the number of bat passes (defined as a sequence of at least
56
57 147 two "search-phase" echolocation calls separated by less than a second) per transect section.
58
59
60

1
2
3 148 However, we chose to analyse our data as bat activity presence or absence per transect section,
4
5 149 per survey visit, because the distribution of observed counts was highly skewed (many zeros and
6
7
8 150 excessive variation) so that count-based statistical models did not provide any reasonable fit.
9
10 151 Thus, throughout the remainder of this paper, bat activity is measured as the probability of at
11
12 152 least one bat pass per unit area. Foraging calls (“buzzes”) were recorded in some cases, but in too
13
14
15 153 small numbers to be included in the analysis presented here. In addition, because the area
16
17 154 covered by each transect section varied slightly (see [Bat data and transects](#)), we here model the
18
19
20 155 probability of detecting bat activity per hectare surveyed within each section.
21

22 23 156 *Statistical analysis*

24
25 157 We used generalized linear mixed effects models (GLMMs) (Gelman & Hill 2007) to analyse
26
27
28 158 our data.
29

30 31 159 *Model structure*

32
33 160 We constructed two models for bat activity; one *P. pipistrellus* and one for *P. pygmaeus* activity.
34
35 161 The structure of both models was the same. In each case, activity was modelled as the probability
36
37 162 of a bat pass per unit area on a given survey visit as the response variable with a GLMM with
38
39 163 binomial errors and a complementary log-log link function (*clog-log*). This link function allowed
40
41 164 us to include transect section area as an offset in the model on the appropriate scale (Agresti
42
43 165 2013). In turn, this offset accounts for slight variation in the size of each transect section (see [Bat](#)
44
45 166 [data and transects](#) and Figure A1), modelling probability of a bat pass per unit area. To account
46
47 167 for the non-independence of repeated measurements from the same site and transect sections, all
48
49 168 models included transect nested within site as a random effect. To test our two predictions (see
50
51 169 [Introduction](#)), we included two focal fixed factors; (1) transect section (distance bands; 0-100m,
52
53 170 100-200m, 200-300m, 300-400m and 400-500m from the SWTs) and (2) turbine number (single
54
55
56
57
58
59
60

1
2
3 171 or multiple SWTs), as well as the interaction between the two. In addition to these focal factors
4
5 172 we included the two selected habitat variables (edge density and proportion of woodland; see
6
7
8 173 [Habitat data and variable selection](#)) as continuous covariates. Moreover, because bat activity is
9
10 174 known to vary with weather conditions (particularly temperature and wind speed), across the
11
12 175 season and through the night, we also included covariates for the minimum temperature over 24
13
14
15 176 hours ($^{\circ}\text{C}$), wind speed (m s^{-1}), day number (1 January = 1), time to midnight (minutes, with
16
17 177 negative and positive values indicating before and after midnight respectively) and time to
18
19
20 178 midnight² (to account for potential non-linear effects of the latter). Finally we included year as a
21
22 179 factorial predictor to account for potentially systematic differences in bat activity between the
23
24 180 two years of the study. Thus, accounting for any confounding effects of habitat and
25
26
27 181 environmental conditions, a transect section effect would indicate a systematic difference in bat
28
29 182 activity as a function of distance from SWTs (Prediction 1), and an interaction between this and
30
31
32 183 turbine number would indicate that the strength of this effect depends on the number of SWTs
33
34 184 installed (Prediction 2).

35
36
37 185 To avoid overparameterisation of the models, interactions other than those specified
38
39 186 above (i.e. between confounding effects) were not considered. All inputs were standardised
40
41
42 187 (centred to 0 and scaled to 1 SD) to improve performance of parameter estimation and to allow
43
44 188 direct interpretation of relative effect sizes. Marginal and conditional R^2 were presented as
45
46 189 measures of model fit (Nakagawa & Schielzeth 2013). We calculated and presented model
47
48
49 190 predictions on the response (measured) scale. Summary statistics of unstandardised model inputs
50
51 191 are given in Table A1.

52
53
54 192 *Model predictions and interpretation*

1
2
3 193 We chose to present full models only, to avoid issues with any form of model simplification
4
5 194 (Cade 2015, Whittingham *et al.* 2006). We interpreted the model outputs through assessment of
6
7
8 195 (1) relative effect sizes using standardised inputs, and (2) 95% quantiles of predictions of the
9
10 196 response at each of the levels of interest (specifically the number of turbines and distance band)
11
12 197 at average values for the remaining covariates. The 95% quantiles of predicted distributions
13
14 198 were made using N = 1000 bootstrap simulations (generating new values of both the “spherical”
15
16 199 random effects and the independent errors in each run) of parameter estimates of each model.
17
18
19

20 200 We used R version 3.1.3 (<http://cran.r-project.org/>) for all statistical analyses including, in
21
22 201 particular package *lme4* (Bates *et al.* 2014), to fit GLMMs, and its `bootMer()` function to produce
23
24 202 parametric bootstrap samples..
25
26
27

28 203

31 204 RESULTS

32
33
34 205 Over the two years of the study, we collected bat activity data in N = 1395 transect sections,
35
36 206 during 78 survey visits. Most sites (N = 30) were surveyed on at least two occasions (2-6 surveys
37
38 207 per site), but four sites were only visited once. Within this sample, N = 20 were single-turbine
39
40 208 sites, and N = 14 were multiple-turbine sites (2-4 turbines). In total, we recorded N = 1867 bat
41
42 209 passes, of which 98.4% (N = 1838) were *Pipistrellus* spp. (N = 969 [51.9%] *Pipistrellus*
43
44 210 *pipistrellus*, N = 714 [38.2%] *P. pygmaeus*, and N = 155 [8.3%] *Pipistrellus* sp.). The remaining
45
46 211 1.6% of total calls recorded were a mixture of *Nyctalus noctula*, *Plecotus auritus* Linnaeus and
47
48 212 *Myotis* spp., but because these species combined comprised N = 29 calls, here we focus on
49
50 213 analyses of *Pipistrellus* spp. activity. Overall, *Pipistrellus* bat activity was detected on 466
51
52 214 section surveys.
53
54
55
56
57
58
59
60

1
2
3 215 Parameter estimates for the models for *P. pipistrellus* and *P. pygmaeus* are presented in Table 1.
4
5
6 216 In terms of confounding effects, there were strong non-zero effects of wind speed (negative),
7
8 217 edge density (positive) on the activity of both species, as well as an effect of time to midnight
9
10 218 (linearly negative for *P. pipistrellus* and quadratic for *P. pygmaeus*). In addition, both the
11
12 219 proportion of woodland present and day number affected *P. pygmaeus* activity positively, and *P.*
13
14 220 *pipistrellus* activity was higher in 2014 compared to 2013.

15
16
17
18 221 Accounting for these effects of environmental and habitat variables, we find some evidence for
19
20 222 an effect of turbine proximity. Specifically, this effect is strongest for *P. pipistrellus* (Table 1a),
21
22 223 for which activity in distance band 1 (0-100m, intercept in Table 1a) is lower than the activity in
23
24 224 distance bands 2-5 (100-500m). Based on parameter estimates and their errors, no such effect of
25
26 225 turbine proximity is evident in the model for *P. pygmaeus* activity (Table 1b). Moreover, again
27
28 226 based on parameter estimates alone, there is no evidence for an interaction between distance
29
30 227 band and number of turbines installed.

31
32
33
34
35 228 This interpretation of coefficient estimates above is largely mirrored by assessment of the
36
37 229 predicted levels of activity, although a strong suggestion of an interaction with the number of
38
39 230 turbines installed is evident (Figure 1). Predicted activity is lowest for both species in distance
40
41 231 band 1 (0-100m). In particular, for both species, at multiple SWT sites the predicted mean
42
43 232 activity (measured as the probability of a bat pass per unit area) in the first distance bands (*P.*
44
45 233 *pipistrellus* (Figure 1a): 0.057; *P. pygmaeus* (Figure 1b): 0.109) falls outside the 95% quantiles
46
47 234 of point estimates for all other distance bands (combined 95% quantiles for bands 2-5, *P.*
48
49 235 *pipistrellus*: 0.062-0.248; *P. pygmaeus*: 0.113-0.399), whereas the means of the distance bands 2-
50
51 236 5 all fall within each other's 95% quantiles. There is no such difference for single SWT sites: the
52
53 237 predicted means in all distance bands fall within each other's 95% quantiles, although in the case
54
55
56
57
58
59
60

1
2
3 238 of *P. pipistrellus* activity tends to be lower in the first distance band as well (Figure 1a, single
4
5 239 SWT).

6
7
8 240

9
10
11 241 **DISCUSSION**

12
13
14
15 242 We show that, when accounting for confounding effects of habitat and environmental conditions,
16
17 243 activity of *P. pipistrellus* activity is lower at 0-100m from installed SWTs, compared to 200-
18
19 244 500m. Although not apparent in estimated model parameters, assessment of predictions suggests
20
21 245 that this may also be the case for *P. pygmaeus* activity but this effect was weaker. Similarly, the
22
23 246 distribution of model predictions suggest that the negative effect of SWT presence in the
24
25 247 “closest” distance band is particularly prominent in sites with multiple SWTs installed, with an
26
27 248 apparent absence of any difference between distance bands in the activity of *P. pygmaeus* in
28
29 249 single SWT sites.

30
31
32
33
34 250 On this basis, we conclude that the adverse effect of SWTs on *Pipistrellus* bat activity shown
35
36 251 previously (Minderman *et al.* 2012) may in some cases persist over longer distances in the
37
38 252 landscape. Although here we show that differences in average predicted activity may be
39
40 253 measured when comparing 0-100m to 200-500m distance bands from SWTs, it should be
41
42 254 stressed that this does not show that adverse effects would remain evident at 100m from SWTs.
43
44 255 Indeed, our previous study (Minderman *et al.* 2012) showed a potential drop in average activity
45
46 256 of up to 50% within 25m from operating SWTs. Even in the absence of strong effects beyond
47
48 257 25m, this alone could explain the difference between 0-100m and 200-500m shown in the
49
50 258 current study. Nevertheless, the present findings do suggest that *Pipistrellus* bats may avoid
51
52 259 SWTs on a wider (landscape) scale than previously demonstrated.

53
54
55
56
57
58
59
60

1
2
3 260 Although the mechanism for any avoidance behaviour is unclear, experimental studies in a
4
5
6 261 laboratory setting have shown that the return of echolocation pulses off spinning turbine blades
7
8 262 may be erratic, potentially causing navigational problems for bats (Long, Flint & Lepper 2010).
9
10 263 Alternatively, some bats may avoid foraging in noisy environments (Schaub, Ostwald & Siemers
11
12 264 2008) but whether noise generated by turbines (including SWTs) has this effect has not been
13
14
15 265 tested. In either case, it would be reasonable to assume that the strength of such interference
16
17 266 would dissipate with distance, especially if it involves high-frequency noise (which attenuates
18
19
20 267 quickly over distance), limiting the spatial scale of any adverse effect of the turbine. However,
21
22 268 neither the extent of acoustic noise generated by SWTs nor how this operates over any distance
23
24 269 has been studied, so it is unclear to what extent this could explain the results show here.
25
26
27
28 270 Furthermore, although the model parameter estimates provide no evidence that such avoidance is
29
30 271 stronger in sites with multiple SWTs installed for either species, model predictions of activity
31
32 272 suggest that the decrease in activity in the “near” distance bands tends to be stronger in multiple
33
34 273 SWT sites, particularly for *P. pygmaeus* activity (Figure 1b). Given known local avoidance of
35
36 274 individual turbines (e.g. Minderman et al. 2012), it is possible that clusters of SWTs installed in
37
38 275 relatively close proximity (in our sample between 12 and 90m apart with mean 46.4m and only
39
40 276 one site >62m and <25m respectively) to each other have adverse effects over longer distances.
41
42
43
44 277 For example, while local avoidance of single turbines may cause local changes of commuting
45
46 278 routes, multiple SWTs may cause the loss of such routes altogether (similar to the effect of
47
48 279 lighting on bat activity; Stone *et al.* 2009). However, given the mixed evidence for the interactive
49
50 280 effect between number of turbines and proximity in the current study, this suggestion needs
51
52 281 further study. Moreover, the possibility of cumulative effects (i.e. the combined effect of
53
54 282 multiple wind farm sites in a wider area, “regionally cumulative effects”) is often raised for
55
56
57
58
59
60

1
2
3 283 larger turbines and wind farms (Masden *et al.* 2010; SNH 2012). Here we aimed to answer a
4
5 284 slightly different question, namely whether the installation of a larger number of SWTs installed
6
7
8 285 in a single site increases the effect on bats (i.e. “locally cumulative effects”). The findings
9
10 286 reported here should thus not be used to draw conclusions on whether the effects of multiple
11
12 287 SWT sites developed in a wider area could combine into a “regional cumulative effect”. Finally,
13
14 288 it should be stressed that in accordance with our previous findings (Minderman *et al.* 2012) we
15
16 289 do not find any evidence for the hypothesis that the species studied here are attracted to small
17
18 290 wind turbines, as suggested for other species by Cryan *et al.* (2014). The suggestion that the
19
20 291 interactive effect between SWT proximity and the number of SWTs installed is stronger for *P.*
21
22 292 *pygmaeus* activity is in itself interesting, as it highlights the possibility that different species vary
23
24 293 in their response to SWT presence. Although morphologically similar, the two species
25
26 294 considered here are different in a number of key ecological aspects, including habitat preferences
27
28 295 (Vaughan 1997, Nicholls and Racey 2006, Lintott *et al.* 2015). Although our own findings reflect
29
30 296 this (e.g. strong positive effects of proportion of woodland on *P. pygmaeus*, but not *P.*
31
32 297 *pipistrellus*), whether such ecological differences between these species affect their response to
33
34 298 SWT presence remains to be tested, and unfortunately we did not have sufficient statistical
35
36 299 power to do so here.
37
38
39
40
41
42
43

44 300 There are a number of important caveats with our findings. First of all, these data should not be
45
46 301 generalized to species other than *Pipistrellus* spp. Although *Pipistrellus* bats are by far the
47
48 302 commonest species in both our wider study area as well as in the specific habitats we studied
49
50 303 (Middleton 2006), it is likely that the collection of sufficient data to draw further species-specific
51
52 304 conclusions would require targeted surveys. The need to collect data at a wide range of sites
53
54 305 prevented us from doing this here. However, the effect of larger turbines in wind farms can vary
55
56
57
58
59
60

1
2
3 306 between species (e.g. collision mortality can vary between species depending on environmental
4
5 307 conditions, Baerwald & Barclay 2011) and we therefore consider this to be a research priority for
6
7
8 308 SWTs. Secondly, the limited number of sites with more than two turbines in our sample means
9
10
11 309 that we cannot exclude the possibility that a larger number of turbines would cause stronger
12
13 310 effects; indeed modelling studies in birds (Schaub 2012) suggest that this may well be the case.
14
15 311 However, based on our own experience and anecdotal data collected during the construction of
16
17 312 our SWT owner database (e.g. Park, Turner & Minderman 2013), we feel that our sample of sites
18
19
20 313 fairly represents the usual number of SWTs installed per site; sites with more than two and in
21
22 314 particular more than three SWTs remain relatively rare in the UK. Finally, the combination of
23
24 315 recording bat activity over a relatively large scale (up to 500m from SWTs) along transects, and
25
26
27 316 a distance-based approach to turbine effects, required that we analysed our data in distance
28
29 317 bands. As noted previously, our choice of relatively 'coarse' 100m distance bands may limit our
30
31
32 318 ability to draw conclusions on the exact distance over which avoidance behaviour might operate,
33
34 319 but small scale effects have already been studied (Minderman et al. 2012) and we here chose to
35
36 320 focus on wider-scale effects. In addition, to allow us to generalise our sampling regime across
37
38
39 321 sites with single and multiple SWTs, we chose to use single starting locations for all transects.
40
41 322 As a result, in some cases where multiple SWTs were present, this may have caused some
42
43 323 imprecision in the allocation of individual passes to distance bands (as exact distance to
44
45
46 324 individual turbines would differ by a small amount). However, our choice of transect placement
47
48
49 325 minimised any such possible bias: in multiple SWT sites, transects were oriented in such a way
50
51 326 that they were as close to perpendicular to the orientation of the SWT "line". In cases where this
52
53 327 was impossible, e.g. one of the transects would pass within 50m of another turbine, the transect
54
55 328 in question was excluded from the analyses presented here (i.e. leading to some sites with <4
56
57
58
59
60

1
2
3 329 transects, see Methods). Moreover, accurately allocating bat passes to even more narrow distance
4
5 330 bands is likely to introduce false precision. Thus, collecting data on behavioural responses to
6
7
8 331 turbines on a finer scale would require alternative methods.
9

10
11 332 In conclusion, we suggest that adverse effects of SWTs on *Pipistrellus* bat activity may be
12
13 333 measurable over a greater spatial scale than previously thought (0-100m), although combined
14
15 334 with our previous work it is likely that much of such avoidance behaviour occurs in relatively
16
17
18 335 close proximity to turbines (<25m). It appears that such effects are stronger when multiple SWTs
19
20 336 are installed, particularly for *P. pygmaeus*, but this requires further study. Finally, it should be
21
22
23 337 stressed that the effects reported here are on bat activity only. Although they may reflect changes
24
25 338 in habitat use, and this may in turn lead to population-level effects, direct effects of SWT
26
27
28 339 developments on the latter remain to be studied and the conclusions drawn here cannot be thus
29
30 340 extrapolated.
31

32
33 341 The SWT sector is expanding rapidly worldwide, and as a result planners and decision makers
34
35 342 are increasingly faced with having to advise or decide on planned developments of multiple
36
37
38 343 SWTs in clusters. To date, there were no data to inform such decisions, and the findings
39
40 344 presented here are therefore vital and timely. We reiterate our recommendation that individual
41
42
43 345 SWTs within the range of sizes considered here (6-25m hub height) should be sited away from
44
45 346 potentially valuable bat habitat (at least >25m, Minderman *et al.* 2012), but based on the current
46
47
48 347 findings we further stress that particular care should be taken in landscapes where limited
49
50 348 alternative habitats are available. This would include known roost, commuting and foraging
51
52 349 sites. In addition, on the basis of the potential for wider landscape scale effects on some species,
53
54
55 350 we suggest that the (at least the potential for) presence of particularly vulnerable or locally rare
56
57
58
59
60

1
2
3 351 species is assessed within 100m from proposed sites, particularly when more than one turbine is
4
5 352 planned.

6
7
8
9 353 **ACKNOWLEDGEMENTS**

10
11 354 We thank all site owners for kindly allowing access to their turbines. Thanks to Kathryn
12
13 355 Hamilton for additional data collection, Sofia Motta Pralon for GIS data entry, to Cerian
14
15 356 Tatchley, Nils Bunnefeld, Chris Pendlebury and Claudia Garratt for useful discussions, and to
16
17 357 two three anonymous reviewers for their help in substantially improving this manuscript from
18
19 358 previous versions. This study was funded by a University of Stirling Impact Fellowship to JM
20
21 359 and a grant from the People's Trust for Endangered Species to KJP.
22
23
24
25
26
27

28 360 **References**

- 29
30 361 Agresti, A. (2013) *Categorical data analysis*, 3rd ed. Wiley, Hoboken, NJ, USA.
31
32
33 362 Baerwald, E.F. & Barclay, R.M.R. (2011) Patterns of activity and fatality of migratory bats at a
34
35 363 wind energy facility in Alberta, Canada. *J Wildlife Manage*, **75**, 1103–1114.
36
37
38 364 Barton, K. (2015) *MuMIn: Multi-Model Inference*. R package version 1.13.4. Available at:
39
40 365 <http://cran.r-project.org/web/packages/MuMIn/index.html> [Accessed 28 April 2015].
41
42
43 366 Bates, D., Maechler, M., Bolker, B. & Walker, S. (2014) *lme4: Linear mixed-effects models*
44
45 367 *using Eigen and S4*. R package version 1.1-7. Available at: [http://cran.r-](http://cran.r-project.org/web/packages/lme4/index.html)
46
47 368 [project.org/web/packages/lme4/index.html](http://cran.r-project.org/web/packages/lme4/index.html) [Accessed 28 April 2015].
48
49
50 369 Cade, B.S. (2015) Model averaging and muddled multimodel inferences. *Ecology*, **96**, 2370–
51
52 370 2382.
53
54
55
56
57
58
59
60

- 1
2
3 371 Cryan, P.M., Gorresen, P.M., Hein, C.D., et al (2014) Behavior of bats at wind turbines. *Proc*
4
5 372 *Natl Acad Sci* **111**:15126–15131. doi: 10.1073/pnas.1406672111
6
7
8 373 Gelman, A. & Hill, J. (2007) *Data Analysis Using Regression and Multilevel/Hierarchical*
9 374 *Models*. Cambridge University Press, New York.
10
11
12 375 Gelman, A. & Su, Y.-S. (2014) *arm: Data Analysis Using Regression and*
13 376 *Multilevel/Hierarchical Models*. R package version 1.7-07. Available at : [http://cran.r-](http://cran.r-project.org/web/packages/arm/arm.pdf)
14 377 [project.org/web/packages/arm/arm.pdf](http://cran.r-project.org/web/packages/arm/arm.pdf) [Accessed 28 April 2015].
15
16
17 378 Kuvlesky, W.P., Brennan, L.A., Morrison, M.L., Boydston, K.K., Ballard, B.M. & Bryant, F.C.
18 379 (2007) Wind Energy Development and Wildlife Conservation: Challenges and
19 380 Opportunities. *J. Wildl. Manage.*, **71**, 2487–2498.
20
21
22 381 Lintott, P.R., Bunnefeld, N., Minderman, J., Fuentes-Montemayor, E., Mayhew, R.J., Olley, L.
23 382 & Park, K.J. (2015) Differential Responses to Woodland Character and Landscape Context
24 383 by Cryptic Bats in Urban Environments ed D. Russo. *PLoS One*, **10**, e0126850.
25
26
27 384 Long, C., Flint, J. & Lepper, P. (2010) Wind turbines and bat mortality: Doppler shift profiles
28 385 and ultrasonic bat-like pulse reflection from moving turbine blades. *J. Acoust. Soc. Am.*, **128**,
29 386 2238–2245.
30
31
32 387 Masden, E.A., Fox, A.D., Furness, R.W., Bullman, R. & Haydon, D.T. (2010) Cumulative
33 388 impact assessments and bird/wind farm interactions: Developing a conceptual framework.
34 389 *Env. Impact Assess. Rev.*, **30**, 1–7.
35
36
37 390 Middleton, N. (2006) *An Introduction to the Bats of Scotland*. ECHOES Ecology Ltd., Polmont,
38 391 UK. Available at:
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 392 [http://www.echoesecology.co.uk/documents/AnIntroductionToTheBatsofScotlandEdition1D](http://www.echoesecology.co.uk/documents/AnIntroductionToTheBatsofScotlandEdition1Dec2006_002.pdf)
4
5 393 [ec2006_002.pdf](http://www.echoesecology.co.uk/documents/AnIntroductionToTheBatsofScotlandEdition1Dec2006_002.pdf) [Accessed 28 April 2015].
6
7
8
9 394 Minderman, J., Fuentes-Montemayor, E., Pearce-Higgins, J.W., Pendlebury, C.J. & Park, K.J.
10
11 395 (2014) Estimates and correlates of bird and bat mortality at small wind turbine sites. *Biodiv.*
12
13 396 *Cons.*, **24**, 467–482.
14
15
16 397 Minderman, J., Pendlebury, C.J., Pearce-Higgins, J.W. & Park, K.J. (2012) Experimental
17
18 398 evidence for the effect of small wind turbine proximity and operation on bird and bat
19
20 399 activity. *PLoS ONE*, **7**, e41177.
21
22
23
24 400 Nakagawa, S. & Schielzeth, H. (2013) A general and simple method for obtaining R² from
25
26 401 generalized linear mixed-effects models. *Methods Ecol. Evol.*, **4**, 133–142.
27
28
29
30 402 Nicholls, B. & Racey, P.A. (2006) Habitat selection as a mechanism of resource partitioning in
31
32 403 two cryptic bat species *Pipistrellus pipistrellus* and *Pipistrellus pygmaeus*. *Ecography*, **29**,
33
34 404 697–708.
35
36
37 405 Park, K.J., Turner, A. & Minderman, J. (2013) Integrating applied ecology and planning policy:
38
39 406 The case of micro-turbines and wildlife conservation. *J. Appl. Ecol.*, **50**, 199–204.
40
41
42
43 407 Pearce-Higgins, J.W., Stephen, L., Langston, R.H.W., Bainbridge, I.P. & Bullman, R. (2009)
44
45 408 The distribution of breeding birds around upland wind farms. *J. Appl. Ecol.*, **46**, 1323–1331.
46
47
48
49 409 Rydell, J., Bach, L., Dubourg-Savage, M., Green, M., Rodrigues, L. & Hedenstrom, A. (2010)
50
51 410 Bat mortality at wind turbines in northwestern Europe. *Acta Chiropterologica*, **12**, 261–274.
52
53
54 411 Schaub, M. (2012) Spatial distribution of wind turbines is crucial for the survival of red kite
55
56 412 populations. *Biol. Cons.*, **155**, 111–118.
57
58
59
60

- 1
2
3 413 Schaub, A., Ostwald, J. & Siemers, B. (2008) Foraging bats avoid noise. *J. Exp. Biol.*, **211**,
4
5 414 3174–3180.
6
7
8
9 415 SNH. (2012) *Assessing the Cumulative Impact of Onshore Wind Energy Developments*. Scottish
10
11 416 Natural Heritage, Inverness, UK. Available at: <http://www.snh.gov.uk/docs/A675503.pdf>
12
13 417 [Accessed 28 April 2015].
14
15
16 418 Stone, E.L., Jones, G. & Harris, S. (2009) Street Lighting Disturbs Commuting Bats. *Curr Biol*,
17
18 419 **19**, 1123–1127.
19
20
21
22 420 Swift, S.M. (2009) Activity patterns of Pipistrelle bats (*Pipistrellus pipistrellus*) in north-east
23
24 421 Scotland. *J. Zool.*, **190**, 285–295.
25
26
27 422 Tuomainen, U. & Candolin, U. (2011) Behavioural responses to human-induced environmental
28
29 423 change. *Biol. Rev.*, **86**, 640–657.
30
31
32
33 424 UK Meteorological Office (2006) *MIDAS Land Surface Stations Data (1853-Current)*. NCAS
34
35 425 British Atmospheric Data Centre, Didcot, UK. Available at:
36
37 426 <http://catalogue.ceda.ac.uk/uuid/220a65615218d5c9cc9e4785a3234bd0> [Accessed 28 April
38
39 427 2015].
40
41
42
43 428 Vaughan, N., Jones, G. & Harris, S. (1997) Habitat use by bats (Chiroptera) assessed by means
44
45 429 of a broad-band acoustic method. *J Appl Ecol*, **34**, 716–730.
46
47
48 430 Walsh, A., Harris, S. (1996) Foraging habitat preferences of vespertilionid bats in Britain. *J Appl*
49
50 431 *Ecol* **33**:508–518.
51
52
53
54 432 Whittingham, M.J., Stephens, P.A., Bradbury, R.B. & Freckleton, R.P. (2006) Why do we still
55
56 433 use stepwise modelling in ecology and behaviour? *J Anim Ecol*, **75**, 1182–1189.
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

434 Zwart, M.C., McKenzie, A.J., Minderman, J., Whittingham, M.J. (2015) Conflicts between birds
435 and on-shore wind farms. In: *Problematic wildlife: a cross-disciplinary approach*. Springer
436 Science+Business Media, New York, NY, pp 489–504. Angelici, F.M. (Ed.)


Review Copy

437 **Table 1.** Parameter estimates (*b*) and their standard errors (S.E.) from Generalized Linear Mixed
 438 Effects Models (GLMMs) for the activity (probability of a pass ha⁻¹) of (a) *P. Pipistrellus* and
 439 (b) *P. pygmaeus*. All models included site and transect within site as a nested random effect
 440 structure (variance estimates at the bottom of the table), had binomial errors and were fitted with
 441 a complimentary log-log link. All coefficients are parameter estimates based on model fits using
 442 standardised (centered and scaled to 1 SD) inputs. Instances where the point estimate ± 2 *S.E.
 443 does not overlap zero are highlighted in bold, indicating evidence that the parameter is different
 444 from zero.

<i>Fixed effects</i>	<i>(a) P. pipistrellus</i>		<i>(b) P. pygmaeus</i>	
	<i>b</i>	S.E.	<i>b</i>	S.E.
(Intercept)	-2.203	0.197	-1.864	0.198
Distance band 2 (100-200m)	0.521	0.208	0.174	0.203
Distance band 3 (200-300m)	0.730	0.207	0.253	0.205
Distance band 4 (300-400m)	0.596	0.213	0.232	0.210
Distance band 5 (400-500m)	0.489	0.223	0.306	0.211
No. turbines (multiple)	-0.080	0.367	-0.263	0.363
Min 24h. Temperature	-0.041	0.150	-0.099	0.162
Day number	0.124	0.203	0.739	0.195
Year (2014)	0.826	0.252	-0.602	0.301
Time to midnight	-0.536	0.156	0.294	0.175
Time to midnight ²	-0.472	0.287	-1.208	0.332
Wind speed (m s ⁻¹)	-0.549	0.172	-0.759	0.191
Edge density (m m ⁻²)	0.670	0.132	0.620	0.144
Proportion woodland	-0.067	0.141	0.493	0.125
D. band 2 * no. turbines	0.260	0.428	0.693	0.412
D. band 3 * no. turbines	0.433	0.422	0.421	0.405
D. band 4 * no. turbines	-0.379	0.424	0.461	0.411
D. band 5 * no. turbines	-0.018	0.443	0.781	0.412
<i>Random effects</i> σ^2				
Transect / Site	0.374		0.579	
Site	0.377		0.447	
<i>Model fit</i>				
Marginal R^2	0.304		0.255	
Conditional R^2	0.451		0.507	

445 **FIGURE CAPTIONS**

1
2
3 446 **Figure 1.** Observed (small grey points, average per site visit) and predicted (dark grey and black
4
5 447 points, and lines) of **(a)** *P. pipistrellus*, and **(b)** *P. pygmaeus* activity (probability of a pass ha⁻¹)
6
7
8 448 in five distance bands from single- and multiple SWT installations. Predictions are based on
9
10 449 model fits using standardized (scaled and centered) bat activity from Generalized Linear Mixed
11
12 450 Effects Models (GLMMs, Table 1), in each of five distance bands around small wind turbines.
13
14
15 451 Dark grey circles and their corresponding lines are predicted median levels for single-turbine
16
17 452 sites, and black circles and their corresponding lines are median levels for multiple-turbine sites.
18
19
20 453 Small grey dots are the observed mean probabilities of obtaining a bat pass across all
21
22 454 observations in each distance class, for each site. The error bars (lines) are the 95% quantiles of
23
24
25 455 predictions calculated from N = 1000 bootstrap samples of estimated parameter distributions.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



456

457 **Figure 1.**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

458

459

Review Copy

1
2
3 **1 Landscape-scale effects of single- and multiple small wind turbines on bat activity**
4

5
6 *2 Jeroen Minderman, Mairi H. Gillis, Helen F. Daly & Kirsty J. Park*
7
8

9
10 3

11
12 **4 Supplementary material**
13
14

15
16 5
17

18
19 **6 Preliminary habitat data analysis and additional tables**
20

21 7 A total of 10 habitat variables were quantified in each transect section in all sites (Table A2),
22
23 8 using 1:1250 UK Ordnance Survey MasterMap Topography digital map (OS) data and QGIS v.
24
25 9 1.8. These variables were chosen on the basis of their known potential effect on bat activity. The
26
27 10 five proportion land cover variables (proportion buildings, road or tracks, rough grassland,
28
29 11 roadside, and woodland) were calculated from the OS polygon area data and were calculated as
30
31 12 the sum of each of these five land cover types in each transect section, divided by the size of the
32
33 13 transect section. These were defined as the total area size of all polygons listed as (1) "Buildings"
34
35 14 in the *Theme* field; (2) "Road Tracks and Paths" in the *Theme* field; (3) "Rough grassland"
36
37 15 (including all lower classifications) in the *descrTerm* field; (4) "Roadside" in the *descrGroup*
38
39 16 field and (5) "Coniferous or Nonconiferous trees" in the *descrGroup* field. The four distance
40
41 17 variables (distance to buildings, linear features, trees and water) were calculated by overlaying
42
43 18 the OS data with a 1x1m 'raster' of grid cells, calculating the distance (m) between each raster
44
45 19 cell and the nearest raster cell containing each of the four land cover types (respectively,
46
47 20 buildings as defined above, any linear feature as contained in the OS "line" data, any woodland
48
49 21 as defined above and any water polygon), and averaging these distances for all cells within the
50
51 22 transect sections.
52
53
54
55
56
57
58
59
60

1
2
3 23 Finally, edge density (m m^{-2}) was expressed as the total length of all line features contained in
4
5 24 the OS Linear data per transect section, divided by the transect section area. Line features
6
7
8 25 include e.g. building-, woodland and water edges, hedgerows, roads and tracks, roadsides, and
9
10 26 field boundaries.

11
12
13 27 To identify a subset of habitat variables that were most descriptive of bat activity (see main text),
14
15
16 28 we ran an initial model selection procedure on a Generalized Linear Mixed Effects (GLMM)
17
18 29 model for the probability of a bat pass per hectare as a function of all 10 habitat variables only.
19
20 30 In addition to these habitat variables this model included transect within site as a random effect,
21
22
23 31 a binomial error distribution with a log-log link function and transect section size (ha) as an
24
25 32 offset. To avoid problems with collinearity, we limited the full model set to include only one of
26
27
28 33 the five 'proportion' habitat variables (because these represent proportion cover, lower cover of
29
30 34 one automatically implies greater cover of another, leading to high collinearity). We further
31
32 35 excluded all models including (1) **both** proportion of woodland and distance to trees, (2) **both**
33
34 36 proportion buildings and distance to buildings, (3) **both** edge density and distance to linear
35
36 37 features **or** proportion of roads or tracks. Again, these latter exclusions were to avoid high
37
38 38 collinearity. For example, a greater proportion of woodland in a given transect section naturally
39
40 39 implies shorter distances to trees; edge density includes linear features such roads and tracks; etc.
41
42 40 Thus, these variables effectively measured very similar things but in slightly different ways. All
43
44 41 model fitting procedures and analyses followed the methodology described in the main text.
45
46
47
48

49
50 42 The resulting full model set comprised of a total of 112 models. The candidate set ($\Delta\text{AICc}<4$)
51
52 43 contained 4 models. The 'top' model retained three of the ten habitat variables: distance to water,
53
54 44 edge density and the proportion of woodland (Table A3). Distance to water was only included in
55
56 45 two out of the four models in the top set, however. By contrast, edge density and the proportion
57
58
59
60

1
2
3 46 of woodland were retained in all four models in the candidate set. Thus, we chose to use edge
4
5 47 density and the proportion of woodland as the key two habitat variables explaining bat activity.
6
7
8 48 As discussed, we chose to limit the number of explanatory variables for the models for bat
9
10 49 activity in this way to avoid problems with overparameterisation. However, we did test whether
11
12 50 the inclusion of distance to water (because of the known importance of riparian habitats to *P.*
13
14 51 *pygmaeus* in particular) made any difference to the models presented in Table 1 in the main text
15
16 52 or whether it would change their interpretation. The estimated effect of distance to water was not
17
18 53 significantly different from zero in either the model for *P. pipistrellus* (-0.259 +/- 0.151) or the
19
20 54 model for *P. pygmaeus* (-0.043 +/- 0.181, although convergence of the estimates of this model
21
22 55 was poor, presumably due to overparameterisation), neither was the interpretation or significance
23
24 56 of any of the other parameters changed by its inclusions.
25
26
27
28
29

30 57 Similarly, we tested whether the inclusion of turbine hub height as an extra predictor, as well as
31
32 58 its interaction with distance from turbines, affected our results. As per the addition of distance to
33
34 59 water, this did lead to estimation problems for some of the models under consideration. Bearing
35
36 60 this in mind, none of the interaction terms between hub height and distance from turbine were
37
38 61 different from zero for either the *P. pipistrellus* or *P. pygmaeus* models, and when this
39
40 62 interaction term was further dropped from both models, the effect of hub height was not different
41
42 63 from zero in either model (*P. pipistrellus*: -0.119 +/- 0.231; *P. pygmaeus*: -0.124 +/- 0.268). The
43
44 64 remaining parameter estimates and their interpretation also remain qualitatively unchanged.
45
46
47
48
49

50 65

51 66

52 67

53

54

55

56

57

58

59

60

68 **Table A1.** Descriptive statistics of predictor variables in the Generalized Linear Mixed Model
 69 (GLMM) for *Pipistrellus* sp. bat activity (Table 1), on their original (unstandardised) scale.

	Mean	Median	Std. dev	Min.	Max.
Minimum 24h temperature (°C)	11.76	12.00	2.75	5.00	17.60
Julian day number	196.59	195.00	28.85	147.00	246.00
Time to midnight (min)	-26.07	-41.00	82.54	-202.30	197.77
Time to midnight ²	7487.47	4669.44	8375.13	0.00	40925.29
Wind speed (m s ⁻¹)	1.07	0.40	1.46	0.00	9.80
Edge density (m m ⁻²)	0.04	0.03	0.02	0.00	0.13
Proportion woodland	0.05	0.00	0.13	0.00	0.92

70

71 **Table A2.** Descriptive statistics of all ten original habitat variables as predictors for bat activity on their original (unstandardised)
 72 scale, split by single- and multiple SWT sites.

73


	Single turbines					Multiple turbines (2-4)				
	Mean	Median	SD	Min.	Max.	Mean	Median	SD	Min.	Max.
Distance to buildings (m)	174.48	152.02	114.83	13.65	596.93	222.76	209.89	135.88	9.2	623.77
Distance to linear features (m)	24.67	17.92	21.86	5.9	132.37	21.09	18.5	16.5	5.37	190.97
Distance to trees (m)	144.88	125.65	110.28	0.19	519.74	211.45	196.33	150.86	1.89	646.47
Distance to water (m)	218.3	174	166.43	13.2	689.27	186.47	153.42	142.33	17.88	652.74
Edge density (m m ⁻²)	0.04	0.03	0.03	0.00	0.13	0.04	0.04	0.02	0.00	0.13
Proportion of buildings	0.01	0.00	0.03	0.00	0.20	0.01	0.00	0.04	0.00	0.29
Proportion of roads or tracks	0.02	0.01	0.03	0.00	0.26	0.02	0.01	0.03	0.00	0.15
Proportion of rough grassland	0.07	0.00	0.18	0.00	1.00	0.02	0.00	0.07	0.00	0.49
Proportion of roadsides	0.02	0.00	0.04	0.00	0.26	0.02	0.00	0.03	0.00	0.14
Proportion of woodland	0.07	0.00	0.15	0.00	0.92	0.04	0.00	0.12	0.00	0.77

74 **Table A3.** Candidate set of Generalized Linear Mixed Models for bat activity (probability of bat
 75 activity $h_{a^{-1}}$, binomial errors, complementary log-log link) within 4 AICc points of the “top”
 76 model, as a function of ten candidate habitat predictors only. Coefficients are parameter
 77 estimates based on models fit using standardized (centered and scaled to 1 SD) inputs. K is the
 78 number of parameters in the models. All models included site and transect within site as a nested
 79 random effect structure.

Model	Intercept	Distance to buildings	Distance to water	Edge density	Proportion woodland	K	Log-likelihood	AICc	Δ AICc	Model weight
1	-1.146		-0.305	0.523	0.41	6	-824.8	1661.7	0.000	0.541
2	-1.146	0.028	-0.301	0.532	0.408	7	-824.8	1663.7	1.962	0.203
3	-1.139			0.549	0.445	5	-826.9	1663.9	2.184	0.181
4	-1.138	0.059		0.568	0.438	6	-826.8	1665.7	3.957	0.075

80

81



82

83 **Figure A1.** Conceptual example of a bat transect layout. The turbine is at the intersection of the
84 transects. The dashed lines show the transects walked, whereas the solid line represents the area
85 for which The habitat surrounding the turbines varied significantly between sites, and this was
86 reflected in transect placement, so that (1) the combination of all transects within each site
87 covered all major habitats present, and (2) overall distance separating each transect was
88 maximized.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy