SUPPLEMENTARY INFORMATION

Benzylic functionalisation of phenyl all *cis*-2, 3, 5, 6 –tetrafluoro - cyclohexane provides access to new organo-fluorine building blocks

Tetiana Bykova, Nawaf Al-Maharik, Michael Bühl, Alexandra M. Z. Slawin,
Tomas Lebl and David O' Hagan*

EaStCHEM School of Chemistry, University of St Andrews, North Haugh, St Andrews,
Fife, KY16 9ST, UK

E.mail; do1@st-andrews.ac.uk

Table of Contents

1. GENERAL INFORMATION.....	2
2. PROCEDURE AND ANALYTICAL DATA.....	3
3. NMR SPECTRA (¹H, ¹³C, ¹⁹F NMR).....	10
4. ¹⁹F NMR VT STUDIES: CONVERSION OF COMPOUND 4 TO 6.	26
5. 2D ¹H, ¹H-DQF-COSY (A), 2D ¹H, ¹⁹F HMBC (B) AND 1D ¹⁹F WITH ¹H DECOUPLING (C) AND 1D SELECTIVE ¹H, ¹⁹F-HOESY (D, E, F) NMR SPECTRA FOR AMINO ALCOHOL 9.....	28
6. (A) ¹H, ¹⁹F NMR HMBC; (B) ¹⁹F OBSERVE WITH ¹H DECOUPLING; (C) 1D SELECTIVE ¹H, ¹⁹F HOESY (C, D, E, F) FOR 4-BROMOBENZYL-1,2,3,5,6- PENTAFLUOROCYCLOHEXANE-1-CARBOXAMIDE (22).....	30
7. (A) ¹H, ¹⁹F NMR HMBC; (B) ¹⁹F OBSERVE WITH ¹H DECOUPLING; (C) 1D SELECTIVE ¹H, ¹⁹F HOESY (C, D, E, F) FOR 1,2,3,5,6-PENTAFLUORO-4-NITROBENZYL CYCLOHEXANE-1-CARBOXAMIDE (24)	32
8. (A) ¹H, ¹⁹F NMR HMBC; (B) ¹⁹F OBSERVE WITH ¹H DECOUPLING; (C) 1D SELECTIVE ¹H, ¹⁹F HOESY (C, D, E, F) FOR 1,4-PHENYLENEBIS(METHYLENE)BIS- 1,2,3,5,6-PENTAFLUOROCYCLOHEXANE-1-CARBOXAMIDE (25)	33
9. DENSITY FUNCTIONAL THEORY COMPUTATIONS	34

1. General information

^1H , ^{13}C and ^{19}F NMR spectra were recorded on Bruker Avance or Avance III spectrometers at 500 MHz (^1H NMR), 125 MHz (^{13}C NMR) and 470 MHz (^{19}F NMR). Chemical shifts data were reported as δ in units of parts per million relative to residual solvent and were referenced to the internal solvent, where appropriate. Coupling constants were reported in Hertz (Hz). Analytical thin layer chromatography (TLC) was carried out on Merck silica gel 60 F254 glass-supported plates. Visualisation was by absorption of UV light (λ max 253 or 365 nm), or by thermal development after dipping in an aqueous solution of potassium permanganate, potassium carbonate and sodium hydroxide. Flash column chromatography was carried out on Merck Geduran silica gel 40-63 micron, eluting with solvents as supplied, under a positive pressure of compressed air. Melting points were recorded on an Electrothermal 9100 melting point apparatus and are uncorrected. Electrospray mass spectra were recorded at the University of St Andrews Mass Spectrometry facility on either a Micromass LC TOF spectrometer or a ThermoFisher Orbitrap Excalibur spectrometer from solutions of the analyte in methanol. Additional spectra were obtained at the EPSRC National Mass Spectrometry Service Centre at Swansea using a ThermoFisher Orbitrap LQT XL spectrometer fitted with an ASAP solids probe operating in atmospheric pressure chemical ionisation (APCI) mode.

2. Procedure and analytical data

1-Bromo-2,3,5,6-tetrafluorocyclohexyl)benzene (4):

N-Bromosuccinimide (2.05 g, 113.79 mmol) was added to a solution of **2** (2.20 g, 94.82 mmol) in CCl_4 (20 ml) and the mixture was heated under reflux for 18 h. After cooling to r.t, the solvent was removed under reduced pressure, dissolved in DCM and washed with H_2O and the aqueous layer was then extracted with DCM (200 mL x 3). The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated to give orange solid, which was subjected to column chromatography using silica gel (petrol ether/DCM 7:3), furnishing compound **4** (2.62 g, 84.39 mmol, 89%): M.p.= 178 °C; ^1H NMR (400 MHz, CDCl_3) δ 7.69 (2H, d, J = 7.2, Ar), 7.54-7.36 (3H, m, Ar), 5.70 (2H, dd, J = 48.4, 5.5, H-2, 6), 5.60-5.36 (2H, m, H-3, 5), 2.80-2.62 (1H, m, H-4a), 2.60-2.46 (1H, m, H-4b); ^{13}C NMR (101 MHz, CDCl_3) δ 136.6 (Ar), 129.6 (Ar), 129.2 (Ar), 126.4 (Ar), 88.0 (dd, J = 199.0, 18.9, C-3, 5), 85.4 (dm, J = 201.5, C-2, 6), 56.8 (C-1), 27.1 (tt, J = 22.3, 3.3, C-4); ^{19}F NMR (376 MHz, CDCl_3) δ -189.3 (AA'XX', F-2, 6), -194.68 (AA'XX', F-3, 5); FTMS (ESI $^-$) m/z calcd for ([M]) 311.11332; found 311.1689; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 2922.16 (C-H), 1392.61 (C-F), 1101.35 (C-F)

2,3,5,6-tetrafluoro-1-phenylcyclohexan-1-ol (6):

Sodium azide (100 mg, 1.54 mmol) was added to a solution of benzyl bromide **4** (300 mg, 0.97 mmol) in DMF (10 mL) and the reaction was stirred for 18 h at 90 °C until TLC analyses revealed the consumption of the starting material. The mixture was poured into water (50 mL), extracted with diethyl ether (2 x 100 mL), washed with brine (3 x 150 mL), dried over Na_2SO_4 , filtered, and the filtrate was concentrated under reduced pressure. The product was purified over silica gel (petrol ether/ethyl acetate 7:3), furnishing compound **6** (2.62 g, 84.39 mmol, 82%): M.p.= 125 °C; ^1H NMR (400 MHz, CDCl_3) δ 7.80-7.69 (2H, m, Ar), 7.59-7.38 (3H, m, Ar), 5.29-4.94 (4H, m, H-2, 3, 5, 6), 2.78-2.62 (1H, m, H-4a), 7.80-7.69 (1H, m, H-4b), 2.07 (1H, s, OH); ^{13}C NMR (126 MHz, CDCl_3) δ 139.5 (Ar), 129.6 (Ar), 129.3 (Ar), 126.1 (Ar), 89.6 (dd, J = 196.8, 15.2, C-2, 6), 86.1 (dm, J = 191.1, C-3, 5), 73.1 (C-1), 26.7 (t, J = 21.9, C-4); ^{19}F NMR (376 MHz, CDCl_3) δ -198.8 (AA'XX', F-2, 6), -207.7 (AA'XX', F-3, 5); FTMS (ESI $^+$) m/z calcd for ([M]-H $^+$) 247.0746; found 247.0751; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3578.01 (O-H), 2925.1 (C-H), 2853.73 (C-H), 1390.7 (-C-H), 1160.2 (C-F), 1013.61 (C-F).

2,3,5,6-Tetrafluoro-1-phenylcyclohexyl)acetamide (7):

Concentrated H_2SO_4 (1 mL) was added to a solution of benzyl bromide **4** (400 mg, 1.29 mmol) in MeCN (200 ml). The reaction mixture was stirred for 10-40 h at 90 °C until TLC analyses indicated the consumption of the starting material. After cooling to r.t., the mixture was poured into water (200 mL), extracted with EtOAc (3 x 200 mL), dried over Na_2SO_4 and filtered. The filtrate was concentrated under reduced pressure and purified over silica gel (petrol ether/ EtOAc 6:4) furnishing compound **7** (254 mg, 0.88 mmol, 68%): M.p.= 282 °C; ^1H NMR (500 MHz, $(\text{CD}_3)_2\text{CO}$) δ 7.73 (2H, dt, J = 7.7, 1.1, Ar), 7.41 (2H, td, J = 8.0, 7.3, 1.8, Ar), 7.34 (1H, tt, J = 7.4, 2.0, Ar), 6.07-5.90 (2H, m, H-2, 6), 5.43-5.20 (2H, m, H-3, 5), 2.57-2.47 (2 H, m, H-4), 1.82 (3H, s, CH_3); ^{13}C

NMR (126 MHz, CD_3CO): δ 169.5 (C=O), 139.3 (Ar), 128.2 (Ar), 127.6 (Ar), 126.9 (Ar), 88.2 (dm, J = 205.7 Hz, C-2, 6), 86.1 (dm, J = 180.5 Hz, C-3, 5), 60.7 (C-1), 26.8 (t, J = 22.3, C-4), 22.9 (CH_3); ^{19}F NMR (376 MHz, CD_3CO): δ -197.0 (AA'), -208.7 (BB'); IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3257 (N-H), 3217 (N-H), 3057 (C-H), 3001 (C-H), 1683 (C=O), 1559 (C-C); FTMS (ESI^-) m/z calcd for $([\text{M}]-\text{H})$ 288.1011; found 288.1018.

1-Amino-3,5,6-trifluoro-2-phenylcyclohexan-2-ol (9) and 2,3,5,6-tetrafluoro-1-phenylcyclohexan-1-amine (8):

Acetamide **7** (50 mg, 0.17 mmol) was heated in aqueous HCl (10 mL of 6 M solution) at 120 °C for 18 h. The reaction was cooled to r.t and the solution concentrated under reduced pressure. The mixture was purified over silica gel (petrol ether/ethyl acetate 3:7), furnishing products **9** and **8**. The residue containing products **9** and **8** was dissolved in MeCN: water (40:60) and purified by semi-preparative HPLC on a Shimadzu system (Phenomenex Luna C18 (250 × 10.00 mm, 5 μ); mobile Phase: A (H_2O + 5% MeCN), B (MeCN); isocratic conditions: 50% B, flow rate: 10 mL. min $^{-1}$; detection 215 nm. Fractions containing products **7** (tR = 21.5 min) and **8** (tR = 23.5 min) were concentrated under reduced pressure to eliminate MeCN and freeze dried from frozen solutions in water, to offer **9** (6 mg, 0.025 mmol, 15 %) as white crystalline solid: M.p.= 170 °C; ^1H NMR (300 MHz, MeOD) δ 7.60 (2H, d, J = 7.9, Ar), 7.37 (2H, dd, J = 8.5, 6.7, Ar), 7.33-7.21 (1H, m, Ar), 5.39-5.19 (1H, m, H-5), 5.03-4.94 (1H, m, H-3), 4.62 (1H, dd, J = 49.9, 7.3, H-6), 4.40 (2 H, dd J = 14.5, 8.4, 2.2, H-2), 2.61-2.41 (1H, m, H-4a), 2.33-2.04 (1H, m, H-4b); ^{13}C NMR (126 MHz, MeOD) δ 143.4 (Ar), 127.8 (Ar), 126.7 (Ar), 126.4 (Ar), 93.7 (dd, J = 182.2, 15.8, C-6), 90.7 (dd, J = 172.7, 16.7, H-3), 87.8-85.2 (m, H-5), 71.9 (d, J = 18.7, H-2), 60.5 (C-1), 30.5 (td, J = 21.2, 4.4, C-4); ^{19}F NMR (282 MHz, MeOD) δ -191.33 (d, J = 5.3, F-3), -198.28 (dd, J = 15.1, 5.2, F-5), -208.17 (d, J = 15.1, F-6); FTMS (ESI^+) m/z calcd for $([\text{M}]+\text{H})$ 246.1105; found 246.1103; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3451.1 (N-H), 3246.20 (O-H), 2866.84 (C-H), 1539.2 (C=C), 1072 (C-N); followed by (**8**) (15 mg, 0.06 mmol, 35%) as white crystalline solid: M.p.= 159-160 °C; ^1H NMR (300 MHz, MeOD) δ 7.65 (2H, d, J = 7.3, Ar), 7.40 (2H, dd, J = 8.4, 6.9, Ar), 7.34-7.26 (1H, tt, J = 7.3, 1.3, Ar), 5.49-5.03 (4H, m, H-2, 3, 5, 6), 2.67-2.31 (2H, m, H-4); ^{13}C NMR (126 MHz, MeOD) δ 143.5 (Ar), 128.1 (Ar), 127.0 (Ar), 125.7 (Ar), 90.0 (dd, J = 194.5, 16.8, C-2, 6), 86.5 (dm, J = 188.6, C-3, 5), 58.5 (C-1), 27.1 (t, J = 23.4, C-4); ^{19}F NMR (282 MHz, MeOD) -198.25 (AA'), -206.95 (BB'); FTMS (ESI^+) m/z calcd for $([\text{M}]+\text{Na})$ 270.0881; found 270.0875; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3580 (N-H), 2916 (C-H), 2849 (C-H), 1653 (N-H), 1111 (C-F), 1028 (C-F).

1-Acetamido-2,3,5,6-tetrafluorocyclohexane-1-carboxylic acid (10):

RuCl₃ (5 mg, 0.026 mmol, 0.05 eq.) was added to a solution of acetamide **7** (150 mg, 0.52 mmol) and periodic acid H₅IO₆ (1 g, 4.4 mmol) in a AcCN/CCl₄/H₂O (14 mL, 2:2:3) solvent system, followed by reflux for 18 h. After cooling, the solvent was concentrated and the product redissolved in EtOAc and washed with H₂O. The aqueous layer was then extracted with EtOAc (200 mL x 2). The organic extracts were dried over NaSO₄, filtered and concentrated under reduced pressure. The mixture was dissolved in a minimum amount of H₂O/MeCN mixture and was loaded onto a reverse phase cartridge (1000 mg Extract Clean C18-HC, preconditioned with H₂O), and washed with H₂O: MeCN (50:50) (2 x 10 mL) resulting in compound **10** (80 mg, 0.31 mmol, 61%): M.p.= 209-210 °C; ¹H NMR (500 MHz, (CD₃)₂CO): δ 4.21 (2H, d, *J* = 41.2, H-2, 6), 3.83- 3.52 (2H, m, H-3, 5), 1.23-1.10 (2H, m, H-4), 0.70 (3H, s, (CH₃)); ¹³C NMR (126 MHz, (CD₃)₂CO): δ 170.9 (C=O), 167.0 (C-1), 85.4 (d, *J* = 146.8, C-2, 6), 83.8 (d, *J* = 131.6, C-3, 5), 25.2 (t, *J* = 22.4, C-4), 20.0 (CH₃); ¹⁹F NMR (376 MHz, (CD₃)₂CO): δ -199.2 (AA'), -211.7 (BB'); IR: ν_{max}/cm⁻¹ 3273 (N-H), 2987 (C-H), 1662 (C=O), 1374 (C-H); FTMS (ESI⁻) *m/z* calcd for ([M]-H) 256.0596; found 256.0601.

1-Amino-2,3,5,6-tetrafluorocyclohexane-1-carboxylic acid (11);

Carboxylic acid **10** (70 mg, 0.27 mmol) was heated in aqueous HCl (10 mL of 6 M solution) at 120 °C for 18 h. The reaction was cooled to r.t and the solution was concentrated under reduced pressure. The mixture was dissolved in minimum amount of H₂O and was then loaded onto a reverse phase cartridge (1000 mg Extract Clean C18-HC, preconditioned with H₂O), washed with H₂O (10 mL), and the product eluted with H₂O: MeCN (70:30) (3 x 10 mL) resulting in amino acid **11** (32 mg, 0.15 mmol, 55%): M.p.= 198-199 °C; ¹H NMR (500 MHz, MeOD): δ 5-20-4.95 (4H, m, H-2, 3, 5, 6), 2.57-2.45 (1H, m, H-4a), 2.41- 2.22 (1H, m, H-4b); ¹³C NMR (126 MHz, MeOD): δ 171.7 (COOH), 88.3 (dd, *J* = 191.8, 17.5), 85.4 (dt, *J* = 182.0, 15.1), 28.8 (C-1), 27.4 (t, *J* = 22.1); ¹⁹F NMR (282 MHz, MeOD): δ -200.3 (F-2, 6), -209.8 (F-3, 5); IR: ν_{max}/cm⁻¹ 3133 (N-H), 3034 (N-H), 1727 (C=O), 1324 (C-O), 1069 (C-N); FTMS (ESI⁻) *m/z* calcd for ([M]-H) 214.0491; found 214.0493.

1,2,3,5,6-Pentafluorocyclohexyl)benzene (13)

A solution of bromide **4** (800 mg, 25.81 mmol) and Ag(II)F (650 mg, 51.61 mmol) in Et₂O was refluxed for 18 h under an argon atmosphere and in light free conditions. The reaction was then cooled to r.t, filtered through a short pad of celite and was washed with Et₂O. The solvent was concentrated under reduced pressure. Purification over silica gel (petrol ether/Et₂O 6:4) gave benzyl fluoride **13** (529 mg, 21.16 mmol, 82%): M.p.= 121 °C; ¹H NMR (400 MHz, CDCl₃): δ 7.71-7.67 (2H, m, Ar), 7.61-7.39 (3H, m, Ar), 5.30-4.79 (4H, m, H-2, 3, 5, 6), 2.89-2.66 (1H, m, H-4a), 2.65-2.41 (1H, m, H-4b); ¹³C NMR (126 MHz, CDCl₃): δ 135.3 (*J* = 18.3, Ar), 129.8 (Ar), 128.7 (Ar), 125.7 (dt, *J* = 8.9, 4.1), 91.2 (d, *J* = 187.8 Hz, C-1), 87.7 (dm, *J* = 193.9, C-3, 5), 85.3 (dm, *J* = 185.9, C-2, -6), 26.3 (t, *J* = 22.0, C-4); ¹⁹F NMR (376 MHz, CDCl₃): δ -166.6 (t, *J* = 13.6 Hz), -199.7 (dt, *J* = 8.2, 4.3), -213.5 (dd, *J* = 13.0, 7.9); IR: ν_{max}/cm⁻¹ 2922.16 (C-H), 1452.4 (C=C), 1390.68 (C-F), 1336.67 (C-F), 1168.86 (C-F); FTMS (ESI⁻) *m/z* calcd for ([M]-H) 249.0702; found 248.9607.

1,2,3,5,6-Pentafluorocyclohexane-1-carboxylic acid (14):

RuCl₃ (6 mg, 0.0295 mmol, 0.05 eq) was added to a solution of pentafluorocyclohexane **13** (150 mg, 0.59 mmol) and periodic acid H₅IO₆ (1.08 g, 4.72 mmol) in AcCN/CCl₄/H₂O (14 mL, 2:2:3). The reaction was heated under reflux for 18 h and was cooled. The solvent mixture was removed under reduced pressure and the product was re-dissolved in EtOAc and washed with H₂O. The aqueous layer was then extracted with EtOAc (200 mL x 2). The organic fraction was dried over NaSO₄, filtered and concentrated under reduced pressure. The product was dissolved in a minimum volume of H₂O/MeCN and then was loaded onto a reverse phase cartridge (1000 mg Extract Clean C18-HC, preconditioned with H₂O), washed with H₂O (10 mL), and the product eluted with H₂O: MeCN (50:50) (2 x 10 mL) resulting in compound **14** (108 mg, 0.44 mmol, 74%): M.p.= 125 °C; ¹H NMR (500 MHz, MeOD): δ 5.29 (2H, dm, *J* = 44.0, H-2, 6), 5.03-4.95 (2H, m, H-3, 5), 2.54- 2.39 (2H, m, H-4^{a,b}); ¹³C NMR (126 MHz, MeOD): δ 165.0 (s, COOH), 124.7 (C-1), 85.0 (dm, *J* = 191.8, C-2, 3, 5, 6), 26.0 (t, *J* = 22.1, C-4); ¹⁹F NMR (376 MHz, MeOD): δ -174.4 (s(br), F-1), -200.8 (dt, *J* = 7.6, 3.9, F-2, 6), -215.8-216.7 (m, F-3, 5); IR: ν_{max}/cm⁻¹ 2935.66 (O-H), 1730 (C=O), 1392.61 (C-F), 1296.16 (C-F); FTMS (ESI⁻) *m/z* calcd for ([M]-H⁺) 217.0288; found 217.0288.

1,2,3,5,6-Pentafluorocyclohexyl)methanol (15):

BH₃.THF (1M in THF, 1 mL, 1 mmol) was added dropwise to a solution of pentafluorocyclohexane-1-carboxylic acid (**14**) (60 mg, 0.28 mmol) in THF (3 mL) at 0 °C. The reaction was stirred at 50 °C for 18 h, after which it was quenched with sat. aq. NaHCO₃ and extracted with EtOAc (2 x 30 mL). The combined organics were washed with brine (2 x 10 mL), dried (Na₂SO₄), and concentrated. The resulting material was purified over silica gel (EtOAc/petrol ether 1:1) to provide the product **15** as white crystalline solid, which was found to sublime under reduced pressure (31 mg, 0.15 mmol, 55%): M.p.= 80 °C; ¹H NMR (400 MHz, CDCl₃): δ 4.99 (2H, d, *J* = 46.7, H-2, 6), 4.89- 4.67 (2H, m, H-3, 5), 4.16 (2H, ddt, *J* = 26.3, 6.4, 3.1, CH₂-OH), 2.74-2.57 (1H, m, H-4a), 2.52-2.41 (1H, m, H-4b); ¹³C NMR (126 MHz, CDCl₃): δ 110.0 (d, *J*=213.5, C-1), 86.7- 82.9 (m, C-2, 3, 5, 6), 61.9 (d, *J* = 18.3, CH₂OH), 26.7 (t, *J* = 22.0, C-4); ¹⁹F NMR (376 MHz, CDCl₃): δ -184.12 (s(br), F-1), -199.8 (d, *J* = 44.0, F-2, 6), -215.7 (s (br), F-3,5); IR: ν_{max}/cm⁻¹ 3290.56 (O-H), 2945.3 (C-H), 1716.66, 1622.13 (C=C), 1388.75 (C-F), 1267.23 (C-F), 1139.93 (C-O), 1026.13 (C-F); FTMS (ESI⁻) *m/z* calcd for ([M]-H⁺) 203.0495; found 203.0500.

1,2,3,5,6-Pentafluorocyclohexyl)methyl methanesulfonate (16):

Et₃N (0.032 mL, 0.23 mmol) and MeSO₂Cl (0.012 mL, 0.14 mmol) were sequentially added to a solution of pentafluoro alcohol **15** (32 mg, 0.12 mmol) in DCM (3 mL) at 0 °C. The reaction was stirred at 0 °C for 30 min, and was then allowed to warm to RT and left stirring for 18 h. The reaction was quenched with H₂O (5 mL) and extracted with DCM (2 x 10 mL). The combined organics were washed with H₂O (10 mL), brine (10 mL), dried (Na₂SO₄) and concentrated. The resulting material was purified over silica gel (petroleum ether/ EtOAc 7:3) to provide the product as a colourless gum (21 mg, 0.076 mmol, 63 %): ¹H NMR (400 MHz, CDCl₃): δ 5.08-4.78 (4H, m, H-2, 3, 5, 6), 4.71 (2H, dt, *J* = 24.7, 2.7, CH₂-O), 3.14 (3H, s, CH₃), 2.72-2.55 (1H, m,

H-4a), 2.54-2.41 (1H, m, H-4b); ^{19}F NMR (376 MHz, CDCl_3): δ -181.4 (s, F-1), -199.9 (s, F-2, 6), -214.5 (s, F-3, 5); ^{13}C NMR (101 MHz, CDCl_3): δ 84.6 (dm, J = 186.8, C-2, 3, 5, 6), 66.8 (d, J = 18.2, $\text{CH}_2\text{-O}$), 37.8 (CH_3), 26.5 (t, J = 22.0, C-4); IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3041.74 (C-H), 1357.89 (S=O), 1174.65 (C-O), 1128.36 (C-F), 1008.77 (C-F); FTMS (ESI^+) m/z calcd for $([\text{M}]+3\text{H}^+)$ 285.0583; found 285.0180.

General coupling procedure for synthesis of compounds 22-25:

EDCI hydrochloride (1.3 eq) and N-methylmorpholine (NMM) (4 eq.) were added to a solution of carboxylic acid **14** (1 eq.) and HOBt (1.3 eq.) in dry DMF and the solution was stirred for 5 min at 0 °C. the appropriate benzylamine (1.3 eq.) was then added and the solution stirred at rt for 16 h. The reaction was diluted by sat ammonium chloride solution, stirred for 1 h and then extracted into ethyl acetate. The organic layer was washed with 10 % NaHCO_3 and brine, dried and the organic solvent was removed under reduced pressure. The products were purified over silica gel (EtOAc/ petrol ether).

1,2,3,5,6-Pentafluoro-N-(4-bromobenzyl)cyclohexane-1-carboxamide (**22**):

The general procedure above was carried out with **14** (100 mg, 0.46 mmol), HOBt (81 mg, 0.6 mmol), NMM (185 mg, 1.83 mmol), EDCI hydrochloride (114 mg, 0.6 mmol) and 4-bromobenzylamine (**18**), (112 mg, 0.6 mmol). After purification over silica (petroleum ether/ EtOAc 6:4) compound **22** (153 mg, 0.4 mmol, 87%) was isolated as a white solid: M.p.= 143-144 °C; ^1H NMR (400 MHz, CDCl_3) δ 7.50 (2H, d, J = 8.4, Ar), 7.19 (2H, d, J = 8.5, Ar), 6.75 (1H, s, NH), 5.11 (2H, d, J = 45.9, H-3, 5), 5.01-4.72 (2H, m, H-2, 6), 4.55 (2H, d, J = 5.8, $\text{CH}_2\text{-Ar}$), 2.77- 2.55 (1H, m, H-4a), 2.58-2.41 (1H, m. H-4b); ^{13}C NMR (101 MHz, CDCl_3) δ 164.0 (CO) , 135.8 (Ar) , 132.0 (Ar) , 129.3 (Ar), 121.9 (Ar), 110.0 (C-1), 85.9 (dm, J = 155.2, C-3, 5), 84.2 (dm, J = 137.7, C-2, 6), 43.3 ($\text{CH}_2\text{-Ar}$), 26.2 (t, J = 22.2, C-4); ^{19}F NMR (376 MHz, CDCl_3) δ -174.77 (s (br), F-1) , -199.52 (dt, J =8.3, 4.5) (F-2,6), - (212.53-213.63) (m, F-3,5); FTMS (ESI^+) m/z calcd for $([\text{M}]+\text{H}^+)$ 386.0178; found 386.0172; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3462.22 (N-H), 2956.87 (C-H), 2306.87, 1697.55 (C=O), 1529.55 (C=C), 1487.12 (C=C), 1114.86 (C-N), 1028.06 (C-F).

1,2,3,5,6-Pentafluoro-N-(4-fluorobenzyl)cyclohexane-1-carboxamide (**23**):

The general procedure above was carried out with **14** (10 mg, 0.046 mmol), HOBt (8 mg, 0.06 mmol), NMM (19 mg, 0.183 mmol), EDCI hydrochloride (11 mg, 0.06 mmol) and 4-fluorobenzylamine (**19**), (10 mg, 0.07 mmol). Purification over silica gel (petroleum ether/ EtOAc 7:3) gave compound **23** (9 mg, 0.026 mmol, 57%) as white solid: M.p.= 181-182 °C; ^1H NMR (400 MHz, CDCl_3) δ 7.30-7.28 (2H, m, Ar) 7.07 (2H, dd, J = 17.8, 8.6, Ar), 6.72 (1H, s, NH), 5.12 (2 H, d, J = 47.2, H-3, 5), 4.98-4.71 (2H, m, H-2, 6), 4.57 (2H, d, J = 5.7, $\text{CH}_2\text{-Ar}$), 2.69-2.61 (1H, m, H-4a), 2.58-2.44 (1H, m, H-4b); ^{13}C NMR (126 MHz, CDCl_3) δ 195.6 (CO), 163.4 (d, J = 240.2, Ar), 132.5 (d, J = 3.1, Ar), 129.4 (d, J = 8.3, Ar), 115.8 (d, J = 21.5, Ar), 85.2 (dm, J = 16.6, C-3, 5), 84.3 (dm, J = 137.4, C-2, 6), 43.3 ($\text{CH}_2\text{-Ar}$), 29.7 (C-4); ^{19}F NMR (377 MHz, CDCl_3) δ -114.17 (F-Ar), -174.8 (F-1), -199.5 (dt, J = 8.2, 4.4, F-2, 6), -(210.2-214.8) (m, C-3, 5); FTMS (ESI^+) m/z calcd for $([\text{M}]+\text{H})$ 326.0979; found 326.0974; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3566.38

(N-H), 2922.16 (C-H), 2852.72 (C-H), 2358.94, 1716. 12(C=O), 1683(C=C), 1515.1 (C=C), 1338.6 (C-F).

1,2,3,5,6-Pentafluoro-N-(4-nitrobenzyl)cyclohexane-1-carboxamide (24):

The general procedure above was carried out with **14** (100 mg, 0.46 mmol), HOBt (81 mg, 0.6 mmol), NMM (185 mg, 1.83 mmol), EDCI hydrochloride (114 mg, 0.6 mmol) and 4-Nitrobenzylamine hydrochloride (**20**), (112 mg, 0.6 mmol). Purification over silica gel (petroleum ether/ EtOAc 6:4) gave compound **25** (130 mg, 0.37 mmol, 82 %) as white solid: M.p.= 156-157 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.25 (2H, d, *J* = 9.2, Ar), 7.49 (2H, d, *J* = 9.2, Ar), 6.90 (1H, s, NH), 5.14 (2H, d, *J* = 47.2, H-3, 5), 5.03-4.76 (2H, m, H-2, 6), 4.72 (2H, d, *J* = 6.0, CH₂-Ar), 2.75-2.63 (1H, m, H-4a), 2.59-2.45 (1H, m, H-4b); ¹³C NMR (101 MHz, CDCl₃) δ 164.3 (CO), 148.3 (Ar), 144.1 (Ar), 128.1 (Ar), 124.1 (Ar), 110.0 (C-1), 88.1-82.1 (m, C-2, 3, 5, 6), 43.2 (CH₂-Ar), 26.3 (t, *J* = 22.6, C-4); ¹⁹F NMR (376 MHz, CDCl₃) δ -174.68 (s(br), F-1), -199.45 (dt, *J* = 8.1, 4.3, F-2, 6), -212.48-212.70 (m, F-3, 5); FTMS (ESI⁺) *m/z* calcd for ([M]+H) 353.0924; found 353.0914; IR: ν_{max}/cm⁻¹ 3296.35 (N-H), 2360.87, 2343.51, 1681 (C=O), 1558.48 (N-O), 1519 (C=C), 1248.24 (C-F).

(1,4-Phenylenebis(methylene))bis(1,2,3,5,6-

pentafluorocyclohexane-1-carboxamide) (25):

The general procedure above was carried out with **14** (30 mg, 0.138 mmol), HOBt (24 mg, 0.18 mmol), NMM (56 mg, 0.55 mmol), EDCI hydrochloride (35 mg, 0.18 mmol) and 1,4-Phenylenedimethanamine (**21**), (25 mg, 0.18 mmol). Purification over silica gel (petroleum ether/ EtOAc 6:4) gave compound **23** (52 mg, 0.097 mmol, 70%) as white solid: M.p.= 178 °C; ¹H NMR (700 MHz, (CD₃)₂CO) δ 8.14 (2H, s(br), NH), 7.31 (4H, s, Ar), 5.45 (4H, d, *J* = 44.1 Hz, H-3, 5), 5.28-4.94 (4H, m, H-2, 6), 4.55 (4H, d, *J* = 6.0, CH₂Ar), 2.59-2.51 (1H, m, H-4a), 2.51-2.41 (1H, m, H-4b); ¹³C NMR (176 MHz, (CD₃)₂CO) δ 163.2 (CO), 163.1 (CO), 137.4 (Ar), 127.5 (Ar), 110.0 (C-1), 85.3 (dm, *J* = 189.1, C-2, 3, 5, 6), 42.8 (CH₂Ar), 26.3 (t, *J* = 43.2, C-4); ¹⁹F NMR (659 MHz, (CD₃)₂CO) δ -171.35 (F-1), -200.53 (d, *J* = 43.7, F-2, 6), -215.26 (F-3, 5); FTMS (ESI⁺) *m/z* calcd for ([M]+H⁺) 537.1599; found 537.1587; IR: ν_{max}/cm⁻¹ 3469.94 (N-H), 3061.03 (C-H), 1716/64 (C=O), 1523.76 (C=C), 1128.36 (C-F), 1060.85 (C-F).

1,2,3,5,6-Pentafluoro-N-(4-aminobenzyl)cyclohexane-1-carboxamide (26):

SnCl₂·2H₂O (38 mg, 0.17 mmol) was added to a solution of carboxamide **24** (20 mg, 0.057 mmol) in EtOAc (10 mL). The reaction mixture was then stirred at 90 °C for 16h. The mixture was then filtered through celite, and the filtrate was washed with 10% NaHCO₃. The layers were separated and the aq layer was further extracted with EtOAc. The combined organics were dried (MgSO₄) and concentrated under reduced pressure to provide product **27** (11 mg, 0.037, 65%) as a yellow solid: M.p.= decomposed at 250 °C; ¹H (400 MHz, MeOD) δ 7.09 (2H, d, *J* = 8.2, Ar), 6.72 (2H, d, *J* = 8.1, Ar), 5.28 (2H, d, *J* = 46.7, H-3, 5), 5.05-4.80 (2H, m, H-2, 6), 4.39 (2H, s, CH₂-Ar), 2.62-2.40 (2H, m, H-4); ¹³C NMR (126 MHz, MeOD)

δ 164.2 (CO), 146.4 (Ar), 128.2 (Ar) , 127.3 (Ar) , 115.4 (Ar), 100 (C-1), 85.0 (dm, J = 185.2, C-2, 3, 5, 6), 42.7 (CH₂-Ar) , 26.1 (t, J = 24.2, C-4); ¹⁹F NMR (376 MHz, MeOD) δ -201.26 (s(br), F-1), -201.69 (s(Br), F-2, 6), -215.93 (s(br), F-3, 5); FTMS (ESI⁺) m/z calcd for ([M]+H) 323.118; found 323.116; IR: $\nu_{\text{max}}/\text{cm}^{-1}$ 3462.22 (N-H), 2956.87 (N-H), 2360.87, 1697.55 (C=O), 1529.55 (C=C), 1114.86 C(-F), 1028.08 (C-F).

3. NMR Spectra (^1H , ^{13}C , ^{19}F NMR)

1-bromo-2,3,5,6-tetrafluorocyclohexyl)benzene (4)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

2,3,5,6-tetrafluoro-1-phenylcyclohexan-1-ol (6)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

2,3,5,6-tetrafluoro-1-phenylcyclohexyl)acetamide (7)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

2,3,5,6-tetrafluoro-1-phenylcyclohexan-1-amine (8)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1-amino-3,5,6-trifluoro-2-phenylcyclohexan-2-ol (9)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1-acetamido-2,3,5,6-tetrafluorocyclohexane-1-carboxylic acid (10):

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1-amino-2,3,5,6-tetrafluorocyclohexane-1-carboxylic acid (11)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1,2,3,5,6-pentafluorocyclohexane-1-carboxylic acid (14)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1,2,3,5,6-pentafluorocyclohexyl)methanol (15)

¹H NMR (400 MHz, CDCl₃)

¹³C NMR (101 MHz, CDCl₃)

¹⁹F NMR (376 MHz, CDCl₃)

MsOH₂C F **1,2,3,5,6-pentafluorocyclohexyl)methyl methanesulfonate (16)**

¹H NMR (400 MHz, CDCl₃)

¹³C NMR (101 MHz, CDCl₃)

¹⁹F NMR (376 MHz, CDCl₃)

1,2,3,5,6-pentafluoro-*N*-(4-bromobenzyl)cyclohexane-1-carboxamide (22)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1,2,3,5,6-pentafluoro-*N*-(4-fluorobenzyl)cyclohexane-1-carboxamide (23)

1,2,3,5,6-pentafluoro-*N*-(4-nitrobenzyl)cyclohexane-1-carboxamide (24)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

1,2,3,5,6-pentafluoro-*N*-(4-aminobenzyl)cyclohexane-1-carboxamide (26)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (101 MHz, CDCl_3)

^{19}F NMR (376 MHz, CDCl_3)

4. ^{19}F NMR VT studies: Conversion of compound 4 to 6.

In order to verify if the reaction undergoes through intermediate involving DMF or directly by an attack of the OH^- on the cationic intermediate, the reaction has been performed in the NMR tube directly in NMR machine at 80°C using DMF as a solvent and a D_2O probe with the ^{19}F [^1H] NMR being recorded every hour.

Figure 4.1: Magnified ^{19}F [^1H decoupled] NMR in D_2O of the reaction, areas (A) and (B), starting from 0 h at r.t (top) followed by 0.5 h, 1 h, 2 h, 3 h, 4 h, 5 h, 6 h at 80°C , sample at r.t after 6 h at 80°C and control sample of final product 6 which has been dissolved in DMF and run with D_2O probe.

The formation of the new signals (Figure 4.1, section A and B) was observed. At the closer look of Figure 4.1 (section A) the new signal could be observed at δ -198.5 ppm after just 1 hour at 80°C and at 2 hours the second signal at δ -197.0 ppm start to appear and increase with the time. Similar effect also observed in the area (B) (Figure 2.4), where the second set of fluorine signals corresponding to the final product 6, starting with appearing signal at δ -207.4 ppm after 1 h and then it starts to split into two peaks after 2 h, with a gradual increase over the time of the second signal (at δ -207.6 ppm).

Once the reaction is cooled down, the signals shift and change from the broad singlets to a more refined AA'BB' pattern with complete disappearance of the intermediate signals at δ -198.5 ppm and δ -207.4 ppm. Figure 4.2 indicates the comparison of the ^{19}F [^1H] NMR of the cooled down reaction to the starting material and to the previously prepared product **6**, which has been dissolved in DMF with D_2O probe, showing ^{19}F [^1H] NMR signals of control product and the product formed during the reaction are sufficiently similar in order to assume that they are the same products.

Figure 4.2: ^{19}F [^1H] NMR signal comparison of the reaction at r.t after 6 hours at 80 °C (top), of starting material **4** (middle) and of previously prepared product **6**.

5. 2D ^1H , ^1H -DQF-COSY (A), 2D ^1H , ^{19}F HMBC (B) and 1D ^{19}F with ^1H decoupling (C) and 1D selective ^1H , ^{19}F -HOESY (D, E, F) NMR spectra for 1-amino-3,5,6-trifluoro-2-phenylcyclohexan-2-ol (9)

(A) 2D ^1H , ^1H -DQF-COSY NMR

(B) 2D ^1H , ^{19}F HMBC NMR

Figure S.1: (A) 2D ^1H , ^1H -DQF-COSY NMR (300 MHz, MeOD) spectrum of 9 at rt in MeOD; (B) 2D ^1H , ^{19}F HMBC NMR spectrum of 9 at rt (377 MHz, MeOD).

Figure 5.2: C) ^1H NMR Spectrum of 9 in MeOD; D) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-6 at rt; E) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-5; F) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-3. NMR spectra were recorded using Bruker AVANCE 500 MHz spectrometer equipped with 5 mm QNP-probe at rt.

6. (A) ^1H , ^{19}F NMR HMBC; (B) ^{19}F observe with ^1H decoupling; (C) 1D selective ^1H , ^{19}F HOESY (C, D, E, F) for 4-bromobenzyl-1,2,3,5,6-pentafluorocyclohexane-1-carboxamide (22)

A) ^1H , ^{19}F (500 MHz, CD_2Cl_2) HMBC NMR

B) ^{19}F (500 MHz, CD_2Cl_2) with ^1H decoupling NMR, SW 80 ppm

Figure 6.1: A) 2D ^{19}F HMBC NMR spectrum of 22; B) ^{19}F with ^1H decoupling NMR spectrum of 22 along with the structure of 13, identifying the 3 fluorine peaks that relate to F-1, 2, 3, 5, 6 fluorine atoms.

Figure 6.2: 1D selective ^1H , ^{19}F -HOESY spectra were recorded using Bruker AVANCE 500 MHz spectrometer equipped with 5 mm QNP-probe at rt; C) ^1H NMR Spectrum of 22.D) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-1; E) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-2 and F-6; F) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-3 and F-5.

7. (A) ^1H , ^{19}F NMR HMBC; (B) ^{19}F observe with ^1H decoupling; (C) 1D selective ^1H , ^{19}F HOESY (C, D, E, F) for 1,2,3,5,6-pentafluoro-4-nitrobenzyl cyclohexane-1-carboxamide (24)

A) ^1H , ^{19}F (500 MHz, CD_2Cl_2) HMBC NMR

B) ^{19}F (500 MHz, CD_2Cl_2) with ^1H decoupling NMR, SW 80 ppm

Figure 7.1: A) ^{19}F HMBC NMR spectrum of 24. B) ^{19}F with ^1H decoupling NMR spectrum of 24 along with the structure of 24, identifying the 3 fluorine peaks that relate to F-1, 2, 3, 5, 6 fluorine atoms.

Figure 7.2: C) ^1H NMR Spectrum; D) 1D selective ^1H , ^{19}F -HOESY with increased number of scans up to 800, irradiation of F-1; E) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-2,-6. F) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-3,-5

8. (A) ^1H , ^{19}F NMR HMBC; (B) ^{19}F observe with ^1H decoupling; (C) 1D selective ^1H , ^{19}F HOESY (C, D, E, F) for 1,4-phenylenebis(methylene)bis-1,2,3,5,6-pentafluorocyclohexane-1-carboxamide (25)

Figure 8.1: A) ^{19}F HMBC NMR spectrum of 25. B) ^{19}F with ^1H decoupling NMR spectrum of 25 along with the structure of 25, identifying the 3 fluorine peaks that relate to F-1, 2, 3, 5, 6 fluorine atoms.

Figure 8.2: C) ^1H NMR Spectrum of 25. D) 1D selective ^1H , ^{19}F -HOESY with increased number of scans up to 800: Irradiation of F-1; E) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-2 and F-6; F) 1D selective ^1H , ^{19}F -HOESY: Irradiation of F-3 and F-5.

9. Density Functional Theory Computations

Coordinates of the DFT-optimised α -fluoroamide 16

Cartesian coordinates in Å, B3LYP/6-311+G** optimised (xyz format)

ax1

F,0,0.9687001115,7.436209174,3.9340216595
F,0,2.664408351,6.8014625287,6.177233602
F,0,1.6219930437,4.4706040042,7.2197199292
F,0,-0.90835014,3.7992726836,5.74120872
F,0,-1.3568685499,5.7330169911,3.8274539268
O,0,-0.5150795221,6.3522242234,6.9684756659
N,0,-0.4224530984,8.3491946315,5.8888072802
C,0,0.6855581596,6.4112198459,4.8827848051
C,0,2.0630552146,5.8873895557,5.3223816603
H,0,2.6730660482,5.8616260576,4.4115871679
C,0,2.1159707985,4.4834923049,5.9267438207
H,0,3.171156527,4.2002903101,5.9983717421
C,0,1.3645643871,3.4771269083,5.0543218768
H,0,1.9448831401,3.3019547365,4.140983936
H,0,1.3081420183,2.5241191351,5.583756497
C,0,-0.0488751048,3.9088415126,4.6615875504
H,0,-0.441386764,3.2411635999,3.8877370028
C,0,-0.0549769937,5.325166946,4.0845758149
H,0,0.4546438599,5.2728743205,3.1150817183
C,0,-0.1518682918,7.0342877689,6.0342007836
C,0,-1.188890387,9.0669434972,6.8974679838
H,0,-2.1223512113,8.5415629492,7.1065195707
H,0,-1.4130519204,10.0645901549,6.5210290255
H,0,-0.0594287869,8.8356599648,5.0857497228
H,0,-0.6284100192,9.1518113955,7.8322360279

ax2

F,0,0.6063631172,1.037763822,2.1005786926
F,0,0.273277177,2.4196218923,-0.4579074744
F,0,-1.0081311473,0.5515410109,-2.0079589305
F,0,-1.1490351734,-1.9217274347,-0.717831749
F,0,0.053391708,-1.8145591667,1.7440258838
O,0,2.6245344837,0.2781450021,0.757224913
N,0,1.4804928735,-0.6304713407,-0.9700758452
C,0,0.2228887813,0.4337527156,0.9009069394
C,0,-0.5862117704,1.5192451156,0.1540788209
H,0,-1.1245196191,2.073845928,0.9315211245
C,0,-1.6307411742,1.0483283277,-0.8542324405
H,0,-2.2113700097,1.9157029407,-1.1770399668
C,0,-2.5357368126,-0.0294185207,-0.2661636853
H,0,-3.157335237,0.4150666379,0.5189466324
H,0,-3.2114505201,-0.3978916504,-1.0404498936
C,0,-1.7535548343,-1.1990044989,0.321954691
H,0,-2.4185750502,-1.9037425742,0.8263350388
C,0,-0.7028038502,-0.7292511649,1.3256241225
H,0,-1.2480815798,-0.365804607,2.2044271715
C,0,1.576170082,0.0100015669,0.2101339678
C,0,2.6790631555,-1.051933763,-1.6831206036
H,0,3.3321856856,-0.1979840272,-1.8740124429

H,0,2.3780628218,-1.4944969856,-2.6321048063
H,0,0.5798291846,-0.8394184142,-1.3696663536
H,0,3.2403347082,-1.7887248116,-1.1033478063

eq1

F,0,-0.547837692,6.0209747952,7.2113418717
F,0,0.0800578561,2.5062639789,7.3156208361
F,0,-0.894045656,2.8795498154,9.8317915304
F,0,3.0792548989,5.4476255642,9.137557586
F,0,2.4078736025,4.0575457119,6.8630105668
O,0,0.1551481443,5.8223032644,4.6568484188
N,0,-0.0090953905,3.5569847275,4.7445319578
C,0,0.170767335,4.8724213854,6.8469830284
C,0,-0.5013868216,3.7384479763,7.6548695512
H,0,-1.5603586262,3.6764621142,7.3961642108
C,0,-0.3238083402,3.9591473592,9.154739927
H,0,-0.8964812632,4.8469762622,9.4366879916
C,0,1.1422186888,4.1060948855,9.5452232867
H,0,1.2216632411,4.2717949112,10.6209403456
H,0,1.6883353669,3.1948564949,9.2929911624
C,0,1.7356019267,5.2931964179,8.7943275095
H,0,1.2292414063,6.2225857613,9.0714469882
C,0,1.6295693575,5.1374281839,7.2805930111
H,0,1.993077437,6.0295578723,6.7671534642
C,0,0.0999473254,4.7862544225,5.2923389187
C,0,0.026122627,3.3797808187,3.2971054988
H,0,-0.6575192963,4.0839434792,2.8234265463
H,0,-0.2838185462,2.3618159747,3.0628948845
H,0,0.0920001249,2.7485138004,5.339756838
H,0,1.0296603239,3.5532601225,2.8979114794

eq2

F,0,-0.5115157648,-0.0616232962,1.7202083571
F,0,0.2468676509,-1.2907378805,-1.5515214178
F,0,2.198955942,-2.5413937879,-0.1121860611
F,0,2.5971410692,2.2008943403,0.3613981013
F,0,0.5573365278,1.5140415939,-1.3182873908
O,0,-2.1335604152,0.6166092962,-1.3466300727
N,0,-2.8663306791,-0.1880920589,0.6540616046
C,0,-0.463553548,0.0569048505,0.3002477316
C,0,0.1976974837,-1.2637926782,-0.1705028842
H,0,-0.4132749572,-2.1094635503,0.1573042887
C,0,1.6138071464,-1.3823664906,0.3992564504
H,0,1.5360662415,-1.5222926765,1.4812822959
C,0,2.4772630857,-0.1705112391,0.0689861298
H,0,3.4699945478,-0.2947716453,0.5051236835
H,0,2.5802285222,-0.0721810621,-1.013494797
C,0,1.8138530758,1.0796395516,0.6380368944
H,0,1.7247997368,1.0196787002,1.7260170754
C,0,0.4229272707,1.3004508434,0.0391392902
H,0,-0.0493015617,2.1849025645,0.4749219635
C,0,-1.9108791699,0.2101629145,-0.2248252498
C,0,-4.2825982798,-0.1015869442,0.3209081494
H,0,-4.6729850585,0.9095328367,0.4714513005
H,0,-4.8380708728,-0.797846267,0.9495349709

H,0,-2.5940557341,-0.3943773858,1.6021028382
H,0,-4.4211102594,-0.3693655293,-0.725875252

ax1 (in water)

F,0,1.0128642494,0.029359834,2.003224109
F,0,0.175880595,2.4032669388,0.547913929
F,0,-1.5366920137,1.4976302468,-1.3853889707
F,0,-1.488833498,-1.4922632027,-1.3833734835
F,0,0.2262655666,-2.3658645543,0.55732803
O,0,0.9671120318,0.0610038708,-1.5426158307
N,0,2.614799167,-0.0131621994,0.0159205236
C,0,0.2905509124,0.0200553237,0.7766543251
C,0,-0.6003398752,1.2744234537,0.8258713964
H,0,-0.9327236447,1.3685527486,1.8637492215
C,0,-1.8589074287,1.2794253487,-0.0389855959
H,0,-2.4680471077,2.1332680363,0.2637335688
C,0,-2.6519731498,-0.0130419099,0.117107351
H,0,-3.111063551,-0.0190003758,1.1104170707
H,0,-3.4703559384,-0.0227542527,-0.6046160534
C,0,-1.8277238131,-1.2858528929,-0.0387656991
H,0,-2.4181534641,-2.1551482864,0.2573469278
C,0,-0.5743343784,-1.2527571528,0.8309061569
H,0,-0.9063105406,-1.3510704543,1.8681652931
C,0,1.3328313846,0.0264329318,-0.3767237932
C,0,3.708989568,-0.0222460384,-0.9479323478
H,0,3.627249269,-0.8817359571,-1.6164561423
H,0,4.6473151428,-0.0831070199,-0.4003916443
H,0,2.823366429,-0.035193038,1.0018119485
H,0,3.7039000876,0.8887836014,-1.5502762906

ax2 (in water)

F,0,0.6433059045,0.8491787873,2.1783674686
F,0,0.2243126736,2.480807881,-0.2088935564
F,0,-1.0210263812,0.7597245116,-1.92345668
F,0,-1.1054489609,-1.8620001155,-0.8851706582
F,0,0.1361242458,-1.9497460048,1.5526855646
O,0,2.6152940648,0.4324363196,0.6411573885
N,0,1.4547652406,-0.5719489476,-1.0181291508
C,0,0.2189184228,0.3668895287,0.9286419955
C,0,-0.6273451033,1.5070783013,0.3214148022
H,0,-1.1609754132,1.9742145558,1.1534937781
C,0,-1.6682658725,1.114331092,-0.7208210634
H,0,-2.2772358259,1.9889216207,-0.9517300017
C,0,-2.5351540189,-0.0438902732,-0.2466361459
H,0,-3.1698761192,0.3012633398,0.5749467123
H,0,-3.2033370758,-0.3518882168,-1.0522268681
C,0,-1.728347028,-1.243995589,0.22702511
H,0,-2.3740736843,-2.0064942389,0.6622649653
C,0,-0.6736882728,-0.8497545256,1.2556781263
H,0,-1.210898735,-0.5958086699,2.1734714122
C,0,1.5545880042,0.0648101134,0.1532463716
C,0,2.6339148646,-0.8729297284,-1.8236394329
H,0,3.1689768252,0.0429938891,-2.0819265093
H,0,2.3064196552,-1.3654659019,-2.7371184044
H,0,0.5539689902,-0.8786265974,-1.3542849258
H,0,3.3157125995,-1.5343721313,-1.2845922982

eq1 (in water)

F,0,-0.1318400542,0.307937789,1.991198065
F,0,-0.011610757,-1.5846070581,-1.0396031143
F,0,2.2736461447,-2.4700040595,0.1246407875
F,0,2.4200433903,2.2461498108,-0.6516457159
F,0,0.027415428,1.2096685911,-1.4578901804
O,0,-2.5271674933,1.0385011323,1.1744133695
N,0,-2.5165871314,-0.4975666088,-0.4956196696
C,0,-0.389355205,0.1166215195,0.625088266
C,0,0.2436490716,-1.2580890409,0.3058042275
H,0,-0.2163883116,-2.03419464,0.9184412263
C,0,1.7551957713,-1.2234794252,0.5096228375
H,0,1.959991826,-1.1153763277,1.5764334538
C,0,2.4258944889,-0.1203729999,-0.3001417778
H,0,3.4986405481,-0.1225658357,-0.1034181915
H,0,2.264944742,-0.2912208289,-1.3666933101
C,0,1.8326879933,1.2222113174,0.1087554812
H,0,2.0488933736,1.4458957868,1.1557130718
C,0,0.3221455992,1.2879081813,-0.0914884088
H,0,-0.0806491566,2.2319748061,0.2780636142
C,0,-1.932755237,0.2456845555,0.4526492727
C,0,-3.9471927396,-0.3997721491,-0.7715015715
H,0,-4.528921794,-0.6779996059,0.1091256938
H,0,-4.1848134358,-1.0798176609,-1.5867926543
H,0,-1.9496678619,-1.1012444136,-1.0723727497
H,0,-4.2159471996,0.6182061641,-1.0611140227

25

eq2 (in water)

F,0,-0.5176987715,0.0070149109,1.7164229509
F,0,0.3868302015,-1.3969797578,-1.4503755882
F,0,2.3645399466,-2.4155339174,0.097053703
F,0,2.4375616356,2.3629486638,0.1497673205
F,0,0.4454177017,1.4238285161,-1.4292222319
O,0,-2.1329273978,0.1043457083,-1.4475002986
N,0,-2.8819789716,-0.0258669238,0.6929584393
C,0,-0.474099677,0.0168585745,0.301687875
C,0,0.2843822196,-1.2844382204,-0.0651707017
H,0,-0.2761337524,-2.1488831762,0.2950745614
C,0,1.690952873,-1.2633638887,0.5349346518
H,0,1.6100047052,-1.3438897824,1.6202740957
C,0,2.4838259608,-0.0275261802,0.126401411
H,0,3.4640032695,-0.0468676527,0.6046095155
H,0,2.6228307082,-0.0181356867,-0.9568901436
C,0,1.7275146795,1.2226623067,0.5605265623
H,0,1.6446515689,1.282062622,1.6469948569
C,0,0.3262603905,1.2978629247,-0.0466784957
H,0,-0.2085087336,2.1757716854,0.3198714074
C,0,-1.9216875648,0.0391510543,-0.2418340062
C,0,-4.2968560732,-0.0226850727,0.3361370923
H,0,-4.57091494,0.9126332831,-0.1566197405
H,0,-4.8805246925,-0.1311158184,1.2478237326
H,0,-2.6203123875,-0.0656081098,1.6662482153
H,0,-4.524021899,-0.8513980626,-0.3369861845