Cumulative effects of cyclones and bleaching on coral cover and species richness at Lizard Island

Authors: Joshua S. Madin^{1,2}, Andrew H. Baird³, Tom C. L. Bridge^{3,4}, Sean R. Connolly³, Kyle J. A. Zawada^{2,5}, Maria Dornelas⁵

¹Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, Kaneohe, Hawai'i, USA

²Department of Biological Sciences, Macquarie University, Sydney NSW 2109, Australia

³Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, QLD 4811, Australia

⁴Biodiversity and Geosciences Program, Museum of Tropical Queensland, Queensland Museum Network, Townsville, QLD 4810, Australia

⁵Centre for Biological Diversity, Scottish Oceans Institute, University of St. Andrews, KY16 9TH, Scotland, UK

Abstract: Coral reefs are being subjected to an increase in the frequency and intensity of disturbance, such as bleaching and cyclones, and it is important to document the effects of such disturbance on reef coral assemblages. Between March 2014 and May 2017, the reefs of Lizard Island in the northern section of the Great Barrier Reef were affected by four consecutive, disturbances: severe tropical cyclones Ita and Nathan in 2014 and 2015, and mass bleaching in 2016 and 2017. Loss of coral cover following the cyclones was patchy and dependent on the direction of the waves generated. In contrast, loss of cover following bleaching was much more uniform. Overall, coral cover declined five-fold from 36% pre-cyclone Ita to 7% post-bleaching in 2017, while mean species richness dropped from 10 to four species per transect. The spatial scale and magnitude of the loss of coral cover in the region suggests that it will be many years before these reefs recover.

27

28 Key words: community ecology, coral reefs, climate change, disturbance, diversity,

29

30 **Introduction**

31 Coral reefs are under threat, with the general consensus that these ecosystems are degrading
32 globally (Hughes et al. 2003, Hughes et al. 2018a). Reef-building corals provide the habitat
33 structure that allows associated coral reef fauna to thrive. Therefore, declines in coral cover lead to
34 loss of fish abundance and species richness (e.g. Jones et al. 2004, Graham et al. 2006). In the
35 context of global change, the intensity and frequency of climate-associated disturbances on coral
36 reefs are increasing (Bender et al. 2010). To predict the fate of coral reefs, we need to document and
37 understand the effects of different types of disturbance on reef coral abundance and diversity.

38 In the Indo-Pacific region, coral cover has declined on average by 0.7% per year since the
39 late 1960s (Bruno & Selig 2007). Coral cover change is not monotonic: typically, declines are
40 episodic and driven by disturbance events such as crown-of-thorn starfish outbreaks, cyclones and,
41 more recently, mass bleaching events. Disturbance-led declines are typically interspersed with
42 periods when, under the right conditions, coral cover increases (Gilmour et al. 2013). Coral reefs
43 are naturally mosaics of communities at different successional stages of recovery from a myriad of
44 disturbances. However, rapid sequences of intense disturbances can result in phase-shifts in coral
45 ecosystems (Hughes 1994, Hughes & Connell 1999) from which recovery is much less likely. Some
46 reefs can recover from large losses of cover, with the abundance of herbivores, topographic
47 complexity, and depth being important predictors of the probability of recovery (Graham et al.
48 2015).

49 Sections of the northern Great Barrier Reef (GBR) centred around Lizard Island were
50 recently affected by a sequence of extreme disturbances—the first disturbances on record to affect
51 these reefs in 20 years (Wakeford et al. 2008, Pratchett 2010, Death et al. 2012). In April 2014,
52 Severe Tropical Cyclone (STC) Ita (Category 4) crossed the island from the north. In March 2015,

53 STC Nathan (Category 4) affected the island through waves generated from the south. The largest
54 mass bleaching event on record affected the island in April 2016, with estimates of bleaching and
55 loss of coral cover in the area of between 50-100% (Hughes et al. 2017), and close to 90% for
56 Lizard Island (Hughes et al. 2018b). We documented a subsequent bleaching event in early 2017.
57 This sequence of disturbance has drastically changed the assemblage structure of reefs around the
58 island when compared to data from 2011 and the mid-1990s. Here, we quantified the cumulative
59 effects of the two cyclones and two bleaching events on the cover, richness and composition of
60 coral assemblages at 19 sites around Lizard Island.

61
62 **Materials and methods**

63 We used line intercept transects (Loya 1972) to estimate coral cover and species diversity at two to
64 19 sites around Lizard Island, in the northern Great Barrier Reef (14.6688° S, 145.4594° E) in 1995,
65 1996, 1997, 2011, 2014, 2015, 2016 and 2017 (see map in Fig. 3). North Reef and South Island
66 were surveyed in 1995; Lizard Head and Washing Machine were added in 1996; North Reef, Lizard
67 Head, Trimodal, Lagoon 1 and 2 and Horseshoe reef sites were sampled on every sampling
68 occasion since 2011, the remaining 13 sites were sampled since 2015. At each site, five or six 10m
69 transects were haphazardly deployed parallel to the reef margin at 1 to 2 m depth and no less than
70 10m apart. All colonies with a maximum diameter greater than 5 cm were identified to genus
71 between 1995 and 1997 and to species between 2011 and 2017 following Veron (1986). We did not
72 transform cover or richness data, and used Welch's t-tests for unequal variances for reporting
73 comparisons with an alpha of 0.05. Analyses were run using the function "t.test" in the statistical
74 software R (R Core Team 2018).

75
76 **Results and Discussion**

77 Coral cover, taxonomic richness and assemblage structure at the start of surveys in 2011, and prior
78 to the sequence of severe disturbances, were very similar to those in the historical surveys in the

1990s (Figs. 1 and 2). The series of disturbances corresponded with in a five-fold loss of coral cover between 2011 and 2017 at the six sites that were surveyed consistently during this time (from 36% to 7% average cover; Fig. 1). At the island scale, the decline in cover was particularly high following cyclone Nathan in 2015, after which cover approximately halved, followed by the 2016 bleaching, after which cover halved again. However, the spatial pattern of loss varied among the different disturbances (Fig. 3). Following cyclones, coral loss was greatest at sites facing the direction from which the cyclone approached. For example, following STC Ita, cover declined proportionally by 85% at North Reef ($t=5.33$, $df=7.47$, $p<0.001$), whereas there was no statistically significant change in cover at sites on the south side of the island (e.g., Trimodal, Lagoon 1 and 2, Lizard Head and Horseshoe). Similarly, south-facing sites suffered major declines following cyclone Nathan. Coral cover declined proportionally by 90% and 66% at Trimodal and Lagoon 2, respectively ($t=9.7$, $df=8.57$, $p<0.001$; $t=2.98$, $df=6.62$, $p=0.02$), whereas there was no further change in cover at North Reef ($t=0.40$, $df=2.82$, $p=0.72$). In contrast, loss of cover following bleaching was more uniform across the six main sites following both bleaching events, with all sites with greater than 10% cover remaining showing significant declines in cover. Nonetheless, coral cover at some of the 19 sites surveyed since 2014 has remained relatively constant or even increased, particularly Resort, Cooks Path and Turtle Beach (Fig. 3), which were dominated by taxa such as *Montipora* and *Porites* that are less susceptible to bleaching. Overall, these patterns are consistent with previous research suggesting that cyclone-driven declines in cover are patchy and mediated by exposure to prevailing winds (Connell et al. 2004, Wakeford et al. 2008, Fabricius et al. 2008), whereas bleaching-driven declines tend to be more uniform and contingent on the structure of the coral assemblage at the time of bleaching (Marshall & Baird 2000, McClanahan et al. 2005).

The cumulative results of these four major disturbances are drastically altered coral assemblages. Loss of cover was accompanied by declines in genus and species richness (Fig. 4). The total number of species across the six consistently surveyed sites decreased from 76 in 2011 to

105 49 in 2017. Furthermore, only 28 of the species recorded in 2011 were recorded again in 2017.
106 Many of the additional 21 species recorded in 2017 were likely hidden within the living structure of
107 the reef, such as underneath tabular-*Acropora* spp. colonies (Baird & Hughes 2000). Therefore, the
108 loss of species richness is likely to be more extensive than the observed net difference before and
109 after the disturbances (i.e., 76 vs. 28 species). The spatial patterns across sites for average species
110 richness per transect (and genus richness, which could be included for 1995-97) were consistent
111 with the patterns of change in cover, but far less pronounced (Fig. 1 and 3).

112 In the historical surveys, and prior to STC Ita, most reefs around Lizard Island were
113 dominated by *Acropora* spp. (Fig. 2). For instance, on Trimodal Reef in 2005 there were 43
114 *Acropora* spp, including the third and fourth most abundant species at the site (Dornelas &
115 Connolly 2008). Only one of these species, *A. hyacinthus*, occurred on the transects in 2017 and
116 was represented by only a few, very small colonies. *Acropora* cover declined by over 95% (Fig. 2,
117 dark blue), more than any other genus. Currently, the dominant benthic taxa at Lizard Island are soft
118 corals and members of the hard coral genus *Porites*, and family *Faviidae*. Here, *Porites*,
119 *Pocillopora*, *Faviidae* and *Stylophora* were the only common taxa that lost less than half their cover
120 across the sequence of disturbances (Fig. 2).

121 The two cyclones and two bleaching events have changed Lizard Island coral assemblages
122 profoundly. The high mortalities resulting from these events have reduced coral cover to below 4%
123 at ten out of the 19 sites. Only two sites have cover above 40% in 2017, which was the norm for
124 Lizard Island's reefs prior to these disturbances (Fig. 1; Wakeford et al. 2008, Pratchett et al. 2010).
125 Moreover, the species most affected were those that created much of the structural complexity of
126 these reefs. The current most common taxa have much simpler morphologies (e.g., massive *Porites*
127 and *Faviidae*) and account for very little coral over on the reef of Lizard Island. These changes are
128 likely to have severe knock-on effects on the abundance and diversity of fishes that rely on
129 structural complexity for habitat (Pratchett et al. 2006, Wilson et al. 2006).

Whether the reefs around Lizard will recover from this degraded state remains to be seen.

Estimates of recovery time following major disturbances range from 5-20 years (Done et al. 2010, Johns et al. 2014). While the effects of the cyclones were dramatic, the spatially clustered nature of their effects mean that multiple source populations remained in the region to potentially supply recruits for recovery. However, the cumulative effects of the cyclones and mass bleaching means that far fewer reproductively active adults remain in the region. In fact, recruitment rates in the area have dropped by two orders of magnitude following the mass bleaching (Woods et al. in prep) and are likely to increase the expected recovery period. In contrast, the abundance of juveniles (i.e., colonies < 5 cm maximum diameter) has been much more stable, and the juveniles of most taxa were less affected by bleaching than the adults (Alvarez-Noriega et al. 2018) offering some promise of a recovery, but only if the region does not experience another severe disturbance event in the next decade.

References

- Alvarez-Noriega M, Baird AH, Bridge T, Dornelas M, Fontoura L, Precoda K, Torres-Pulliza D, Woods RM, Zawada K, Madin JS (2018) Contrasting patterns of bleaching susceptibility between juvenile and adult corals. *Coral Reefs* 37:527-532
- Baird AH, Hughes TP (2000) Competitive dominance by tabular corals: an experimental analysis of recruitment and survival of understorey assemblages. *Journal of Experimental Marine Biology and Ecology* 251:117-132
- Bender MA, Knutson TR, Tuleya RE, Sirutis JJ, Vecchi GA, Garner ST, Held IM (2010) Modeled Impact of Anthropogenic Warming on the Frequency of Intense Atlantic Hurricanes. *Science* 327:454-458
- Bruno JF, Selig ER (2007) Regional decline of coral cover in the Indo-Pacific: Timing, extent, and subregional comparisons. *Plos One* 2:e711

156 De'ath G, Fabricius KE, Sweatman H, Puotinen M (2012) The 27-year decline of coral cover on the
157 Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the*
158 *United States of America* 109:17995-17999

159 Done T, DeVantier L, Turak E, Fisk D, Wakeford M, van Woesik R (2010) Coral growth on three
160 reefs: development of recovery benchmarks using a space for time approach. *Coral Reefs*
161 29:815-833

162 Dornelas M, Connolly SR (2008) Multiple modes in a coral species abundance distribution.
163 *Ecology Letters* 11:1008-1016

164 Fabricius KE, De'ath G, Puotinen ML, Done T, Cooper TF, Burgess SC (2008) Disturbance
165 gradients on inshore and offshore coral reefs caused by a severe tropical cyclone. *Limnology*
166 *and Oceanography* 53:690-704

167 Gilmour JP, Smith LD, Heyward AJ, Baird AH, Pratchett MS (2013) Recovery of an isolated coral
168 reef system following severe disturbance. *Science* 340:69-71

169 Graham NAJ, Jennings S, MacNeil MA, Mouillot D, Wilson SK (2015) Predicting climate-driven
170 regime shifts versus rebound potential in coral reefs. *Nature* 518:94-97

171 Graham NAJ, Wilson SK, Jennings S, Polunin NVC, Bijoux JP, Robinson J (2006) Dynamic
172 fragility of oceanic coral reef ecosystems. *Proceedings of the National Academy of Sciences*
173 *of the United States of America* 103:8425-8429

174 Hughes TP (1994) Catastrophes, Phase-Shifts, and Large-Scale Degradation of a Caribbean Coral-
175 Reef. *Science* 265:1547-1551

176 Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK,
177 Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H,
178 Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett
179 M, Schoepf V, Torda G, Wilson SK (2018a) Spatial and temporal patterns of mass bleaching
180 of corals in the Anthropocene. *Science* 359:80-83

181 Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-
 182 Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nystrom M, Palumbi SR,
 183 Pandolfi JM, Rosen B, Roughgarden J (2003) Climate change, human impacts, and the
 184 resilience of coral reefs. *Science* 301:929-933

185 Hughes TP, Connell JH (1999) Multiple stressors on coral reefs: A long-term perspective.
 186 *Limnology and Oceanography* 44:932-940

187 Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock
 188 RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE,
 189 Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF,
 190 Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs J-PA, Hoogenboom MO, Kennedy
 191 EV, Kuo C-y, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ,
 192 Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda
 193 G, Wachenfeld DR, Willis BL, Wilson SK (2017) Global warming and recurrent mass
 194 bleaching of corals. *Nature* 543:373-377

195 Hughes TP, Kerry JT, Baird AH, Connolly SR, Dietzel A, Eakin CM, Heron SF, Hoey AS,
 196 Hoogenboom MO, Liu G, McWilliam MJ, Pears RJ, Pratchett MS, Skirving WJ, Stella JS,
 197 Torda G (2018b) Global warming transforms coral reef assemblages. *Nature* 556:492-496

198 Johns KA, Osborne KO, Logan M (2014) Contrasting rates of coral recovery and reassembly in
 199 coral communities on the Great Barrier Reef. *Coral Reefs* 33:553-563

200 Jones GP, McCormick MI, Srinivasan M, Eagle JV (2004) Coral decline threatens fish biodiversity
 201 in marine reserves. *Proceedings of the National Academy of Sciences of the United States of*
 202 *America* 101:8251-8253

203 Loya Y (1972) Community structure and species diversity of hermatypic corals at Eilat, Red Sea.
 204 *Marine Biology* 13:100-112

205 Marshall PA, Baird AH (2000) Bleaching of corals on the Great Barrier Reef: differential
 206 susceptibilities among taxa. *Coral Reefs* 19:155-163

207 McClanahan TR, Maina J, Moothien-Pillay R, Baker AC (2005) Effects of geography, taxa, water
 208 flow, and temperature variation on coral bleaching intensity in Mauritius. *Marine Ecology-
 209 Progress Series* 298:131-142
 210 Pratchett MS, Wilson SK, Baird AH (2006) Long-term monitoring of the Great Barrier Reef. *J Fish
 211 Biol* 69:1269-1280
 212 Pratchett MS (2010) Changes in coral assemblages during an outbreak of *Acanthaster planci* at
 213 Lizard Island, northern Great Barrier Reef (1995–1999). *Coral Reefs* 29:717-725
 214 R Core Team (2018) R: A language and environment for statistical computing. R Foundation for
 215 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
 216 Veron JEN (1986) Veron, J. E. N. 1986. Corals of Australia and the Indo-Pacific. Angus &
 217 Robertson, Sydney.
 218 Wakeford M, Done TJ, Johnson CR (2008) Decadal trends in a coral community and evidence of
 219 changed disturbance regime. *Coral Reefs* 27:1–13
 220 Wilson SK, Graham NAJ, Pratchett MS, Jones GP, Polunin NVC (2006) Multiple disturbances and
 221 the global degradation of coral reefs: are reef fishes at risk or resilient? *Glob Change Biol*
 222 12:2220-2234
 223 Woods RM, Baird AH, Madin JS (unpublished) A collapse in larval supply following mass coral
 224 bleaching.
 225
 226
 227
 228
 229

230 **Acknowledgments**

231 We thank the Lizard Island Research Station staff for their support. Funding was provided by the
 232 Australian Research Council Centre of Excellence for Coral Reef Studies (AHB COE140100020)
 233 and the Templeton Foundation (60501). MD is grateful to the Scottish Funding Council (MASTS,

234 grant reference HR09011), and the European Research Council (grant BioTIME, and grant
235 BioCHANGE). MD and JSM thank the John Templeton Foundation for funding.

236

237 **Figure captions**

238 **Figure 1.** Change in average (A) coral cover, (B) genus richness and (C) species richness per
239 transect ($\pm$ SE) for the six consistently surveyed reefs at Lizard Island. Black arrows: tropical
240 cyclones; red arrows: bleaching events.

241

242 **Figure 2.** Compositional change in cover of the ten most dominant taxa: nine coral genera and a
243 single category for soft corals. Arrows correspond with tropical cyclones (black) and bleaching
244 events (red). Soft coral was not surveyed in the 1990s. Due to low densities, *Seriatopora* was not
245 recorded in some years. Sample sizes reported refer to number of sites surveyed (see Methods for
246 details of within-site replication).

247

248 **Figure 3.** Change in coral cover (grey bars $\pm$ SE) and per transect species richness (white bars $\pm$ SE)
249 across the 19 Lizard Island sites (6 surveyed since 2011, 13 since 2015). Grey shading land and
250 light grey shading is coral reef.

251

252 **Figure 4.** Temporal trajectories of changes in coral cover and species richness per transect for the
253 six consistently surveyed sites. Grey background points show data for all sites and years.

254

255

256

257

258

259

261

262

263

264