

# **Trends in sleeping difficulties among European adolescents: are these associated with physical inactivity and excessive screen time?**

Ariane Ghekiere<sup>1</sup>, Jelle Van Cauwenberg<sup>1,2</sup>, Ann Vandendriessche<sup>1</sup>, Joanna Inchley<sup>3</sup>, Margarida Gaspar de Matos<sup>4</sup>, Alberto Borraccino<sup>5</sup>, Inese Gobina<sup>6</sup>, Jorma Tynjälä<sup>7</sup>, Benedicte Deforche<sup>1,8</sup>, Bart De Clercq<sup>1</sup>

<sup>1</sup> Department of Public Health, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

<sup>2</sup> Fund for Scientific Research Flanders (FWO), Brussels, Belgium

<sup>3</sup> Child and Adolescent Health Research Unit, School of Medicine, University of St Andrews, St Andrews, United Kingdom

<sup>4</sup> Faculdade de Motricidade Humana / University of Lisbon, Lisboa, Portugal

<sup>5</sup> Department of Public Health and Paediatrics, University of Torino, Torino, Italy

<sup>6</sup> Department of Public Health and Epidemiology, Riga Stradins University, Riga, Latvia

<sup>7</sup> Faculty of Sport and Health Sciences, Research Centre for Health Promotion, University of Jyväskylä, Jyväskylä, Finland

<sup>8</sup> Physical Activity, Nutrition and Health Research Unit, Faculty of Physical Education and Physical Therapy, Brussels, Belgium

Corresponding author:

Bart De Clercq

[b.declercq@ugent.be](mailto:b.declercq@ugent.be)

T. +32 (0)9 332 36 18

F. +32 (0)9 332 31 20

## **Conflict of Interest**

The authors declare no conflicts of interest.

## **Acknowledgements**

The International Coordinator of the 2002 – 2014 HBSC surveys was Candace Currie. Data Bank Manager was Oddrun Samdal.

## **Keywords**

Adolescents – 24-hour approach – sleep – sedentary behaviour – physical activity – trend

## Abstract

**Objectives.** We examined changes in sleep-onset difficulties over time, and associations with physical activity and screen time behavior among adolescents.

**Methods.** We used data from last four survey waves of the Health Behavior of School-aged Children (HBSC) study (2002-2006-2010-2014). Multilevel logistic regression analyses were conducted to explore associations between regular sleeping difficulties, excessive screen time exposure, and being insufficiently physically active (i.e. < 60 min daily) among 33 European and non-European countries.

**Results.** Findings indicate an increase in the prevalence of sleep-onset difficulties and in excessive screen time exposure, and a small but significant increase in physical activity levels. Additionally, adolescents exceeding 2h daily screen time had 20% higher odds of reporting sleep-onset difficulties, while no association was found for physical activity. The strength of the association between screen time and sleep-onset difficulties increased over time, which may reflect a change in type of screen time use (e.g. the increased use of easy accessible screens such as smartphones and tablets).

**Conclusions.** Effective strategies to reduce screen time are key to reverse the detrimental trend in sleep-onset difficulties among adolescents.

**Trends in sleeping difficulties among European adolescents: are these associated with physical inactivity and excessive screen time?**

## Introduction

Adolescents, defined as young people between the age of 10 to 19 years old, have often been considered as having a low burden of disease and relatively good physical and mental health compared with other stages of the life course (Gore et al. 2011). Nevertheless, the period of adolescence is characterized by many changes, such as physical changes in weight, height and muscle mass, as well as changes in life style behaviours (Christie and Viner 2005). A number of health-related behaviours (e.g. physical activity, sedentary behaviour, nutrition and sleep) are shaped during adolescence and impact health during adolescence, but also have profound consequences for cardio-metabolic health during adulthood (Ekelund et al. 2012).

In 2014, Canadian researchers emphasized the importance of using a 24-hour approach to study the link between several health behaviours and actual health status (Irish et al. 2014). This 24-hour approach stresses the importance of all health behaviours performed in a day, including sleep, physical activity and sedentary behaviour, to explain health outcomes. Sleep is an important health behaviour as humans sleep approximately one third of a day (Irish et al. 2014), it impacts the circadian metabolic effects and allows the brain to rest and consolidate day experiences (Brand and Kirov 2011). Furthermore, insufficient sleep has been associated with bad mood, high blood pressure, increased risk of type II diabetes, higher engagement in risky behaviour and drowsy driving and lower academic performances (Hysing et al. 2016; Owens 2014; Short et al. 2013).

However, sleep has been largely neglected in the past as an important health behaviour. The 24-hour approach also emphasizes the need to better understand how waking behaviours such as physical activity and sedentary behaviour are associated with sleep, as it is hypothesized that these behaviours are interrelated (Irish et al. 2014). For example, an association between screen time (a highly prevalent type of sedentary behaviour) and sleep difficulties can partly be explained by the exposure to blue light, which increases the alertness of the individual and may induce difficulties in falling asleep (Boniel-Nissim et al. 2015; Nuutinen et al. 2013). The association between sleep and physical activity has been hypothesized to be bi-directional, i.e. engagement in physical activity is associated with better sleep, but evidence is also accumulating that poor sleep may also lead to lower physical activity levels (Kline

2014). The association between physical activity and sedentary behaviour is not univocal, with some studies reporting an inverse association (Pearson et al. 2014; Tanaka et al. 2017), while others reported no association (Borraccino et al. 2009; Serrano-Sanchez et al. 2011). Recently, the interrelation between sleep, screen time and physical activity was tested within a large sample of English adults and showed that low physical activity levels, high screen time levels and poor sleep duration tend to cluster. The clustering of these three behaviours was much more prevalent in overweight or obese adults compared to a normal weight group, showing that these health behaviours are linked with obesity Cassidy et al. (2017). One recent U.S. study found that more daily screen time was associated with worse sleep quality, while lower physical activity was associated with more awakenings at night and higher sleep fragmentation among girls aged 7- to 12 years with a low socio-economic status (Greever et al. 2017). Also, Chilean girls aged 12 years old with low screen time and high physical activity levels reported lower prevalence of sleeping difficulties, compared to those with high screen time and low physical activity levels (Aguilar et al. 2015). Finally, a study among Chinese college students showed a positive association between screen time and sleeping difficulties, but no association was found between physical activity and prevalence of sleeping difficulties (Wu et al. 2015). However, large scale studies examining the association between these three behaviours in an international context are currently lacking. Additionally, no studies have explored this association among adolescents in a European context.

International trend analyses showed a strong increase in screen time among adolescents during the past decade, while physical activity levels generally remain low and efforts to increase physical activity seem to have small but positive effects on the prevalence of adolescents being sufficiently physically active (Bucksch et al. 2016; Inchley et al. 2017; Kalman et al. 2015; Nelson et al. 2006). Only limited information is available on how the prevalence of sleeping difficulties has changed during the past decade (Kronholm et al. 2015; Pallesen et al. 2008b). Since physical activity and screen time have been shown to be related to sleeping difficulties, it would be informative to describe the trends in sleeping difficulties together with the trends in physical activity and screen time.

Sleeping difficulties is a term which has been used in a wide range of contexts, and can cover many problems related to bad sleep. The current study focusses on difficulties in falling asleep. Limited

information is available on the prevalence of difficulties of falling asleep among adolescents. To the authors knowledge, there is only one paper which describes the changes in difficulties on falling asleep among a sample of Norwegian adolescents, and found that prevalence increased significantly between 1983 and 2005 (Pallesen et al. 2008a). No evidence is available on how this prevalence might have changed during the last decade, how this might differ across countries, and limited information is available on factors associated with these problems with falling asleep. Therefore, the aim of the current study was two-fold. First, we aimed to describe the prevalence of sleep-onset difficulties, insufficient physical activity and excessive screen time in 33 European countries between 2002 and 2014. Secondly, the aim was to examine the associations of physical activity and screen time with sleep-onset difficulties and whether these associations have changed over time and differ between countries.

## Methods

Data from the four most recent waves of the Health Behaviour of School-aged Children (HBSC) (2002-2006-2010-2014) were used for the current analysis. The HBSC study is a large international study developed by an international network of researchers, in collaboration with the World Health Organization (WHO) Regional Office for Europe. The study aims to obtain insights into health behaviours and well-being among school children via a self-reported questionnaire which is repeated every four years (World Health Organisation 2016).

The current study includes only those 33 countries/regions which participated in the four waves (Belgium – Flemish and French part; United Kingdom – Scotland, Wales, England). Sample selection protocols were described previously (World Health Organisation 2016). Questionnaires, which were in the national language of the participating areas, were completed in schools by adolescents aged 11, 13 and 15 years old. For the purposes of the current study, questions regarding physical activity, screen time and sleep-onset difficulties were included in the analyses, along with age, gender and material socio-economic position, measured using the Family Affluence Scale (Currie et al. 2008). Physical activity was self-reported by the adolescents as the number of days of being physically active at moderate-to-vigorous-intensity for 60 minutes in the past 7 days (range 0 to 7) (Prochaska et al. 2001). Screen time behaviour was measured by responses on three different questions (“How many hours a

day, in your free time, do you usually spend watching TV, videos, DVDs, and other entertainment on a screen?”, “How many hours a day, in your free time, do you usually spend playing games on a computer, games console, tablet (like iPad), smartphone or other electronic device (not including moving or fitness games)?” and “How many hours a day, in your free time, do you usually spend using electronic devices such as computers, tablets (like iPad) or smart phones for other purposes, for example, homework, emailing, tweeting, Facebook, chatting, surfing the internet?”) separately for week and weekend days. Nine response categories were possible, ranging from none at all to up to more than 7 hours daily. These responses were treated as continuous, and were thus recoded as follows: none at all = 0, half an hour a day = 0.5, about an hour a day = 1.0 etc. Average daily screen time was calculated by summing these three behaviours for week- and weekend days separately, which were used to create a weighted average mean ( $= (5 \times \text{weekday} + 2 \times \text{weekend day}) / 7 \text{ days}$ ) (Rey-López et al. 2010). Sleep-onset difficulties were assessed by a single item: “how often in the past six months did you have difficulties in falling asleep?”. Response categories were “about every day”, “more than once a week”, “about every week”, “about every month” and “rarely or never”. The three behaviours were dichotomized to conform with the current international health guidelines for physical activity and screen time, i.e. being physically active for 7 days (1) or not (0) (World Health Organization 2011), exceeding 2hour daily screen time (1) or not (0) (American Academy of Pediatrics: Children, adolescents, and television 2001) and having difficulties in falling asleep ‘about every day/more than once a week’ (1) or less (0) (Irish et al. 2014).

## Analysis

First, prevalence (%) was calculated for each country/region ( $n=33$ ) for each survey year (2002, 2006, 2010, 2014) for not meeting the physical activity guidelines ( $=$ less than 7 days physically active for at least 60 minutes in last 7 days), exceeding screen time guidelines ( $=$ more than 2h per day) and having regular sleep-onset difficulties ( $=$ about every day and more than once a week). Twelve year differences were calculated (prevalence of behaviour in 2014 - prevalence in 2002) and mapped on geographical areas (i.e. countries) to illustrate geographical differences in the changes over time, with the use of Tableau Software. A mean score was calculated per country if region information was available (i.e. Belgium and United Kingdom). Positive trends were marked green, while detrimental (unfavourable) changes were marked red. Secondly, a multilevel logistic regression model was fitted in R studio (lme4



package) to assess whether sleep-onset difficulties could be explained by exceeding the screen time guidelines and not reaching the physical activity guidelines. In order to examine if the observations vary between clusters (i.e. countries), we calculated the Intraclass Correlation Coefficient (ICC). The ICC indicated that 3.1% of the variance in sleep-onset difficulties is attributable to contextual differences (i.e. differences between countries). In addition, the likelihood ratio test showed that the multilevel logistic model fitted the data better than the single-level logistic model. Therefore, adolescents (level 1) were clustered within year by country units (level 2), which in turn were clustered within countries (level 3) (Fairbrother 2013). A stepwise model fitting approach was used, by first fitting a time effect model (model 1), then age, gender and FAS were added (model 2). Age and FAS were centered around their mean. In order to be able to assess the relative rank within the socioeconomic hierarchy across different time points and cross-national contexts, FAS-scores were transformed to ridsits corrected for survey year, country, gender and age. A ridity is a proportional rank score ( $j$ ) that represents the proportion ( $P$ ) of observations with lower scores ( $\sum_{n=1}^{j-1} P_{2n}$ ) plus one-half the proportion with equal scores ( $\frac{P_{2j}}{2}$ ) so that  $Ridity_j = \sum_{n=1}^{j-1} P_{2n} + \frac{P_{2j}}{2}$ . To calculate the ridity-based FAS score, we simply add the cumulative proportion of the adjacent lower value ( $\sum_{n=1}^{j-1} P_{2n}$ ) to one-half the proportion of the value itself ( $\frac{P_{2j}}{2}$ ). When added to a regression model of health, the coefficient for the ridity-based FAS score represents the slope index of health inequality (i.e. the difference in health between the highest and lowest FAS groups). Physical activity was added to model 2 (model 3a), followed by screen time (model 3b), and physical activity and screen time simultaneously (model 3c). In model 4a and model 4b, a random slope over country was allowed to test whether associations differed across countries, for physical activity and screen time respectively. Finally, interactions between physical activity and time (model 5a), and screen time and time (model 5b) were tested to examine whether associations of physical activity and screen time with sleep-onset difficulties have changed over time. Model fit was compared using Akaike and Bayesian Information Criterion (AIC and BIC, respectively) values. Loglikelihood ratio tests were conducted to examine significant improvements/deteriorations in model fits. Significance was set at  $\alpha < 0.05$ .

## Results

### Sample descriptives

Table 1 shows an overview of the sample descriptives across the four years. Age and sex distributions are comparable across survey years. Prevalence of sleep-onset difficulties and excessive screen time increased over time, while a small increase in the prevalence of sufficient physical activity can be observed. Additionally, an increase in mean daily screen time can be observed.

### Descriptive trends in sleep-onset difficulties, physical activity and screen time

Different time trends can be observed across countries (see Table 2). Figures 1 to 3 geographically illustrate the magnitude of the 12 year changes in sleep-onset difficulties, screen time and physical activity. In most countries, an increase in the prevalence of sleep-onset difficulties was observed over 12 years, except for Greece (-4.3%), Spain (-4.1%), Norway (-2.0%), England (-1.7%) and Portugal (-1.3%). In 2014, on average around 20% of the adolescents had regular difficulties in falling asleep, but prevalence ranged from 9.6% in Ukraine to 37.4% in France.

For physical activity, 19 out of 33 countries (58%) showed a positive change over 12 years, i.e. decreases in the proportion of adolescents not being moderately or vigorously physically active every day (see Figure 2). However, prevalence of insufficient physical activity was still very high across all countries with, on average, 80% of adolescents being insufficiently physically active in 2014 ranging from 89.9% (Italy) to 72.2% (Finland). It can also be observed that most countries improved their physical activity levels from 2002 to 2006, followed by a small decrease in physical activity in 2010 and again small increases in physical activity levels from 2010 to 2014.

Prevalence of exceeding the 2h screen time/day has increased in almost all countries, except for Israel (-1.6%) and Lithuania (-27.7%) (see Figure 3). Around 90% of the adolescents in the participating countries exceeded the guidelines of 2h of daily screen time in 2014, ranging from 56.7% in Lithuania to 95.4% in the Netherlands. It can also be observed that countries with lower prevalence in 2002 had larger increases in their prevalence for excessive screen time (and vice versa, e.g. Israel), and differences between countries decreased over time. In 2014, prevalence of excessive ST was over 90% in 12 out of 33 countries.

## Change in sleep-onset difficulties over time and the association with screen time and physical activity

Results of the multilevel logistic regression models are shown in Table 3. Model 1 shows a significant time trend for the odds of having regular sleep-onset difficulties. No differences were observed between 2002 and 2006, but a significant increase in odds was found for 2010 (OR=1.102, 95% CI 1.043-1.164) and 2014 (OR=1.208, 95% CI 1.145-1.275) compared to 2002. Adding socio-demographic variables to the model increased the model fit (model 2), i.e. age, gender and FAS were significantly associated with the odds of having regular sleep-onset difficulties. Adolescents aged one year older had 2.3% higher odds (95% CI=1.019-1.027) of having sleep-onset difficulties. Higher affluent groups had 3.3% (95% CI=0.964-0.970) lower odds of having sleep-onset difficulties compared to the lower affluent groups. Girls had 44.0% (95% CI= 1.421-1.458) higher odds to report sleep-onset difficulties. Adding conformity to the physical activity guidelines into the model (3a) only slightly improved the model fit, and physical activity was not significantly associated with sleep-onset difficulties. Adding conformity to the screen time guidelines to model 2 increased the model fit, and illustrated a positive association between excessive screen time and the odds of sleep-onset difficulties (model 3b). Adolescents exceeding the guidelines for screen time had 20.1% (95% CI=1.178-1.225) higher odds of reporting regular sleep-onset difficulties. In model 3c, both conformity to the physical activity and screen time guidelines were added simultaneously to the model, and showed a slightly better model fit compared to model 3b in which only screen time was added. This model confirms that adolescents who exceeded the screen time guidelines had higher odds of reporting regular sleep-onset difficulties even after adjustment for physical activity. In model 4a and 4b, a random country slope was added for physical activity and screen time respectively, and each showed a significant increase in model fit (model 4a vs 3a  $\chi^2$  (2df)=72.777,  $p<0.001$ ; model 4b vs 3b  $\chi^2$  (2df)=54.22,  $p<0.001$ ). This implies that the strength of the association between physical activity and screen time with sleep-onset difficulties differs significantly between countries. Finally, an interaction term for physical activity (model 5a) and screen time (model 5b) by time was added to the model. Model 5a showed a significant interaction between physical activity and time showing that the magnitude of the association between physical activity and sleep-onset changes over time, although in none of the years, the magnitude differed significantly from 1. The

association between physical activity and sleep-onset difficulties was significantly larger in 2014 compared to 2002 (OR 2002=0.992, 95% CI=0.976-1.010; OR 2006=0.991, 95% CI=0.958-1.026; OR 2010=1.015, 95% CI=0.981-1.095; OR 2014=0.971, 95% CI=0.939-1.004). Model 5b illustrates that the association between screen time and sleep-onset difficulties got stronger over time (OR 2002 =1.201, 95% CI=1.178-1.225; OR 2006 =1.163, 95% CI=1.118-1.211; OR 2010=1.219, 95% CI=1.170-1.270, OR 2014=1.419, 95% CI=1.360-1.480). The magnitude of the association between screen time and sleep-onset difficulties increased each wave and all waves did significantly differ from 2002 (see Table 3, model 5b). Country specific effect sizes for screen time and physical activity are available in a supplementary file (Supplementary Table 1 & 2).

## Discussion

The current study was the first to describe co-occurrence of trends in sleep-onset difficulties, physical activity and screen time, and to examine the association between sleep-onset difficulties, physical activity and screen time among a large sample of mainly European adolescents.

Across the majority of countries, unfavourable changes were observed for screen time and sleep-onset difficulties, while physical activity levels slightly increased over time. Results indicated increases in the prevalence of sleep-onset difficulties over time (odds ranged between 1.208 (model 1) and 1.279 (model 4a)). The prevalence of sleep-onset difficulties differed across gender (girls had 44% higher odds of reporting sleep-onset difficulties), socio-economic groups (lower SES was associated with more sleep-onset difficulties), and age groups (older adolescents reported more sleep-onset difficulties). These differences in sleep-onset difficulties between socio-demographic groups were found in some previous studies (Lundqvist 2014; Wiklund et al. 2012). However, the existence of these differences needs further investigation. As sleep-onset difficulties may lead to the development of more serious sleeping disorders (e.g. impairment of sleep onset, sleep maintenance, premature final morning awakening, non-restorative sleep) and delayed sleep time disorders, it is important to identify appropriate strategies to decrease the prevalence of sleep-onset difficulties. For example, adolescent could be stimulated to develop consistent sleep schedules, given that adolescents with a schedule on both weekdays and weekends experience less difficulties in falling asleep than their counterparts with inconsistent sleep schedules (Mindell and Owens 2010).

Excessive screen time gets more prevalent over time, and in many countries, more than 90% of the adolescents exceeded the recommendations of 2h screen time daily. This increase in excessive screen time was recently described in the HBSC International report 2014 (Inchley et al. 2017), and can be attributed to a rapid increase in computer/tablet/smartphone use for both gaming and non-gaming purposes. Although not investigated within the current study, it can be hypothesized that this increase is mainly explained by the increased use of smartphones among adolescents during the past few years (Škařupová et al. 2016). Smartphones are portable and accessible screen time devices, and especially when used before going to bed, they negatively influence sleep quality and quantity (Christensen et al.

2016). This shift in content of screen time may explain our finding that the strength of the association between screen time and sleep-onset difficulties increased over time.

The prevalence of sleep-onset difficulties was related to the prevalence of excessive screen time, but no association was found with physical activity, which is in contrast to findings of a previous study conducted in Switzerland, which found that high physical activity groups had less sleep-onset difficulties compared to low physical activity group (Lang et al. 2013). It might be that no association was found in our study due to our crude measure of physical activity. In depth analyses illustrated that in some countries significant associations were found, with most of them being negatively associated with sleep-onset difficulties, but in some regions also positive associations were found (results not shown). More research including more detailed assessment of physical activity (e.g. time of the day, intensity and duration of physical activity) is needed to explore the association between physical activity and sleep-onset difficulties. It could be hypothesized that sleep quality or sleep quantity improves the night following the physical activity session, if it is not performed just before sleeping time (Kline 2014). However, such information is not available within the HBSC study. Similarly, no information was available on the timing of screen time, although it could be hypothesized that screen time right before bedtime may have the largest implications on difficulties in falling asleep (Hale and Guan 2015). In the current study, only difficulties in falling asleep were examined, while other elements related to sleep such as sleep duration, sleep quality, differences in sleep duration between week and weekend days, sleep hygiene and other sleep problems are also important to examine in future studies. Furthermore, more insights are needed in how associations might differ across countries and regions, as no overall association between physical activity and sleep-onset difficulties was found, but sub-analyses showed significant associations in some regions. Therefore, there is need for more in depth study on physical activity and sleep, in order to prevent careless made conclusions that physical activity is not related to sleep/sleep problems.

### Strengths and limitations

This study pioneered to present an international overview of trends over time in sleep-onset difficulties in a large international sample, highlighting an emerging health problem which has often been neglected

in past health behaviour studies. This study was limited to mainly European countries, and future studies should explore whether these trends are also observed in other regions such as the U.S. and Australia, and how associations may differ for in for example regions which are characterized by high mobile device use. Additionally, more insights are needed in the relationship between physical activity, screen time and other sleep related health behaviours or outcomes, as this information was not available in the current study. A second limitation was the use of self-reported measures, which are prone to social desirability and recall bias. Also, screen time levels may have been somewhat overestimated in the current study, due to the co-occurrence of screen time activities. Third, we did not consider the context and content of screen time activities. Although we hypothesized that the increase in screen time is generally explained by the increased use of smartphones during the past few years (Škařupová et al. 2016), this does not automatically imply that this increase is driven by leisure activities. In turn, educational demands such as obligatory electronic homework and tests may have contributed to this increase. In that sense, the association between sleeping difficulties and screen-time may be confounded by un-observed factors. Unfortunately, no such information was available in the data. The use of objective measures which can assess both physical activity, screen time or sedentary behaviour in general and sleep would overcome the above mentioned limitations and allow researchers to further examine the 24-hour approach. This may generate better understanding of the associations between these three behaviours as information on the specific moment of the day, intensity and duration of the activities can be obtained.

## Conclusions

To conclude, findings of the current study indicate that sleep-onset difficulties are an emerging health problem. Additionally, findings suggest that interventions are needed to lower the prevalence of excessive screen time, which could lead to less difficulties in falling asleep among adolescents. However, more insights into the determinants of sleep-onset difficulties (and sleep in general) are needed, as screen time is only one pathway associated with sleep-onset difficulties, and little is known on other potential influencing factors.

## **Compliance with Ethical Standards**

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

HBSC is a WHO collaborative study and as such fulfils all WHO ethical requirements. The study is approved in each country separately by national or ministry ethical committees. The Flemish study was approved by the ethical committee of the University hospital of Ghent (project EC/2013/1145).

## **Informed consent**

Informed consent was obtained from all individual participants included in the study.

## **Funding**

The study was funded by the government of Flanders (grant number GEO-1GDD2A-WT) and the Research Foundation Flanders (grant number 12I117N).

## **Conflict of Interest**

The authors declare no conflicts of interest.



## References

- Aguilar MM, Vergara FA, Velasquez EJA, Garcia-Hermoso A (2015) Physical activity, screen time and sleep patterns in Chilean girls *An Pediatr* 83:304-310 doi:10.1016/j.anpedi.2014.12.006
- American Academy of Pediatrics: Children, adolescents, and television (2001) *Pediatrics* 107:423-426
- Boniell-Nissim M, Lenzi M, Zsiros E et al. (2015) International trends in electronic media communication among 11- to 15-year-olds in 30 countries from 2002 to 2010: association with ease of communication with friends of the opposite sex *European Journal of Public Health* 25:41-45 doi:10.1093/eurpub/ckv025
- Borraccino A, Lemma P, Iannotti, RJ et al. (2009) Socioeconomic effects on meeting physical activity guidelines: comparisons among 32 countries *Med Sci Sports Exerc* 41:749-756 doi:10.1249/MSS.0b013e3181917722
- Brand S, Kirov R (2011) Sleep and its importance in adolescence and in common adolescent somatic and psychiatric conditions *International Journal of General Medicine* 4:425-442 doi:10.2147/IJGM.S11557
- Bucksch J, Sigmundova D, Hamrik Z et al. (2016) International Trends in Adolescent Screen-Time Behaviours From 2002 to 2010 *Journal of Adolescent Health* 58:417-425 doi:<https://doi.org/10.1016/j.jadohealth.2015.11.014>
- Cassidy S, Chau JY, Catt M, Bauman A, Trenell MI (2017) Low physical activity, high television viewing and poor sleep duration cluster in overweight and obese adults; a cross-sectional study of 398,984 participants from the UK Biobank *International Journal of Behavioural Nutrition and Physical Activity* 14:57 doi:10.1186/s12966-017-0514-y
- Christensen MA, Bettencourt L, Kaye L et al. (2016) Direct Measurements of Smartphone Screen-Time: Relationships with Demographics and Sleep *PLOS ONE* 11:e0165331 doi:10.1371/journal.pone.0165331
- Christie D, Viner R (2005) Adolescent development *BMJ : British Medical Journal* 330:301-304
- Currie C, Molcho M, Boyce W, Holstein B, Torsheim T, Richter M (2008) Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) family affluence scale *Social science & medicine* (1982) 66:1429-1436 doi:10.1016/j.socscimed.2007.11.024
- Ekelund U, Luan J, Sherar LB, Esliger DW, Griew P, Cooper A (2012) Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents *Jama* 307:704-712 doi:10.1001/jama.2012.156
- Fairbrother M (2013) Two Multilevel Modeling Techniques for Analyzing Comparative Longitudinal Survey Datasets *Political Science Research and Methods* 2:119-140 doi:10.1017/psrm.2013.24

- Gore FM, Bloem PJN, Patton GC et al. (2011) Global burden of disease in young people aged 10-24 years: a systematic analysis *The Lancet* 377:2093-2102 doi:10.1016/S0140-6736(11)60512-6
- Greever CJ, Ahmadi M, Sirard J, Alhassan S (2017) Associations among physical activity, screen time, and sleep in low socioeconomic status urban girls *Preventive medicine reports* 5:275-278 doi:10.1016/j.pmedr.2017.01.014
- Hale L, Guan S (2015) Screen time and sleep among school-aged children and adolescents: a systematic literature review *Sleep Med Rev* 21 doi:10.1016/j.smr.2014.07.007
- Hysing M, Harvey AG, Linton SJ, Askeland KG, Sivertsen B (2016) Sleep and academic performance in later adolescence: results from a large population-based study *J Sleep Res* 25:318-324 doi:10.1111/jsr.12373
- Inchley J, Currie D, Jewell J, Breda J, Barnekow V (2017) Adolescent obesity and related behaviours: trends and inequalities in the WHO European Region, 2002–2014.
- Irish LA, Kline CE, Rothenberger SD et al. (2014) A 24-hour approach to the study of health behaviours: temporal relationships between waking health behaviours and sleep *Annals of behavioural medicine : a publication of the Society of Behavioural Medicine* 47:189-197 doi:10.1007/s12160-013-9533-3
- Kalman M, Inchley J, Sigmundova D et al. (2015) Secular trends in moderate-to-vigorous physical activity in 32 countries from 2002 to 2010: a cross-national perspective *Eur J Public Health* 25 Suppl 2:37-40 doi:10.1093/eurpub/ckv024
- Kline CE (2014) The bidirectional relationship between exercise and sleep: Implications for exercise adherence and sleep improvement *American journal of lifestyle medicine* 8:375-379 doi:10.1177/1559827614544437
- Kronholm E, Puusniekka R, Jokela J et al. (2015) Trends in self-reported sleep problems, tiredness and related school performance among Finnish adolescents from 1984 to 2011 *J Sleep Res* 24:3-10 doi:10.1111/jsr.12258
- Lang C, Brand S, Feldmeth AK, Holsboer-Trachsler E, Pühse U, Gerber M (2013) Increased self-reported and objectively assessed physical activity predict sleep quality among adolescents *Physiology & Behaviour* 120:46-53 doi:<https://doi.org/10.1016/j.physbeh.2013.07.001>
- Lundqvist L (2014) Socio-demographic variation in sleep difficulties among adolescents in Sweden.
- Mindell JA, Owens JA (2010) *A Clinical Guide to Pediatric Sleep: Diagnosis and Management of Sleep Problems*. Lippincott Williams & Wilkins,
- Nelson MC, Neumark-Stzainer D, Hannan PJ, Sirard JR, Story M (2006) Longitudinal and Secular Trends in Physical Activity and Sedentary Behaviour During Adolescence *Pediatrics* 118:e1627-e1634 doi:10.1542/peds.2006-0926
- Nuutinen T, Ray C, Roos E (2013) Do computer use, TV viewing, and the presence of the media in the bedroom predict school-aged children's sleep habits in a longitudinal study? *BMC Public Health* 13:684 doi:10.1186/1471-2458-13-684

- Owens J (2014) Insufficient sleep in adolescents and young adults: an update on causes and consequences *Pediatrics* 134:e921-932 doi:10.1542/peds.2014-1696
- Pallesen S, Hetland J, Sivertsen B, Samdal O, Torsheim T, Nordhus IH (2008a) Time trends in sleep-onset difficulties among Norwegian adolescents: 1983-2005 *Scand J Public Health* 36:889-895 doi:10.1177/1403494808095953
- Pallesen S, Hetland J, Sivertsen B, Samdal O, Torsheim T, Nordhus IH (2008b) Time trends in sleep-onset difficulties among Norwegian adolescents: 1983--2005 *Scand J Public Health* 36:889-895 doi:10.1177/1403494808095953
- Pearson N, Braithwaite RE, Biddle SJ (2014) Associations between sedentary behaviour and physical activity in children and adolescents: a meta-analysis *Obes Rev* 15 doi:10.1111/obr.12188
- Prochaska JJ, Sallis JF, Long B (2001) A physical activity screening measure for use with adolescents in primary care *Archives of Pediatrics & Adolescent Medicine* 155:554-559 doi:10.1001/archpedi.155.5.554
- Rey-López JP, Vicente-Rodriguez G, Ortega FB et al. (2010) Sedentary patterns and media availability in European adolescents: The HELENA study *Preventive Medicine* 51:50-55 doi:<http://dx.doi.org/10.1016/j.ypmed.2010.03.013>
- Serrano-Sanchez JA, Martí-Trujillo S, Lera-Navarro A, Dorado-García C, González-Henríquez JJ, Sanchis-Moysi J (2011) Associations between Screen Time and Physical Activity among Spanish Adolescents *PLoS ONE* 6:e24453 doi:10.1371/journal.pone.0024453
- Short MA, Gradisar M, Lack LC, Wright HR (2013) The impact of sleep on adolescent depressed mood, alertness and academic performance *Journal of adolescence* 36:1025-1033 doi:10.1016/j.adolescence.2013.08.007
- Škařupová K, Ólafsson K, Blinka L (2016) The effect of smartphone use on trends in European adolescents' excessive Internet use *Behaviour & Information Technology* 35:68-74 doi:10.1080/0144929X.2015.1114144
- Tanaka C, Tanaka M, Okuda M, Inoue S, Aoyama T, Tanaka S (2017) Association between objectively evaluated physical activity and sedentary behaviour and screen time in primary school children *BMC Research Notes* 10:175 doi:10.1186/s13104-017-2495-y
- Wiklund M, Malmgren-Olsson E-B, Öhman A, Bergström E, Fjellman-Wiklund A (2012) Subjective health complaints in older adolescents are related to perceived stress, anxiety and gender – a cross-sectional school study in Northern Sweden *BMC Public Health* 12:993-993 doi:10.1186/1471-2458-12-993
- World Health Organisation (2016) About HBSC - World Health Organization collaborative cross-national survey. <http://www.hbsc.org/about/index.html>.
- World Health Organization (2011) Global Recommendations on Physical Activity for Health 5-17 years old

Wu XY, Tao SM, Zhang YK, Zhang SC, Tao FB (2015) Low Physical Activity and High Screen Time Can Increase the Risks of Mental Health Problems and Poor Sleep Quality among Chinese College Students Plos One 10:10 doi:10.1371/journal.pone.0119607

**Table 1: Sample descriptives across four measurement periods (33 countries), Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014**

	2002	2006	2010	2014
n	155300	167656	172729	175399
<b>Age (mean ± SD, years)</b>	13.5 ± 1.7	13.6 ± 1.7	13.6 ± 1.6	13.6 ± 1.6
<b>Sex (% girls)</b>	51.2	51.0	50.9	50.8
<b>Sleep-onset difficulties (%)</b>				
About every day	8.7	9.0	9.9	11.0
More than once a week	9.1	9.2	9.9	10.5
About every week	10.0	10.0	11.2	11.5
About every month	15.2	15.5	16.7	16.7
Rarely or never	57.0	56.3	52.3	50.3
<b>Physical Activity (% , number of days MVPA)</b>				
0	4.7	4.4	4.2	4.1
1	9.3	7.7	7.4	6.8
2	16.0	13.6	13.7	12.6
3	17.9	16.7	16.7	16.4
4	14.7	15.2	15.9	15.6
5	12.7	14.0	14.4	15.1
6	7.9	9.0	9.5	10.0
7	16.7	19.4	18.1	19.4
<b>Screen time (mean ± SD, hours per day)</b>	4.0 ± 2.4	5.5 ± 3.7	6.0 ± 3.9	6.5 ± 4.4
<b>Screen time (% exceeding 2h daily)</b>	79.8	87.1	88.5	88.2
<b>FAS (mean ridit score)</b>	0.50 ± 0.28	0.50 ± 0.28	0.50 ± 0.28	0.50 ± 0.28

MVPA = Moderate to vigorous physical activity, FAS = Family Affluence Scale

**Table 2: Prevalence of sleep-onset difficulties, insufficient physical activity and excessive screen time across four measurement periods in 33 countries, Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014.**

	Sleep on-set difficulties					Physical activity					Screen time				
	% reporting to have regular difficulties in falling asleep					% not obtaining the PA guidelines of 60 min daily					% exceeding 2h screen time/day				
Country	2002	2006	2010	2014	12 yr evolution	2002	2006	2010	2014	12 yr evolution	2002	2006	2010	2014	12 yr evolution
Austria	16.7	16.9	17.5	17.6	0.9	78.5	80.8	76.6	80.3	1.8		85.5	83.6	85.3	
Belgium (Flemish)	18.1	18.5	18.9	22.2	4.1	89.4	84.6	84.0	85.0	-4.4	80.4	89.4	88.1	89.2	8.8
Belgium (French)	23.8	25.7	29.1	30.4	6.6	100.0	76.0	83.2	82.2	-17.8	72.6	83.9	84.0	87.0	14.4
Canada	23.3	25.7	26.5	27.2	3.9	78.7	76.9	78.0	75.4	-3.3	83.5	90.1	88.2	90.8	7.3
Croatia	12.1	12.7	13.4	13.1	1.0	80.5	77.3	79.9	75.3	-5.2	82.8	88.0	92.3	86.2	3.4
Czech Republic	17.0	18.3	20.9	22.0	5.0	73.4	78.4	77.2	78.7	5.3	78.5	88.6	90.6	89.2	10.7
Denmark	19.6	22.6	25.1	24.7	5.1	84.1	78.1	88.4	86.9	2.8	82.3	90.5	92.9	93.6	11.3
Estonia	13.8	15.1	17.8	18.2	4.4	88.5	82.3	85.8	83.7	-4.8	91.8	95.0	94.5	93.5	1.7
Finland	14.5	17.3	16.4	18.8	4.3	85.3	77.9	76.7	72.2	-13.1	79.3	86.5	90.9	91.2	11.9
France	28.0	29.5	30.7	37.4	9.4	91.4	86.8	88.4	87.6	-3.8	68.7	82.8	83.3	86.9	18.2
Germany	13.4	16.6	17.6	19.9	6.5	88.7	83.4	83.5	84.9	-3.8	75.2	85.4	86.7	86.7	11.5
Greece	18.1	14.4	13.5	13.8	-4.3	82.1	84.4	86.1	86.6	4.5	75.3	87.7	90.9	88.8	13.5
Greenland	20.1	15.8	17.9	24.7	4.6	71.6	72.5	76.1	83.1	11.5	73.8	74.5	69.2		
Hungary	15.2	14.8	15.6	17.1	1.9	84.6	81.1	81.3	77.9	-6.7	79.0	86.2	89.6	88.5	9.5
Ireland	16.4	17.1	18.1	21.9	5.5	72.7	69.9	73.7	77.2	4.5	61.9	75.5	78.7	85.4	23.5
Israel	19.1	23.1	20.1	19.8	0.7	85.3	83.5	83.7	89.3	4.0	90.3	91.9	91.5	88.7	-1.6
Italy	15.5	14.6	16.4	19.8	4.3	89.5	85.0	92.0	89.9	0.4	73.8	79.5	85.6	86.2	12.4
Latvia	16.4	14.3	18.4	18.0	1.6	85.6	77.4	80.4	81.7	-3.9	88.8	91.6	92.8	92.8	4.0
Lithuania	12.6	17.4	18.7	17.9	5.3	74.4	80.9	83.6	79.5	5.1	84.4	90.9	85.1	56.7	-27.7
Macedonia	12.9	15.0	13.4	15.5	2.6	83.2	77.7	78.6	73.7	-9.5	81.2	87.6	89.7	87.5	6.3
Netherlands	22.8	15.6	20.8	26.3	3.5	83.2	78.8	81.3	82.0	-1.2	84.4	94.1	93.7	95.4	11.0
Norway	18.4	19.8	21.6	16.4	-2.0	88.8	85.9	85.2	82.4	-6.4	83.7	86.3	83.5	89.0	5.3
Poland	14.6	14.1	15.9	19.4	4.8	83.1	82.7	80.1	76.1	-7.0	86.7	90.8	92.8	90.9	4.2
Portugal	14.8	13.1	15.9	13.5	-1.3	88.2	85.5	87.0	85.1	-3.1	83.6	91.8	91.9	88.4	4.8
Russia	8.9	11.0	16.6	12.2	3.3	86.3	87.2	87.6	83.0	-3.3	85.1	88.5	92.7	90.6	5.5
Slovenia	16.1	16.5	15.0	22.3	6.2	77.9	82.6	79.9	81.6	3.7	79.2	86.9	89.3	85.5	6.3
Spain	16.7	14.1	15.0	12.6	-4.1	83.5	79.6	77.8	75.3	-8.2	76.7	81.0	86.6	87.0	10.3
Sweden	23.8	20.8	20.9	30.4	6.6	87.8	84.6	87.0	86.4	-1.4	79.2	87.8	92.1	94.0	14.8
Switzerland	17.5	21.3	21.9	24.2	6.7	85.1	87.3	88.1	85.7	0.6	60.8	71.1	75.3	81.4	20.6
Ukraine	9.2	7.9	7.9	9.6	0.4	83.7	78.4	77.5	74.0	-9.7	88.2	89.5	87.2		
UK (England)	23.4	21.1	21.9	21.7	-1.7	78.3	82.3	80.7	82.3	4.0	70.8	86.5	91.7	91.7	20.9
UK (Scotland)	22.3	21.5	23.1	22.7	0.4	80.7	78.5	85.7	82.8	2.1	84.4	91.3	92.0	94.3	9.9
UK (Wales)	22.2	22.5	20.3	24.3	2.1	83.2	79.5	81.3	83.8	0.6	85.3	91.6	91.9	92.5	7.2
<b>Total</b>	<b>17.5</b>	<b>17.8</b>	<b>19.4</b>	<b>20.8</b>	<b>3.3</b>	<b>84.1</b>	<b>81.1</b>	<b>82.2</b>	<b>81.1</b>	<b>-3.0</b>	<b>79.8</b>	<b>87.1</b>	<b>88.5</b>	<b>88.2</b>	<b>8.4</b>

*Detrimental evolutions are marked red, positive changes are marked green.*


**Table 3: Results of the logistic regression analyses on the odds of reporting sleep-onset difficulties, Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014.**

	Model 1		Model 2		Model 3a		Model 3b		Model 3c		Model 4a		Model 4b		Model 5a		Model 5b	
n	671084		635133		635133		635133		615130		635133		615130		615130		615130	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<b>Fixed effects</b>																		
intercept	0.205***	0.186-0.226	0.164***	0.146-0.185	0.164***	0.146-0.185	0.140***	0.124-0.158	0.140***	0.124-0.158	0.164***	0.146-0.185	0.141***	0.126-0.158	0.140***	0.124-0.157	0.152***	0.135-0.172
<b>Level 2: time*country</b>																		
Time (ref=2002)																		
2006	1.018	0.959-1.081	1.038	0.973-1.106	1.038	0.973-1.106	1.027	0.963-1.094	1.027	0.963-1.095	1.041	0.976-1.110	1.027	0.962-1.095	1.033	0.968-1.102	0.974	0.901-1.053
2010	1.102***	1.043-1.164	1.152***	1.081-1.230	1.152***	1.081-1.230	1.140***	1.069-1.216	1.140***	1.069-1.216	1.154***	1.082-1.230	1.140***	1.069-1.215	1.143***	1.071-1.219	1.039	0.960-1.124
2014	1.208***	1.145-1.275	1.273***	1.194-1.358	1.273***	1.194-1.358	1.261***	1.181-1.346	1.261***	1.181-1.346	1.279***	1.199-1.366	1.258***	1.176-1.343	1.273***	1.192-1.360	1.002	0.924-1.086
<b>Level 1: individual</b>																		
Age			1.023***	1.019-1.027	1.023***	1.019-1.027	1.019***	1.015-1.023	1.019***	1.015-1.023	1.023***	1.019-1.027	1.019***	1.015-1.023	1.019***	1.015-1.023	1.018***	1.014-1.022
FAS			0.967***	0.964-0.970	0.967***	0.964-0.970	0.965***	0.962-0.969	0.965***	0.962-0.969	0.967***	0.964-0.970	0.966***	0.962-0.969	0.965***	0.962-0.969	0.965***	0.961-0.969
Gender (ref=boys)			1.440***	1.421-1.458	1.440***	1.420-1.457	1.453***	1.434-1.472	1.453***	1.434-1.472	1.440***	1.421-1.458	1.454***	1.435-1.473	1.453***	1.434-1.472	1.452***	1.433-1.471
PA (ref=less than 7days MVPA/week)					0.992	0.976-1.010			0.999	0.982-1.017	1.002	0.966-1.039			1.029	0.990-1.070	1.000	0.983-1.018
ST (ref=less than 2h/day)							1.201***	1.178-1.225	1.201***	1.178-1.225			1.192***	1.145-1.240	1.201***	1.177-1.225	1.087***	1.050-1.125
PA * Time (2006)																0.963	0.915-1.014	
PA* Time (2010)																0.986	0.937-1.038	
PA * Time (2014)																0.943***	0.897-0.992	
ST * Time (2006)																		1.070*
ST * Time (2010)																		1.121***
ST * Time (2014)																		1.305***
																		1.015-1.128
																		1.106-1.182
																		1.256-1.379
<b>Random effects</b>																		
	Variance		Variance		Variance		Variance		Variance		Variance		Variance		Variance		Variance	
Country variance	0.0892		0.1037		0.1036		0.1046		0.1046		0.1049		0.088		0.1046		0.104	
Time*country variance	0.0162		0.0164		0.0164		0.0158		0.0158		0.0162		0.0159		0.0157		0.0159	
PA											0.0085							
ST													0.0095					
<b>Model fit parameters</b>																		
AIC	639009.2		602357.2		602356.8		584848.7		584850.7		602289.7		584798.4		584850.5		584762.3	
BIC	639077.7		602459.5		602447.7		584962		584975.3		602426.1		584934.4		585009.1		584920.9	
Loglikelihood	-319498.6		-301169.6		-301169.3		-292414.3		-292414.3		-301132.9		-292387.2		-292411.2		-292367.2	

\*p<0.05; \*\*p<0.01;\*\*\*p<0.001; MVPA = Moderate to vigorous physical activity; FAS = Family Affluence Scale




**Figure 1: Twelve year changes in the prevalence of sleep-onset difficulties, Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014**



*Red color indicates detrimental evolutions. Green color indicates positive evolutions.  
The darker the color, the larger the twelve year change.*




**Figure 2. Twelve year changes in the prevalence of excessive screen time, Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014**



*Red color indicates detrimental evolutions. Green color indicates positive evolutions.  
The darker the color, the larger the twelve year change.*

**Figure 3. Twelve year change in the prevalence of insufficient physical activity, Health Behaviour in School-Aged Children (HBSC) survey, 33 European countries, 2002–2014**



*Red color indicates detrimental evolutions. Green color indicates positive evolutions.  
The darker the color, the larger the twelve year change.*

**Supplementary Table 1: Country specific effect sizes for the relation between screen time and sleep-onset difficulties**

	<b>B</b>	<b>B standardized</b>	<b>OR</b>	<b>Significance</b>
Austria	0,281	0,268	1,31	***
Belgium (Flemish)	0,142	0,125	1,13	*
Belgium (French)	0,137	0,117	1,12	**
Canada	0,315	0,226	1,25	***
Croatia	0,004	0,004	1,00	
Czech Republic	0,059	0,050	1,05	
Denmark	0,305	0,221	1,25	***
England	0,162	0,144	1,16	**
Estonia	0,157	0,103	1,11	(*)
Finland	0,402	0,358	1,43	***
France	0,250	0,218	1,24	***
Germany	0,231	0,229	1,26	***
Greece	0,196	0,191	1,21	**
Greenland	0,071	0,080	1,08	
Hungary	0,142	0,136	1,15	*
Ireland	0,178	0,194	1,21	***
Israel	0,080	0,058	1,06	
Italy	0,104	0,109	1,12	(*)
Latvia	0,164	0,121	1,13	*
Lithuania	0,006	0,007	1,01	
Macedonia	0,307	0,300	1,35	***
Netherlands	0,279	0,185	1,20	***
Norway	0,215	0,196	1,22	***
Poland	0,081	0,069	1,07	0
Portugal	0,008	0,007	1,01	
Russia	-0,066	-0,065	0,94	
Scotland	0,236	0,163	1,18	***
Slovenia	0,404	0,373	1,45	***
Spain	0,112	0,122	1,13	*
Sweden	0,364	0,258	1,29	***
Switzerland	0,290	0,312	1,37	***
Ukraine	0,094	0,089	1,09	
Wales	0,203	0,142	1,15	**

(\*)= p<0,1; \*=p<0,05; \*\*=p<0,01; \*\*\*=p<0,001

**Supplementary Table 2: Country specific effect sizes for the relation between physical activity and sleep-onset difficulties**

	<b>B</b>	<b>B standardized</b>	<b>OR</b>	<b>Significance</b>
Austria	-0,056	-0,060	0,94	
Belgium (Flemish)	0,064	0,056	1,06	
Belgium (French)	0,090	0,072	1,07	(*)
Canada	-0,066	-0,063	0,94	*
Croatia	0,056	0,069	1,07	
Czech Republic	0,115	0,122	1,13	**
Denmark	0,006	0,005	1,00	
England	-0,072	-0,068	0,93	
Estonia	0,024	0,023	1,02	
Finland	-0,046	-0,051	0,95	
France	0,129	0,088	1,09	**
Germany	0,041	0,039	1,04	
Greece	0,042	0,004	1,00	
Greenland	0,154	0,168	1,18	
Hungary	0,165	0,177	1,19	**
Ireland	0,037	0,042	1,04	
Israel	0,201	0,175	1,19	***
Italy	0,011	0,009	1,01	
Latvia	0,073	0,076	1,08	
Lithuania	0,014	0,015	1,02	
Macedonia	-0,109	-0,129	0,88	(*)
Netherlands	-0,086	-0,082	0,92	(*)
Norway	-0,146	-0,129	0,88	*
Poland	-0,015	-0,017	0,98	
Portugal	0,058	0,057	1,06	
Russia	0,179	0,193	1,21	**
Scotland	-0,021	-0,021	0,98	
Slovenia	-0,009	-0,009	0,99	
Spain	-0,246	-0,289	0,75	***
Sweden	-0,069	-0,055	0,95	
Switzerland	0,003	0,003	1,00	
Ukraine	0,033	0,040	1,04	
Wales	-0,039	-0,039	0,96	

(\*)= p<0,1; \*=p<0,05; \*\*=p<0,01; \*\*\*=p<0,001