

Supplementary Material for “A balance of winners and losers in the Anthropocene”

Maria Dornelas, Nicholas J. Gotelli, Hideyasu Shimadzu, Faye Moyes, Anne E. Magurran, Brian McGill

Data sources

Metadata for the data sources is available at Figshare: 10.6084/m9.figshare.7687778

	Reference
18	(Zachmann <i>et al.</i> 2010)
33	(Widdicombe <i>et al.</i> 2010a; Widdicombe <i>et al.</i> 2010b)
39	(Holmes & Sturges 1975; Holmes <i>et al.</i> 1986; Holmes & Sherry 1988, 2001)
42	(Beven 1976; Williamson 1987; Gibbons <i>et al.</i> 1993; Standley <i>et al.</i> 1996; Stone <i>et al.</i> 1997; Gaston & Blackburn 2000; Lack 2010)
44	(Halpern & Dyrness 2010; Halpern & Lutz 2013a, b)
46	(Williamson 1983)
47	(Vickery & Nudds 1984)
51	(NERC ; Lindén & Rajala 1981)
52	(NERC ; Pulliainen 1981)
54	(Willig 2016)
56	(Friggens 2008)
57	(LTER)
58	(Waide 2010b, a)
59	(Ernest <i>et al.</i> 2009)
63	(Moore 1991b; Moore 1991a)
67	(CWAC)
68	(Management)
70	(Vanholder 1997)
78	(Zettler 2005)
81	(Robinson <i>et al.</i> 2007; Robinson 2010)
84	(Milchakova 2011)
85	(Addinck & de Kluijver 2003)
92	(Naumov)
97	(Markhaseva <i>et al.</i> 1985)
100	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
101	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
108	(Woehler)
110	(Ostler)
112	(NIWA)
113	(Pugh)
117	(NODC 2011)
119	(Clark & Branton 2007)
121	(CRED 2011)
125	(Reichert 2009)
147	(SWP)
152	(CMarZ)
162	(EPA)
163	(NPRB)
166	(PIROP ; Brown 1975; Diamond <i>et al.</i> 1986; Huettmann 1998; Halpin <i>et al.</i> 2009; Read <i>et al.</i> 2009)

169	(Yen <i>et al.</i> 2004; Jahncke 2006; Rintoul <i>et al.</i> 2006; Yen <i>et al.</i> 2006)
171	(BMMROOS)
172	(POPA ; Machete & Santos 2007; Amorim <i>et al.</i> 2008; Morato <i>et al.</i> 2008)
173	(JAMSTEC)
176	(Kennedy & Spry 2011)
178	(Boutillier 2007)
180	(ECNA)
182	(Wade 2011)
183	(Tremblay & Branton 2007)
187	(NCCOS 2003)
191	(NEFSC 2010)
192	(AADC)
193	(Henderson ; Henderson & Magurran 2014b; Henderson & Magurran 2014a)
195	(USGS)
196	(Moore & Howson)
197	(DATRAS 2013b)
198	(DATRAS 2013a)
200	(NEFSC 2010)
204	(Degraer <i>et al.</i> 2006)
206	(DATRAS 2010d)
208	(DATRAS 2010b)
209	(DATRAS 2010a)
210	(DATRAS 2010c)
213	(NMFS 2005)
214	(Harmon 2012)
215	(HMANA)
217	(USFS)
219	(NatureCounts 2012b)
220	(NatureCounts 2012b)
221	(Viereck <i>et al.</i> 2005)
225	(Shochat <i>et al.</i> 2004)
229	(McLarney <i>et al.</i> 2010)
232	(Williams)
236	(Gido)
237	(Lathrop 2000)
238	(LTER 2011)
240	(Muldavin)
243	(Day <i>et al.</i> 2004; Day 2010)
244	(NatureCounts 2012a)
246	(Chen <i>et al.</i> 2015)
247	(Rudstam 2008)
248	(Shi <i>et al.</i> 2015)
249	(Thomsen <i>et al.</i> 2016)
252	(Reichert 2010)
253	(LTER)
254	(LTER)
256	(DATRAS 2010a)
271	(Reed 2014a)
272	(Reed 2014b)
273	(Reed 2014c)
278	(NIWA)
285	(REVIZEE)

288	(Clark & Branton 2007)
290	(CMAR)
291	(CMAR)
292	(CMAR)
297	(Carpenter 2015)
300	(Landis & Gage 2014)
301	(Jonas & Joern 2007; Joern 2016)
305	(Wiley)
306	(Brook <i>et al.</i> 2006; NERC 2010)
308	(Merritt 1999)
310	(Sal <i>et al.</i> 2013)
311	(Kaufman)
313	(Knops & Tilman)
316	(Lightfoot)
318	(SANParks 2011)
319	(Wilgers <i>et al.</i> 2006)
321	(Lightfoot 2007)
327	(Kelt <i>et al.</i> 2013)
329	(Scott <i>et al.</i>)
330	(Davies <i>et al.</i> 2014)
332	(Grossman 2007)
333	(Sandercock)
335	(Grossman <i>et al.</i> 1982)
336	(Ernest <i>et al.</i> 2009)
337	(Vermont Center for Ecostudies <i>et al.</i> 2015)
339	(Svensson <i>et al.</i> 2010)
340	(Lightfoot 2011)
350	(Billett <i>et al.</i> 2001; Billett <i>et al.</i> 2010)
351	(Ruhl & Smith 2004; Kuhnz <i>et al.</i> 2014)
354	(Davies <i>et al.</i> 2016)
356	(Bradford <i>et al.</i> 2014)
357	(Stapp 2013)
358	(Dickson <i>et al.</i> 1993)
359	(Reed 2014)
361	(Hall 1984)
363	(Enemar <i>et al.</i> 2004)
366	(Lightfoot 2013)
374	(Monitoring Site 1000 Project <i>et al.</i> 2013)
375	(Monitoring Site 1000 Project <i>et al.</i> 2014)
378	(Benedetti-Cecchi)
379	(Benedetti-Cecchi)
382	(Krefting & Ahlgren 1974; NERC 2010)
413	(Kendeigh 1982)
414	(Kendeigh 1982)
416	(Kendeigh 1982)
418	(Sweatman <i>et al.</i> 2001)
419	(Fraser 2014)
420	(Svensson 2006)
427	(Pomati <i>et al.</i> 2012; Pomati <i>et al.</i> 2015)
428	(Stenseth <i>et al.</i> 1999; Olsen <i>et al.</i> 2009; Rogers <i>et al.</i> 2011; Barceló <i>et al.</i> 2016)
430	(NIWA)
431	(NIWA)

432	(NIWA)
433	(Budy <i>et al.</i> ; Budy <i>et al.</i>)
439	(Zakharov 1998)
440	(Zakharov 1998)
441	(Zakharov 1998)
466	(Neat & Campbell 2011)
468	(Kenner <i>et al.</i> 2013)
469	(Kushner <i>et al.</i> 2013)
473	(Anderson <i>et al.</i> 2011)
475	(Hogstad 1993)
477	(Douglass <i>et al.</i> 2010; Lefcheck 2015)
478	(Wagner <i>et al.</i> 2011)
499	(Institute of Agricultural and Fisheries research (ILVO) 2016a)
500	(Institute of Agricultural and Fisheries research (ILVO) 2016b)
505	(Edelist <i>et al.</i> 2013)

References

1.
AADC. "Whale Catches in Southern Ocean". OBIS - Australian Antarctic Data Centre. Available at: <http://www.iobis.org>, accessed 2013.
2.
Addinck, W. & de Kluijver, M. (2003). North Sea observations of Crustacea, Polychaeta, Echinodermata, Mollusca and some other groups between 1986 and 2003. Expert Centre for Taxonomic Idenditification (ETI), the Netherlands. Available at: <http://www.emodnet-biology.eu/data-catalog?module=dataset&dasid=1037>, accessed 2012.
3.
Amorim, P., Figueiredo, M., Machete, M., Morato, T., Martins, A. & Serrão Santos, R. (2008). Spatial variability of seabird distribution associated with environmental factors: a case study of marine Important Bird Areas in the Azores. *ICES Journal of Marine Science*, 66, 29-40.
4.
Anderson, J., Vermeire, L. & Adler, P.B. (2011). Fourteen years of mapped, permanent quadrats in a northern mixed prairie, USA. *Ecology*, 92, 1703-1703.
5.
Barceló, C., Ciannelli, L., Olsen, E.M., Johannessen, T. & Knutsen, H. (2016). Eight decades of sampling reveal a contemporary novel fish assemblage in coastal nursery habitats. *Global Change Biology*, 22, 1155-1167.
6.
Benedetti-Cecchi, L. "Calafuria Mid-shore Intertidal Dataset (1991-2014)". Department of Biology, University of Pisa. Accessed 2016.
- 7.

- Beven, G. (1976). Changes in breeding bird populations of an oak-wood on Bookham Common, Surrey, over twenty-seven years. *London Naturalist*, 55, 23-42.
- 8.
- Billett, D., Bett, B., Reid, W., Boorman, B. & Priede, I. (2010). Long-term change in the abyssal NE Atlantic: The 'Amperima Event'revisited. *Deep Sea Research Part II: Topical Studies in Oceanography*, 57, 1406-1417.
- 9.
- Billett, D., Bett, B., Rice, A., Thurston, M., Galéron, J., Sibuet, M. et al. (2001). Long-term change in the megabenthos of the Porcupine Abyssal Plain (NE Atlantic). *Progress in Oceanography*, 50, 325-348.
- 10.
- BMMROOS. "Bahamas Marine Mammal Research Organisation Opportunistic Sightings - OBIS SEAMAP". Available at: <http://www.iobis.org>, accessed 2012.
- 11.
- Boutillier, J.A. (2007). "Pacific Shrimp Trawl Survey". In: ShrimpTrawl Bio database, Fisheries and Oceans Canada PBS Shellfish Data Unit. OBIS Canada Digital Collections, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada. Available at: <http://www.iobis.org>, accessed 2012.
- 12.
- Bradford, M.G., Murphy, H.T., Ford, A.J., Hogan, D.L. & Metcalfe, D.J. (2014). Long-term stem inventory data from tropical rain forest plots in Australia. *Ecology*, 95, 2362-2362.
- 13.
- Brook, B.W., Traill, L.W. & Bradshaw, C.J. (2006). Minimum viable population sizes and global extinction risk are unrelated. *Ecology letters*, 9, 375-382.
- 14.
- Brown, R.G. (1975). Atlas of eastern Canadian seabirds.
- 15.
- Budy, P., Luecke, C. & O'Brien, J. "Zooplankton density for lake samples collected near Toolik Lake Arctic LTER in the summer from 1983 to 1992". Arctic Long-Term Ecological Research. Available at: <http://arc-lter.ecosystems.mbl.edu/1983-1992arclterzoops>, accessed 2016.
- 16.
- Budy, P., Luecke, C. & O'Brien, J. "Zooplankton density for lake samples collected near Toolik Lake Arctic LTER in the summers between 1993-2002". Arctic Long-Term Ecological Research. Available at: <http://arc-lter.ecosystems.mbl.edu/1993-2002arcticlterzoops>, accessed 2016.
- 17.
- Carpenter, R. (2015). "MCR LTER: Coral Reef: Long-term Population and Community Dynamics: Other Benthic Invertebrates, ongoing since 2005". Moorea Coral Reef LTER, knb-lter-mcr.7.28. Available at: doi:10.6073/pasta/8e7b3a0c7a8bf315739921861cc79d10, accessed 2016.
- 18.
- Chen, H., Liao, Y.-C., Chen, C.-Y., Tsai, J.-I., Chen, L.-S. & Shao, K.-T. (2015). Long-term monitoring dataset of fish assemblages impinged at nuclear power plants in northern Taiwan. *Scientific data*, 2, 150071.
- 19.

- Clark, D. & Branton, B. (2007). DFO Maritimes Research Vessel Trawl Surveys, OBIS Canada Digital Collections. *Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada, OBIS Canada*.
- 20.
- CMAR. "CSIRO Marine Data Warehouse - OBIS Australia", CSIRO Division of Marine and Atmospheric Research (CMAR), Australia. Available at <http://www.iobis.org>, accessed 2012.
- 21.
- CMarZ. "CMarZ (Census of Marine Zooplankton)-Asia Database". Accessed through OBIS- SCAR-MarBIN., Available at: <http://www.iobis.org/mapper/?dataset=1500>, accessed 2012.
- 22.
- CRED (2011). "Towed-Diver Fish Biomass Surveys in the Pacific Ocean 2000-2010". Coral Reef Ecosystem Division (CRED), Pacific Island Fisheries Sciences Center, National Marine Fisheries Service. Available at: <http://www.iobis.org/mapper/?dataset=1581>, accessed 2012.
- 23.
- CWAC. "Coordinated Waterbird Counts (CWAC) - AfrOBIS". Available at: <http://www.iobis.org/mapper/?dataset=603>, accessed 2012.
- 24.
- DATRAS (2010a). "Fish trawl survey: ICES Beam Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2761>, accessed 2013.
- 25.
- DATRAS (2010b). "Fish trawl survey: ICES French Southern Atlantic Bottom Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2759>, accessed 2013.
- 26.
- DATRAS (2010c). "Fish trawl survey: ICES North Sea International Bottom Trawl Survey for commercial fish species. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2763>, accessed 2013.
- 27.
- DATRAS (2010d). "Fish trawl survey: Northern Irish Ground Fish Trawl Survey. ICES Database of trawl surveys (DATRAS)." The International Council for the Exploration of the Sea, Copenhagen. Available at: <http://www.emodnet-biology.eu/data-catalog?%3Fmodule=dataset&dasid=2764>, accessed 2013.
- 28.
- DATRAS (2013a). ICES Baltic International Trawl Survey For Commercial Fish Species (1991 - 2013). Available at <https://datras.ices.dk>, accessed 2013.
- 29.
- DATRAS (2013b). ICES Scottish West Coast Survey For Commercial Fish Species 1985-2013. Available at <https://datras.ices.dk>, accessed 2014.
- 30.

- Davies, C.H., Armstrong, A.J., Baird, M., Coman, F., Edgar, S., Gaughan, D. *et al.* (2014). Over 75 years of zooplankton data from Australia. *Ecology*, 95, 3229-3229.
- 31.
- Davies, C.H., Coughlan, A., Hallegraeff, G., Ajani, P., Armbrecht, L., Atkins, N. *et al.* (2016). A database of marine phytoplankton abundance, biomass and species composition in Australian waters. *Scientific data*, 3.
- 32.
- Day, F. (2010). "Long-term N-fertilized vegetation plots on Hog Island, Virginia Coastal Barrier Islands, 1992-2014." Virginia Coast Reserve Long-Term Ecological Research Project. Available at: <http://www.vcrler.virginia.edu/cgi-bin/showDataset.cgi?docid=knb-lter-vcr.106>, accessed 2013.
- 33.
- Day, F.P., Conn, C., Crawford, E. & Stevenson, M. (2004). Long-term effects of nitrogen fertilization on plant community structure on a coastal barrier island dune chronosequence. *Journal of Coastal Research*, 722-730.
- 34.
- Degraer, S., Wittoeck, J., Appeltans, W., Cooreman, K., Deprez, T., Hillewaert, H. *et al.* (2006). "Macrobel: Long term trends in the macrobenthos of the Belgian Continental Shelf." Oostende, Belgium. Available at: <http://www.emodnet-biology.eu/catalog?module=dataset&dasid=145>, accessed 2013.
- 35.
- Diamond, A., Gaston, A. & Brown, R. (1986). Converting PIROP Counts of Seabirds at Sea to Absolute Densities. Progress Notes No 164. *Canadian Wildlife Service, Ottawa*.
- 36.
- Dickson, J.G., Conner, R.N. & Williamson, J.H. (1993). Neotropical migratory bird communities in a developing pine plantation. *Proceedings on the Annual Conference. SEAFWA*, 47, 439-446.
- 37.
- Douglass, J.G., France, K.E., Richardson, J.P. & Duffy, J.E. (2010). Seasonal and interannual change in a Chesapeake Bay eelgrass community: Insights into biotic and abiotic control of community structure. *Limnology and Oceanography*, 55, 1499-1520.
- 38.
- ECNA. "East Coast North America Strategic Assessment Project, Groundfish Atlas for the East Coast of North America". Available at: <http://www.iobis.org>, accessed 2012.
- 39.
- Edelist, D., Rilov, G., Golani, D., Carlton, J.T. & Spanier, E. (2013). Restructuring the Sea: profound shifts in the world's most invaded marine ecosystem. *Diversity and Distributions*, 19, 69-77.
- 40.
- Enemar, A., Sjöstrand, B., Andersson, G. & von Proschwitz, T. (2004). The 37-year dynamics of a subalpine passerine bird community, with special emphasis on the influence of environmental temperature and *Epirrita autumnata* cycles. *Ornis Svecica*, 14, 63-106.
- 41.
- EPA. "EPA'S EMAP Database". U.S. Environmental Protection Agency through its Environmental Monitoring and Assessment Program (EMAP). Available at: <http://www.iobis.org>, accessed

2012. Also available at <https://archive.epa.gov/emap/archive-emap/web/html/index-2021.html>

42.

Ernest, S., Valone, T.J. & Brown, J.H. (2009). Long-term monitoring and experimental manipulation of a Chihuahuan Desert ecosystem near Portal, Arizona, USA. *Ecology*, 90, 1708-1708.

43.

Fraser, W. (2014). "At-sea seabird censuses. Data on the species encountered (including marine mammals), their abundance, distribution and behavior. Data collected aboard cruises off the coast of the Western Antarctic Peninsula, 1993 - present". Palmer Station Antarctica LTER. Available at: <http://dx.doi.org/10.6073/pasta/e3871e749fa737dd94d5a269ac90e8ce>, accessed 2016.

44.

Friggins, M. (2008). "Sevilleta LTER Small Mammal Population Data", Albuquerque, NM: Sevilleta Long Term Ecological Research Site Database: SEV008. Available at: <http://sev.lternet.edu/data/sev-8>, accessed 2012.

45.

Gaston, K.J. & Blackburn, T.M. (2000). Pattern and process in macroecology. Wiley-Blackwell Oxford, England.

46.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993). *The new atlas of breeding birds in Britain and Ireland: 1988-1991*. T & AD Poyser London.

47.

Gido, K.B. "Fish population on selected watersheds at Konza Prairie - CFP01." Konza Prairie LTER Program. Available at: <http://www.konza.ksu.edu/KNZ/pages/data/Knzdsdetail.aspx?datasetCode=CFP01>, accessed 2012.

48.

Grossman, G.D. (2007). "Stream fish assemblage stability in a southern Appalachian stream at the Ceweeta Hydrologic Laboratory from 1984 to 1995". Ceweeta Long Term Ecological Research Program. Available at: <http://dx.doi.org/10.6073/pasta/f0baf5f59c89f670e04f537f5cc05290>, accessed 2016.

49.

Grossman, G.D., Moyle, P.B. & Whitaker Jr, J.O. (1982). Stochasticity in structural and functional characteristics of an Indiana stream fish assemblage: a test of community theory. *American Naturalist*, 423-454.

50.

Hall, G.A. (1984). A long-term bird population study in an Appalachian spruce forest. *The Wilson Bulletin*, 228-240.

51.

Halpern, C.B. & Dyrness, C. (2010). "Plant succession and biomass dynamics following logging and burning in the Andrews Experimental Forest Watersheds 1 and 3, 1962-Present". Long-Term Ecological Research. Forest Science Data Bank, Corvallis. Available at: <http://andrewsforest.oregonstate.edu/data/abstract.cfm?dbcode=TP073>, accessed 2012.

52.

- Halpern, C.B. & Lutz, J.A. (2013a). Canopy closure exerts weak controls on understory dynamics: a 30-year study of overstory–understory interactions. *Ecological Monographs*, 83, 221-237.
- 53.
- Halpern, C.B. & Lutz, J.A. (2013b). "Canopy closure exerts weak controls on understory dynamics: a 30-year study of overstory–understory interactions". Available at: Dryad DigitalRepository, doi:10.5061/dryad.1q88j, accessed 2013.
- 54.
- Halpin, P.N., Read, A.J., Fujioka, E., Best, B.D., Donnelly, B., Hazen, L.J. et al. (2009). OBIS-SEAMAP: The world data center for marine mammal, sea bird, and sea turtle distributions. *Oceanography*, 22, 104-115.
- 55.
- Harmon, M.F., J. (2012). "Long-term growth, mortality and regeneration of trees in permanent vegetation plots in the Pacific Northwest, 1910 to present." Long-Term Ecological Research. Forest Science Data Bank, Corvallis. Available at: <http://andrewsforest.oregonstate.edu/data/abstract.cfm?dbcode=TV010>, accessed 2012.
- 56.
- Henderson, P.A. The long-term study of the fish and crustacean community of the Bristol Channel. Available at <http://www.pisces-conservation.com/>, accessed 2013.
- 57.
- Henderson, P.A. & Magurran, A.E. (2014a). Data from: Direct evidence that density-dependent regulation underpins the temporal stability of abundant species in a diverse animal community. Dryad Data Repository.
- 58.
- Henderson, P.A. & Magurran, A.E. (2014b). Direct evidence that density-dependent regulation underpins the temporal stability of abundant species in a diverse animal community. *Proceedings of the Royal Society B: Biological Sciences*, 281.
- 59.
- HMANA. "Hawk Migration Association of North America (HMANA)". Available at: <http://www.hmana.org/>, accessed 2012.
- 60.
- Hogstad, O. (1993). Structure and dynamics of a passerine bird community in a spruce-dominated boreal forest. A 12-year study. In: *Annales Zoologici Fennici*. JSTOR, pp. 43-54.
- 61.
- Holmes, R. & Sturges, F. (1975). Bird community dynamics and energetics in a northern hardwoods ecosystem. *The Journal of Animal Ecology*, 175-200.
- 62.
- Holmes, R.T. & Sherry, T.W. (1988). Assessing population trends of New Hampshire forest birds: local vs. regional patterns. *The Auk*, 756-768.
- 63.
- Holmes, R.T. & Sherry, T.W. (2001). Thirty-year bird population trends in an unfragmented temperate deciduous forest: importance of habitat change. *The Auk*, 118, 589-609.
- 64.

- Holmes, R.T., Sherry, T.W. & Sturges, F.W. (1986). Bird Community Dynamics in a Temperate Deciduous Forest: Long-Term Trends at Hubbard Brook. *Ecological Monographs*, 56, 201-220.
- 65.
- Huettmann, F. (1998). An ecological GIS research application for the northern Atlantic-The PIROP database software, environmental data sets and the role of the internet. *Riekert/Tochtermann*.
- 66.
- Institute of Agricultural and Fisheries research (ILVO), B. (2016a). Macrofauna monitoring at long-term monitoring stations in the Belgian part of the North Sea between 1979 and 1999. Available at: <http://dx.doi.org/10.14284/201>, accessed 2016.
- 67.
- Institute of Agricultural and Fisheries research (ILVO), B. (2016b). Macrofauna monitoring at long-term monitoring stations in the Belgian part of the North Sea from 2001 on. Available at: <http://dx.doi.org/10.14284/202>, accessed 2016.
- 68.
- Jahncke, J.R., C. (2006). "CalCOFI and NMFS Seabird and Marine Mammal Observation Data, 1987-2006". California Cooperative Oceanic Fisheries Investigations (CalCOFI) and National Marine Fisheries Service (NMFS) cruises, 1987-2006 - OBIS SEAMAP. Available at: <http://www.iobis.org>, accessed 2012.
- 69.
- JAMSTEC. "Marine Biological Sample Database, JAMSTEC" (OBIS_JAPAN). Available at: <http://www.iobis.org/mapper/?dataset=2289>, accessed 2012.
- 70.
- Joern, A. (2016). CGR02 Sweep Sampling of Grasshoppers on Konza Prairie LTER watersheds (1982-present). Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/7060b2c244229a37e3bfc8c18f14ad02>, accessed 2016.
- 71.
- Jonas, J.L. & Joern, A. (2007). Grasshopper (Orthoptera: Acrididae) communities respond to fire, bison grazing and weather in North American tallgrass prairie: a long-term study. *Oecologia*, 153, 699-711.
- 72.
- Kaufman, D.W. Seasonal summary of numbers of small mammals on 14 LTER traplines in prairie habitats at Konza Prairie. Konza Prairie Long-Term Ecological Research. . Available at: <http://lter.konza.ksu.edu/content/csm01-seasonal-summary-numbers-small-mammals-14-lter-traplines-prairie-habitats-konza>, accessed 2016.
- 73.
- Kelt, D., Meserve, P., Gutiérrez, J., Milstead, W.B. & Previtali, M. (2013). Long-term monitoring of mammals in the face of biotic and abiotic influences at a semiarid site in north-central Chile. *Ecology*, 94, 977-977.
- 74.
- Kendeigh, S.C. (1982). Bird populations in east central Illinois: Fluctuations, variations, and development over a half-century. *University of Illinois Press*.
- 75.

- Kennedy, M. & Spry, J. (2011). Atlantic Zone Monitoring Program Maritimes Region plankton datasets. *Fisheries and Oceans Canada-BioChem archive. OBIS Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada.*
- 76.
- Kenner, M.C., Estes, J.A., Tinker, M.T., Bodkin, J.L., Cowen, R.K., Harrold, C. et al. (2013). A multi-decade time series of kelp forest community structure at San Nicolas Island, California (USA). *Ecology*, 94, 2654-2654.
- 77.
- Knops, J. & Tilman, D. Successional Dynamics on a Resampled Chronosequence - Experiment 014. Cedar Creek Ecosystem Science Reserve. Available at <http://www.cedarcreek.umn.edu/research/data/dataset?ghe014>, accessed 2016.
- 78.
- Krefting, L.W. & Ahlgren, C.E. (1974). Small Mammals and Vegetation Changes After Fire in a Mixed Conifer-Hardwood Forest. *Ecology*, 55, 1391-1398.
- 79.
- Kuhnz, L.A., Ruhl, H.A., Huffard, C.L. & Smith, K.L. (2014). Rapid changes and long-term cycles in the benthic megafaunal community observed over 24 years in the abyssal northeast Pacific. *Progress in Oceanography*, 124, 1-11.
- 80.
- Kushner, D.J., Rassweiler, A., McLaughlin, J.P. & Lafferty, K.D. (2013). A multi-decade time series of kelp forest community structure at the California Channel Islands. *Ecology*, 94, 2655-2655.
- 81.
- Lack, P. (2010). *The atlas of wintering birds in Britain and Ireland*. A&C Black.
- 82.
- Landis, D. & Gage, S. (2014). Insect Populations via Sticky Traps at KBS-LTER. Available at: <http://lter.kbs.msu.edu/datatables/67>, accessed 2016.
- 83.
- Lathrop, R. (2000). "Madison Wisconsin Lakes Zooplankton 1976 - 1994." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <http://lter.limnology.wisc.edu/dataset/madison-wisconsin-lakes-zooplankton-1976-1994>, accessed 2013.
- 84.
- Lefcheck, J.S. (2015). The use of functional traits to elucidate the causes and consequences of biological diversity. The College of William & Mary, PhD thesis. Available at: <http://gradworks.umi.com/36/62/3662989.html>, accessed 2016.
- 85.
- Lightfoot, D. "Small Mammal Exclosure Study (SMES)". Sevilleta Long Term Ecological Research Program. Available at: <http://sev.lternet.edu/content/small-mammal-exclosure-study-smes-0>, accessed 2016.
- 86.
- Lightfoot, D. (2007). "Jornada Grasshopper Data". Jornada Basin LTER. Available at: <http://jornada.nmsu.edu/lter/dataset/49712/view>, accessed 2016.
- 87.

- Lightfoot, D. (2011). "Small Mammal Exclosure Study (SMES) Vegetation Data from the Chihuahuan Desert Grassland and Shrubland at the Sevilleta National Wildlife Refuge, New Mexico (2006-2009)". Long Term Ecological Research Network. Available at: <http://dx.doi.org/10.6073/pasta/d80d5e2196cd11ef79df23ebe5a77c19>, accessed 2016.
- 88.
- Lightfoot, D. (2013). "Lizard pitfall trap data (LTER-II, LTER-III)". Jornada Basin LTER. Available at: <http://jornada.nmsu.edu/lter/dataset/49821/view>, accessed 2016.
- 89.
- Lindén, H. & Rajala, P. (1981). Fluctuations and long-term trends in the relative densities of tetraonid populations in Finland, 1964-77. *Finnish Game Research*, 39, 13-34.
- 90.
- LTER, N. "North Temperate Lakes LTER: Phytoplankton - Madison Lakes Area 1995 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-phytoplankton-madison-lakes-area-1995-current>, accessed 2013.
- 91.
- LTER, N. "North Temperate Lakes LTER: Zooplankton - Trout Lake Area 1982 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-zooplankton-trout-lake-area-1982-current>, accessed 2013.
- 92.
- LTER, N. "NTLFI02 North Temperate Lakes LTER: Fish Abundance 1981 - current". North Temperate Lakes Long Term Ecological Research program, NSF. Center for Limnology, University of Wisconsin-Madison. Available at: <https://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-fish-abundance-1981-current>, accessed 2012.
- 93.
- LTER, N. (2011). "North Temperate Lakes LTER: Zooplankton - Madison Lakes Area 1997 - current." North Temperate Lakes Long Term Ecological Research Program, Center for Limnology, University of Wisconsin-Madison. Available at: <http://lter.limnology.wisc.edu/dataset/north-temperate-lakes-lter-zooplankton-madison-lakes-area-1997-current>, accessed 2013.
- 94.
- Machete, M. & Santos, R. (2007). Azores Fisheries Observer Program (POPA): a case study of the multidisciplinary use of observer data. In: *Proceedings of the 5th International Fisheries Observer Conference*, pp. 15-18.
- 95.
- Management, M.a.C. "Marine and Coastal Management - Copepod Surveys - AfrOBIS". Available at: <http://www.iobis.org/>, accessed 2012.
- 96.
- Markhaseva, E.L., Golikov, A.A., Agapova, T.A. & Beig, A.A. (1985). Archives of the Arctic Seas Zooplankton., Available at: <http://www.iobis.org/mapper/?dataset=4470>, accessed 2012.
- 97.

- McLarney, W.O., Meador, J. & Chamblee, J. (2010). "Upper Little Tennessee River Biomonitoring Program Database." Ceweeta Long Term Ecological Research Program. Available at: https://cweeta.uga.edu/dbpublic/dataset_details.asp?accession=4045, accessed 2012.
- 98.
- Merritt, J. (1999). Long Term Mammal Data from Powdermill Biological Station 1979-1999. Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/83c888854e239a79597999895bb61cf>, accessed 2016.
- 99.
- Milchakova, N.A., Ryabogina, V. G. & Chernyshova, E. B. (2011). "Macroalgae of the Crimean coastal zone (the Black Sea, 1967-2007)". Sevastopol, IBSS. Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&dasid=2690>, accessed 2012.
- 100.
- Monitoring Site 1000 Project, Biodiversity Center & Japan, M.o.E.o. (2013). "Monitoring site 1000 Shorebird Survey" (ShorebirdsDatapackage2012.zip, downloaded from <http://www.biodic.go.jp/moni1000/> findings/data/index.html), Accessed 2016.
- 101.
- Monitoring Site 1000 Project, Biodiversity Center & Japan, M.o.E.o. (2014). "Monitoring site 1000 Forest and grassland research - Surface wandering beetles survey data". (GBDataPackage2014ver1.zip, downloaded from <http://www.biodic.go.jp/moni1000/findings/data/index.html>), Accessed 2016.
- 102.
- Moore, J.J. & Howson, C.M. "Survey of the rocky shores in the region of Sullom Voe, Shetland, A report to SOTEAG from Aquatic Survey & Monitoring Ltd", Cosheston, Pembrokeshire. 29 p. Available at: <http://www.soteag.org.uk>, accessed 2013.
- 103.
- Moore, N.W. (1991a). "The development of dragonfly communities and the consequences of territorial behaviour: A 27-year study on small ponds at Woodwalton Fen, Cambridgeshire, United Kingdom". NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database Version 2.0. Available at: <http://www3.imperial.ac.uk/cpb/databases/gpdd>, accessed 2012.
- 104.
- Moore, N.W. (1991b). The development of dragonfly communities and the consequences of territorial behaviour: a 27 year study on small ponds at Woodwalton Fen, Cambridgeshire, United Kingdom. *Odonatologica*, 20, 203-231.
- 105.
- Morato, T., Varkey, D.A., Damaso, C., Machete, M., Santos, M., Prieto, R. et al. (2008). Evidence of a seamount effect on aggregating visitors. *Marine Ecology Progress Series*, 357, 23-32.
- 106.
- Muldavin, E. "Pinon-Juniper (Core Site) Quadrat Data for the Net Primary Production Study at the Sevilleta National Wildlife Refuge, New Mexico (2003-Present)." Sevilleta Long Term Ecological Research Program., Available at: <http://sev.lternet.edu/node/1718>, accessed 2013.
- 107.

NatureCounts (2012a). Bird Studies Canada "BC Coastal Waterbird Survey (2004)." NatureCounts, a node of the Avian Knowledge Network. Available at: <http://www.birdscanada.org/birdmon/>, accessed 2012.

108.

NatureCounts (2012b). Bird Studies Canada "Marsh Monitoring Program." NatureCounts, a node of the Avian Knowledge Network. Available at: <http://www.birdscanada.org/birdmon/>, accessed 2012.

109.

Naumov, A. "Benthos of the White Sea. A database". White Sea Biological Station, Zoological Institute RAS. Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&dasid=2769>, accessed 2012.

110.

NCCOS (2003). "National Benthic Infaunal Database (NBID)". NOAA/NOS/NCCOS/CCEHBR/Coastal Ecology Program. NOAA's Ocean Service, National Centers for Coastal Ocean Science (NCCOS). Available at: <https://data.noaa.gov/dataset/national-benthic-infaunal-database-nbid>, accessed 2012.

111.

Neat, F. & Campbell, N. (2011). Demersal fish diversity of the isolated Rockall plateau compared with the adjacent west coast shelf of Scotland. *Biological Journal of the Linnean Society*, 104, 138-147.

112.

NEFSC (2010). "Benthic Database (OBIS-USA)", Northeast Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Department of Commerce., Available at: <http://www.iobis.org/mapper/?dataset=1694>, accessed 2012.

113.

NERC. "Fluctuations and long-term trends in the relative densities of tetraonid populations in Finland, 1964-77." NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database v2.0. . Available at: <https://www.imperial.ac.uk/cpb/gpdd2/secure/register.aspx>, accessed 2012.

114.

NERC. "A transect survey of small land carnivore and red fox populations on a subarctic fell in Finnish Forest Lapland over 13 winters". NERC Centre for Population Biology, Imperial College, The Global Population Dynamics Database v2.0., Available at: <http://www3.imperial.ac.uk/cpb/databases/gpdd>, accessed 2012.

115.

NERC (2010). "The Global Population Dynamics Database Version 2". Centre for Population Biology, Imperial College. Available at: <http://www.sw.ic.ac.uk/cpb/cpb/gpdd.html>, accessed 2016.

116.

NIWA. "The New Zealand Freshwater Fish Database". Available at: <https://www.niwa.co.nz/our-services/online-services/freshwater-fish-database>, accessed 2016.

117.

NIWA. "South Western Pacific Regional OBIS Data Asteroid Subset", NIWA (National Institute of Water and Atmospheric Research - New Zealand) MBIS (Marine Biodata Information System)

- accessed through South Western Pacific OBIS., Available at:
<http://www.iobis.org/mapper/?dataset=219>, accessed 2012.
- 118.
- NIWA. "South Western Pacific Regional OBIS Data provider for the NIWA Marine Biodata Information System". Ocean Biogeographic Information System. Occurrence Dataset., Available at:
<https://doi.org/10.15468/zuiyu>, accessed 2012.
- 119.
- NMFS (2005). "Northeast Fisheries Science Center Bottom Trawl Survey Data (OBIS-USA)." NOAA's National Marine Fisheries Service (NMFS) Northeast Fisheries Science Center. Woods Hole, Massachusetts, USA., Available at: <http://www.iobis.org/mapper/?dataset=1435>, accessed 2013.
- 120.
- NODC (2011). "South TX Outer Continental Shelf and MI, AL, and FL Outer Continental Shelf benthic organism sampling 1974-1978". US National Oceanographic Data Center, Silver Spring, Maryland, USA. Available at http://www.usgs.gov/obis-usa/data_search_and_access/participants.html, accessed 2012.
- 121.
- NPRB. "The Observer Program database", accessed through the OBIS-USA North Pacific Groundfish Observer (North Pacific Research Board). Available at: <http://www.iobis.org>, accessed 2012.
- 122.
- Olsen, E.M., Carlson, S.M., Gjøsæter, J. & Stenseth, N.C. (2009). Nine decades of decreasing phenotypic variability in Atlantic cod. *Ecology Letters*, 12, 622-631.
- 123.
- Ostler, R. Marine Nature Conservation Review (MNCR) and associated benthic marine data held and managed by JNCC-EurOBIS. *Joint Nature Conservation Committee, Centre for Ecology and hydrology, Aberdeenshire, UK.*(Available at <http://www.emodnet-biology.eu/datalatalog>.
- 124.
- PIROP. "PIROP Northwest Atlantic 1965-1992 - OBIS SEAMAP". Available at:
<http://www.iobis.org/mapper/?dataset=2245>, accessed 2012.
- 125.
- Pomati, F., Matthews, B., Jokela, J., Schildknecht, A. & Ibelings, B.W. (2012). Effects of re-oligotrophication and climate warming on plankton richness and community stability in a deep mesotrophic lake. *Oikos*, 121, 1317-1327.
- 126.
- Pomati, F., Tellenbach, C., Matthews, B., Venail, P., Ibelings, B.W. & Ptacnik, R. (2015). Challenges and prospects for interpreting long-term phytoplankton diversity changes in Lake Zurich (Switzerland). *Freshwater Biology*, 60, 1052-1059.
- 127.
- POPA. "cetacean, seabird, and sea turtle sightings in the Azores area 1998-2009 - OBIS SEAMAP" . Available at: <http://www.iobis.org/mapper/?dataset=4257>, accessed 2012.
- 128.

- Pugh, P. Discovery Collections Midwater Database. National Oceanography Centre, Southampton, UK., Available at: <http://www.iobis.org/mapper/?dataset=12>, accessed 2012.
- 129.
- Pulliainen, E. (1981). A transect survey of small land carnivore and red fox populations on a subarctic fell in Finnish Forest Lapland over 13 winters. *Annales Zoologici Fennici*, 270-278.
- 130.
- Read, A., Halpin, P., Crowder, L., Best, B. & Fujioka, E. (2009). OBIS-SEAMAP: mapping marine mammals, birds and turtles. *World Wide Web electronic publication*. <http://seamap.env.duke.edu>. Accessed, 15.
- 131.
- Reed, D.C. (2014). "SBC LTER: Reef: Kelp Forest Community Dynamics: Fish abundance". Santa Barbara Coastal LTER. Available at:
doi:10.6073/pasta/e37ed29111b2fddffc08355252b8b8c7, accessed 2016.
- 132.
- Reed, D.C. (2014a). "SBC LTER: Reef: Kelp forest community dynamics: Abundance and size of giant kelp (*Macrocystis pyrifera*), ongoing since 2000". Santa Barbara Coastal LTER. Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.18>
doi:10.6073/pasta/d90872297e30026b263a119d4f5bca9f, accessed 2016.
- 133.
- Reed, D.C. (2014b). "SBC LTER: Reef: Kelp forest community dynamics: Fish abundance". Santa Barbara Coastal LTER. Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.17>
doi:10.6073/pasta/e37ed29111b2fddffc08355252b8b8c7, accessed 2016.
- 134.
- Reed, D.C. (2014c). "SBC LTER: Reef: Kelp forest community dynamics: Invertebrate and algal density". Santa Barbara Coastal LTER. . Available at: <http://sbc.lternet.edu/cgi-bin/showDataset.cgi?docid=knb-lter-sbc.19>
doi:10.6073/pasta/cd4cf864efecd69891dfe1d73b9ac9c3, accessed 2016.
- 135.
- Reichert, M. (2009). "MARMAP Chevron Trap Survey 1990-2009". SCDNR/NOAA MARMAP Program, SCDNR MARMAP Aggregate Data Surveys, The Marine Resources Monitoring, Assessment, and Prediction (MARMAP) Program, Marine Resources Research Institute, South Carolina Department of Natural Resources U.S.A., Available at: http://www.usgs.gov/obis-usa/data_search_and_access/participants.html, accessed 2012.
- 136.
- Reichert, M. (2010). "MARMAP Blackfish Trap Survey 1990-2009". SCDNR/NOAA MARMAP Program. SCDNR MARMAP Aggregate Data Surveys. The Marine Resources Monitoring. Assessment. and Prediction (MARMAP) Program. Marine Resources Research Institute. South Carolina Department of Natural Resources USA. Available at: <http://www.usgs.gov/obis-usa/>, accessed 2013.
- 137.
- REVIZEE. "Previous_fisheries_REVIZEE_Program", accessed through Tropical and Subtropical Western South Pacific OBIS., Available at: <http://www.iobis.org/mapper/?dataset=2411>, accessed 2012.
- 138.

Rintoul, C., Langabeer-Schlagenhauf, B., Hyrenbach, K., Morgan, K. & Sydeman, W. (2006). Atlas of California Current marine birds and mammals: Version 1. *Unpublished Report, PRBO Conservation Science, Petaluma, CA.*

139.

Robinson, K.P. (2010). "CRRU (Cetacean Research and Rescue Unit) Cetacean sightings in Scotland waters". Available at: <http://www.emodnet-biology.eu/component/imis/?module=dataset&dasid=2819>, accessed 2012.

140.

Robinson, K.P., Baumgartner, N., Eisfeld, S.M., Clark, N.M., Culloch, R.M., Haskins, G.N. et al. (2007). The summer distribution and occurrence of cetaceans in the coastal waters of the outer southern Moray Firth in northeast Scotland (UK). *Lutra*, 50, 19.

141.

Rogers, L.A., Stige, L.C., Olsen, E.M., Knutsen, H., Chan, K.-S. & Stenseth, N.C. (2011). Climate and population density drive changes in cod body size throughout a century on the Norwegian coast. *Proceedings of the National Academy of Sciences*, 108, 1961-1966.

142.

Rudstam, L. (2008). "Zooplankton survey of Oneida Lake, New York, 1964 – 2012", KNB Data Repository. Available at: <https://knb.ecoinformatics.org/#view/kgordon.17.56>, accessed 2016.

143.

Ruhl, H.A. & Smith, K.L. (2004). Shifts in deep-sea community structure linked to climate and food supply. *Science*, 305, 513-515.

144.

Sal, S., López-Urrutia, Á., Irigoien, X., Harbour, D.S. & Harris, R.P. (2013). Marine microplankton diversity database. *Ecology*, 94, 1658-1658.

145.

Sandercock, B.K. "Variable distance line-transect sampling of bird population numbers in different habitats on Konza Prairie (1981 - 2009)". Konza Prairie Long Term Ecological Research Program. Available at: <http://www.konza.ksu.edu/knz/pages/data/KnzEntity.aspx?id=CBP011>, accessed 2016.

146.

SANParks (2011). "Karoo National Park Census Data. 1994 - 2009". Available at: <http://datadryad.org/handle/10255/dryad.13079?show=full>, accessed 2016.

147.

Scott, D., Metts, B. & Lance, S. "The Rainbow Bay Long-term Study". Available at: <http://srelherp.uga.edu/projects/rbay.htm>, accessed 2016.

148.

Shi, Z., Sherry, R., Xu, X., Hararuk, O., Souza, L., Jiang, L. et al. (2015). Evidence for long-term shift in plant community composition under decadal experimental warming. *Journal of Ecology*, 103, 1131-1140.

149.

Shochat, E., Katti, M. & Warren, P. (2004). "Point count bird censusing: long-term monitoring of bird distribution and diversity in central Arizona-Phoenix: period 2000 to 2011". Central Arizona-

- Phoenix Long-Term Ecological Research. Global Institute for Sustainability, Arizona State University. Available at: <https://caplter.asu.edu/data/data-catalog/?id=46>, accessed 2012.
- 150.
- Standley, P., Bucknell, N., Swash, A. & Collins, I. (1996). The Birds of Berkshire. Berkshire Atlas Group. *Reading, UK.*
- 151.
- Stapp, P. (2013). SGS-LTER Long-Term Monitoring Project: Small Mammals on Trapping Webs on the Central Plains Experimental Range, Nunn, Colorado, USA 1994 -2006, ARS Study Number 118. Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/2e311b4e40fea38e573890f473807ba9>, accessed 2017.
- 152.
- Stenseth, N.C., Bjørnstad, O.N., Falck, W., Fromentin, J.M., Gjøsæter, J. & Gray, J.S. (1999). Dynamics of coastal cod populations: intra- and intercohort density dependence and stochastic processes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266, 1645-1654.
- 153.
- Stone, B., Sears, J., Cranswick, P., Gregory, R., Gibbons, D., Rehfisch, M. et al. (1997). Population estimates of birds in Britain and in the United Kingdom. *British Birds*, 90, 1-22.
- 154.
- Svensson, S. (2006). Species composition and population fluctuations of alpine bird communities during 38 years in the Scandinavian mountain range. *Ornis Svecica*, 16, 183-210.
- 155.
- Svensson, S., Thorner, A. & Nyholm, N. (2010). Species trends, turnover and composition of a woodland bird community in southern Sweden during a period of fifty-seven years. *Ornis Svecica*, 20, 31-44.
- 156.
- Sweatman, H.P.A., Cheal, A.J., Coleman, G.J., Delean, S., Fitzpatrick, B.M., Miller, I.R. et al. (2001). Long-Term Monitoring of the Great Barrier Reef. Status Report Number 5. Available at: <http://www.iobis.org/mapper/?dataset=233>, accessed 2017.
- 157.
- SWP. "South Western Pacific Regional OBIS Data Bryozoan Subset". Accessed through South Western Pacific OBIS., Available at: <http://www.iobis.org/mapper/?dataset=221>, accessed 2012.
- 158.
- Thomsen, P.F., Jørgensen, P.S., Bruun, H.H., Pedersen, J., Riis-Nielsen, T., Jonko, K. et al. (2016). Resource specialists lead local insect community turnover associated with temperature – analysis of an 18-year full-seasonal record of moths and beetles. *Journal of Animal Ecology*, 85, 251-261.
- 159.
- Tremblay, J.M. & Branton, B. (2007). DFO Maritimes Research Vessel Trawl Surveys, OBIS Canada Digital Collections. *Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada, OBIS Canada.*
- 160.

- USFS. "Landbird Monitoring Program (UMT-LBMP)." US Forest Service. Available at: <http://www.avianknowledge.net/>, accessed 2012.
- 161.
- USGS. Patuxent Wildlife Research Center "North American Breeding Bird Survey" ftp data set, version 2014.0. Available at: <ftp://ftpext.usgs.gov/pub/er/md/laurel/BBS/DataFiles/>, accessed 2013.
- 162.
- Vanholder, B. (1997). "Belgian Migrating Lepidoptera". NERC Centre for Population Biology, Imperial College. The Global Population Dynamics Database v2.0. . Available at: <https://www.imperial.ac.uk/cpb/gpdd2/secure/register.aspx>, accessed 2012.
- 163.
- Vermont Center for Ecostudies, Lambert, J.D. & Hart, J. (2015). "Mountain Birdwatch 1.0". KNB Data Repository, 10.5063/F1DN430G. accessed 2016.
- 164.
- Vickery, W.L. & Nudds, T.D. (1984). Detection of Density-Dependent Effects in Annual Duck Censuses. *Ecology*, 65, 96-104.
- 165.
- Viereck, L.A., Van Cleve, K., Chapin, F.S., Ruess, R.W. & Hollingsworth, T.N. (2005). Vegetation Plots of the Bonanza Creek LTER Control Plots: Species Count (1975 - 2004). Environmental Data Initiative., Available at: <http://dx.doi.org/10.6073/pasta/8dd0e1ac48e2f82b51adabfb3c62ae2>, accessed 2012.
- 166.
- Wade, E. (2011). Snow crab research trawl survey database (Southern Gulf of St. Lawrence, Gulf region, Canada) from 1988 to 2010. *OBIS Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada*.
- 167.
- Wagner, R., Marxsen, J., Zwick, P. & Cox, E.J. (2011). *Central European Stream Ecosystems: The Long Term Study of the Breitenbach*. John Wiley & Sons.
- 168.
- Waide, R.B. (2010a). "Bird abundance - point counts. El Verde Field Station, Puerto Rico: Luquillo Long Term Ecological Research Site Database: Data Set 23". Available at: <http://luq.lter.net.edu/data/luqmetadata23>, accessed 2012.
- 169.
- Waide, R.B. (2010b). Bird abundance - point counts. Long Term Ecological Research Network. Available at: <http://dx.doi.org/10.6073/pasta/0d96957379936a038ebbbcc6135b2fab>, accessed 2012.
- 170.
- Widdicombe, C., Eloire, D., Harbour, D., Harris, R. & Somerfield, P. (2010a). Long-term phytoplankton community dynamics in the Western English Channel. *Journal of Plankton Research*, fbp127.
- 171.
- Widdicombe, C.E., Harbour, D. & Harris, R. (2010b). Time series of phytoplankton abundance and composition at station L4 in the English Channel from 1988 to 2009. *PANGAEA*, 754335, 0-0.
- 172.

- Wiley, R.H. "Population estimates of Appalachian salamanders". Ceweeta LTER. Available at:
<http://ceweeta.uga.edu/eml/1044.xml>, accessed 2016.
- 173.
- Wilgers, D.J., Horne, E.A., Sandercock, B.K. & Volkmann, A.W. (2006). Effects of rangeland management on community dynamics of the herpetofauna of the tallgrass prairie. *Herpetologica*, 62, 378-388.
- 174.
- Williams, D. "Pelagic Fish Observations 1968-1999." Australian Antarctic Data Centre. Available at:
<http://www.gbif.org/dataset/85b0a82a-f762-11e1-a439-00145eb45e9a>, accessed 2012.
- 175.
- Williamson, M. (1983). The land-bird community of Skokholm: ordination and turnover. *Oikos*, 378-384.
- 176.
- Williamson, M. (1987). Are communities ever stable? In: *Symposium of the British Ecological Society*. 177.
- Willig, M.R.B., C. P. (2016). "El Verde Grid long-term invertebrate data: Luquillo Long Term Ecological Research Site Database: Data Set 107". Available at:<http://luq.lternet.edu/data/luqmetadata107/7427> accessed 2016.
- 178.
- Woehler, E. "Seabirds of the Southern and South Indian Ocean - Australian Antarctic Data Centre". Available at: <http://www.iobis.org>, accessed 2012.
- 179.
- Yen, P., Sydeman, W., Bograd, S. & Hyrenbach, K. (2006). Spring-time distributions of migratory marine birds in the southern California Current: Oceanic eddy associations and coastal habitat hotspots over 17 years. *Deep Sea Research Part II: Topical Studies in Oceanography*, 53, 399-418.
- 180.
- Yen, P.P., Sydeman, W.J. & Hyrenbach, K.D. (2004). Marine bird and cetacean associations with bathymetric habitats and shallow-water topographies: implications for trophic transfer and conservation. *Journal of Marine Systems*, 50, 79-99.
- 181.
- Zachmann, L., Moffet, C. & Adler, P. (2010). Mapped quadrats in sagebrush steppe: long-term data for analyzing demographic rates and plant-plant interactions. *Ecology*, 91, 3427-3427.
- 182.
- Zakharov, V.D. (1998). Biodiversity of bird population of terrestrial habitats in Southern Ural. Miass: IGZ. *Ural Branch of Russian Academy of Sciences*, 158 p.
- 183.
- Zettler, M.L. (2005). Macrozoobenthos baltic sea (1980-2005) as part of the IOW-Monitoring. Institut für Ostseeforschung Warnemünde, Germany., Available at:
<http://www.iobis.org/mapper/?dataset=2289>, accessed 2012.