

1
2
3
4
5
6
7
8
9

Adaptation of sperm whales to open-boat whalers: rapid social learning on a large scale?

Hal Whitehead^{1*}, Tim D. Smith², and Luke Rendell³

¹Department of Biology, Dalhousie University, Halifax, Nova Scotia, B3H 4R2, Canada.

²World Whaling History, Redding, California, USA.

³Sea Mammal Research Unit and Centre for Social Learning and Cognitive Evolution, School of Biology, University of St Andrews, Fife KY16 8LB, UK.

*Correspondence to: hwhitehe@dal.ca.

10 **Abstract:**

11 Animals can mitigate human threats, but how do they do this, and how fast can they adapt?
12 Hunting sperm whales was a major 19th Century industry. Analysis of data from digitized
13 logbooks of American whalers in the North Pacific found that the rate at which whalers
14 succeeded in harpooning (striking) sighted whales fell by about 58% over the first few years of
15 exploitation in a region. This decline cannot be explained by the earliest whalers being more
16 competent, as their strike rates outside the North Pacific, where whaling had a longer history,
17 were not elevated. The initial killing of particularly vulnerable individuals would not have
18 produced the observed rapid decline in strike rate. It appears that whales swiftly learned effective
19 defensive behaviour. Sperm whales live in kin-based social units. Our models show that social
20 learning, in which naïve social units, when confronted by whalers, learned defensive measures
21 from grouped social units with experience, could lead to the documented rapid decline in strike
22 rate. This rapid, large-scale adoption of new behaviour enlarges our concept of the spatio-
23 temporal dynamics of non-human culture.

24
25 **Keywords:** social learning, culture, sperm whale, whaling, defensive measures

26
27 **1. Introduction**

28 Social learning can change the behaviour of whole populations, sometimes over large spatial
29 scales [1–3]. However, beyond humans and beyond dialect evolution [4] there is little evidence
30 for large-scale cultural change in behaviour that is rapid (over less than about half a generation).

31 In the context of a predator-prey interaction that was particularly significant for economic
32 development [5] and marine ecology [6], we investigated a hypothesis that the prey—sperm
33 whales (*Physeter macrocephalus*)—quickly learned effective defensive measures from one
34 another when faced with a novel predator—the open-boat whaler. In the 18th and 19th Centuries,
35 pelagic whalers from Europe and North America spread around the world, exploiting new
36 “grounds” and new species [5,7]. They searched for whales from sail-powered whaleships, with
37 sightings of groups of sperm whales typically interspersed by days or more of searching. Once
38 whales were sighted, whalers rowed whaleboats towards the whales, aiming to “strike” them
39 with hand-thrown harpoons, a process typically lasting tens of minutes to hours (Fig. 1). If
40 whales were killed, they would be towed back to the whaleship for processing (“trying out”).

41 Some historians have suggested that the success rate of open-boat whalers in harpooning sighted
42 whales—the second stage in these whaling operations—dropped substantially during the initial
43 years of industrial exploitation, and that this was due to socially-learned changes in whale
44 behaviour [8,9]. While quantitative data for the earliest years of any new exploitation of wildlife
45 are usually few or absent, the development of pelagic whaling operations in the North Pacific by
46 American whalers during the middle 19th Century is well chronicled in digitized whalers’
47 logbooks [10].

48 We used these data to quantify the success of whalers in striking (i.e. harpooning) sighted sperm
49 whales during the early years of exploitation. We assessed whether this decline was caused by
50 socially-learned changes in the defensive behaviour of the whales [8,9] (hypothesis HX),
51 evaluating support for several alternative hypotheses using causal models: the first whalers on a
52 ground were particularly proficient (H1); the whalers initially captured especially vulnerable

53 whales (H2); or the whales learned to escape whalers from their own individual experience of
54 encountering them (H3).

56 2. Methods

57 Detailed methods, together with justifications and a consideration of assumptions, as well as raw
58 data and computer code for analysis, are given in the Supplementary Material. Data come from a
59 compilation and digitization of logbooks of American whalers working between 10-50°N in the
60 North Pacific by Maury [11] and the Census of Marine Life [10], giving for each voyage-day (a
61 logged day of searching for whales by a particular vessel; e.g. “Ship-Alexander_06-May-1822”)
62 noon position, whether sperm whales were sighted, and how many whales were struck
63 (harpooned) or tried (processed). For this study we needed to determine the date of first
64 exploitation in an area with reasonable accuracy and this was only possible for the North Pacific,
65 with other oceans being exploited earlier, or largely by non-American whalers [5,7,12,13].

66 We delineated areas by calculating, for each voyage-day d on which sperm whales were sighted,
67 the time lag since the first sighting within $h=1,000$ km (based on displacement of photo-identified
68 female sperm whales over lags of years in Southeast Pacific [14]). Then, for each time lag since
69 first sighting in the region, T years, we calculated the mean number of whales struck per sighting
70 day, the “strike rate”, $y(T)$: for voyage-days with time lag T since the first sighting in the region,
71 the total number of sperm whales struck divided by the number of voyage-days with sperm
72 whale sightings. To model the observed decrease in strike rate with time lag since first
73 exploitation and estimate the magnitude and temporal scale of the initial decline, we fit a
74 heuristic model of an initial exponential decline in strike rate to a plateau. We also fit causal
75 models corresponding to hypotheses H2, H3 and HX. Equations are in Table 1, and detailed
76 derivations are in the Supplementary Material.

77 Under H2, we assume that the population, of size $P(0)$ at the commencement of whaling, has a
78 proportion q of vulnerable individuals which are struck at a rate f_1 when the group containing the
79 individual is sighted, while the less vulnerable animals are struck at a rate f_2 . If s is the rate of
80 groups (or individuals) being sighted by whalers per voyage-day, then the number of vulnerable
81 animals in the population t days after the commencement of exploitation is $N_1(t)=P(0).q.(1-s.f_1)^t$,
82 and the number of less vulnerable individuals $N_2(t)=P(0).(1-q).(1-s.f_2)^t$, leading to equation H2
83 in Table 1.

84 In order to assess learning at the individual or within-unit level as a driver of an initial decline in
85 strike rate (H3), we estimated the proportion of whales that had experience of whalers at
86 different numbers of years after the initiation of whaling, $Z(T)$. $Z(T)$ increases at a rate of $s(1-$
87 $Z(T))$ per day. Integrating over T gives $Z(T)=1-e^{-s.365.T}$, leading to equation H3 in Table 1.

88 Studies of living animals in the Pacific Ocean have found groups to be made up of on average
89 two kin-based social units of females and their young, that generally remain grouped for around
90 7-14 days [15]. To model a rapid form of social learning (HX), we assumed that a group being
91 approached by whalers would act like an experienced group if any of the units that made it up
92 had previous experience with whalers. Then, the probability that at least one unit in the group
93 had experience of whaling becomes $Z'(T)=1-e^{-(g/u).s.365.T}$, where g is the group size and u the unit
94 size. This leads to equation HX in Table 1.

95 We estimated $g=21.75$, $u=10.5$ from literature and $s=0.000313$ sightings.group⁻¹.day⁻¹ using
96 population estimates (see Detailed Methods in Supplementary Material). We assessed the fit of
97 each model using adjusted R², and the relative support of the causal models using the Akaike
98 information criterion [16]. To evaluate the results' robustness, we systematically changed g , u ,
99 and s to outer plausible values, varied h , and alternatively used whaler-defined grounds as
100 regions (Supplementary Material).

102 3. Results

103 The data set included 77,749 operational voyage-days, with sperm whale sightings on 2,405
104 voyage-days. The heuristic model indicated a 58% drop in strike rates over time lag scales of
105 2.4yr after which they stabilized (Fig. 2; Table 1).

106 To assess hypothesis H1, whaler competency, we used the full global data set. When outside the
107 North Pacific, whalers that entered a North Pacific region 0-4 years after the initial sighting had
108 similar rates of trying out whales per sighting day (0.80, SE=0.08) as those entering the North
109 Pacific 5-9 years after the initial sighting (0.77, SE=0.09; strike rates were not available outside
110 the North Pacific, so we used try rates). That the earliest whalers in the North Pacific and those
111 that followed later had similar success when in other oceans, where the sperm whales had
112 typically been exploited for longer, argues against decreasing whaler competence driving the fall
113 in strike rate.

114 Under hypothesis H2, differential vulnerability, an initial decline in strike rates will occur if a
115 population is stratified by vulnerability to whaling and the most vulnerable animals are taken
116 first. These vulnerable individuals might include the young, the old, the sick, the foolhardy, the
117 belligerent, or mothers attempting to protect calves. Our model of H2 produces a decline in strike
118 rate, but it is not as steep as indicated by the real data (Fig. 2), fitting considerably worse than the
119 social learning model (HX) (Table 1). Furthermore, the parameters of the best-fitting model
120 (Table 1) indicated that all vulnerable individuals (2.2% of population) would have been struck
121 when first sighted, while non-vulnerable individuals were struck at a rate of 0.016 per sighting.
122 This extreme contrast in vulnerability seems unrealistic. Alternative formulations of the
123 vulnerable individual model that we can imagine (e.g. more than 2 categories of vulnerability,
124 continuous measure of vulnerability) would tend to reduce the rate of initial decline in strike rate.
125 Thus, a stratification of the population by vulnerability to whalers is not supported as the sole
126 cause of the initial decline in strike rate.

127 The final two potential causes of the fall in strike rate that we considered (H3, HX) assume that
128 whales, individually or socially, learn behaviour that is adaptive for avoiding being struck. A
129 model in which strike rates fell following the first experience of a social unit to whalers (H3)
130 produces virtually the same decline as the vulnerable individual model (H2; Fig. 2). However, in
131 common with the vulnerable individual model, the decline in strike rate due to within-unit
132 learning seems insufficient (with a poorer fit than HX). Realistic modifications to the within-unit
133 learning model that we can imagine, such as imperfect learning from the first encounter with
134 whalers, will tend to reduce the rate of initial decline in strike rate. Thus, learning as individuals
135 or within social units is not supported as the sole cause of the initial decline in strike rate. In
136 contrast the between-unit social learning model (HX) fits the data better than all other causal
137 models (Fig. 2; Table 1), producing a rapid decline in strike rate as units learn defensive
138 measures from one another.

139 In the Supplementary Material we examine the effects of changing our definition of “region” and
140 of variation in the parameters of the models within plausible limits. In all scenarios the between-
141 unit social learning model fit better than other causal models. With a few exceptions, considered
142 below in the discussion of the between-unit learning hypothesis, violations of the assumptions of
143 the causal models would tend to decrease the steepness of the initial decline in strike rates, and
144 thus the fit of the model to the data.

146 4. Discussion

147 While a combination of H1-H3 might produce a steep decline in strike rate, social learning of
148 defensive measures between social units (HX) is the best-supported explanation for the rapid
149 decline in strike rate following the first sperm whale sighting within a region. The whalers
150 themselves wrote of defensive methods that they believed the whales were adopting, including
151 communicating danger within the social group, fleeing—especially upwind—or attacking the
152 whalers [17,18] (Fig. 1). Deep dives would also have been effective. But, perhaps the most
153 straightforward change would be for sperm whales to cease their characteristic defensive
154 behaviour against their most serious previous predator, the killer whale, *Orcinus orca*. Gathering
155 in slow-moving groups at the surface and fighting back with jaws or flukes often works against
156 killer whales [19,20], but will have only assisted the relatively slow-moving, surface-limited,
157 harpoon-bearing open-boat whalers.

158 There are other behavioural changes that the whales may have made in response to whaling, but
159 their impact on strike rates is less clear. There is some evidence that sperm whales formed larger
160 groups in response to whaling [15], but this would likely have increased rather than decreased
161 strike rates. They may have learned to avoid the whalers before the whalers detected them, but
162 this should generally have reduced the mean detection range of the whalers and so increased the
163 strike rate. However, if whales fleeing at long ranges made themselves more visible by blowing
164 hard and showing their bodies forcefully, so increasing the number of sightings with groups that
165 were not easily struck, this might have additionally decreased the strike rate.

166 Thus, there are learned behavioural changes that the sperm whales could have made to reduce
167 strike rates, and some anecdotal witness that they did so. However, learning as individuals or
168 within social units is not supported as the sole cause of the initial decline in strike rate. To
169 achieve the observed reduction in strike rate through behavioural change, some mechanism must
170 have allowed naïve whales without experience of whalers to receive the benefits of experience.

171 We suggest that naïve social units learned defensive measures from grouped experienced social
172 units and adopted them. Encounters with whalers typically lasted hours, and sperm whales
173 through their echolocation and communication systems can probably sense and coordinate
174 behaviour over ranges of several km. Other processes could have enhanced the social learning
175 process. If groups were particularly likely to split between or within social units after experience
176 with whalers, and then to join other units, this will have increased the probability that a naïve
177 animal was grouped with an experienced individual during its first encounter with whalers.

178 Our analysis provides substantial support for rapid (<20% generation time) social learning over
179 large spatial scales. The ability of sperm whales, or potentially other species, to rapidly change
180 behaviour in the face of a new anthropogenic threat by making use of social learning has

181 implications for the population significance of new threats, and their assessment. Data from the
182 earliest exposures may not generalize to later periods, and vice versa.

183
184 **Acknowledgments:** Thanks to Trevor Kenchington, David Lusseau, Matthew Zippel and three
185 anonymous reviewers for useful suggestions.

186 **Funding:** no funding was obtained for this work.

187 **Author contributions:** HW conceived the study and carried out the analyses; TDS curated the
188 data; TDS and LR provided input to the analyses; all authors wrote and revised the manuscript.

189 190 **References**

- 191
192 1. Fisher J, Hinde RA. 1949 The opening of milk bottles by birds. *Br. Birds* **42**, 347–357.
- 193 2. Allen J, Weinrich M, Hoppitt W, Rendell L. 2013 Network-based diffusion analysis
194 reveals cultural transmission of lobtail feeding in humpback whales. *Science* **340**, 485–
195 488.
- 196 3. Jesmer BR *et al.* 2018 Is ungulate migration culturally transmitted? Evidence of social
197 learning from translocated animals. *Science* **361**, 1023–1025.
198 (doi:10.1126/science.aat0985 [doi])
- 199 4. Garland EC, Goldizen AW, Rekdahl ML, Constantine R, Garrigue C, Hauser ND, Poole
200 MM, Robbins J, Noad MJ. 2011 Dynamic horizontal cultural transmission of humpback
201 whale song at the ocean basin scale. *Curr. Biol.* **21**, 687–691.
202 (doi:10.1016/j.cub.2011.03.019)
- 203 5. Starbuck A. 1878 *History of the American whale fishery from its earliest inception to the*
204 *year 1876*. Washington, DC: Government Printing Office.
- 205 6. Lavery TJ, Roudnew B, Gill P, Seymour J, Seuront L, Johnson G, Mitchell JG, Smetacek
206 V. 2010 Iron defecation by sperm whales stimulates carbon export in the Southern Ocean.
207 *Proc. R. Soc. B Biol. Sci.* **277**, 3527–3531.
- 208 7. Jackson G. 2017 *The British whaling trade*. Oxford University Press.
- 209 8. Demuth B. 2019 *Floating coast: an environmental history of the Bering Strait*. New York:
210 W.W. Norton & Company.
- 211 9. Jones RT. 2019 A whale of a difference: southern right whale culture and the Tasman
212 world’s living terrain of encounter. *Environ. Hist. Camb.* **25**, 185–218.
- 213 10. Smith TD, Reeves RR, Josephson EA, Lund JN. 2012 Spatial and seasonal distribution of
214 American whaling and whales in the age of sail. *PLoS One* **7**, e34905.
- 215 11. Maury MF. 1851 *Whale chart*. Washington.
- 216 12. Colnett J. 1798 *A voyage to the South Atlantic and round Cape Horn into the Pacific*
217 *Ocean, for the purpose of extending the spermaceti whale fisheries*. London: W. Bennett.
- 218 13. Wray P, Martin KR. 1983 Historical whaling records from the western Indian Ocean.

- 219 *Reports Int. Whal. Comm. (Special Issue)* **5**, 218–242.
- 220 14. Whitehead H, Coakes A, Jaquet N, Lusseau S. 2008 Movements of sperm whales in the
221 tropical Pacific. *Mar. Ecol. Prog. Ser.* **361**, 291–300.
- 222 15. Whitehead H, Antunes R, Gero S, Wong SNP, Engelhaupt D, Rendell L. 2012 Multilevel
223 societies of female sperm whales (*Physeter macrocephalus*) in the Atlantic and Pacific:
224 why are they so different? *Int. J. Primatol.* **33**, 1142–1164.
- 225 16. Burnham KP, Anderson DR. 2002 *Model selection and multimodel inference: a practical*
226 *information-theoretic approach*. New York: Springer-Verlag.
- 227 17. Philbrick N. 2000 *In the heart of the sea: the tragedy of the whaleship Essex*. New York:
228 Penguin.
- 229 18. Caldwell DK, Caldwell MC, Rice DW. 1966 Behavior of the sperm whale *Physeter*
230 *catodon* L. In *Whales, dolphins and porpoises* (ed KS Norris), pp. 677–717. Berkeley:
231 University of California Press.
- 232 19. Pitman RL, Ballance LT, Mesnick SIL, Chivers SJ, Balance LT, Mesnick SIL, Chivers SJ.
233 2001 Killer whale predation on sperm whales: Observations and implications. *Mar.*
234 *Mammal Sci.* **17**, 494–507.
- 235 20. Arnbohm T, Papastavrou V, Weilgart LS, Whitehead H. 1987 Sperm whales react to an
236 attack by killer whales. *J. Mammal.* **68**, 450–453.

237

238




240
241

242
243

244 **Fig. 1.** Sperm whaling in the North Pacific from the ship *Canton* (by Oliver Wilcox, whaler;
245 New Bedford Whaling Museum).

246
247



248
249
250
251
252
253
254

Fig. 2. A: Sperm whale sightings with estimated date of first sighting in region (scale of 1,000 km). **B:** Strike rate with time lag since first sighting in region (number of voyage-days above each lag; standard errors from Poisson approximation may be biased narrow due to dependencies), with fitted models (Table 1).

255
256
257
258
259

Table 1. Models fit to strike rates (y strikes/sighting day) with time lag (T years) after first sighting in region (s = rate of groups being sighted by whalers per day; g = mean size of group; u = mean size of social unit; see Supplemental Material for derivation of models, and parameter estimates).

Model	AIC	Adjusted R ²	Parameter estimates (SE):		
			Strike rate on vulnerable/naïve animals (f_1)	Strike rate on less vulnerable/experienced animals (f_2)	Proportion of vulnerable animals (q)
<i>Heuristic:</i>					
Exponential					
	$y = a_1 + a_2 \cdot e^{-a_3 T}$	-17.55	0.42	-	-
<i>Causal:</i>					
Vulnerable individuals (H2)					
	$y = g \frac{f_1 q (1 - s f_1)^{365 \cdot T} + f_2 (1 - q) (1 - s f_2)^{365 \cdot T}}{q (1 - s f_1)^{365 \cdot T} + (1 - q) (1 - s f_2)^{365 \cdot T}}$	-9.95	0.28	1.000 (0.000)	0.016 (0.003)
Within-unit learning (H3)					
	$y = g [f_1 e^{-s \cdot 365 \cdot T} + f_2 (1 - e^{-s \cdot 365 \cdot T})]$	-12.14	0.28	0.038 (0.005)	0.016 (0.003)
Between-unit learning (HX)					
	$y = g \left[f_1 e^{-\left(\frac{g}{u}\right) \cdot s \cdot 365 \cdot T} + f_2 \left(1 - e^{-\left(\frac{g}{u}\right) \cdot s \cdot 365 \cdot T} \right) \right]$	-16.71	0.40	0.044 (0.005)	0.019 (0.002)

260