

# Spatiotemporal Variability of Dimethylsulphoniopropionate on a Fringing Coral Reef: The Role of Reefal Carbonate Chemistry and Environmental Variability

Heidi L. Burdett<sup>1\*</sup>, Penelope J. C. Donohue<sup>1</sup>, Angela D. Hatton<sup>2</sup>, Magdy A. Alwany<sup>3</sup>, Nicholas A. Kamenos<sup>1</sup>

**1** School of Geographical and Earth Sciences, University of Glasgow, Glasgow, United Kingdom, **2** Scottish Association for Marine Science, Oban, Argyll, United Kingdom, **3** Department of Marine Science, Suez Canal University, Ismailia, Egypt

## Abstract

Oceanic pH is projected to decrease by up to 0.5 units by 2100 (a process known as ocean acidification, OA), reducing the calcium carbonate saturation state of the oceans. The coastal ocean is expected to experience periods of even lower carbonate saturation state because of the inherent natural variability of coastal habitats. Thus, in order to accurately project the impact of OA on the coastal ocean, we must first understand its natural variability. The production of dimethylsulphoniopropionate (DMSP) by marine algae and the release of DMSP's breakdown product dimethylsulphide (DMS) are often related to environmental stress. This study investigated the spatiotemporal response of tropical macroalgae (*Padina* sp., *Amphiroa* sp. and *Turbinaria* sp.) and the overlying water column to natural changes in reefal carbonate chemistry. We compared macroalgal intracellular DMSP and water column DMSP+DMS concentrations between the environmentally stable reef crest and environmentally variable reef flat of the fringing Suleman Reef, Egypt, over 45-hour sampling periods. Similar diel patterns were observed throughout: maximum intracellular DMSP and water column DMS/P concentrations were observed at night, coinciding with the time of lowest carbonate saturation state. Spatially, water column DMS/P concentrations were highest over areas dominated by seagrass and macroalgae (dissolved DMS/P) and phytoplankton (particulate DMS/P) rather than corals. This research suggests that macroalgae may use DMSP to maintain metabolic function during periods of low carbonate saturation state. In the reef system, seagrass and macroalgae may be more important benthic producers of dissolved DMS/P than corals. An increase in DMS/P concentrations during periods of low carbonate saturation state may become ecologically important in the future under an OA regime, impacting larval settlement and increasing atmospheric emissions of DMS.

**Citation:** Burdett HL, Donohue PJC, Hatton AD, Alwany MA, Kamenos NA (2013) Spatiotemporal Variability of Dimethylsulphoniopropionate on a Fringing Coral Reef: The Role of Reefal Carbonate Chemistry and Environmental Variability. PLoS ONE 8(5): e64651. doi:10.1371/journal.pone.0064651

**Editor:** Tilmann Harder, University of New South Wales, Australia

**Received:** February 7, 2013; **Accepted:** April 17, 2013; **Published:** May 28, 2013

**Copyright:** © 2013 Burdett et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This research was financially supported by a NERC studentship (NE/H525303/1) to HB, a University of Glasgow Mobility Scholarship to HB and PD and a Royal Society of Edinburgh/Scottish Government Fellowship (RSE 48701/1) and Carnegie Trust for the Universities of Scotland grant to NK. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: h.burdett.1@research.gla.ac.uk

## Introduction

The carbonate chemistry of the oceans is regulated by a carbonate equilibrium that is driven by the dissolution of atmospheric CO<sub>2</sub> into the oceans. Since the Industrial Revolution, atmospheric CO<sub>2</sub> concentrations have increased from 280 ppm to ~390 ppm; CO<sub>2</sub> concentrations have been increasing by ~2 ppm yr<sup>-1</sup> since 2000 [1]. It is projected that continued anthropogenic emissions of CO<sub>2</sub> will cause the pH of the oceans to drop by 0.3–0.5 units by 2100 [2], a process known as ocean acidification (OA). This rate of change has been estimated to be ~100 times faster than during glacial terminations [3], raising concerns over the future survival of calcifying organisms.

Coastal habitats such as coral reefs are characterised by more extreme natural variations in carbonate saturation state compared to the open ocean, reflecting diurnal and seasonal cycles driven by biological and physical processes [3]. Thus, in order to accurately project the impact of an OA scenario on coastal ecosystems, one

must first understand contemporary natural variability. Spatial heterogeneity of carbonate chemistry has been observed in coral reef systems, with reef flats and lagoons more variable than fringing forereefs due to biological (benthos composition) and physical (wave action and residence time) variables [4,5,6,7,8]. Similarly, surface pCO<sub>2</sub> in atoll and barrier reef lagoons may be higher than offshore waters [9]. Where the benthos is dominated by filamentous turf algae, pH may be high (i.e. less acidic) [7]. Short-term and longer-term temporal heterogeneity in reefal carbonate chemistry has also been observed. During the day, net calcification may be observed, whilst net dissolution may be observed at night [8], influencing the carbonate chemistry of the overlying water column. Seasonal patterns in carbonate chemistry have been observed [10] but this may be de-coupled from reef calcification due to the maintenance of threshold saturation levels during the winter [11].

The response of calcified organisms to reduced carbonate saturation state is varied depending on their carbonate structure,

their method of calcification and their ability to benefit from an increase in CO<sub>2</sub> (e.g. for photosynthesis) [12]. This has led to the suggestion that, under an OA scenario, coral reefs may become dominated by macroalgae (fleshy and coralline) [13]. The negative effect of low carbonate saturation state on the calcification and growth of red coralline algae (which deposit high-magnesium calcite, the most thermodynamically unstable carbonate polymorph) has been well reported. During the winter, net calcification in the temperate/polar coralline alga *Lithothamnion glaciale* was significantly reduced at high CO<sub>2</sub> (750, 950 and 1500 ppm), whilst only the very high CO<sub>2</sub> treatment (1500 ppm) induced a decrease in calcification in the summer [14]. Growth rates and structural integrity of *L. glaciale* may also be reduced under high CO<sub>2</sub> (589–1080  $\mu$ atm) [15,16]. After a 1 year exposure to high CO<sub>2</sub> conditions (700 ppm), net dissolution exceeded net calcification in *Lithophyllum cabiochae* [17], and when combined with high temperature (+3°C) more algal necroses, death and dissolution were observed [17]. Spore production, growth and recruitment of the Corallinaceae are also inhibited by high CO<sub>2</sub> conditions (550–760 ppm) [18,19,20]. Additionally, high CO<sub>2</sub> (1000 ppm) and UV radiation (particularly UVB) may act synergistically to inhibit growth, photosynthetic O<sub>2</sub> evolution and calcification in the geniculate coralline alga *Corallina sessilis* [21].

In the field, calcifying organisms may decrease in abundance under high, variable  $p$ CO<sub>2</sub> conditions [22]. Despite this, *Padina* spp. (Ochrophyta: Dictyoaceae), one of only two known calcifying brown algae, appear to thrive in CO<sub>2</sub> vent systems, albeit with reduced calcification in the more acidified areas [23], attributed to the lower abundance of grazing sea urchins in acidified areas and enhanced photosynthesis from higher CO<sub>2</sub> availability [23]. Experimental reef studies (conducted in 2650 m<sup>3</sup> mesocosms dominated by macroalgae) suggest that although calcification appears to decline under high CO<sub>2</sub> conditions, net organic production does not change [24,25]. However, current studies are not wholly conclusive and more detailed investigations into the biochemical and morphological effects of low carbonate saturation state on calcifying benthic macroalgae are still required.

Dimethylsulphoniopropionate (DMSP) is a sulphur compound produced by many marine algae and is the major precursor to dimethylsulphide (DMS), a gas that may be linked to local climate regulation through aerosol production and cloud formation [26,27]. A number of cellular functions have been described for DMSP in macro- and microalgae, including as a compatible solute [28], a cryoprotectant [29], an antioxidant [30] and a herbivore deterrent [31] and attractant [32]. A general reduction in intracellular DMSP concentrations with increasing latitude has been suggested for macroalgae in the northern hemisphere [31], perhaps in response to the cryoprotective properties of DMSP [29]. However, this suggestion was based primarily on Chlorophyta species, the abundance of which also increases with increasing latitude. In contrast, other macroalgal secondary metabolites (e.g. terpenes) tend to increase in low latitudes, perhaps due to increased grazing pressure [33]. The principal functions of DMSP have not been extensively studied and the lack of data available on macroalgal DMSP concentrations currently prevents such assessments to be made.

Information on the effect of reduced carbonate saturation state on intracellular DMSP concentrations in algae is limited and has provided variable results. Intracellular DMSP concentrations in the green macroalgae *Ulva lactuca* and *U. clathrata* was not affected by elevated  $p$ CO<sub>2</sub> conditions (up to 1514  $\mu$ atm) [34]. However, in the non-geniculate red coralline alga *L. glaciale*, reduced pH (pH 7.7), particularly when variable, led to an increase in intracellular DMSP concentrations [16]. Species-specific responses

have also been observed in phytoplankton. An up-regulation of intracellular DMSP has been observed in *Emiliania huxleyi* under high temperature and high CO<sub>2</sub> (+4°C/1000 ppm) [35]; (+6°C/790 ppm) [36]. In contrast, intracellular DMSP concentrations were reduced in *Thalassiosira pseudonana* and *Phaeodactylum tricornutum* under high temperature and CO<sub>2</sub> (+6°C/790 ppm) [36].

In tropical reef environments, intracellular DMSP may be important as an antioxidant, grazing deterrent and/or compatible solute. Recent studies also suggest that intracellular DMSP may play a role in improving tolerance to variable carbonate chemistry conditions [16]. Corals from environmentally variable conditions (e.g. reef lagoons and flats) may be more thermally tolerant than those from more stable conditions (e.g. reef crests) [37]. Thus, it may be expected that organisms (e.g. corals and macroalgae) on reef flats have also developed mechanisms (e.g. up-regulation of DMSP) to become more tolerant of carbonate system variability compared to those on reef crests. This study assessed natural spatiotemporal variability of intracellular DMSP and water column DMSP+DMS (DMS/P) concentrations in the fringing reef environment of the Red Sea and related this to the carbonate chemistry of the overlying water column. It was hypothesised that (1) intracellular DMSP and water column DMS/P concentrations would fluctuate more on the environmentally variable reef flat compared to the environmentally stable reef crest and (2) corals would be the primary source of DMS/P in the reef system, as was observed on the Great Barrier Reef, Australia [38,39].

## Materials and Methods

### Study Site and Field Sampling Design

Research was conducted on the fringing Suleman Reef, Gulf of Aqaba, northern Red Sea, Egypt (28°28'N, 34°30'W) in August 2011. This reef is characterised by four distinct zones: seagrass beds nearest the shore, the reef flat 40–60 m from the shore, the reef crest 100–120 m from the shore and the reef slope which extends away from the reef crest to a depth of ~10 m. On average, water depth was uniform across the reef platform (~0.8 m). All research was approved by the Dahab Marine Research Center and the Marine Environmental Center, Suez Canal University, provided the permit to conduct the research.

Water and macroalgal samples were taken at the reef flat (dominated by fleshy and coralline macroalgae) and the reef crest (dominated by small branching corals, macroalgae and encrusting coralline algae) over two 45-hour experimental periods, at T14, T21, T29, T38 and T45 hours (Where T0 was 00:00 on day one). *In situ* water temperature, salinity and dissolved oxygen were recorded at each timepoint using a YSI Pro 2030 instrument. *In situ* photosynthetically active radiation (PAR) at the sea bed was recorded hourly at the reef flat and reef crest using an Apogee QSO-E underwater quantum sensor and Gemini voltage data logger.

### Carbonate Chemistry

At each timepoint, water samples were preserved for carbonate chemistry with MgCl<sub>2</sub> [40] and stored in the dark for subsequent total alkalinity (TA) and dissolved inorganic carbon (DIC) analysis. DIC was determined using a CO<sub>2</sub> Coulometer (CM5014 v.3, UIC Ltd.) with acidification module (CM5130 v.2, UIC Ltd.) and methods described by Dickson et al. [40]. TA was determined using the 2-stage open-cell titration method [40]. *In situ* TA, DIC, temperature and salinity measurements were used to calculate pH,  $p$ CO<sub>2</sub>, HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> concentrations and calcite ( $\Omega$ Ca) and aragonite ( $\Omega$ Ar) saturation states using CO2SYS [41] with

dissociation constants from Mehrbach et al. [42] refit by Dickson and Millero [43] and  $[KSO_4]$  using Dickson [44].

### Intracellular DMSP

Three macroalgal species were analysed for intracellular DMSP ( $n=10$  per species, per timepoint, per location): *Padina* sp. (Ochrophyta: Dictyotales), *Amphiroa* sp. (Rhodophyta: Corallinales) and *Turbinaria* sp. (Ochrophyta: Fucales). Both *Padina* sp. and *Amphiroa* sp. were found on the reef crest and reef flat; *Turbinaria* sp. was found only on the reef crest. Macroalgal samples ( $\sim 0.2$  g) were stored in 10 M NaOH in gas-tight glass vials sealed with Pharma-Fix (Grace Alltech) crimp lids to hydrolyse all intracellular DMSP into DMS. Vials were stored in the dark until DMS quantification.

### Water Column DMSP

At each sampling timepoint, water samples were collected from the reef flat and reef crest ( $n=5$  per location per timepoint), from 15 cm above the seabed. 50 ml water samples were filtered using a 0.7  $\mu\text{m}$  depth filter (Millepore). The filtrate and filter paper were stored in glass vials sealed with Pharma-Fix septa at a final concentration of 0.33 M NaOH. These samples represented the total dissolved and particulate DMS+DMSP pool (DMSPt). Logistical constraints limited DMSPt sample collections to the first 45-hour sampling period only.

### Reef Transect – Spatial Variation

In addition to the reef flat/reef crest diel comparison, water samples ( $n=3$  per location) were also taken at one timepoint (16:00) along a transect perpendicular to the shore at five locations: at the seagrass bed (10 m from the shore), on the reef flat (50 m from the shore), crest (100 m from the shore) and slope (120 m from the shore) and 270 m offshore (sandy bottom, no discernible macro-primary production). Samples were prepared for DMSPt using the methods described above.

### DMSP Analysis

All DMSP samples were analysed using a Shimadzu 2014 gas chromatograph (GC) fitted with a 25 m capillary column (Restek RTx-5MS 30 m column, 0.25 mm ID) and a sulphur-specific FPD detector (injector port and column oven temperature: 45°C, detector: 200°C). Samples were analysed within 2 months of collection. Intracellular DMSP samples were analysed by direct injection of the vial headspace into the GC injector port. Water column DMSPt samples were pre-concentrated using the purge and cryotrap technique [45] before injection into the GC. Sample concentrations were quantified from DMSP standard calibration curves (DMSP standard from Research Plus Inc.). The limit of detection was 960 ng S per 100  $\mu\text{l}$  headspace injection and 0.64 ng S per injection using the cryotrap system; standard and sample precision was within 3%.

### Proportion Calcified

The proportion of *Padina* sp. and *Amphiroa* sp. from the reef crest and flat that was calcified was determined by recording the mass of samples before and after overnight storage in 10 M NaOH to break down the organic material. The mass remaining represented the calcified portion, which was expressed as a percentage of the original mass. The proportion of *Amphiroa* sp. from the reef crest that was calcified could not be calculated as the carbonate structure broke down in the NaOH preventing final mass assessment.

### Statistics

Intracellular DMSP measurements could not be transformed to achieve normality and homogeneity of variance, thus a non-parametric multi-comparison Kruskal-Wallis test was performed to compare the data sets. Transect DMSPt data were analysed using general linear models (test assumptions met). Analyses were conducted using Minitab V15.

Generalised Additive Models (GAMs) were used to assess the relationship between the reefal abiotic parameters and measured macroalgal and water column DMSP concentrations. GAMs were conducted in R (V2.14.2) using the ‘mgcv’ package [46]. Thin plate regression splines were used as the smoothing basis, allowing multiple predictors to be considered and reducing subjectivity as knot locations (i.e. where splines join) are not manually specified [46]. Abiotic parameter interactions were identified using the ‘tree’ package in R. The most parsimonious GAM for each DMSP dataset was determined using generalised cross validation techniques (GCV) [46].

### Results


#### Water Chemistry

Both the reef crest and reef flat exhibited a diel trend in water temperature: highest water temperatures were in the afternoon and lowest temperatures were at dawn (Figure 1A). Whilst minimum water temperature on the reef flat and reef crest were the same (26.0°C), maximum temperatures were higher on the reef flat than the reef crest (29.1°C compared to 27.5°C), resulting in the reef flat having a wider temperature range than the reef crest (3.15°C and 1.5°C respectively, Figure 1A). The range in dissolved oxygen was also wider on the reef flat (60–160%) than the reef crest (110–70%, Figure S1A, supporting information). No difference was observed between the reef crest and reef flat in terms of PAR (0.8 m depth, Figure S1B, supporting information), with a maximum of  $>1500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  at 14:00 on day one and day two of the 45-hour experimental periods.

TA did not exhibit a strong diurnal oscillation on the reef crest (2422–2681  $\mu\text{mol kg}^{-1}$ ) or flat (2433–2574  $\mu\text{mol kg}^{-1}$ , Figure 1B). DIC on the reef flat ( $<2000 \mu\text{mol kg}^{-1}$ ) was lower than the reef crest ( $\sim 2100 \mu\text{mol kg}^{-1}$ ) at all daytime sampling points; at night (21 h and 45 h) DIC on the flat and crest were similar (2100–2200  $\mu\text{mol kg}^{-1}$ , Figure 1B). Saturation states were highest in the day (maximum  $\Omega_{\text{Ca}}$ : 9.3 on the crest and 7.8 on the flat;  $\Omega_{\text{Ar}}$ : 5.9 on the crest and 4.9 on the flat) and lowest at night/dawn (minimum  $\Omega_{\text{Ca}}$ : 3.9 on the crest and 3.7 on the flat;  $\Omega_{\text{Ar}}$ : 2.4 on the crest and 2.3 on the flat, Figure 1C). pH,  $\text{CO}_3^{2-}$  and  $\text{HCO}_3^-$  concentrations and  $p\text{CO}_2$  diel trends (derived from TA and DIC) are available in Figure S1C–F (supporting information).

#### Intracellular and Water Column DMSP

A general diurnal oscillation was observed for most DMSP measurements: high DMSP concentrations at 21:00 (T21+T45) and low concentrations at 14:00 (T14+T38, Figure 2). An exception to this was *Padina* sp. on the crest (no diurnal trend, Figure 2A). No statistical difference was observed between the intracellular DMSP concentrations of *Padina* sp. from the reef flat and crest except at T29 ( $Z=3.41$ ,  $p<0.001$ , Figure 2A). Intracellular DMSP in *Amphiroa* sp. on the flat ( $12.6 \pm 5.5 \text{ mg S g}^{-1}$ , mean  $\pm$  SD) was significantly lower than the crest ( $24.9 \pm 8.2 \text{ mg S g}^{-1}$ ,  $Z=4.73$ ,  $p<0.001$ , Figure 2B), with a significant diurnal trend of low intracellular DMSP at 14:00 (T14+T38) and high DMSPi at 21:00 (T21+T45,  $Z=4.63$ ,



**Figure 1. Diel pattern of abiotic parameters on Suleman Reef, Egypt.** *In situ* A: water temperature (°C), B: TA and DIC ( $\mu\text{mol kg}^{-1}$ ) and C: calcite ( $\Omega\text{Ca}$ ) and aragonite ( $\Omega\text{Ar}$ ) saturation states at the reef crest (black symbols) and reef flat (open symbols) over a 45-hour period on Suleman Reef, Egypt. Grey shading indicates night-time (sunset – sunrise). Data presented as mean  $\pm$  0.5 range of two experimental runs. doi:10.1371/journal.pone.0064651.g001

$p < 0.001$ , Figure 2B). A diurnal trend was also evident in the intracellular DMSP concentrations of *Turbinaria* sp. from the reef crest ( $< 0.3 \text{ mg S g}^{-1}$  at 14:00,  $> 0.5 \text{ mg S g}^{-1}$  at 21:00 [T21+T45]), although this was not significant ( $Z < 3.40$ ) and was characterised by a relatively high variability at 21:00 (T21+T45) and 07:00 (T29, Figure 2C). DMSPt did not differ significantly between the crest and flat ( $T_{22} = -0.18$ ,  $p = 0.862$ , Figure 2D), but there was suggestion of a diurnal trend, with elevated concentrations at 21:00 (T21+T45) compared to 14:00 (T14+T38), particularly on the reef flat.

#### Associations between Environmental Conditions and DMSP

Carbonate chemistry parameters were present in the most parsimonious GAM models for all DMSP datasets (Table 1). Water temperature was significantly associated with intracellular DMSP concentrations in *Amphiroa* sp. (reef crest and flat, Table 1). Dissolved oxygen was significantly associated with intracellular DMSP concentrations in *Padina* sp. (reef crest), *Turbinaria* sp. (reef crest) and water column DMSPt (reef flat, Table 1). The % deviance explained by the GAM models was generally higher for reef flat datasets, except *Padina* sp. (reef crest: 66%, reef flat: 44% deviance explained, Table 1).

#### Transect DMSPt

A significant difference in DMSPt was observed between the five transect sampling sites ( $F_4 = 10.38$ ,  $p = 0.001$ ) (Figure 3). DMSPt on the reef crest ( $17.29 \pm 0.53 \text{ nmol L}^{-1}$ , mean  $\pm$  SE) and slope ( $14.67 \pm 1.11 \text{ nmol L}^{-1}$ ) were significantly lower than the seagrass ( $23.38 \pm 2.04 \text{ nmol L}^{-1}$ ) and offshore ( $23.89 \pm 1.29 \text{ nmol L}^{-1}$ ) sites. The reef flat exhibited an intermediate DMSPt concentration ( $18.32 \pm 0.66 \text{ nmol L}^{-1}$ ).


#### Proportion Calcified

No significant difference in the proportion of thallus that was calcified was observed between *Padina* sp. samples from the reef crest ( $56.5 \pm 16.0\%$ , mean  $\pm$  SD) and flat ( $51.5 \pm 4.7\%$ ,  $T_1 = 0.30$ ,  $p = 0.816$ ). The proportion for *Amphiroa* sp. on the reef flat that was calcified was  $43.6 \pm 5.0\%$ . The proportion of *Amphiroa* sp. on the reef crest that was calcified could not be determined, although visually these samples appeared to be more heavily calcified.

#### Discussion

##### Biological Control of Carbonate Chemistry

Biological processes, particularly photosynthesis/respiration and calcification/dissolution can strongly influence the carbonate chemistry of tropical reefs [4,7,8]. The Suleman reef platform was, on average, only 0.8 m deep, thus biological processes are




**Figure 2. Diel pattern of DMS/P measurements on Suleman Reef, Egypt.** Intracellular DMSP ( $\text{mg S g}^{-1}$  algae as intracellular DMSP, DMSPi) of A: *Padina* sp., B: *Amphiroa* sp. and C: *Turbinaria* sp. and, in the water column, D: total DMS+DMSP (DMSPt,  $\text{nmol L}^{-1}$ ) on the reef crest (black circles) and reef flat (open circles) over a 45-hour period on Suleman Reef, Egypt. Grey shading indicates night-time (sunset – sunrise). Data presented as mean  $\pm$  SE. Note the different y-axes on all graphs. doi:10.1371/journal.pone.0064651.g002

**Table 1. Most parsimonious GAM models for intracellular DMSP (DMSPi) for *Padina* sp., *Amphiroa* sp. and *Turbinaria* sp. and water column DMSP on the reef crest and flat at Suleman Reef, Egypt.**

Location	Sample	GAM formula	Est. df	GCV score	Adj. R <sup>2</sup>	Dev. exp. (%)
Crest	<i>Padina</i>	DMSPi ~ s(TA)+s(DO)	6.92	0.084	0.64	66.4
	<i>Amphiroa</i>	DMSPi ~ s(TA)+s(Temp:TA)	5.16	0.074	0.278	31.3
	<i>Turbinaria</i>	DMSPi ~ s(DO)+s(DO:DIC)	4.00	0.150	0.536	56.2
Water column		DMSPt ~ s( $\Omega$ Ca)	2.57	0.009	0.169	27
Flat	<i>Padina</i>	DMSPi ~ s(TA)+s(TA: $\Omega$ Ca)	5.02	0.100	0.408	43.7
	<i>Amphiroa</i>	DMSPi ~ s(Temp)+s( $\Omega$ Ca)+s(pCO <sub>2</sub> )	5.99	0.034	0.739	75.4
	Water column	DMSPt ~ s(DO)+s(pH)	3.98	0.005	0.745	80.9

Statistics presented: estimated degrees of freedom (Est. df), GCV score, Adjusted R<sup>2</sup> (Adj. R<sup>2</sup>), % deviance explained (Dev. exp.).  
 TA: Total alkalinity, DO: dissolved oxygen, Temp: water temperature, DIC: dissolved inorganic carbon,  $\Omega$ Ca: calcite saturation state, pCO<sub>2</sub>: partial pressure of CO<sub>2</sub>.  
 doi:10.1371/journal.pone.0064651.t001



**Figure 3. DMSPt measurements along a transect across Suleman Reef, Egypt.** Measurements taken from the five zones of Suleman reef: seagrass beds nearest the shore, the reef flat, reef crest, reef slope and 270 m offshore. Data presented as mean  $\pm$  SE. doi:10.1371/journal.pone.0064651.g003

likely to have strongly influenced the observed diel patterns in carbonate chemistry, probably due to the low surface area to volume ratio [47]. In general, carbonate parameters were more variable on the reef flat compared to the reef crest. The diel difference between CO<sub>2</sub> utilisation and release was likely higher on the reef flat than the crest due to the high proportion of macroalgal cover on the reef flat. CO<sub>2</sub> uptake for photosynthesis during the day (which reduces DIC) and release by respiration at night (which increases DIC) were the probable drivers of the diel pattern in DIC concentrations. This proposed biological control mechanism was also reflected in the dissolved oxygen concentrations, which were  $\sim$ 160% saturated during the day (release from photosynthesis) and  $<$ 60% saturated during the night (uptake for respiration) on the reef flat (see Figure S1A, supporting information). The TA and DIC measurements observed in Suleman Reef are within the range cited for other tropical reef systems [47]. Such large diel changes in oxygen saturation have also been reported for Puerto Rican reefs (62–138%) [6].

### Species-specific Intracellular DMSP Concentrations

Intracellular DMSP concentrations of Red Sea tropical macroalgae in this study were higher than intracellular DMSP concentrations reported for the same genera from Australia [39] and the Caribbean [48]. Restricted water circulation within the Gulf of Aqaba results in more extreme conditions (salinity, temperature and carbonate chemistry) compared to other tropical reefs [49], which may necessitate an increase in intracellular DMSP relative to other tropical reef environments. Despite this, inter- and intra-species differences in intracellular DMSP concentrations were observed in the macroalgae of Suleman Reef, with *Amphiroa* sp. (crest)  $>$  *Amphiroa* sp. (flat)  $>$  *Padina* sp. (crest and flat)  $>$  *Turbinaria* sp. (crest). Grazing pressure may have contributed to the species-specific and diel patterns in intracellular DMSP concentrations, as grazing pressure is typically highest at night on tropical reefs [50] and DMSP may be an important grazing defence mechanism in macroalgae [51,52,53].

### Diel Regulation of DMSP

The production of DMSP by algae is energetically costly [54], thus the regulation of DMSP in response to environmental change (e.g. hypersalinity) may be relatively slow [55]. However, the principal functions of DMSP in algae under varying environmental conditions are not well understood. Overall, a clear diel pattern in macroalgal intracellular DMSP and water column DMSPt concentrations was observed, with DMSP concentrations typically highest at night, supporting the macroalgal grazing defence hypothesis [51,52,53]. Our results also suggest that DMSP concentrations were driven, at least in part, by the reefal carbonate chemistry. Thus, intracellular DMSP may play a role in maintaining macroalgal cellular function whilst under low carbonate saturation conditions, although other factors may also play a role (e.g. nutrients, grazing pressure). In temperate red coralline algae, DMSP may be up-regulated in response to acute changes in saturation state [16], thus it may be proposed that the algae sampled in this study also respond to decreases in saturation state by up-regulating intracellular DMSP concentrations. DMSPt will have consisted of benthic, detrital, land and phytoplankton material. DMSPt on the reef flat was well described by dissolved oxygen and pH (81% deviance explained); whilst DMSPt on the reef crest was not well described by any of the measured abiotic parameters (most parsimonious parameter:  $\Omega$ Ca, 27% deviance explained), suggesting that DMSPt on the crest was influenced by offshore waters more than DMSPt on the reef flat (different phytoplanktonic community for example).

DMSP and its breakdown products have been proposed as an ‘antioxidant cascade’ [30], able to ‘mop up’ reactive oxygen species produced by photosynthesis or elevated temperature. Water temperature was a significant factor contributing to observed intracellular DMSP concentrations in *Amphiroa* sp. This suggests that either DMSP was utilised during the day and replenished at night or there was a lag in the response of intracellular DMSP in *Amphiroa* sp. to ROS production (perhaps due to the energy outlay required) [54]. However, such a long lag

in DMSP response (seven hours or more) may not be an effective mechanism against oxidative damage.

Dissolved oxygen levels remained well above the suggested level for hypoxic impact (~30% saturation) [56] and the effect of reduced oxygen saturation on the intracellular DMSP concentrations of macroalgae is not known. However, dissolved oxygen did appear to influence the intracellular DMSP concentrations of *Padina* sp. and *Turbinaria* sp., suggesting that DMSP may be up-regulated during periods of low oxygen saturation.

### Benthic Producers of DMSP

Tropical corals have been described as one of the most important benthic sources of DMSP and DMS in the coastal zone [39]. Most data are from the Great Barrier Reef (GBR), Australia but the sources of water column DMSP and DMS are undefined. The GBR is also morphologically and hydrographically different to the fringing reef of this study. Results from this study suggest that other macro-primary producers may be more important sources of DMSP than corals. Nearshore DMSP concentrations were highest in areas dominated by seagrass beds and macroalgae (on the reef flat) rather than corals (reef crest and slope). DMSP concentrations were also elevated at the offshore site, where the primary source is likely to have been phytoplankton. Importantly, DMSP in the water column is available to microorganisms for potential breakdown into DMS, which is then available for atmospheric flux. Given the shallow water depth of the Suleman Reef platform (~0.8 m), the inshore reef areas may be important sources of atmospheric DMS in the region.

### Structural Tolerance to Variable Conditions

Calcification in *Amphiroa* sp. (high-magnesium calcite) was visibly reduced on the reef flat, suggesting that the algae were, relatively speaking, partially decalcified in the more variable conditions. This could not be quantitatively confirmed but changes to the calcite structure of red coralline algae in response to variable carbonate saturation has been observed in the temperate coralline alga *Lithothamnion glaciale* [16]. Grazing pressure may have also contributed to the observed differences in calcification. In contrast, the calcification of *Padina* sp. (aragonite) was not significantly different between the reef flat and reef crest, despite previous reports of decalcified *Padina* sp. in naturally low pH environments in Italy and Papua New Guinea [23].

### References

- NOAA. Trends in atmospheric carbon dioxide. Available: <http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed 2012.
- Caldeira K, Wickett ME (2005) Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *Journal of Geophysical Research* 110: C09S04.
- Pelejero C, Calvo E, Hoegh-Guldberg O (2010) Paleo-perspectives on ocean acidification. *Trends in Ecology & Evolution* 25: 332–344.
- Anthony KRN, A. Kleypas J, Gattuso J-P (2011) Coral reefs modify their seawater carbon chemistry – implications for impacts of ocean acidification. *Global Change Biology* 17: 3655–3666.
- Kleypas JA, Anthony KRN, Gattuso J-P (2011) Coral reefs modify their seawater carbon chemistry – case study from a barrier reef (Moorea, French Polynesia). *Global Change Biology* 17: 3667–3678.
- Gray SEC, DeGrandpre MD, Langdon C, Corredor JE (2012) Short-term and seasonal pH, pCO<sub>2</sub> and saturation state variability in a coral-reef ecosystem. *Global Biogeochem Cycles* 26: GB3012.
- Gagliano M, McCormick M, Moore J, Depczynski M (2010) The basics of acidification: baseline variability of pH on Australian coral reefs. *Marine Biology* 157: 1849–1856.
- Zhang Z, Falter J, Lowe R, Ivey G (2012) The combined influence of hydrodynamic forcing and calcification on the spatial distribution of alkalinity in a coral reef system. *Journal of Geophysical Research* 117: C04034.
- Suzuki A, Kawahata H (2003) Carbon budget of coral reef systems: an overview of observations in fringing reefs, barrier reefs and atolls in the Indo-Pacific regions. *Tellus B* 55: 428–444.
- Bates NR, Amat A, Andersson AJ (2010) Feedbacks and responses of coral calcification on the Bermuda reef system to seasonal changes in biological processes and ocean acidification. *Biogeosciences* 7: 2509–2530.
- Falter JL, Lowe RJ, Atkinson MJ, Cuet P (2012) Seasonal coupling and decoupling of net calcification rates from coral reef metabolism and carbonate chemistry at Ningaloo Reef, Western Australia. *Journal of Geophysical Research* 117: C05003.
- Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed response to CO<sub>2</sub>-induced ocean acidification. *Geology* 37: 1131–1134.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, et al. (2007) Coral Reefs Under Rapid Climate Change and Ocean Acidification. *Science* 318: 1737–1742.
- Büdenbender J, U R, A F (2011) Calcification of the Arctic coralline red alga *Lithothamnion glaciale* in response to elevated CO<sub>2</sub>. *Marine Ecology Progress Series* 441: 79–87.
- Ragazzola F, Foster LC, Form A, Anderson PSL, Hansteen TH, et al. (2012) Ocean acidification weakens the structural integrity of coralline algae. *Global Change Biology* 18: 2804–2812.
- Burdett HL, Aloisio E, Calosi P, Findlay HS, Widdicombe S, et al. (2012) The effect of chronic and acute low pH on the intracellular DMSP production and epithelial cell morphology of red coralline algae. *Marine Biology Research* 8: 756–763.
- Martin S, Gattuso JP (2009) Response of Mediterranean coralline algae to ocean acidification and elevated temperature. *Global Change Biology* 15: 2089–2100.

### Environmental Implications and Conclusions

In coastal regions, the effect of OA will be superimposed on the naturally variable environment, resulting in new carbonate extremes [3]. Our study suggests that intracellular DMSP concentrations may be regulated in response to variability in reefal carbonate chemistry, helping to maintain cellular function during periods of lower carbonate saturation. Macroalgal intracellular DMSP regulation may also impact the function of macroalgae within the reefal ecosystem. Coralline algae are important settlement cues for invertebrate larvae, particularly corals, e.g. [57,58], a function which may be reduced under high-CO<sub>2</sub> conditions [59] and which may be driven by intracellular DMSP concentrations [60]. It is projected that other calcifying reef organisms such as urchins may be detrimentally affected by OA, e.g. [22,61,62], impacting the food web dynamics of the reef ecosystem and perhaps allowing a phase shift towards macroalgal dominance to occur [63]. A shift towards macroalgal dominance may subsequently impact DMSP biogeochemistry, affecting ecosystem function and perhaps resulting in higher water column DMSP concentrations. This will increase the potential atmospheric emissions of DMS from the area, with subsequent impacts on local climate regulation [26,27].

### Supporting Information

**Figure S1 Diel pattern in abiotic parameters of Suleman Reef, Egypt.** *In situ* A: dissolved oxygen (%), B: photosynthetically active radiation (PAR,  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), C:  $p\text{CO}_2$  ( $\mu\text{atm}$ ), D:  $\text{HCO}_3^-$  concentration ( $\mu\text{mol kg}^{-1}$ ), E:  $\text{CO}_3^{2-}$  concentration ( $\mu\text{mol kg}^{-1}$ ), F: pH. (TIF)

### Acknowledgments

Thanks extend to the staff at the Dahab Marine Research Centre for their provision of research facilities.

### Author Contributions

Conceived and designed the experiments: HB PD NK. Performed the experiments: HB PD NK. Analyzed the data: HB. Wrote the paper: HB PD AH MA NK.

18. Cumani F, Bradassi F, Di Pascoli A, Bressan G (2010) Marine Acidification Effects on Reproduction and Growth Rates of Corallinaceae Spores (Rhodophyta). *CIESM Congress Proceedings* 39: 735.
19. Kuffner IB, Andersson AJ, Jokiel PL, Rodgers KuS, Mackenzie FT (2008) Decreased abundance of crustose coralline algae due to ocean acidification. *Nature Geoscience* 1: 114–117.
20. Jokiel P, Rodgers K, Kuffner I, Andersson A, Cox E, et al. (2008) Ocean acidification and calcifying reef organisms: a mesocosm investigation. *Coral Reefs* 27: 473–483.
21. Gao K, Zheng Y (2010) Combined effects of ocean acidification and solar UV radiation on photosynthesis, growth, pigmentation and calcification of the coralline alga *Corallina sessilis* (Rhodophyta). *Global Change Biology* 16: 2388–2398.
22. Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, et al. (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454: 96–99.
23. Johnson VR, Russell BD, Fabricius KE, Brownlee C, Hall-Spencer JM (2012) Temperate and tropical brown macroalgae thrive, despite decalcification, along natural CO<sub>2</sub> gradients. *Global Change Biology*: n/a-n/a.
24. Langdon C, Broecker WS, Hammond DE, Glenn E, Fitzsimmons K, et al. (2003) Effect of elevated CO<sub>2</sub> on the community metabolism of an experimental coral reef. *Global Biogeochem Cycles* 17: 1011.
25. Langdon C, Takahashi T, Sweeney C, Chipman D, Goddard J, et al. (2000) Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochem Cycles* 14: 639–654.
26. Charlson RJ, Lovelock JE, Andreae MO, Warren SG (1987) Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Nature* 326: 655–661.
27. Ayers GP, Caine JM (2007) The CLAW hypothesis: a review of the major developments. *Environmental Chemistry* 4: 366–374.
28. Kirst G (1996) Osmotic adjustment in phytoplankton and macroalgae: the use of dimethylsulfoniopropionate (DMSP). In: Kiene RP, Visscher PT, Keller Maureen D, Kirst G, editors. *Biological and Environmental Chemistry of DMSP and Related Sulfonium Compounds*. New York: Plenum Press. 121–129.
29. Karsten U, Kuck K, Vogt C, Kirst GO (1996) Dimethylsulfoniopropionate production in phototrophic organisms and its physiological function as a cryoprotectant. In: Kiene RP, Visscher PT, Keller MD, Kirst GO, editors. *Biological Chemistry of DMSP and Related Sulfonium Compounds*. New York: Plenum Press. 143–153.
30. Sunda W, Kieber DJ, Kiene RP, Huntsman S (2002) An antioxidant function for DMSP and DMS in marine algae. *Nature* 418: 317–320.
31. Van Alstyne K, Puglisi M (2007) DMSP in marine macroalgae and macroinvertebrates: Distribution, function, and ecological impacts. *Aquatic Sciences - Research Across Boundaries* 69: 394–402.
32. Seymour JR, Simó R, Ahmed T, Stocker R (2010) Chemoattraction to Dimethylsulfoniopropionate Throughout the Marine Microbial Food Web. *Science* 329: 342–345.
33. Dethier M, Van Alstyne K, Duggins D (2001) Spatial Patterns in Macroalgal Chemical Defenses. *Marine Chemical Ecology*: CRC Press. 301–324.
34. Kerrison P, Suggett D, Hepburn L, Steinke M (2012) Effect of elevated  $\mu\text{CO}_2$  on the production of dimethylsulphoniopropionate (DMSP) and dimethylsulphide (DMS) in two species of *Ulva* (Chlorophyceae). *Biogeochemistry* 110: 5–16.
35. Arnold HE, Kerrison P, Steinke M (2012) Interacting effects of ocean acidification and warming on growth and DMS-production in the haptophyte coccolithophore *Emiliania huxleyi*. *Global Change Biology*: n/a-n/a.
36. Spielmeier A, Pohnert G (2012) Influence of temperature and elevated carbon dioxide on the production of dimethylsulfoniopropionate and glycine betaine by marine phytoplankton. *Marine Environmental Research* 73: 62–69.
37. Oliver T, Palumbi S (2011) Do fluctuating temperature environments elevate coral thermal tolerance? *Coral Reefs* 30: 429–440.
38. Broadbent AD, Jones GB (2004) DMS and DMSP in mucus ropes, coral mucus, surface films and sediment pore waters from coral reefs in the Great Barrier Reef. *Marine and Freshwater Research* 55: 849–855.
39. Broadbent AD, Jones GB, Jones RJ (2002) DMSP in corals and benthic algae from the Great Barrier Reef. *Estuarine Coastal and Shelf Science* 55: 547–555.
40. Dickson AG, Sabine CL, Christian JR, editors (2007) *Guide to best practices for ocean CO<sub>2</sub> measurements*. 191 p.
41. Pierrot D, Lewis E, Wallace DWR (2006) CO<sub>2</sub>SYS DOS Program developed for CO<sub>2</sub> system calculations. ORNL/CDIAC-105. Carbon Dioxide Information analysis Center. Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN.
42. Mehrbach C, Culberson CH, Hawley JE, Pytkowicz RM (1973) Measurement of the Apparent Dissociation Constants of Carbonic Acid in Seawater at Atmospheric Pressure. *Limnology and Oceanography* 18: 897–907.
43. Dickson AG, Millero FJ (1987) A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research Part A Oceanographic Research Papers* 34: 1733–1743.
44. Dickson AG (1990) Standard potential of the reaction:  $\text{AgCl(s)} + 1/2\text{H}_2(\text{g}) = \text{Ag(s)} + \text{HCl(aq)}$ , and the standard acidity constant of the ion  $\text{HSO}_4^-$  in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics* 22: 113–127.
45. Turner SM, Malin G, Bagander LE, Leck C (1990) Interlaboratory calibration and sample analysis of dimethyl sulfide in water. *Marine Chemistry* 29: 47–62.
46. Wood SN (2006) *Generalised Additive Models. An Introduction with R*. Boca Raton: Taylor & Francis.
47. Kleypas JA, Langdon C (2006) *Coral reefs and changing seawater carbonate chemistry*. Coral Reefs and Climate Change: Science and Management. Washington, DC: AGU. 73–110.
48. Dacey JWH, King GM, Lobel PS (1994) Herbivory by reef fishes and the production of dimethylsulfide and acrylic-acid. *Marine Ecology Progress Series* 112: 67–74.
49. Chiffings A (2003) *Marine Region 11: Arabian Seas. A global representative system of Marine Protected Areas*.
50. Lewis SM, Wainwright PC (1985) Herbivore abundance and grazing intensity on a Caribbean coral reef. *Journal of Experimental Marine Biology and Ecology* 87: 215–228.
51. Van Alstyne KL, Houser LT (2003) Dimethylsulphide release during macroinvertebrate grazing and its role as an activated chemical defense. *Marine Ecology Progress Series* 250: 175–181.
52. Van Alstyne KL, Pelletreau KN, Kirby A (2009) Nutritional preferences override chemical defenses in determining food choice by a generalist herbivore, *Littorina sitkana*. *Journal of Experimental Marine Biology and Ecology* 379: 85–91.
53. Van Alstyne KL, Wolfe GV, Freidenburg TL, Neill A, Hicken C (2001) Activated defense systems in marine macroalgae: evidence for an ecological role for DMSP cleavage. *Marine Ecology Progress Series* 213: 53–65.
54. Stefels J (2000) Physiological aspects of the production and conversion of DMSP in marine algae and higher plants. *Journal of Sea Research* 43: 183–197.
55. Kirst G (1989) Salinity tolerance of eukaryotic marine algae. *Annu Rev Plant Physiol Plant Mol Biol* 40: 21–53.
56. Doney SC (2010) The Growing Human Footprint on Coastal and Open-Ocean Biogeochemistry. *Science* 328: 1512–1516.
57. Steller D, Cáceres-Martínez C (2009) Coralline algal rhodoliths enhance larval settlement and early growth of the Pacific calico scallop *Argopecten ventricosus*. *Marine Ecology Progress Series* 396: 49–60.
58. Huggett M, Williamson J, de Nys R, Kjelleberg S, Steinberg P (2006) Larval settlement of the common Australian sea urchin *Heliocidaris erythrogramma* in response to bacteria from the surface of coralline algae. *Oecologia* 149: 604–619.
59. Webster NS, Uthicke S, Botté ES, Flores F, Negri AP (2013) Ocean acidification reduces induction of coral settlement by crustose coralline algae. *Global Change Biology* 19: 303–315.
60. Steinberg PD, De Nys R (2002) Chemical Mediation of Colonization of Seaweed Surfaces. *Journal of Phycology* 38: 621–629.
61. Stumpp M, Dupont S, Thorndyke MC, Melzner F (2011) CO<sub>2</sub> induced seawater acidification impacts sea urchin larval development II: Gene expression patterns in pluteus larvae. *Comparative Biochemistry and Physiology - Part A: Molecular Integrative Physiology* 160: 320–330.
62. Stumpp M, Wren J, Melzner F, Thorndyke MC, Dupont ST (2011) CO<sub>2</sub> induced seawater acidification impacts sea urchin larval development I: Elevated metabolic rates decrease scope for growth and induce developmental delay. *Comparative Biochemistry and Physiology - Part A: Molecular Integrative Physiology* 160: 331–340.
63. Dudgeon S, Aronson R, Bruno J, Precht W (2010) Phase shifts and stable states on coral reefs. *Marine Ecology Progress Series* 413: 201–216.