Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

A planar dianionic ditelluride and a cyclic tritelluride supported by P₂N₂ rings†Andreas Nordheider,^{a,b} Tristram Chivers,^{a*} Ramalingam Thirumoorthi,^a Kasun S. Athukorala Arachchige,^b Alexandra M. Z. Slawin,^b J. Derek Woollins^b and Ignacio Vargas-Baca^c⁵ Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Oxidation of alkali metal derivatives of [Te(^tBuN)P(μ -N^tBu)₂P(N^tBu)Te]²⁻ with I₂ produces the intermediate ditelluride dianion [Te(^tBuN)P(μ -N^tBu)₂P(N^tBu)Te]₂²⁻ with a planar PTeTeP conformation and, subsequently, the cyclic tritelluride [(^tBuN)P(μ -N^tBu)₂P(N^tBu)(μ -TeTeTe)].

In their pioneering studies of inorganic macrocycles that incorporate P₂N₂ rings,¹ Wright and co-workers have employed cyclocondensation to generate NH- or O-bridged systems of various sizes involving P^{III}P^VN₂ rings^{2,3} and a reductive method for the synthesis of a hexamer in which monoselenido (–Se–) units span P^{III}P^VN₂ rings.⁴ Recently, we have applied a mild oxidative approach to generate the trimers [(^tBuN)P(μ -N^tBu)₂P(N^tBu)(μ -E–E–)]₃ with a planar P₆E₆ (E = S, Se) framework in which dichalcogenido (–E–E–) groups are linked by perpendicular P^VN₂ rings.⁵ The synthesis of these polychalcogen macrocycles involves the two-electron oxidation of the dianions **1a** and **1b** with I₂. We now report a detailed investigation of the oxidation of the tellurium analogue **1c** that provides important insights into the initial oxidation process, as well as a notable difference in the final outcome of the oxidation compared to that observed for **1a** and **1b**. Specifically, we have identified and structurally characterised (a) the dianionic ditelluride [Te(^tBuN)P(μ -N^tBu)₂P(N^tBu)Te]₂²⁻ (**2**), with an unusual planar conformation, as the product of one-electron oxidation of **1c** and (b) the cyclic tritelluride [(^tBuN)P(μ -N^tBu)₂P(N^tBu)(μ -TeTeTe)] (**3**) as the result of further oxidation. DFT calculations were carried out in order to determine the reason(s) for the unusual planarity of the PTeTeP unit in the ditelluride **2**.

The oxidation of the dianion **1c** (as its dilithium derivative)^{6a} with a one-half molar equivalent of I₂ led to the isolation of extremely sensitive black crystals, which were identified by X-ray crystallography as [Li(tmeda)]₂**2** (Eq. 1).⁷ As depicted in Fig. 1, the dianion [Te(^tBuN)P(μ -N^tBu)₂P(N^tBu)Te]₂²⁻ (**2**) is comprised of a central ditellurido unit that links two P^VN₂ rings, each of which has exocyclic Te and N^tBu substituents that are *N*,*Te*-chelated to the tmeda-solvated Li⁺ counterions. The ³¹P NMR spectrum of [Li(tmeda)]₂**2** in d₈-toluene exhibits singlets at δ = –78.5 and –117.0 with ¹J(P,Te) = 1670 and 1219 Hz, respectively, reflecting the disparity in the P–Te bond lengths [d(P2–Te2) = 2.396(2) and d(P1–Te1) = 2.497(2) Å]. The Te1–Te1' bond length in **2** is 2.777(3) Å, cf. 2.70–2.71 Å for typical aryl ditellurides,⁸ 2.75–2.76 Å for ditellurides supported by an intramolecular Te⋯N interaction,⁹ and 2.77 Å for the bulky derivative (TpsiTe)₂ (Tpsi = tris(phenyldimethylsilyl)methyl).¹⁰

Atypically, the ditelluride [Li(tmeda)]₂**2** is perfectly planar (< P–Te–Te–P = 180.0°). The only other examples of antiperiplanar ditellurides are bis(chloro-2-pyridyl) ditelluride¹¹ and the diacyl ditelluride (2-MeOC₆H₄COTe)₂,¹² which exhibit intramolecular heteroatom-tellurium interactions and (TpsiTe)₂ for which the antiperiplanar conformation is imposed by the bulky substituents.¹⁰ The Te1⋯Te2' distance in [Li(tmeda)]₂**2** (3.884 Å) is shorter than the sum of van der Waals radii for Te (4.12 Å).¹³ Moreover, the closely related neutral ditelluride [(^tBuNH)P(μ -N^tBu)₂P(N^tBu)Te]₂ (**4**), which has a similar steric environment for the Te–Te linkage, exhibits a dihedral angle of –123.8° and a Te–Te bond length of 2.7204(9) Å (Fig. S1).¹⁴ In view of these observations, DFT calculations were carried out to probe whether the conformation of [Li(tmeda)]₂**2** is stabilized by intramolecular Te1⋯Te2' secondary bonding interactions (3.884 Å). Satisfactory structures were obtained from geometry optimizations (PBE-D3, TZP, ZORA) for planar and synclinal models of [Li(tmeda)]₂**2** and **4** simplified using Me groups in lieu of ^tBu. In both instances the planar conformations were higher in energy, by 8 and 37 kJ mol^{–1} for the models of [Li(tmeda)]₂**2** and **4**, respectively. Preferred P–Te–Te–P torsion angles are 98° for the former and 90° for the latter. The small difference of energy suggests that the conformation observed in the crystal structure of [Li(tmeda)]₂**2** is imposed by packing forces. Further analysis on the electronic structures of the [Li(tmeda)]₂**2** models failed to identify a particular orbital interaction or contribution from dispersion that would help stabilize the antiperiplanar geometry.

In order to provide a direct comparison of the oxidation of **1c** with that of **1a** and **1b**⁵, we have developed a synthesis of the sodium salt $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ via the metallation-telluration sequence illustrated in Scheme 1. Yellow crystals of

Scheme 1 Synthesis of the cyclic tritelluride **3**

$[\text{Na}(\text{tmeda})]_2\textbf{1c}$ suitable for an X-ray analysis were obtained from *n*-hexane at -30°C and the molecular structure is illustrated in Fig. 2.¹⁵ In contrast to the lithium analogue $[\text{Li}(\text{thf})]_2\textbf{1c}$ in which the dianionic ligand is coordinated to the Li^+ cations asymmetrically (*Te*, *Te'* and *N*, *Te* chelation),^{6a} the sodium salt $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ adopts a symmetrical structure (*Te*, *Te'* and *N*, *N'* coordination).¹⁷ The mean P–Te distance of $2.420(2)\text{ \AA}$ in $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ is shorter by ca. $0.03\text{ \AA}$ than that involving the two-coordinate Te in $[\text{Li}(\text{thf})]_2\textbf{1c}$.^{6a} Consistently, the $^1J(\text{P}, \text{Te})$ coupling constant of 1583 Hz for $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ is substantially larger than the values of 1309 and 1467 Hz found for the lithium analogue.^{6a} A doublet is observed in the ^{125}Te NMR spectrum of $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ at -148.7 ppm , *cf.* -289 and -87 ppm for the two inequivalent Te environments in $[\text{Li}(\text{thf})]_2\textbf{1c}$.^{6a}

Fig. 1. Molecular structure of $[\text{Li}(\text{tmeda})]_2$. Hydrogen atoms have been omitted for clarity. Selected bond distances ($\text{\AA}$) and angles ($^\circ$): $\text{Te1}-\text{Te1}'$ $2.7768(8)$, $\text{Te1}-\text{P1}$ $2.4970(18)$, $\text{Te2}-\text{P2}$ $2.3957(18)$, $\text{N4}-\text{Li1}$ $2.008(9)$, $\text{Te2}-\text{Li1}$ $2.848(12)$, $\text{P1}-\text{N1}$ $1.510(6)$, $\text{P2}-\text{N4}$ $1.566(5)$, $\text{P}-\text{N}_{\text{bridging}}$ (range) $1.679(4)-1.712(4)$, $\text{N5}-\text{Li1}$ $2.171(9)$, $\text{P1}-\text{Te1}-\text{Te1}'$ $104.41(4)$.

Treatment of $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ with an equimolar quantity of I_2 in toluene produced a dark red-black solid, which was recrystallised from pentane to give black crystals that were identified as the tritelluride $[(\text{tBuN})\text{P}(\mu\text{-N}'\text{Bu})_2\text{P}(\text{N}'\text{Bu})(\mu\text{-TeTeTe})] (\textbf{3})$ by single-crystal X-ray analysis.¹⁸ Higher yields (41 %) of **3** are obtained via metathesis of $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ with $\text{TeCl}_2\cdot\text{tmtu}$ in toluene (Scheme 1).

As shown in Fig. 3, the molecular structure of **3** is comprised of a tritelluride ligand bridging the $\text{P}^{\text{V}}_2\text{N}_2$ ring. Compound **3** is a rare example of a cyclic tritelluride, the only other representative being the Sn^{IV} complex $[\text{ArSn}(\mu\text{-Te})_2(\mu\text{-TeTeTe})\text{SnAr}]$ ($\text{Ar} = 2,6\text{-bis}(2,4,6\text{-triisopropylphenyl})\text{phenyl}$).¹⁹ Structurally characterised

Fig. 2. Molecular structure of $[\text{Na}(\text{tmeda})]_2\textbf{1c}$. *tert*-Butyl groups and hydrogen atoms have been omitted for clarity. Selected bond distances ($\text{\AA}$) and angles ($^\circ$): $\text{P1}-\text{Te1}$ $2.418(2)$, $\text{P2}-\text{Te2}$ $2.422(2)$, $\text{Na1}-\text{Te1}$ $3.047(4)$, $\text{Na2}-\text{Te2}$ $3.049(3)$, $\text{Na}-\text{N}$ (range) $2.451(8)-2.750(9)$, $\text{P1}-\text{N3}$ $1.576(7)$, $\text{P2}-\text{N4}$ $1.566(7)$, $\text{P}-\text{N}_{\text{bridging}}$ (range) $1.697(7)-1.719(7)$; $\text{P1}-\text{Te1}-\text{Na2}$ $80.41(8)$, $\text{P2}-\text{Te2}-\text{Na2}$ $80.35(8)$, $\text{Te1}-\text{Na2}-\text{Te2}$ $125.1(1)$, $\text{N3}-\text{Na1}-\text{N4}$ $101.2(2)$, $\text{N5}-\text{Na1}-\text{N6}$ $72.6(3)$, $\text{N7}-\text{Na2}-\text{N8}$ $76.5(3)$.

Fig. 3. Molecular structure of **3**. Hydrogen atoms have been omitted for clarity. Selected bond distances ($\text{\AA}$) and angles ($^\circ$): $\text{P1}-\text{Te1}$ $2.5317(10)$, $\text{P2}-\text{Te2}$ $2.5405(10)$, $\text{Te1}-\text{Te3}$ $2.7155(4)$, $\text{Te2}-\text{Te3}$ $2.7158(4)$, $\text{P1}-\text{N1}$ $1.508(3)$, $\text{P2}-\text{N4}$ $1.507(4)$, $\text{P}-\text{N}_{\text{bridging}}$ $1.679(3)-1.694(3)$; $\text{P1}-\text{Te1}-\text{Te3}$ $95.63(2)$, $\text{P2}-\text{Te2}-\text{Te3}$ $96.06(3)$, $\text{Te1}-\text{Te3}-\text{Te2}$ $104.50(1)$.

acyclic tritellurides incorporate either bulky substituents RTeTeTeR ($\text{R} = \text{C}(\text{SiMe}_3)_3$ ²⁰ or intramolecular $\text{N}\cdots\text{Te}$ coordination ($2\text{-pyridyl-C}_6\text{H}_5$,²¹ $8\text{-Me}_2\text{NC}_{10}\text{H}_7$ ²²); the thermally unstable dication $[\text{Bu}_3\text{PTeTeTeP}^+\text{Bu}_3]^{2+}$ has also been identified.²³ The mean Te–Te bond length of $2.716(1)\text{ \AA}$ and Te–Te–Te bond angle of $104.50(1)^\circ$ in **3** are comparable to the corresponding values of $2.705(2)\text{ \AA}$ and $104.02(5)^\circ$ found for $[\text{ArSn}(\mu\text{-Te})_2(\mu\text{-TeTeTe})\text{SnAr}]$.¹⁹ The P–Te bond length of $2.536(1)\text{ \AA}$ is notably longer (by ca. $0.12\text{ \AA}$) than the mean value for $[\text{Na}(\text{tmeda})]_2\textbf{1c}$ indicating a weak P–Te bond in **3**. Consistently, the $^1J(\text{P}, \text{Te})$ coupling of 1030 Hz observed in both

the ^{31}P and ^{125}Te NMR spectra of **3** in d_8 -toluene is among the lowest reported for P–Te compounds.²⁴ The ^{125}Te NMR spectrum of **3** is comprised of a well-separated pseudo-doublet of doublets at -442.8 ppm ($^1J(\text{P},\text{Te}) = 1031$ Hz, $^3J(\text{P},\text{Te}) = 41$ Hz), and a pseudo-triplet centred at $+361.9$ (t, $^2J(\text{P},\text{Te}) = 35$ Hz); these resonances are assigned to the equivalent pair of tellurium atoms Te1, Te2 and the unique tellurium centre Te3, respectively (Fig. 3). The heterocycle **3** is the first structurally characterised neutral P–Te ring, although the cation $[\text{N}(\text{P}^i\text{Pr}_2\text{Te})_2]^+$ embodying a five-membered NP^V_2Te_2 ring is known.²⁵

In summary, the formation of the ditellurido linkage in **2** is comparable to the generation of neutral dichalcogenides upon one-electron oxidation of the PNP-bridged monoanions $[\text{EPR}_2\text{NPR}_2\text{PE}]^-$ (E = S, Se, Te; R = ^iPr , ^tBu).²⁶ The identification of the acyclic intermediate **2** provides an important discernment into the process involved in the construction of P^V_2N_2 -bridged polychalcogen macrocycles via the oxidation pathway.²⁷ It also supplies an incentive for investigations of the sulfur and selenium analogues, which are potentially versatile building blocks for the construction of P^V_2N_2 -stabilised macrocycles that incorporate main group elements or transition metals in addition to dichalcogenido linkages.

Notes and references

^aDepartment of Chemistry, University of Calgary, Calgary, AB, Canada T2N 1N4. E-mail: chivers@ucalgary.ca; Fax: +1 403-289-9488; Tel: +1 403-220-5741

^bDepartment of Chemistry, University of St Andrews, St Andrews UK, KY16 9ST

^cDepartment of Chemistry and Chemical Biology, McMaster University, Hamilton, ON, Canada L8S 4M1

[†]Electronic supplementary information (ESI) available: Experimental and crystallographic data in pdf format. CCDC references numbers 909218–909222 for $[\text{Na}(\text{tmeda})_2]_2$, **1c**, $[\text{Li}(\text{tmeda})_2]_2$, **3**, **4** and $[\text{Na}_4(\text{P}_2(\mu\text{-N}^t\text{Bu})_2(\text{N}^i\text{Bu})_2)_2]$

- For a review, see E. L. Doyle, L. Riera and D. S. Wright, *Eur. J. Inorg. Chem.*, 2003, 3279.
- (a) A. Bashall, A. D. Bond, E. L. Doyle, F. Garcia, S. Kidd, G. T. Lawson, M. C. Parry, M. McPartlin, A. D. Woods and D. S. Wright, *Chem. Eur. J.*, 2002, **8**, 3377; (b) F. Garcia, J. M. Goodman, R. A. Kowenicki, I. Kuzu, M. McPartlin, M. A. Silva, L. Riera, A. D. Woods and D. S. Wright, *Chem. Eur. J.*, 2003, **10**, 6066.
- S. G. Calera, D. J. Eisler, J. M. Goodman, M. McPartlin, S. Singh and D. S. Wright, *Dalton Trans.*, 2009, 1293.
- S. Gonzles-Calera, D. J. Eisler, J. V. Morey, M. McPartlin, S. Singh and D. S. Wright, *Angew. Chem. Int. Ed.*, 2008, **47**, 1111.
- A. Nordheider, T. Chivers, R. Thirumoorthi, I. Vargas-Baca and J. D. Woollins, *Chem. Commun.*, 2012, **48**, 6346.
- (a) G. G. Briand, T. Chivers and M. Parvez, *Angew. Chem. Int. Ed.*, 2002, **41**, 3468; (b) G. G. Briand, T. Chivers, M. Parvez, and G. Schatte, *Inorg. Chem.*, 2003, **42**, 525.
- Crystal data for $[\text{Li}(\text{tmeda})_2]_2$: $\text{C}_{50}\text{H}_{110}\text{N}_{12}\text{Li}_2\text{P}_4\text{Te}_4$, $M = 1818.12$, triclinic, space group $P\bar{1}$, $a = 13.760(3)$, $b = 14.301(3)$, $c = 14.481(3)$ Å, $\alpha = 60.533(5)^\circ$, $\gamma = 65.043(5)^\circ$, $\beta = 69.151(5)^\circ$, $V = 2209.3(7)$ Å³, $Z = 1$, $\rho_{\text{calcd}} = 1.366$ g cm^{−3}, $\mu = 1.423$ mm^{−1}, $T = 93$ K, 14563 reflections collected (θ range $3.04\text{--}25.37^\circ$), 7868 unique ($R_{\text{int}} = 0.0517$), $R_1 = 0.0424$ for 5839 reflections with $I > 2\sigma(I)$ and $wR_2 = 0.1182$ (for all data).
- (a) G. Llabres, O. Dideberg and L. Dupont, *Acta Crystallogr.*, 1972, **B28**, 2438; (b) M. Soirlet, G. Van den Bossche, O. Dideberg and L. Dupont, *Acta Crystallogr.*, 1979, **B35**, 1727; (c) E. S. Lang, R. A. Burrow and E. T. Silveira, *Acta Crystallogr.*, 2002, **C58**, o397; (d) A.

- Fuller, L. A. S. Scott-Hayward, Y. Li, M. Bhl, A. M. Z. Slawin and J. D. Woollins, *J. Am. Chem. Soc.*, 2010, **132**, 5799.
- (a) S. C. Menon, H. B. Singh, J. M. Jasinski, J. P. Jasinski and R. J. Butcher, *Organometallics*, 1996, **15**, 1707; (b) J. E. Drake, M. B. Hursthouse, M. Kulcsar, M. E. Light and A. Silvestru, *J. Organomet. Chem.*, 2001, **623**, 153; (c) G. Mugesch, A. Panda, S. Kumar, S. Apte, H. B. Singh and R. Butcher, *Organometallics*, 2002, **21**, 884.
- T. M. Klaptke, B. Krumm, H. Nth, J. C. Glvez-Ruiz, K. Polborn, I. Schwab and M. Suter, *Inorg. Chem.*, 2005, **44**, 5254.
- The acute $\angle\text{C}\text{--}\text{Te}\text{--}\text{Te}$ angle of $85.8(1)^\circ$ for this derivative suggests a significant intramolecular $\text{N}\cdots\text{Te}$ interaction, but this parameter is not discussed in the paper. T. Junk, K. J. Irgolic and E. A. Meyers, *Acta Crystallogr.*, 1993, **C49**, 975.
- The planarity of this derivative is attributed to intramolecular non-bonded interactions between the carbonyl oxygen atoms (rather than the methoxy oxygen atoms) and the tellurium atoms. O. Niyomura, S. Kato and S. Inagaki, *J. Am. Chem. Soc.*, 2000, **122**, 2132.
- N. W. Alcock, *Adv. Inorg. Chem. Radiochem.*, 1972, **15**, 1.
- Red crystals of the neutral ditelluride $[(\text{BuNH})\text{P}(\mu\text{N}^t\text{Bu})_2\text{P}(\text{N}^i\text{Bu})\text{Te}]_2$ (**4**) were obtained from the decomposition of $[\text{Li}(\text{tmeda})_2]_2$ and identified by an X-ray structural determination (see ESI).
- Crystal data for $[\text{Na}(\text{tmeda})_2]_2$: **1c**: $\text{C}_{28}\text{H}_{68}\text{N}_8\text{P}_2\text{Te}_2\text{Na}_2$, $M = 880.02$, monoclinic, space group $P2_1/n$, $a = 11.1830(5)$, $b = 30.2640(12)$, $c = 12.5750(5)$ Å, $\alpha = \gamma = 90.00^\circ$, $\beta = 95.151(1)^\circ$, $V = 4238.7(3)$ Å³, $Z = 4$, $\rho_{\text{calcd}} = 1.379$ g cm^{−3}, $\mu = 1.500$ mm^{−1}, $T = 173(2)$ K, 12267 reflections collected (θ range $1.4\text{--}27.5^\circ$), 7193 unique ($R_{\text{int}} = 0.0436$), $R_1 = 0.0644$ for 6017 reflections with $I > 2\sigma(I)$ and $wR_2 = 0.1771$ (for all data).
- The new reagent $[\text{Na}_2\{\text{BuNP}(\mu\text{-N}^t\text{Bu})_2\text{PN}^i\text{Bu}\}]_2$ was shown to be a dimer by a single-crystal X-ray structural determination (see ESI).
- The sulfur and selenium analogues of $[\text{Na}(\text{tmeda})_2]_2$, **1c**, in which tmeda is replaced by two thf ligands on the Na sites, adopt a similar symmetrical bonding arrangement. (a) T. Chivers, M. Krahn, and M. Parvez, *Chem. Commun.*, 2000, 463; (b) T. Chivers, M. Krahn, M. Parvez and G. Shatte, *Inorg. Chem.*, 2001, **40**, 2547; (c) G. G. Briand, T. Chivers and M. Krahn, *Coord. Chem. Rev.*, 2002, **233**, 237.
- Crystal data for **3**: $\text{C}_{16}\text{H}_{36}\text{N}_4\text{P}_2\text{Te}_3$, $M = 729.23$, monoclinic, space group $C2/c$, $a = 33.1490(2)$, $b = 14.2630(3)$, $c = 11.5420(3)$ Å, $\alpha = \gamma = 90.00^\circ$, $\beta = 107.354(1)^\circ$, $V = 5208.7(2)$ Å³, $Z = 8$, $\rho_{\text{calcd}} = 1.860$ g cm^{−3}, $\mu = 3.469$ mm^{−1}, $T = 173(2)$ K, 20105 reflections collected (θ range $1.6\text{--}27.4^\circ$), 5894 unique ($R_{\text{int}} = 0.0408$), $R_1 = 0.0358$ for 5360 reflections with $I > 2\sigma(I)$ and $wR_2 = 0.0742$ (for all data).
- M. Saito, H. Hashimoto and T. Tajima, *Acta Crystallogr. Section E*, 2010, **E66**, m885.
- F. Sladky, B. Bildstein, C. Rieker, A. Gieren, H. Betz and T. Hbner, *J. Chem. Soc., Chem. Commun.*, 1985, 1800.
- T. A. Hamor, N. Al-Salim, A. A. West and W. R. McWhinnie, *J. Organomet. Chem.*, 1986, **310**, C5.
- J. Beckmann, J. Bolsinger and A. Duthie, *Organometallics*, 2009, **28**, 4610.
- N. Kuhn, H. Schumann and R. Boese, *J. Chem. Soc., Chem. Commun.*, 1987, 1257.
- A P–Te bond distance of $2.605(1)$ Å with $^1J(\text{P},\text{Te}) = 780$ Hz has been reported for the tricyclohexylphosphine adduct of a 1,2,5-telluradiazolium cation. J. L. Dutton and P. J. Ragona, *Inorg. Chem.*, 2009, **48**, 1722.
- (a) J. Konu, T. Chivers and H. M. Tuononen, *Chem. Commun.*, 2006, 1634; (b) J. Konu, T. Chivers and H. M. Tuononen, *Inorg. Chem.*, 2006, **45**, 10678.
- (a) T. Chivers, J. S. Ritch, S. D. Robertson, J. Konu and H. M. Tuononen, *Acc. Chem. Res.*, 2010, **43**, 1053; (b) T. Chivers, D. J. Eisler, J. S. Ritch and H. M. Tuononen, *Angew. Chem. Int. Ed.*, 2005, **44**, 4953; (c) J. S. Ritch, T. Chivers, D. J. Eisler and H. M. Tuononen, *Chem.–Eur. J.*, 2007, **13**, 4643.
- Wright and co-workers have invoked the involvement of an acyclic intermediate in the formation of NH -bridged $\text{P}^{\text{III}}_2\text{N}_2$ macrocycles on the basis of detailed ^{31}P NMR studies. F. Garcia, J. M. Goodman, R.