

RARE EVENTS FOR THE MANNEVILLE-POMEAU MAP

ANA CRISTINA MOREIRA FREITAS, JORGE MILHAZES FREITAS, MIKE TODD,
AND SANDRO VAIENTI

ABSTRACT. We prove a dichotomy for Manneville-Pomeau maps $f : [0, 1] \rightarrow [0, 1]$: given any point $\zeta \in [0, 1]$, either the Rare Events Point Processes (REPP), counting the number of exceedances, which correspond to entrances in balls around ζ , converge in distribution to a Poisson process; or the point ζ is periodic and the REPP converge in distribution to a compound Poisson process. Our method is to use inducing techniques for all points except 0 and its preimages, extending a recent result [HWZ14], and then to deal with the remaining points separately. The preimages of 0 are dealt with applying recent results in [AFV14]. The point $\zeta = 0$ is studied separately because the tangency with the identity map at this point creates too much dependence, which causes severe clustering of exceedances. The Extremal Index, which measures the intensity of clustering, is equal to 0 at $\zeta = 0$, which ultimately leads to a degenerate limit distribution for the partial maxima of stochastic processes arising from the dynamics and for the usual normalising sequences. We prove that using adapted normalising sequences we can still obtain non-degenerate limit distributions at $\zeta = 0$.

1. INTRODUCTION

One of the standard ways to investigate the statistical properties of a dynamical system $f : \mathcal{X} \rightarrow \mathcal{X}$ with respect to some measure \mathbb{P} is to look at its recurrence to certain points ζ in the system. This can be connected to Extreme Value theory: supposing that $\varphi : \mathcal{X} \rightarrow \mathbb{R}$ is an observable taking its unique maximum u_φ at ζ , one can look at the behaviour of the iterates $x, f(x), f^2(x), \dots$ via the observations

$$X_i = X_i(x) = \varphi \circ f^i(x). \quad (1)$$

If \mathbb{P} is an f -invariant probability measure then X_0, X_1, \dots is a stationary stochastic process. We furthermore assume that \mathbb{P} is ergodic in order to isolate specific statistical behaviour. So Birkhoff's Ergodic Theorem implies that these random variables will satisfy the law of large numbers. We can now consider the random variable given by the maximum of this process:

$$M_n = M_n(x) := \max\{X_0(x), \dots, X_{n-1}(x)\}.$$

Date: April 12, 2016.

2000 *Mathematics Subject Classification.* 37A50, 60G55, 60G70, 37B20, 60G10, 37C25.

ACMF was partially supported by FCT grant SFRH/BPD/66174/2009. JMF was partially supported by FCT grant SFRH/BPD/66040/2009. Both these grants are financially supported by the program POPH/FSE. ACMF, JMF and SV are partially supported by FCT (Portugal) project PTDC/MAT/120346/2010, which is funded by national and European structural funds through the programs FEDER and COMPETE. ACMF, JMF and MT are also supported by CMUP (UID/MAT/00144/2013), which is funded by FCT (Portugal) with national (MEC) and European structural funds through the programs FEDER, under the partnership agreement PT2020. SV has been supported by the ANR grant PERTURBATIONS and by the project MOD TER COM of the french Region PACA.

Again by the ergodic theorem, if ζ is in the support of \mathbb{P} then for any small ball around ζ , $f^n(x)$ must hit the ball for some n and typical x . Hence if φ is sufficiently regular, then we expect $M_n \rightarrow u_\varphi$. Therefore, to obtain a non-trivial limit law, we need to rescale $\{M_n\}_n$. Indeed, we say that we have an *Extreme Value Law* (EVL) for M_n if there is a non-degenerate distribution function $H : \mathbb{R} \rightarrow [0, 1]$ with $H(0) = 0$ and a sequence of levels $u_n = u_n(\tau)$ such that

$$n\mathbb{P}(X_0 > u_n) \rightarrow \tau \text{ as } n \rightarrow \infty, \quad (2)$$

and for which the following holds:

$$\mathbb{P}(M_n \leq u_n) \rightarrow \bar{H}(\tau) = 1 - H(\tau) \text{ as } n \rightarrow \infty,$$

where the convergence is meant at the continuity points of $H(\tau)$.

In recent years, there has been a great deal of work on EVLs in the context of dynamical systems (see for example [Col01, FF08a, VHF09, FFT10, GHN11, HNT12, LFW12, Kel12, FHN14, AFV14]), the standard form of the observable φ being a function of the distance to ζ , for example $\varphi(x) = -\log d(x, \zeta)$ for d a metric on \mathcal{X} . Note that instead of the log, different functions can be composed with the distance (see [FFT10, page 679]); moreover, φ need not depend on the distance to ζ , see [FFT11, Section 5]. However, for the purposes of this paper, as in [FFT10], we assume that φ is a function of the distance to ζ and is invertible in a vicinity of ζ , so that $\varphi^{-1}(u_n)$ is well defined for u_n sufficiently close to $\varphi(\zeta)$, which can possibly be infinite (see Section 4.2.3 for further comments on this).

In many cases it has been shown that for \mathbb{P} -a.e. $\zeta \in \mathcal{X}$, this setup gives an EVL with $\bar{H} = e^{-\tau}$. More recently it has been shown that if ζ is a periodic point of period p then $\bar{H} = e^{-\theta\tau}$ where $\theta \in (0, 1)$ depends on the Jacobean of the measure for f^p , and is referred to as the *Extremal Index* (EI). The EI is known to measure the intensity of clustering of exceedances of the levels u_n . In fact, in many cases, the EI is equal to the inverse of the average cluster size, so that the EI is equal to 1 when there is no clustering. In the case of a class of uniformly hyperbolic dynamical systems, a stronger property, a dichotomy, has been shown: either ζ is periodic and we have an EVL with some extremal index $\theta \in (0, 1)$, or there is an EVL $\bar{H} = e^{-\tau}$. This was shown for f some uniformly expanding interval map with a finite number of branches in [FP12] (see also [FFT12, Section 6]) and with a countable number of branches in [AFV14]; here, depending on the precise form of the map, the measure can be absolutely continuous with respect to Lebesgue (acip), or an equilibrium state for some Hölder potential. Inducing methods have been used to extend some of these results to non-uniformly hyperbolic dynamical systems (see [FFT13] which built on [BSTV03]), but the results have not thus far extended to such a complete dichotomy.

We note that from a more probabilistic direction, such as processes under certain mixing conditions, which can be related to some symbolic dynamical systems, there is related work on Hitting Time Statistics, which as in [FFT10] can be seen as Extreme Value Laws. For a review on early work in this direction see [AG01]. For recent work, including some analysis of *all* points, including periodic ones, see [AV09, AS11].

We can further enrich our process by considering the point process formed by entries into the regions $\{X > u_n\}$, which in good cases gives rise to a Poisson process. An analogous dichotomy can often be shown there also: in the case of a periodic point ζ , we obtain a compound Poisson process. We leave the details of this construction to later.

In this note, we extend the dichotomy to a simple non-uniformly hyperbolic dynamical system, the *Manneville-Pomeau* (MP) map equipped with an absolutely continuous invariant probability measure. The particular form of these maps we will use is, for $\alpha > 0$,

$$f = f_\alpha(x) = \begin{cases} x(1 + 2^\alpha x^\alpha) & \text{for } x \in [0, 1/2) \\ 2x - 1 & \text{for } x \in [1/2, 1] \end{cases}$$

Members of this family of maps are often referred to as Liverani-Saussol-Vaianti maps since they were defined in [LSV99]. If $\alpha \in (0, 1)$ then there is an acip μ_α : we will restrict our attention to this case. As can be seen for example in [LSV99, You99, Hu04], for each $\alpha \in (0, 1)$, the system $([0, 1], f_\alpha, \mu_\alpha)$ has polynomial decay of correlations. That is, letting \mathcal{H}_β denote the space of Hölder continuous functions φ with exponent β equipped with the norm $\|\varphi\|_{\mathcal{H}_\beta} = \|\varphi\|_\infty + |\varphi|_{\mathcal{H}_\beta}$, where

$$|\varphi|_{\mathcal{H}_\beta} = \sup_{x \neq y} \frac{|\varphi(x) - \varphi(y)|}{|x - y|^\beta},$$

there exists $C > 0$ such that for each $\varphi \in \mathcal{H}_\beta$, $\psi \in L^\infty$ and all $t \in \mathbb{N}$,

$$\left| \int \varphi \cdot (\psi \circ f^t) d\mu_\alpha - \int \varphi d\mu_\alpha \int \psi d\mu_\alpha \right| \leq C \|\varphi\|_{\mathcal{H}_\beta} \|\psi\|_\infty \frac{1}{t^{\frac{1}{\alpha}-1}}. \quad (3)$$

Let $h_\alpha = \frac{d\mu_\alpha}{dx}$. From [Hu04] we know that $h_\alpha \in L^{1+\epsilon}$, with $\epsilon < 1/\alpha - 1$, and moreover $\lim_{x \rightarrow 0} \frac{h(x)}{x^{-\alpha}} = C_0 > 0$. Hence, for small $s > 0$ we have that

$$\mu_\alpha([0, s)) \sim_c s^{1-\alpha}, \quad (4)$$

where the notation \sim_c is used in the sense that there is $c > 0$ such that $\lim_{s \rightarrow \infty} \frac{\mu_\alpha([0, s))}{s^{1-\alpha}} = c$.

In this case there are canonical induced maps which capture all but a countable number of points in the phase space, so with some extra consideration for those points not captured, we can prove the full dichotomy, where for ζ a periodic point of period p , the extremal index is $\theta = 1 - 1/|Df^p(\zeta)|$.

For the special case in which ζ is the indifferent fixed point, we prove that there exists an EI equal to zero, which corresponds to a degenerate limit law, when the usual normalising sequences are used. Moreover, for a particular range of α we show that by changing the definition of $(u_n)_n$ given by (2) in a suitable way, we recover a non-degenerate EVL. This latter result relies on information on the transfer operator in [HSV99] as well as a refinement of the techniques for proving EVLs at periodic points developed in [FFT15].

The results that we present here for the Liverani-Saussol-Vaianti maps should be extendable to more general models of intermittent maps with neutral fixed (or periodic) points as long as they admit a first return time induced map which falls in the category of the uniformly expanding maps studied by Rychlik in [Ryc83]. We chose not to treat more general models because we believe that in this way the ideas are presented in a much easier way, without an unnecessary overload of notation and length.

1.1. Point process of hitting times. We will use our observations on our dynamical system to generate point processes. Here we adopt the approach and notation of [Zwe07]. Let $\mathcal{M}_p([0, \infty))$ be the space of counting measures on $([0, \infty), \mathcal{B}_{[0, \infty)})$. We equip this space with

the vague topology, i.e., $\nu_n \rightarrow \nu$ in $\mathcal{M}_p([0, \infty))$ whenever $\nu_n(\psi) \rightarrow \nu(\psi)$ for any continuous function $\psi : [0, \infty) \rightarrow \mathbb{R}$ with compact support. A *point process* N on $[0, \infty)$ is a random element of $\mathcal{M}_p([0, \infty))$. We will be interested in point processes $N_n : X \rightarrow \mathcal{M}_p([0, \infty))$. If we have a fixed measure μ on X , we say that $(N_n)_n$ *converges in distribution* to N if $\mu \circ N_n^{-1}$ converges weakly to $\mu \circ N^{-1}$. We write $N_n \xrightarrow{\mu} N$.

So given X_0, X_1, X_2, \dots and some $u \in \mathbb{R}$, we begin the construction of our point process $\mathbb{R} \rightarrow \mathcal{M}_p([0, \infty))$ as follows. Given $A \subset \mathbb{R}$ we define

$$\mathcal{N}_u(A) := \sum_{i \in A \cap \mathbb{N}_0} \mathbb{1}_{X_i > u}.$$

So $\mathcal{N}_u[0, n)$ counts the number of exceedances of the parameter u among the first n observations of the process X_0, X_1, \dots, X_{n-1} or, in other words, the number of entrances in $U(u) := \{X_0 > u\}$ up to time n .

We next re-scale time using the factor $v := 1/\mathbb{P}(X > u)$ suggested by Kac's Theorem. However, before we give the definition, we need some formalism. Let \mathcal{S} denote the semi-ring of subsets of \mathbb{R}_0^+ whose elements are intervals of the type $[a, b)$, for $a, b \in \mathbb{R}_0^+$. Let \mathcal{R} denote the ring generated by \mathcal{S} . Recall that for every $J \in \mathcal{R}$ there are $k \in \mathbb{N}$ and k intervals $I_1, \dots, I_k \in \mathcal{S}$ such that $J = \cup_{i=1}^k I_i$. In order to fix notation, let $a_j, b_j \in \mathbb{R}_0^+$ be such that $I_j = [a_j, b_j) \in \mathcal{S}$. For $I = [a, b) \in \mathcal{S}$ and $\alpha \in \mathbb{R}$, we denote $\alpha I := [\alpha a, \alpha b)$ and $I + \alpha := [a + \alpha, b + \alpha)$. Similarly, for $J \in \mathcal{R}$ define $\alpha J := \alpha I_1 \cup \dots \cup \alpha I_k$ and $J + \alpha := (I_1 + \alpha) \cup \dots \cup (I_k + \alpha)$.

We suppose that $\tau > 0$ and $(u_n)_n$ is defined so that (2) holds. We let $U(u_n) = \{X_0 > u_n\}$ and let v_n be the corresponding scaling factor defined above.

Definition. We define the *rare event point process* (REPP) by counting the number of exceedances (or hits to $U(u_n)$) during the (re-scaled) time period $v_n J \in \mathcal{R}$, where $J \in \mathcal{R}$. To be more precise, for every $J \in \mathcal{R}$, set

$$N_n(J) := \mathcal{N}_{u_n}(v_n J) = \sum_{j \in v_n J \cap \mathbb{N}_0} \mathbb{1}_{X_j > u_n}.$$

As will see below the REPP just defined converges in distribution to either to standard Poisson process or to a compound Poisson process N with intensity θ and a geometric multiplicity d.f. For completeness, we define here what we mean by a compound Poisson process. (See [Kal86] for more details).

Definition. Let T_1, T_2, \dots be an i.i.d. sequence of random variables with common exponential distribution of mean $1/\theta$. Let D_1, D_2, \dots be another i.i.d. sequence of random variables, independent of the previous one, and with d.f. π . Given these sequences, for $J \in \mathcal{R}$, set

$$N(J) = \int \mathbb{1}_J d \left(\sum_{i=1}^{\infty} D_i \delta_{T_1 + \dots + T_i} \right),$$

where δ_t denotes the Dirac measure at $t > 0$. Whenever we are in this setting, we say that N is a compound Poisson process of intensity θ and multiplicity d.f. π .

Remark 1. In this paper, the multiplicity will always be integer valued which means that π is completely defined by the values $\pi_k = \mathbb{P}(D_1 = k)$, for every $k \in \mathbb{N}_0$. Note that, if $\theta = 1$ and

$\pi_1 = 1$, then N is the homogenous standard Poisson process and, for every $t > 0$, the random variable $N([0, t))$ has a Poisson distribution of mean t .

Remark 2. At periodic points we will see that π is actually a geometric distribution of parameter $\theta \in (0, 1]$, i.e., $\pi_k = \theta(1 - \theta)^{k-1}$, for every $k \in \mathbb{N}_0$. This means that, as in [HV09], here, the random variable $N([0, t))$ follows a Pólya-Aeppli distribution, i.e.:

$$\mathbb{P}(N([0, t)) = k) = e^{-\theta t} \sum_{j=1}^k \theta^j (1 - \theta)^{k-j} \frac{(\theta t)^j}{j!} \binom{k-1}{j-1},$$

for all $k \in \mathbb{N}$ and $\mathbb{P}(N([0, t)) = 0) = e^{-\theta t}$.

Remark 3. By [Kal86, Theorem 4.2], the sequence of point processes $(N_n)_{n \in \mathbb{N}}$ converges in distribution to the point process N iff the sequence of vector r.v. $(N_n(I_1), \dots, N_n(I_k))$ converges in distribution to $(N(I_1), \dots, N(I_k))$, for every $k \in \mathbb{N}$ and all $I_1, \dots, I_k \in \mathcal{S}$ such that $N(\partial I_i) = 0$ a.s., for $i = 1, \dots, k$.

A few examples of studies of REPPs in a dynamical context can be found in [DGS04, HSV99, FFT10, CC13, FHN14], and in a more probabilistic context in [Pit91, AV08, KR14].

1.2. Main results. Let X_0, X_1, \dots be as in (1), with φ as specified above.

Theorem 1. Given $\zeta \in (0, 1]$, consider the REPP N_n defined above. Then either

- (a) ζ is not periodic and N_n converges in distribution to a homogeneous Poisson process N with intensity 1.
- (b) ζ is periodic with period p and N_n converges in distribution to a compound Poisson process N with intensity $\theta = 1 - |D(f^{-p})(\zeta)|$ and multiplicity distribution function π given by $\pi_\kappa = \theta(1 - \theta)^{\kappa-1}$, for every $\kappa \in \mathbb{N}_0$.

Theorem 2. For $\zeta = 0$, consider the maximum function $M_n = M_n(x)$ defined above.

- (a) Let $(u_n)_n = (u_n(\tau))_n$ be chosen as in (2), then $\mathbb{P}(M_n \leq u_n) \rightarrow 1$ as $n \rightarrow \infty$ for any $\tau > 0$.
- (b) For $\alpha \in (0, \sqrt{5} - 2)$ and $\tau \geq 0$, consider the sequence of thresholds $(u_n)_n = (u_n(\tau))_n$ satisfying:

$$\lim_{n \rightarrow \infty} n \mu_\alpha([x_n, \varphi^{-1}(u_n))) = \tau, \tag{5}$$

where x_n is such that $x_n < \varphi^{-1}(u_n)$ and $f_\alpha(x_n) = \varphi^{-1}(u_n)$. Then we have

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_n \leq u_n) \rightarrow e^{-\tau}.$$

Remark 4. Observe that the normalising sequence $(u_n)_{n \in \mathbb{N}}$ given by (5) does not satisfy condition (2). In fact, for such a sequence we have $\lim_{n \rightarrow \infty} n \mu_\alpha(X_0 > u_n) = \infty$. This is coherent with [LLR83, Corollary 3.7.4]. In classical Extreme Value Theory (as well as in our Theorems 1 and 4), the normalising constants are usually taken to be linear families depending on a parameter y , i.e., $u_n = a_n^{-1}y + b_n$, with $a_n > 0$ for all $n \in \mathbb{N}$, so that the limiting distribution for such linear normalisation is an extremal type distribution with $\tau = \tau(y)$ assuming three different types, see [Fre13a] and Section 4.2.3 for more details. When the extremal index is equal to 0, the types of limiting law may be different from the expected

ones if the stochastic process was independent, as noted in [LLR83, Page 69]. It is interesting to observe that the normalising sequences proposed here may lead to the same extremal limiting law as if the process was independent, or not (see Section 4.2.3).

Remark 5. We note that in the case $\zeta = 0$, while it is possible to rescale the thresholds to recover an EVL as in Theorem 2 (2), the corresponding REPP remains degenerate. This result will form part of a forthcoming work [FFR].

Remark 6. The choice $\alpha \in (0, \sqrt{5} - 2)$ is for technical reasons. This upper bound on α depends on the rate of decay of correlations, on the corresponding parameter β and also on ϵ , determining the regularity of the density. We would expect the result to hold for at least $\alpha \in (0, 1/2)$ but the current knowledge about these maps does not allow it. The obstructions appear due to the restrictions on the space of observable functions for which the statement (3) about decay of correlations holds. The underlying reasoning is a blocking argument which requires a mixing condition (see condition $\mathbb{D}_q(u_n)$ below) needed to give asymptotic independence between the blocks of random variables. This condition would follow immediately (for $\alpha < 1/2$) if one could plug indicator functions into (3). Since the existing information about decay of correlations requires the use of Banach spaces such as Hölder functions, one has to replace the indicator functions by suitable Hölder continuous approximations, which ultimately leads to further restrictions on α . These are the same technical limitations experienced for example in [HNT12] where $\alpha \in (0, \omega_0)$ for $\omega_0 \approx 1/13$.

1.3. Comments on history and strategy. Before discussing our approach we introduce some notation. For a dynamical system $f : \mathcal{X} \rightarrow \mathcal{X}$ and a subset $A \subset \mathcal{X}$, for $x \in \mathcal{X}$ define

$$r_A(x) := \inf\{n \in \mathbb{N} : f^n(x) \in A\},$$

the *first hitting time* to A . Note that there is a connection with the behaviour of the variable $r_{U(u_n)}$ and our REPP since we can break that process down into a sequence of first hits to $U(u_n)$. This gives a connection with our REPP and the asymptotics of $r_{U(u_n)}$, the *Hitting Time Statistics* (HTS). One basic difference is that here we are concerned with all hits to $U(u_n)$, not just the first.

Our main result for the case of \mathbb{P} -typical points and for periodic points in $(0, 1)$ follows quickly from previous works, including works already mentioned above, and indeed in some of these papers mention MP explicitly. We also remark that some of the earliest works on HTS for dynamical systems considered the case of MP maps with $\alpha \geq 1$, see for example [CG93, CGS92, CI95], with a focus on the behaviour at 0. In these cases, the sets A_n considered were formed from dynamically defined cylinder sets and the analysis was done at $1/2$, the preimage of 0, so that finite measure sets could be used. In this paper we consider the case $\alpha \in (0, 1)$, so f has an acip, and we also consider more general points and sets A_n .

We will first consider all points in $(0, 1)$, using inducing methods. This will require us to generalise the already very flexible result of [HWZ14] to point processes. Finally we use the approach which goes back to Leadbetter [Lea74] of proving some short range and long range recurrence conditions to prove that we have a degenerate law at 0 (the extremal index is 0).

2. INDUCED POINT PROCESSES

Here we aim to generalise [HWZ14] to point processes. In that paper, they use [Zwe07, Corollary 5] as one of their key tools. In our, fairly analogous, setting we use [Zwe07, Corollary 6] instead. Note that previous results here include [BSTV03, Theorem 2.1], where they proved that for balls around typical points, the HTS of first return maps are the same as that for the original map - they also remarked, without details, that this can be extended to successive return times. Also in [FFT13], we extended this idea to periodic points. The strengths of the approach in [HWZ14] to HTS are that it covers *all* points, and that the proof is rather short.

We will give our result comparing the point process of the induced system to that coming from the original system in a general setting and then later apply this to our MP example. In this section, we take a dynamical system $f : \mathcal{X} \rightarrow \mathcal{X}$ with an ergodic f -invariant probability measure μ , choose a subset $Y \subset \mathcal{X}$, recalling that $r_Y : Y \rightarrow \mathbb{N}$ is the first return time to Y , consider $F = F_Y : Y \rightarrow Y$ to be the first return map f^{r_Y} to Y (note that r_Y and thus F may be undefined at a zero Lebesgue measure set of points which do not return to Y , but most of these points are not important, so we will abuse notation here). Let $\mu_Y(\cdot) = \frac{\mu(\cdot \cap Y)}{\mu(Y)}$ be the conditional measure on Y . By Kac's Theorem μ_Y is F_Y -invariant.

Setting $v_n^Y = 1/\mu_Y(X_0 > u_n)$, for the induced process $X_i^Y = \varphi \circ F_Y^i$,

$$N_n^Y(J) := \mathcal{N}_{u_n}^Y(v_n^Y J) = \sum_{j \in v_n^Y J \cap \mathbb{N}_0} \mathbb{1}_{X_j^Y > u_n}.$$

In keeping with [HWZ14], we denote our inducing domain by Y . Denote the speeded up return time by $r_{A,Y}$ (i.e., $r_{A,Y}(x) = \inf\{n \in \mathbb{N} : F_Y^n(x) \in A\}$) and the induced measure on Y by μ_Y . For each $k \geq 2$ and $x \in \mathcal{X}$, we also define $r_A^k(x) = r_A\left(f^{r_A^1(x)+\dots+r_A^{k-1}(x)}(x)\right)$ and, for $x \in Y$ and $A \subset Y$, the corresponding speeded up version $r_{A,Y}^k(x) = r_A\left(F_Y^{r_{A,Y}^1(x)+\dots+r_{A,Y}^{k-1}(x)}(x)\right)$. Moreover, for $\kappa > 0$ and $I \in \mathcal{S}$, set $I^\eta := \cup_{s \in I} B_\eta^+(s)$ where $B_\eta^+(s) = (s - \eta, s + \eta) \cap [0, \infty)$. For $J \in \mathcal{R}$ such that $J = \cup_{j=1}^\ell I_j$, set $J^\eta = \cup_{j=1}^\ell I_j^\eta$.

Theorem 3. *For every $J \in \mathcal{R}$, assume that $N(J^\eta)$ is continuous in η , for all small η . That is to say that if $J = \cup_{j=1}^\ell I_j$, then for every k_1, \dots, k_ℓ , the map $\eta \mapsto \mu_Y(N(I_1^\eta) \geq k_1, \dots, N(I_\ell^\eta) \geq k_\ell)$ is continuous for $\eta \in [0, \eta_0)$. Then*

$$N_n^Y \xrightarrow{\mu_Y} N \text{ as } n \rightarrow \infty \text{ implies } N_n \xrightarrow{\mu} N \text{ as } n \rightarrow \infty.$$

Proof. By [Zwe07, Corollary 6], for hitting times point processes such as $(N_n)_n$ and an ergodic reference measure m , if $P \ll m$ then $N_n \xrightarrow{P} N$ in $\mathcal{M}_p([0, \infty))$ implies $N_n \xrightarrow{Q} N$ in $\mathcal{M}_p([0, \infty))$ for any $Q \ll m$. So replacing both m and Q with μ and replacing P with μ_Y we see that for our sequence of processes, if $N_n \xrightarrow{\mu_Y} N$ in $\mathcal{M}_p([0, \infty))$, then $N_n \xrightarrow{\mu} N$ in $\mathcal{M}_p([0, \infty))$. Thus, by Remark 3, it suffices to show that for every $J \in \mathcal{R}$, such that $J = \cup_{j=1}^\ell I_j$, and all

$k_1, \dots, k_\ell \in \mathbb{N}$, we have

$$\begin{aligned} \mu_Y(N_n^Y(I_1) \geq k_1, \dots, N_n^Y(I_\ell) \geq k_\ell) &\xrightarrow{n \rightarrow \infty} \mu_Y(N(I_1) \geq k_1, \dots, N(I_\ell) \geq k_\ell) \\ &\text{implies } \mu(N_n(I_1) \geq k_1, \dots, N_n(I_\ell) \geq k_\ell) \xrightarrow{n \rightarrow \infty} \mu(N(I_1) \geq k_1, \dots, N(I_\ell) \geq k_\ell). \end{aligned}$$

For $\delta > 0$ and $M \in \mathbb{N}$, let

$$E_M = E_M^\delta := \left\{ \left(\frac{1-\delta}{\mu(Y)} \right) j \leq r_Y^j \leq \left(\frac{1+\delta}{\mu(Y)} \right) j \text{ for all } j \geq M \right\} \text{ and } G_M := \{r_{U(u_n), Y} \geq M\}.$$

As in [HWZ14], $\mu_Y(G_M^c) \leq N \mu_Y(U(u_n)) \rightarrow 0$ as $n \rightarrow \infty$. Also the ergodic theorem says that $\mu_Y((E_M^\delta)^c) \rightarrow 0$ as $M \rightarrow \infty$. Hence we may restrict our focus to $G_M \cap E_M^\delta$.

For $x \in E_M^\delta$,

$$r_{U(u_n), Y}^k(x) \left(\frac{1-\delta}{\mu(Y)} \right) \leq r_{U(u_n)}^k(x) = \sum_{j=0}^{r_{U(u_n), Y}^k(x)-1} r_Y \circ F_Y^j(x) = r_Y^{r_{U(u_n), Y}^k(x)}(x) \leq r_{U(u_n), Y}^k(x) \left(\frac{1+\delta}{\mu(Y)} \right)$$

We can deduce that for $x \in G_M \cap E_M^\delta$,

$$\mu(Y) r_{U(u_n)}^k(x) \in B_{\delta r_{U(u_n), Y}^k(x)}(r_{U(u_n), Y}^k(x)),$$

where we use the notation $B_\varepsilon(y) = (y - \varepsilon, y + \varepsilon)$.

So if $r_{U(u_n), Y}^k(x) \in v_n^Y J$ then $\mu(Y) r_{U(u_n)}^k(x) \in v_n^Y J^\delta$ and so $r_{U(u_n)}^k(x) \in v_n J^\delta$. Therefore,

$$\begin{aligned} \mu_Y \left(\left\{ N_n(I_1^\delta) \geq k_1, \dots, N_n(I_\ell^\delta) \geq k_\ell \right\} \cap (E_M^\delta \cap G_M) \right) &\geq \\ &\mu_Y \left(\left\{ N_n^Y(I_1) \geq k_1, \dots, N_n^Y(I_\ell) \geq k_\ell \right\} \cap (E_M^\delta \cap G_M) \right). \end{aligned}$$

Setting $\delta' := \frac{\delta}{1+\delta}$, we also obtain that

$$\frac{1}{\mu(Y)} r_{U(u_n), Y}^k(x) \in B_{\delta' r_{U(u_n), Y}^k(x)}(r_{U(u_n), Y}^k(x))$$

for $x \in G_M \cap E_M^\delta$. Analogously to above, this leads us to

$$\begin{aligned} \mu_Y \left(\left\{ N_n^Y(I_1^{\delta'}) \geq k_1, \dots, N_n^Y(I_\ell^{\delta'}) \geq k_\ell \right\} \cap (E_M^\delta \cap G_M) \right) &\geq \\ &\mu_Y \left(\left\{ N_n(I_1) \geq k_1, \dots, N_n(I_\ell) \geq k_\ell \right\} \cap (E_M^\delta \cap G_M) \right). \end{aligned}$$

So since $\varepsilon, \delta > 0$ were arbitrary, we are finished. \square

3. APPLICATION OF INDUCING TO MANNEVILLE-POMEAU

In this section we prove our main theorem for all points $\zeta \in (0, 1)$.

Let \mathcal{P} be the *renewal partition*, that is the partition defined inductively by $Z \in \mathcal{P}$ if $Z = [1/2, 1)$ or $f(Z) \in \mathcal{P}$. Now let $Y \in \mathcal{P}$ and let F_Y be the first return map to Y and μ_Y be the conditional measure on Y . It is well-known that (Y, F_Y, μ_Y) is a Rychlik system (see [Ryc83] or [AFV14, Section 3.2.1] for the essential information about such systems) and so the REPP is understood as in [FFT13, Corollary 3]. Hence by [AFV14] we have the following theorem.

Theorem 4. *Given $\zeta \in Y$, consider the REPP N_n^Y defined above. Then either*

- (a) ζ is not periodic and $N_n^Y \xrightarrow{\mu_Y} N$, where N is a homogeneous Poisson process with intensity 1.
- (b) ζ is periodic with period p and $N_n^Y \xrightarrow{\mu_Y} N$, where N is a compound Poisson process with intensity $\theta = 1 - \left| D(F_Y^{-p})(\zeta) \right|$ and multiplicity d.f. π given by $\pi_\kappa = \theta(1-\theta)^{\kappa-1}$, for every $\kappa \in \mathbb{N}_0$.

For points in $Y \setminus \cup_{n \geq 1} f^{-n}(0)$, this theorem is Proposition 3.2 of [AFV14]. For the boundary points $\cup_{n \geq 1} f^{-n}(0)$, in the language of [AFV14], any such point is called aperiodic non-simple. Hence by Proposition 3.4(1) of that paper, we have a standard extremal index of 1 at all such points. Varying Y means that we have considered all points in $(0, 1)$. So combining Theorems 3 and 4 completes the proof of Theorem 1 for $\zeta \neq 0$.

4. ANALYSIS OF THE INDIFFERENT FIXED POINT

The tangency of the graph of the MP map with the identity map, creates an intensive clustering of exceedances of levels $(u_n)_{n \in \mathbb{N}}$, when they are chosen as in (2), that leads to the existence of an EI equal to 0, which leads to a degenerate limit distribution for M_n . However, if we choose the levels $(u_n)_{n \in \mathbb{N}}$ not in the classical way, but rather a sequence of lower thresholds, so that the exceedances that escape the clustering effect have more weight, then we can recover the existence of a non-degenerate distribution for the maxima.

The proof of an EI equal to 0 for the usual normalising sequences follows easily from the existing connections between Return Times Statistics (RTS), Hitting Times Statistics (HTS) and EVL, which we briefly recall in the next subsection. The proof of the existence of a non degenerate limit, under a different normalising sequence of thresholds, is more complicated and requires some new results from [FFT15], which we will recall below.

4.1. The usual normalising sequences case. For any $\zeta \in [0, 1]$, let $B_\varepsilon(\zeta) = (\zeta - \varepsilon, \zeta + \varepsilon) \cap [0, 1]$. Combining the main result from [FFT10] and [HLV07], if there exists a non degenerate d.f. \tilde{G} such that for all $t \geq 0$,

$$\lim_{\varepsilon \rightarrow 0} \mu_\alpha \left(r_{B_\varepsilon(\zeta)} \leq \frac{t}{\mu_\alpha(B_\varepsilon(\zeta))} \mid B_\varepsilon(\zeta) \right) = \tilde{G}(t),$$

then for G defined by

$$G(t) = \int_0^t (1 - \tilde{G}(s)) ds, \tag{6}$$

it can be shown that H exists and equals G .

Let $U = [0, b)$ and $A = [a, b)$, where a is such that $f(a) = b$, i.e., $b = a + 2^\alpha a^{1+\alpha}$. Using (4) we easily get $\mu_\alpha(U) \sim_c a^{1-\alpha} + (1-\alpha)2^\alpha a + o(a)$ and $\mu_\alpha([0, a)) \sim_c a^{1-\alpha}$.

¹We note that there is an error in [FFT13, Theorem 1], propagated throughout the main results there: the κ should be replaced by $\kappa - 1$.

Next we compute the RTS distribution, which we denote by $\tilde{G}(s)$. For $s \leq 0$, we easily have that $\tilde{G}(s) = 0$, since $r_U \geq 1$, by definition of hitting time. Let $s > 0$ then

$$\begin{aligned} \tilde{G}(s) &= \lim_{b \rightarrow 0} \mu_U \left(r_U \leq \frac{s}{\mu(U)} \right) = \lim_{b \rightarrow 0} \frac{1}{\mu(U)} \mu \left(\left\{ r_U \leq \frac{s}{\mu(U)} \right\} \cap U \right) \\ &\geq \lim_{b \rightarrow 0} \frac{\mu(U \setminus A)}{\mu(U)} = \lim_{b \rightarrow 0} \frac{\mu([0, a])}{\mu([0, b])} = 1 \end{aligned}$$

Then by (6), $G(t) = \int_0^t 1 - \tilde{G}(s) ds = 0$, which, by [FFT10], corresponds to an EI equal to 0. Recall that $\bar{H}(\tau) = e^{-\theta\tau} = 1$, which means that, in this case, $H(\tau) = 0$.

4.2. Adjusted choice of thresholds. In order to prove the existence of EVLs in a dynamical systems context, there are a couple of conditions on the dependence structure of the stochastic process that if verified allow us to obtain such distributional limits. These conditions are motivated by the conditions $D(u_n)$ and $D'(u_n)$ of Leadbetter but were adapted to the dynamical setting and further developed both in the absence of clustering, such as in [Col01, FF08b, HNT12], and in the presence of clustering in [FFT12]. Very recently, in [FFT15], the authors provided some more general conditions, called $\mathbb{D}(u_n)$ and $\mathbb{D}'_q(u_n)$, which subsumed the previous ones and allowed them to address both the presence ($q \geq 1$) and the absence ($q = 0$) of clustering. To distinguish these conditions the authors used a Cyrillic D to denote them. We recall these conditions here.

Given a sequence $(u_n)_{n \in \mathbb{N}}$ of real numbers satisfying (2) and $q \in \mathbb{N}_0$, set

$$A_n^{(q)} := \{X_0 > u_n, X_1 \leq u_n, \dots, X_q \leq u_n\}.$$

For $s, \ell \in \mathbb{N}$ and an event B , let

$$\mathscr{W}_{s, \ell}(B) = \bigcap_{i=s}^{s+\ell-1} f^{-i}(B^c). \quad (7)$$

Condition ($\mathbb{D}_q(u_n)$). We say that $\mathbb{D}(u_n)$ holds for the sequence X_0, X_1, \dots if, for every $\ell, t, n \in \mathbb{N}$

$$\left| \mathbb{P} \left(A_n^{(q)} \cap \mathscr{W}_{t, \ell} \left(A_n^{(q)} \right) \right) - \mathbb{P} \left(A_n^{(q)} \right) \mathbb{P} \left(\mathscr{W}_{0, \ell} \left(A_n^{(q)} \right) \right) \right| \leq \gamma(q, n, t), \quad (8)$$

where $\gamma(q, n, t)$ is decreasing in t and there exists a sequence $(t_n)_{n \in \mathbb{N}}$ such that $t_n = o(n)$ and $n\gamma(q, n, t_n) \rightarrow 0$ when $n \rightarrow \infty$.

For some fixed $q \in \mathbb{N}_0$, consider the sequence $(t_n)_{n \in \mathbb{N}}$ given by condition $\mathbb{D}_q(u_n)$ and let $(k_n)_{n \in \mathbb{N}}$ be another sequence of integers such that

$$k_n \rightarrow \infty \quad \text{and} \quad k_n t_n = o(n). \quad (9)$$

Condition ($\mathbb{D}'_q(u_n)$). We say that $\mathbb{D}'_q(u_n)$ holds for the sequence X_0, X_1, \dots if there exists a sequence $(k_n)_{n \in \mathbb{N}}$ satisfying (9) and such that

$$\lim_{n \rightarrow \infty} n \sum_{j=1}^{\lfloor n/k_n \rfloor} \mathbb{P} \left(A_n^{(q)} \cap f^{-j} \left(A_n^{(q)} \right) \right) = 0. \quad (10)$$

We note that, when $q = 0$, condition $\mathbb{D}'_q(u_n)$ corresponds to condition $D'(u_n)$ from [Lea74].

Now let

$$\vartheta = \lim_{n \rightarrow \infty} \vartheta_n = \lim_{n \rightarrow \infty} \frac{\mathbb{P}(A_n^{(q)})}{\mathbb{P}(U_n)}. \quad (11)$$

From [FFT15, Corollary 2.4], it follows that if the stochastic process X_0, X_1, \dots satisfies both conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$ and the limit in (11) exists then

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_n \leq u_n) = e^{-\vartheta\tau}.$$

Now, we consider the fixed point $\zeta = 0$. For every $n \in \mathbb{N}$, we require $\mu_\alpha(U_n) \sim \tau/n$. Set y_n to be such that $U_n = \{X_0 > u_n\} = [0, y_n)$ and set $x_n \in U_n$ so that $f_\alpha(x_n) = y_n$, *i.e.*, $y_n = x_n + 2^\alpha x_n^{1+\alpha}$. Using (4) we easily get

$$\mu_\alpha(U_n) \sim_c x_n^{1-\alpha} + (1-\alpha)2^\alpha x_n + o(x_n) \quad (12)$$

$$\mu_\alpha([0, x_n)) \sim_c x_n^{1-\alpha} \quad (13)$$

$$\mu_\alpha([x_n, y_n)) \sim_c (1-\alpha)2^\alpha x_n + o(x_n) \quad (14)$$

Now, since we are assuming that $\mu_\alpha(U_n) \sim \tau/n$, then $x_n \sim_c 1/n^{1/(1-\alpha)}$. Observe that $\mu_\alpha(U_n \cap f_\alpha^{-1}(U_n)) = \mu_\alpha([0, x_n)) \sim_c x_n^{1-\alpha} \sim_c 1/n$. Hence, if we consider $q = 0$, the periodicity of ζ implies that $\mathbb{D}'_q(u_n)$ does not hold since

$$n \sum_{j=1}^{\lfloor n/k_n \rfloor} \mathbb{P}(U_n \cap f^{-j}(U_n)) \geq n\mu_\alpha(U_n \cap f_\alpha^{-1}(U_n)) > 0,$$

for all $n \in \mathbb{N}$. Hence, here, given that ζ is a periodic point of period 1 the natural candidate for q is $q = 1$. From here on we always assume that $q = 1$.

In this case, $A_n^{(q)} = [x_n, y_n) =: Q_n$. However, if we plug (14) and (12) into (11), we obtain that $\vartheta = 0$, which means that the natural candidate for a limit distribution for $\mu_\alpha(M_n \leq u_n)$ is degenerate.

The problem is that the indifferent fixed point creates too much dependence. In [FFT12], under a condition called *SP*, we have seen that when ζ is periodic, the probability of having no entrances in U_n , among the first n observations, is asymptotically equal to the probability of having no entrances in Q_n , among the first n observations, *i.e.*,

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_n \leq u_n) = \lim_{n \rightarrow \infty} \mathbb{P}(\mathcal{W}_{0,n}(U_n)) = \lim_{n \rightarrow \infty} \mathbb{P}(\mathcal{W}_{0,n}(Q_n)).$$

In [FFT15], it was shown that it is possible to replace U_n by Q_n even without the *SP* condition (see [FFT15, Proposition 2.7]). Making use of this upgraded result, we can now change the normalising sequence of levels $(u_n)_{n \in \mathbb{N}}$ so that we can still obtain a non-degenerate limit for $\mathbb{P}(M_n \leq u_n)$. To understand the need to change the normalising sequence in order to obtain a non-degenerate limit, recall that condition (2) guaranteed that M_n was normalised by a sequence of levels that kept the average of exceedances among the first n observations at an (almost) constant value $\tau > 0$. When $\vartheta > 0$, condition (2) also guarantees that the average number of entrances in Q_n among the first n observations is kept at an (almost) constant value

$\theta\tau > 0$. Here, since $\vartheta = 0$, we need to change u_n so that the average number of entrances in Q_n is controlled, *i.e.*,

$$\lim_{n \rightarrow \infty} n\mathbb{P}(A_n^{(q)}) = \tau > 0. \quad (15)$$

From equations (2.15) and (2.16) from [FFT15] one gets:

$$\left| \mathbb{P}(\mathcal{W}_{0,n}(A_n^{(q)})) - \left(1 - \left\lfloor \frac{n}{k_n} \right\rfloor \mathbb{P}(A_n^{(q)})\right)^{k_n} \right| \leq 2k_n t_n \mathbb{P}(U_n) + 2n \sum_{j=1}^{\lfloor n/k_n \rfloor - 1} \mathbb{P}(A_n^{(q)} \cap f^{-j} A_n^{(q)}) + \gamma(q, n, t_n) \quad (16)$$

Note that since by (15) we have $\lim_{n \rightarrow \infty} \left(1 - \left\lfloor \frac{n}{k_n} \right\rfloor \mathbb{P}(A_n^{(q)})\right)^{k_n} = e^{-\tau}$, then if both conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$ hold, then all the terms on the left of (16) vanish, as $n \rightarrow \infty$, and consequently:

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_n \leq u_n) = \lim_{n \rightarrow \infty} \mathbb{P}(\mathcal{W}_{0,n}(A_n^{(q)})) = e^{-\tau}. \quad (17)$$

Hence, in order to show that we can still obtain a non-degenerate limiting law for the distribution of M_n when $\zeta = 0$, we start by taking a sequence $(u_n)_{n \in \mathbb{N}}$ so that (15) holds. Note that this implies that by (14) and (12) we have that $x_n \sim_c 1/n$ and $\mu_\alpha(U_n) \sim_c 1/n^{1-\alpha}$. In particular, this means that $\lim_{n \rightarrow \infty} n\mu_\alpha(U_n) = \infty$, which contrasts with the usual case where condition (2) holds.

To prove the existence of the limit in (17) we need to verify conditions $\mathbb{D}_q(u_n)$ and $\mathbb{D}'_q(u_n)$, where $q = 1$. We start by the latter, which is more complicated.

4.2.1. *Proof of $\mathbb{D}'_q(u_n)$.* We will next focus on the proof of $\mathbb{D}'_q(u_n)$ in the case of part (2) of Theorem 2. That is, $(x_n)_n$ will be chosen so that $x_n \sim_c 1/n$, as described above. Later we will note that we can change $(x_n)_n$ to recover a degenerate law as in part (1) of that theorem.

We have to estimate the quantity

$$\Delta'_n := n \sum_{j=1}^{\lfloor n/k_n \rfloor} \mu_\alpha(Q_n \cap f^{-j} Q_n)$$

where $Q_n = [x_n, y_n)$, for $x_n \sim_c \frac{1}{n}$ and $y_n = f(x_n)$. We follow the proof of [HSV99, Lemma 3.5]. By denoting by P the transfer operator and by $\tau_n \in \mathbb{N}$ the *first return time* of the set Q_n into itself, we have:

$$\Delta'_n \leq n \lfloor n/k_n \rfloor \mu_\alpha(Q_n) \sup_{j=\tau_n, \dots, \lfloor n/k_n \rfloor} \sup_{Q_n} \frac{P^j(\mathbf{1}_{Q_n} h)}{h}$$

where h is the density of μ_α . In order to compute $P^{\tau_n}(\mathbf{1}_{Q_n} h)$ we need to know how many branches of f^{τ_n} have their domain intersecting Q_n . If ξ_0 is the original partition into the sets $[0, 1/2)$, $[1/2, 1]$, we denote with ξ_k the join $\xi_k := \xi_0 \vee f^{-1}\xi_0 \vee \dots \vee f^{k-1}\xi_0$.

We begin by observing that Q_n contains at most one boundary point of the partition ξ_{τ_n-1} , otherwise one point of Q_n should be sent into the same set, being f^{τ_n-1} onto on each domain of injectivity. Then when we move to ξ_{τ_n} , the interval Q_n will be covered by at most 4 cylinders

of monotonicity of the partition ξ_{τ_n} . By denoting them from left to right by $C_{\tau_n,1}, \dots, C_{\tau_n,4}$ we have

$$P^{\tau_n}(\mathbf{1}_{Q_n} h) = \sum_{i=1}^4 \frac{h \circ f_i^{-\tau_n} \mathbf{1}_{f_i^{\tau_n} Q_n}}{Df^{\tau_n} \circ f_i^{-\tau_n}}$$

where $f_i^{\tau_n}$ denotes the branch of f^{τ_n} restricted to $C_{\tau_n,i}$. Notice that the density is computed in Q_n whose left boundary point is $1/n$, so h is bounded from above by a constant times n^α . We have now to estimate the derivative $Df^{\tau_n} \circ f_i^{-\tau_n}$ on the sets $Q_n \cap C_{\tau_n,i}$. Let us define r_m as the m -left preimage of 1, $r_m := f_1^{-m}(1)$ and define $m(n)$ as $r_{m(n)} \leq x_n \leq r_{m(n)-1}$. Then the interval $[x_n, y_n)$ will intersect the two cylinders $(r_{m(n)}, r_{m(n)-1})$ and $(r_{m(n)-1}, r_{m(n)-2})$ and the first return of Q_n will be larger than the first returns of those two cylinders; on the other hand the first return of $(r_{m(n)}, r_{m(n)-1})$ is $m(n)$. The derivative Df^{τ_n} will be computed at a point ι_n which will be in one of those two cylinders; suppose without any restriction that $\iota_n \in (r_{m(n)}, r_{m(n)-1})$. Since we need to bound from below the derivatives, we begin to replace $Df^{\tau_n}(\iota_n)$ with $Df^{m(n)}(\iota_n)$; then we observe that the map $f^{m(n)} : [r_{m(n)}, r_{m(n)-1}] \rightarrow [0, 1]$ is onto and we use the distortion bound given, for instance, in [LSY, Lemma 5] which states that there exists a constant C such that for any $m \geq 1$ and any $x, y \in [r_m, r_{m-1}]$ we have $\left| \frac{Df^m(x)}{Df^m(y)} \right| \leq C$. We finally note that $m(n) \sim_c n^\alpha$. This implies immediately that

$$\frac{1}{Df^{m(n)}(\iota_n)} \leq C |r_{m(n)-1} - r_{m(n)}| \sim_c C \frac{1}{m(n)^{\frac{1}{\alpha}+1}} \sim_c C \frac{1}{n^{1+\alpha}}.$$

Consequently (C will continue to denote a constant which could vary from one bound to another)

$$P^{\tau_n}(\mathbf{1}_{Q_n} h) \sim_c \frac{1}{n}$$

We now continue as in [HSV] by getting for the other powers of the transfer operator:

$$\frac{P^j(\mathbf{1}_{Q_n} h)}{h} \leq \frac{P^{j-\tau_n} \mathbf{1}}{h} \sup P^{\tau_n}(\mathbf{1}_{Q_n} h) \leq \frac{P^{j-\tau_n} \frac{h}{\inf h}}{h} \sup P^{\tau_n}(\mathbf{1}_{Q_n} h) \leq \frac{C}{\inf h} \frac{1}{n}$$

and finally

$$\Delta'_n \leq n [n/k_n] \mu_\alpha(Q_n) \frac{C}{\inf h} \frac{1}{n}$$

We now know that $\mu_\alpha(Q_n) \sim_c \frac{1}{n}$; hence

$$\Delta'_n \leq C n [n/k_n] \mu_\alpha(Q_n)^2 \frac{1}{\mu_\alpha(Q_n)} \frac{1}{n} \sim_c [n^2 \mu_\alpha(Q_n)^2] \frac{1}{k_n}.$$

So letting $n \rightarrow \infty$, we see that $\mathbb{D}'_q(u_n)$ holds.

4.2.2. *Proof of $\mathbb{D}_q(u_n)$.* This follows since, as in (3), we have decay of correlations of Hölder functions against bounded measurable functions and condition $\mathbb{D}_q(u_n)$ was designed to follow from sufficiently fast decay of correlations, as shown in [Fre13b, Proposition 5.2]. In order to compute the required rate of decay of correlations, which will impose a restriction on the domain of the parameter α , we recall here the above-mentioned result so that we can follow the computations closely.

Proposition 1 ([Fre13b, Proposition 5.2]). *Assume that \mathcal{X} is a compact subset of \mathbb{R}^d and $f : \mathcal{X} \rightarrow \mathcal{X}$ is a system with an acip \mathbb{P} , such that $\frac{d\mathbb{P}}{dLeb} \in L^{1+\epsilon}$. Assume, moreover, that the system has decay of correlations for all $\varphi \in \mathcal{H}_\beta$ against any $\psi \in L^\infty$ so that there exists some $C > 0$ independent of φ, ψ and t , and a rate function $\varrho : \mathbb{N} \rightarrow \mathbb{R}$ such that*

$$\left| \int \varphi \cdot (\psi \circ f^t) d\mathbb{P} - \int \varphi d\mathbb{P} \int \psi d\mathbb{P} \right| \leq C \|\varphi\|_{\mathcal{H}_\beta} \|\psi\|_\infty \varrho(t), \quad (18)$$

and $n^{1+\beta(1+\max\{0, (\epsilon+1)/\epsilon-d\}+\delta)} \varrho(t_n) \rightarrow 0$, as $n \rightarrow \infty$ for some $\delta > 0$ and $t_n = o(n)$. Then condition $\mathbb{D}_q(u_n)$ holds.

Remark 7. *We note that during the proof, in order to obtain the condition on the rate of decay of correlations, it is assumed that $\mathbb{P}(A_n^{(q)}) \sim_c 1/n$.*

Observe that since we are working in dimension 1, which means $d = 1$, then $\max\{0, (\epsilon + 1)/\epsilon - d\} = 1/\epsilon$. Also, from [Hu04], we may assume that the decay of correlations is written for Lipschitz functions, which allows us to take $\beta = 1$. Hence, for condition $\mathbb{D}_q(u_n)$ hold, we need that the rate of decay of correlations ϱ is sufficiently fast so that there exists some $\delta > 0$ such that

$$\lim_{n \rightarrow \infty} n^{2+1/\epsilon+\delta} \varrho(t_n) = 0, \quad (19)$$

where $t_n = o(n)$. From (4), in order that the density $h_\alpha \in L^{1+\epsilon}$, we need that $\epsilon < 1/\alpha - 1$. Since by (3), we have that $\varrho(t) = t^{-(1/\alpha-1)}$, then by (19) it is obvious that we must have $\alpha < 1/2$, which implies that $1/\epsilon < \alpha + 2\alpha^2$. Taking $t_n = n^{1-\alpha}$, we obtain:

$$n^{2+1/\epsilon+\delta} (n^{1-\alpha})^{-(1/\alpha-1)} = n^{2+1/\epsilon+\delta} n^{-1/\alpha+2-\alpha} < n^{4+2\alpha^2+\delta-1/\alpha}.$$

Hence, if $\alpha < \sqrt{5} - 2$ we can always find $\delta > 0$ so that (19) holds and consequently condition $\mathbb{D}_q(u_n)$ is verified.

4.2.3. Linear normalisation and extremal types limiting laws. As discussed in Remark 4, in classical extreme value theory (independent case), typically, the limiting laws are obtained for linear normalising sequences $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}}$, with $a_n > 0$ so that

$$\lim_{n \rightarrow \infty} \mathbb{P}(a_n(M_n - b_n) \leq y) = e^{-\tau(y)},$$

where $\tau(y)$ is of one of three types specified in [Fre13a, Equation (2.8)], where the tail of the distribution function of X_0 determines the type of limit. Note that $u_n = y/a_n + b_n$ and equation (2) imply that in the usual applications to dynamical systems, when no clustering occurs, the type of limit law is determined by the shape of φ near ζ as well as the shape of the invariant density at that point.

We note that the stochastic process defined as in (1), with $\zeta = 0$, may or may not have the same limiting extremal distribution (under linear normalisation) as the corresponding independent sequence, i.e., consider Z_0, Z_1, \dots i.i.d. and such that Z_0 as the same distribution as X_0 .

To illustrate the possibility of having different extremal limiting distributions, we give an example of φ for which $\mathbb{P}(a_n(M_n - b_n) \leq y) \rightarrow H(y)$ and $\mathbb{P}(a_n^*(M_n^* - b_n^*) \leq y) \rightarrow H^*(y)$ but $H(y) \neq H^*(y)$, where $M_n^* = \max\{Z_0, \dots, Z_{n-1}\}$.

Let $\varphi = 1 - x$. Then $\varphi^{-1}(x) = 1 - x$. Using (15) to define u_n , having in mind (14) and recalling that $Q_n = [x_n, y_n)$, where $y_n = \varphi^{-1}(u_n)$ and $x_n < y_n$ is such that $f_\alpha(x_n) = y_n$, we see that for a well chosen constant $c > 0$, we can take $u_n = 1 - c\tau/n$. Using Theorem 2 and writing $y = -\tau$, we get

$$\lim_{n \rightarrow \infty} \mu_\alpha(M_n \leq 1 - c\tau/n) = e^{-\tau} \iff \lim_{n \rightarrow \infty} \mu_\alpha\left(\frac{n}{c}(M_n - 1) \leq y\right) = e^{-(-y)},$$

which means we have Weibull limiting distribution with exponent equal to 1 and normalising sequences given by $a_n = \frac{n}{c}$ and $b_n = 1$.

If we are now to determine the limiting law for the sequence the i.i.d. sequence Z_0, Z_1, \dots , we have to use equation (2) to define u_n . Now since $U_n = \{X_0 > u_n\} = [0, y_n)$, where $y_n = \varphi^{-1}(u_n)$, then by (13), for a well chosen $c^* > 0$, we can take $u_n = 1 - \left(\frac{c^*\tau}{n}\right)^{1/(1-\alpha)}$. Then using [LLR83, Theorem 1.5.1] and letting $y = -\tau^{1/(1-\alpha)}$, we would get

$$\lim_{n \rightarrow \infty} \mathcal{P}\left(M_n^* \leq 1 - \left(\frac{c^*\tau}{n}\right)^{1/(1-\alpha)}\right) = e^{-\tau} \iff \lim_{n \rightarrow \infty} \mathbb{P}\left(\left(\frac{n}{c^*}\right)^{1/(1-\alpha)}(M_n^* - 1) \leq y\right) = e^{-(-y)^{1-\alpha}},$$

which means we have Weibull limiting distribution with exponent equal to $1 - \alpha$ and normalising sequences given by $a_n = \left(\frac{n}{c^*}\right)^{1/(1-\alpha)}$ and $b_n = 1$.

The same computations would allow us to verify that if $\varphi(x) = -\log(x)$ then both the extremal limiting distributions under linear normalisation for both M_n and M_n^* would be the Gumbel distribution, *i.e.*, $H(y) = H^*(y) = e^{-e^{-y}}$.

REFERENCES

- [AFV14] Hale Aytaç, Jorge Milhazes Freitas, and Sandro Vaienti, *Laws of rare events for deterministic and random dynamical systems*, Trans. Amer. Math. Soc. (2014), Published online with DOI: 10.1090/S0002-9947-2014-06300-9.
- [AG01] M. Abadi and A. Galves, *Inequalities for the occurrence times of rare events in mixing processes. The state of the art*, Markov Process. Related Fields **7** (2001), no. 1, 97–112, Inhomogeneous random systems (Cergy-Pontoise, 2000). MR 1835750
- [AS11] Miguel Abadi and Benoit Saussol, *Hitting and returning to rare events for all alpha-mixing processes*, Stochastic Process. Appl. **121** (2011), no. 2, 314–323. MR 2746177
- [AV08] Miguel Abadi and Nicolas Vergne, *Sharp errors for point-wise Poisson approximations in mixing processes*, Nonlinearity **21** (2008), no. 12, 2871–2885. MR 2461044 (2009m:60050)
- [AV09] ———, *Sharp error terms for return time statistics under mixing conditions*, J. Theoret. Probab. **22** (2009), no. 1, 18–37. MR 2472003 (2010f:60068)
- [BSTV03] H. Bruin, B. Saussol, S. Troubetzkoy, and S. Vaienti, *Return time statistics via inducing*, Ergodic Theory Dynam. Systems **23** (2003), no. 4, 991–1013. MR 1997964 (2005a:37004)
- [CC13] J.-R. Chazottes and P. Collet, *Poisson approximation for the number of visits to balls in non-uniformly hyperbolic dynamical systems*, Ergodic Theory Dynam. Systems **33** (2013), no. 1, 49–80. MR 3009103
- [CG93] P. Collet and A. Galves, *Statistics of close visits to the indifferent fixed point of an interval map*, J. Statist. Phys. **72** (1993), no. 3-4, 459–478. MR 1239564 (94i:58058)
- [CGS92] P. Collet, A. Galves, and B. Schmitt, *Unpredictability of the occurrence time of a long laminar period in a model of temporal intermittency*, Ann. Inst. H. Poincaré Phys. Théor. **57** (1992), no. 3, 319–331. MR 1185337 (94a:58064)
- [CI95] Massimo Campanino and Stefano Isola, *Statistical properties of long return times in type I intermittency*, Forum Math. **7** (1995), no. 3, 331–348. MR 1325560 (96c:60050)

- [Col01] P. Collet, *Statistics of closest return for some non-uniformly hyperbolic systems*, Ergodic Theory Dynam. Systems **21** (2001), no. 2, 401–420. MR MR1827111 (2002a:37038)
- [DGS04] Manfred Denker, Mikhail Gordin, and Anastasya Sharova, *A Poisson limit theorem for toral automorphisms*, Illinois J. Math. **48** (2004), no. 1, 1–20. MR 2048211 (2005a:37015)
- [FF08a] Ana Cristina Moreira Freitas and Jorge Milhazes Freitas, *Extreme values for Benedicks-Carleson quadratic maps*, Ergodic Theory Dynam. Systems **28** (2008), no. 4, 1117–1133. MR MR2437222 (2010a:37013)
- [FF08b] ———, *On the link between dependence and independence in extreme value theory for dynamical systems*, Statist. Probab. Lett. **78** (2008), no. 9, 1088–1093. MR MR2422964 (2009e:37006)
- [FFR] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, and Fagner Bernardini Rodrigues, *The speed of convergence of rare events point processes in non-uniformly hyperbolic systems*, in preparation.
- [FFT10] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, and Mike Todd, *Hitting time statistics and extreme value theory*, Probab. Theory Related Fields **147** (2010), no. 3-4, 675–710. MR 2639719 (2011g:37015)
- [FFT11] ———, *Extreme value laws in dynamical systems for non-smooth observations*, J. Stat. Phys. **142** (2011), no. 1, 108–126. MR 2749711 (2012a:60149)
- [FFT12] ———, *The extremal index, hitting time statistics and periodicity*, Adv. Math. **231** (2012), no. 5, 2626–2665. MR 2970462
- [FFT13] ———, *The compound Poisson limit ruling periodic extreme behaviour of non-uniformly hyperbolic dynamics*, Comm. Math. Phys. **321** (2013), 483–527.
- [FFT15] ———, *Speed of convergence for laws of rare events and escape rates*, Stochastic Process. Appl. **125** (2015), no. 4, 1653–1687.
- [FHN14] Jorge Milhazes Freitas, Nicolai Haydn, and Matthew Nicol, *Convergence of rare event point processes to the Poisson process for planar billiards*, Nonlinearity **27** (2014), no. 7, 1669–1687. MR 3232197
- [FP12] Andrew Ferguson and Mark Pollicott, *Escape rates for Gibbs measures*, Ergodic Theory Dynam. Systems **32** (2012), no. 3, 961–988.
- [Fre13a] Jorge Milhazes Freitas, *Extremal behaviour of chaotic dynamics*, Dyn. Syst. **28** (2013), no. 3, 302–332. MR 3170619
- [Fre13b] ———, *Extremal behaviour of chaotic dynamics*, Dyn. Syst. **28** (2013), no. 3, 302–332. MR 3170619
- [GHN11] Chinmaya Gupta, Mark Holland, and Matthew Nicol, *Extreme value theory and return time statistics for dispersing billiard maps and flows, Lozi maps and Lorenz-like maps*, Ergodic Theory Dynam. Systems **31** (2011), no. 5, 1363–1390. MR 2832250
- [HLV07] N. Haydn, E. Lunedei, and S. Vaienti, *Averaged number of visits*, Chaos **17** (2007), no. 3, 033119, 13. MR 2356973 (2008k:37005a)
- [HNT12] Mark Holland, Matthew Nicol, and Andrei Török, *Extreme value theory for non-uniformly expanding dynamical systems*, Trans. Amer. Math. Soc. **364** (2012), no. 2, 661–688. MR 2846347 (2012k:37064)
- [HSV99] Masaki Hirata, Benoît Saussol, and Sandro Vaienti, *Statistics of return times: a general framework and new applications*, Comm. Math. Phys. **206** (1999), no. 1, 33–55. MR 1736991 (2001c:37007)
- [Hu04] Huyi Hu, *Decay of correlations for piecewise smooth maps with indifferent fixed points*, Ergodic Theory Dynam. Systems **24** (2004), no. 2, 495–524. MR 2054191 (2005a:37064)
- [HV09] Nicolai Haydn and Sandro Vaienti, *The compound Poisson distribution and return times in dynamical systems*, Probab. Theory Related Fields **144** (2009), no. 3-4, 517–542. MR MR2496441
- [HWZ14] Nicolai Haydn, Nicole Winterberg, and Roland Zweimüller, *Return-time statistics, hitting-time statistics and inducing*, Ergodic theory, open dynamics, and coherent structures, Springer Proc. Math. Stat., vol. 70, Springer, New York, 2014, pp. 217–227. MR 3213501
- [Kal86] Olav Kallenberg, *Random measures*, fourth ed., Akademie-Verlag, Berlin, 1986. MR 854102 (87k:60137)
- [Kel12] Gerhard Keller, *Rare events, exponential hitting times and extremal indices via spectral perturbation*, Dynamical Systems **27** (2012), no. 1, 11–27.
- [KR14] Yuri Kifer and Ariel Rapaport, *Poisson and compound Poisson approximations in conventional and nonconventional setups*, Probab. Theory Related Fields **160** (2014), no. 3-4, 797–831. MR 3278921
- [Lea74] M. R. Leadbetter, *On extreme values in stationary sequences*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **28** (1973/74), 289–303. MR 0362465 (50 #14906)
- [LFW12] Valerio Lucarini, Davide Faranda, and Jeroen Wouters, *Universal behavior of extreme value statistics for selected observables of dynamical systems*, J. Stat. Phys. **147** (2012), no. 1, 63–73.

- [LLR83] M. R. Leadbetter, Georg Lindgren, and Holger Rootzén, *Extremes and related properties of random sequences and processes*, Springer Series in Statistics, Springer-Verlag, New York, 1983. MR MR691492 (84h:60050)
- [LSV99] Carlangelo Liverani, Benoît Saussol, and Sandro Vaienti, *A probabilistic approach to intermittency*, Ergodic Theory Dynam. Systems **19** (1999), no. 3, 671–685. MR MR1695915 (2000d:37029)
- [Pit91] B. Pitskel', *Poisson limit law for Markov chains*, Ergodic Theory Dynam. Systems **11** (1991), no. 3, 501–513. MR 1125886
- [Ryc83] Marek Rychlik, *Bounded variation and invariant measures*, Studia Math. **76** (1983), no. 1, 69–80. MR MR728198 (85h:28019)
- [VHF09] Renato Vitolo, Mark P. Holland, and Christopher A. T. Ferro, *Robust extremes in chaotic deterministic systems*, Chaos **19** (2009), no. 4, 043127.
- [You99] Lai-Sang Young, *Recurrence times and rates of mixing*, Israel J. Math. **110** (1999), 153–188. MR MR1750438 (2001j:37062)
- [Zwe07] Roland Zweimüller, *Mixing limit theorems for ergodic transformations*, J. Theoret. Probab. **20** (2007), no. 4, 1059–1071. MR 2359068 (2008h:60119)

ANA CRISTINA MOREIRA FREITAS, CENTRO DE MATEMÁTICA & FACULDADE DE ECONOMIA DA UNIVERSIDADE DO PORTO, RUA DR. ROBERTO FRIAS, 4200-464 PORTO, PORTUGAL

E-mail address: amoreira@fep.up.pt

JORGE MILHAZES FREITAS, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL

E-mail address: jmfreira@fc.up.pt

URL: <http://www.fc.up.pt/pessoas/jmfreira>

MIKE TODD, MATHEMATICAL INSTITUTE, UNIVERSITY OF ST ANDREWS, NORTH HAUGH, ST ANDREWS, KY16 9SS, SCOTLAND,

E-mail address: m.todd@st-andrews.ac.uk

URL: <http://www.mcs.st-and.ac.uk/~miket/>

SANDRO VAIENTI, AIX MARSEILLE UNIVERSITÉ, CNRS, CPT, UMR 7332, 13288 MARSEILLE, FRANCE AND UNIVERSITÉ DE TOULON, CNRS, CPT, UMR 7332, 83957 LA GARDE, FRANCE

E-mail address: vaienti@cpt.univ-mrs.fr