

# Deglaciation chronology of the Donegal Ice Centre, north-west Ireland

PETER WILSON,<sup>1\*</sup> COLIN K. BALLANTYNE,<sup>2</sup> SARA BENETTI,<sup>1</sup> DAVID SMALL,<sup>3</sup> DEREK FABEL<sup>4</sup> and CHRIS D. CLARK<sup>5</sup>

<sup>1</sup>School of Geography and Environmental Sciences, Ulster University, Coleraine BT52 1SA, UK

<sup>2</sup>School of Geography and Sustainable Development, University of St. Andrews, St Andrews KY16 9AL, UK

<sup>3</sup>Department of Geography, University of Durham, Durham, DH1 3LE, UK

<sup>4</sup>Scottish Universities Environmental Research Centre, East Kilbride G75 0QF, UK

<sup>5</sup>Department of Geography, University of Sheffield, Sheffield S10 2TN, UK

Received 25 June 2018; Revised 1 October 2018; Accepted 18 October 2018

**ABSTRACT:** During the Last Glacial Maximum, Donegal in north-west Ireland functioned as an independent centre of ice dispersal that separated and fed into the Donegal Bay Ice Lobe (sourced in the Irish Midlands) to the south and the Hebrides/Malin Sea Ice Stream to the north. We report geochronological data that demonstrate marked contrasts in the timing and rate of deglaciation in northern and southern Donegal. In northern Donegal, which occupied an inter-ice-stream/lobe location, decoupling from the Hebrides/Malin Sea Ice Stream resulted in formation of a marine embayment along the north coast by  $\sim 22\text{--}21$  ka, and subsequent slow ( $\sim 4 \pm 1$  m a<sup>-1</sup>) climatically driven inland retreat of the ice margin to mountain source areas by  $\sim 17$  ka. By contrast, in southern Donegal, which lay near the axis of the Donegal Bay Ice Lobe, deglaciation was delayed until  $\sim 18$  ka following readvance of ice to a moraine in outer Donegal Bay. The ice margin subsequently underwent net retreat, apparently uninterrupted by readvances, at a net rate of  $\sim 18 \pm 6$  m a<sup>-1</sup>. A mean terrestrial cosmogenic nuclide age of  $\sim 15.0$  ka obtained for samples from the foothills of the Blue Stack Mountains in south-east Donegal indicates that ice persisted in valley heads and cirques at the beginning of the Lateglacial Interstadial, suggesting that these and nearby mountains supported the last remnants of the Irish Ice Sheet before complete deglaciation of Ireland, and that almost all the shrinkage of the ice sheet in this sector occurred under stadial conditions before the onset of interstadial warming at  $\sim 14.7$  ka.

© 2018 The Authors. *Journal of Quaternary Science* Published by John Wiley & Sons Ltd.

**KEYWORDS:** British–Irish Ice Sheet; deglaciation; north-west Ireland; terrestrial cosmogenic nuclide surface exposure dating.

## Introduction

During the last (Late Devensian/Late Midlandian) ice-sheet glaciation of Britain and Ireland ( $\sim 32\text{--}15$  ka), the mountains of Donegal in north-west Ireland formed an independent centre of ice dispersal within the more extensive British–Irish Ice Sheet (BIIS). Ice radiating from the Donegal Ice Centre fed north and north-west into a major ice stream (the Hebrides/Malin Sea Ice Stream) on the adjacent Malin Shelf, west and south-west into Donegal Bay, and to the east was confluent with the ice occupying the Irish Midlands. The Donegal ice dome was therefore pivotal in separating ice flows from the Irish Midlands and western Scotland (Fig. 1). Radial ice flow over Donegal for at least part of the last glaciation is demonstrated by the distribution of local erratics, absence of allochthonous erratics, and the alignments of drumlins, moraines, roches moutonnées, striae and meltwater channels (Charlesworth, 1924; Dury, 1957, 1958, 1964; Stephen and Synge, 1965; Colhoun, 1973; McCabe *et al.*, 1993; Knight and McCabe, 1997; Ballantyne *et al.*, 2007; Smith and Knight, 2011; Knight, 2012). Flowsets reconstructed by Greenwood and Clark (2009a,b) show that ice moving south from the Donegal Ice Centre was confluent with west-flowing ice from the Irish Midlands in Donegal Bay, forming an ice lobe (the Donegal Bay Ice Lobe; Ó Cofaigh *et al.*, 2012) that extended north-westwards towards the shelf edge.


Geophysical data obtained for the adjacent offshore shelves indicate that at the global Last Glacial Maximum

(gLGM; 26.5–19 ka, Clark *et al.*, 2009a) grounded ice extended as far as the shelf break,  $\sim 100$  km to the west, where it terminated in a marine setting (Benetti *et al.*, 2010; Dunlop *et al.*, 2010; Ó Cofaigh *et al.*, 2012). These data also indicate that Donegal ice coalesced with ice from western Scotland  $\sim 60$  km north of the present Donegal coastline. Donegal is therefore the key location for determining the timing of decoupling of Irish- and Scottish-sourced ice during the last deglaciation, and is also important for establishing the chronology of ice retreat after the ice margin had retreated to the present coastline.

As part of the wider BRITICE-CHRONO project (<http://www.sheffield.ac.uk/geography/research/britice-chrono/home>), designed to establish a detailed deglaciation chronology of the last BIIS, we present 20 new <sup>10</sup>Be and <sup>36</sup>Cl terrestrial cosmogenic nuclide (TCN) surface exposure ages from six sites in Donegal that were selected to complement and extend the existing deglaciation chronology. The aims of this paper are: (i) to establish the timing of the decoupling of Scottish and Irish ice flowing west across the Malin Shelf; (ii) to reconstruct the chronology of ice margin retreat in Donegal Bay; (iii) to determine the net rate of ice margin recession inland from the north coast of Donegal and in Donegal Bay; (iv) to establish for how long ice persisted locally in the Donegal mountains following its retreat from coastal lowlands; and (v) to explore the wider implications of our results for the interpretation of the deglaciation chronology of the western sector of the last BIIS. The chronology of offshore ice margin retreat from the shelf edge towards the present coastline is considered in a separate paper based on new

\*Correspondence: Peter Wilson, as above.

E-mail: p.wilson@ulster.ac.uk



**Figure 1.** Location map of Donegal and Mayo with terrestrial legacy ages (TCN and  $^{14}\text{C}$ ), BRITICE-CHRONO  $^{14}\text{C}$  ages for the Donegal Bay Moraine, with core numbers, and sites of new TCN ages reported in this paper. Only those legacy ages of relevance to deglaciation are shown. The first TCN age for each site was calculated using version 3.0 of the online exposure age calculator formerly known as the CRONUS-Earth online exposure age calculator with the LLPR; the age in parentheses was calculated using CRONUS-calc 2.0. Ages are mean values of two or more ages for each site except for Glencolumbkille, which is represented by a single age. Terrestrial  $^{14}\text{C}$  ages are minimum ages for deglaciation. Ice divides and generalized ice flow directions are from Greenwood and Clark (2009a). Inset shows location of Donegal and the LGM limit of the BIIS.

radiocarbon ages obtained from marine microfauna retrieved from sediment cores along a transect from the shelf edge to the outer part of Donegal Bay (Ó Cofaigh *et al.*, 2019).

## Donegal

### Regional setting and ice dome extent

County Donegal ( $54^{\circ}28'–55^{\circ}22' \text{ N}$ ,  $06^{\circ}55'–08^{\circ}46' \text{ W}$ ) is predominantly underlain by granites, quartzites and schists with a pronounced NE–SW structural grain that has been accentuated by repeated episodes of Quaternary glacial erosion (Long and McConnell, 1997, 1999). The north and west of the county are mountainous; many summits exceed 500 m OD with the highest point (Errigal) at 751 m OD. From detailed mapping of erosional and depositional landforms, Charlesworth (1924) proposed that the Donegal mountains had nourished and maintained an independent ice dome during the last glaciation, and placed the former ice divide along a line running approximately north–south from the Derryveagh Mountains to the Blue Stack Mountains, close to the present watershed (Fig. 1). He showed that ice-flow from this elongated dome was essentially radial with a focus along pre-existing structurally controlled valleys, but argued that during maximum ice extent topography was probably less of a constraining influence on ice-flow directions than during build-up and retreat phases. Subsequent work is generally supportive of this ice dome hypothesis (e.g. Ballantyne *et al.*, 2007; Greenwood and Clark, 2009a, b; Smith and Knight, 2011).

In contrast, the thickness attained by the ice dome has been a contentious issue. Charlesworth (1924) claimed that all summits lay beneath the ice, although it is not clear if he was referring to the local Last Glacial Maximum (ILGM), which is placed at  $\sim 26.3–24.8$  ka at the shelf break to the west of Donegal (Ó Cofaigh *et al.*, 2019). Complete burial of the mountains by the last ice sheet was favoured by McCabe (1995), while Sellier (1995) maintained that areas above  $\sim 550$  m OD in the Derryveagh Mountains had remained ice free. On the basis of geomorphological evidence and clay-fraction mineralogy, Ballantyne *et al.* (2007) argued for an

ice-shed altitude in excess of 700 m OD, but also reported an absence of evidence for glacial modification on six peripheral summits, including Errigal, and regarded these as being either nunataks during the ILGM or buried beneath a cover of non-erosive cold-based ice.

Conflicting interpretations also concern the lateral extent of the Donegal ice dome. Although Charlesworth (1924) envisaged ice extending offshore to the north and west, others have placed the limit onshore in the north of the county (Stephen and Synge, 1965; Bowen *et al.*, 2002), and a limited offshore extent of  $\sim 10–30$  km to the west has been suggested (e.g. McCabe, 1985; Bowen *et al.*, 1986; Knight, 2003; Ballantyne *et al.*, 2007). More recent work utilizing geophysical techniques to image seabed topography has demonstrated that a concentric sequence of nested moraine ridges indicative of deposition by a grounded ice mass extends westwards to the shelf break 90–100 km from the west coast of Donegal (Sejrup *et al.*, 2005; Benetti *et al.*, 2010; Dunlop *et al.*, 2010; Ó Cofaigh *et al.*, 2012). Radiocarbon dates obtained for marine microfauna in cores retrieved from sediments on the Atlantic shelf north-west of Ireland confirm that these moraines were deposited at the margin of the last ice sheet, and indicate that ice nourished in Donegal began to retreat from the shelf edge in the interval between 26.3 and 24.8 ka cal BP (Ó Cofaigh *et al.*, 2019).

Recognition that the last ice sheet extended to the edge of the Malin Shelf strongly suggests that the Donegal ice dome was of sufficient thickness to have buried all mountain summits during the ILGM, a proposition also supported by climate-proxy-driven thermomechanical models of ice-sheet build-up and decay (Hubbard *et al.*, 2009). Support for this premise comes from south-west Ireland where Ballantyne *et al.* (2011) have argued that the Kerry–Cork Ice Cap attained an altitude of at least 1200 m OD,  $>200$  m above the highest summits, and from north-west Scotland where Fabel *et al.* (2012) have demonstrated that the last ice sheet overtopped all mountain summits. It is therefore extremely unlikely that any of the mountain summits in Donegal formed palaeonunataks during the ILGM (Ballantyne and Ó Cofaigh, 2017).

## Legacy ages, related BRITICE-CHRONO ages and deglaciation

Several previous studies have used either TCN (cosmogenic  $^{10}\text{Be}$  or  $^{36}\text{Cl}$ ) surface exposure dating or  $^{14}\text{C}$  dating to establish the timing of ice retreat and/or readvance from sites in Donegal. Collectively these ages provide the foundation of a deglaciation chronology (Fig. 1; Table 1). Cosmogenic  $^{10}\text{Be}$  exposure ages cited here have been recalculated using the local Loch Lomond production rate (LLPR), and are followed in parentheses by ages obtained from the CRONUScalc online calculator using a global reference production rate; details of these procedures are given in the next section. Insufficient information was available to recalculate the two  $^{36}\text{Cl}$  ages, and we cite them as published by the original authors. The  $^{14}\text{C}$  ages have been (re)calibrated using OxCal 4.2 and, for marine-derived samples, the Marine-13 curve with a marine reservoir correction of 400 years (Bronk Ramsey, 2009; Reimer *et al.*, 2013). All  $^{14}\text{C}$  ages are reported to two decimal places as cal ka BP. TCN ages are reported to one decimal place as ka. Age uncertainties are  $\pm 1\sigma$ . Mean ages reported for two or more TCN ages below and in Tables 1 and 3 are uncertainty-weighted means.

Bowen *et al.* (2002) obtained  $^{36}\text{Cl}$  ages of  $25.1 \pm 1.1$  ka, from glacially smoothed quartzite bedrock at Malin Head, and  $31.0 \pm 17.0$  ka, from either a glacially transported granite boulder or bedrock at Bloody Foreland, but the large uncertainty on the latter age prevents meaningful interpretation, and the former is probably compromised by nuclide inheritance (Ballantyne and Ó Cofaigh, 2017). For Corvish, at the head of Trawbreaga Bay on the north coast, McCabe and Clark (2003) reported  $^{14}\text{C}$  ages for marine microfossils within *in situ* and deformed marine sediments. The basal *in situ* laminated muds yielded ages of  $20.68 \pm 0.16$  and  $18.24 \pm 0.13$  cal ka BP; the older date implies initial deglaciation before  $\sim 20.7$  cal ka BP. Overlying deformed sands and muds gave ages of  $19.50 \pm 0.50$ ,  $18.32 \pm 0.18$  and  $19.03 \pm 0.19$  cal ka BP, and were interpreted by McCabe and Clark (2003) as evidence for reworking of the underlying laminated muds by ice readvance at  $\sim 18$  ka. An age of  $17.06 \pm 0.18$  cal ka BP from *in situ* rhythmically bedded marine muds overlying the deformed muds was regarded as minimal for final deglaciation of the bay.

Seven consistent  $^{10}\text{Be}$  exposure ages from glacially transported boulders on a lateral moraine at Bloody Foreland, the north-westernmost point of Donegal, have given an uncertainty-weighted mean age of  $21.6 \pm 0.7$  ka ( $21.7 \pm 1.8$  ka) (Ballantyne *et al.*, 2007; Clark *et al.*, 2009b; Ballantyne and Ó Cofaigh, 2017; Fig. 2a). Two  $^{10}\text{Be}$  exposure ages from bedrock and a glacially transported boulder on Aran Island, 20 km south-west of Bloody Foreland, have yielded an uncertainty-weighted mean age of  $21.7 \pm 0.8$  ka ( $21.5 \pm 1.8$  ka) (Cullen, 2013). The consistency of these two mean ages provides strong support for retreat of the ice-sheet margin across the Irish sector of the Malin Shelf between  $\sim 26$ – $25$  and  $\sim 22$ – $21$  ka (Clark *et al.*, 2012a; Ó Cofaigh *et al.*, 2012, 2019), decoupling of Malin Shelf ice from Donegal Bay ice at  $\sim 22$ – $21$  ka and the beginning of ice retreat at that time from the present coast towards the mountains. At Glencolumbkille in south-west Donegal,  $^{10}\text{Be}$  exposure ages of  $17.8 \pm 0.6$  ka ( $17.9 \pm 1.5$  ka) and  $19.6 \pm 0.7$  ka ( $19.8 \pm 1.7$  ka) from vein quartz in, respectively, a glacially transported boulder and a roche moutonnée were reported by Ballantyne *et al.* (2007). The latter age overlaps within  $1\sigma$  uncertainties with the mean values from Bloody Foreland and Aran Island, but may be compromised by nuclide inheritance (see below).

The timing of deglaciation of the mountains of Donegal is indicated by  $^{10}\text{Be}$  exposure ages for two sites. Glacially plucked bedrock at 405–430 m OD on a col to the east of Errigal in north Donegal has produced three consistent  $^{10}\text{Be}$  exposure ages averaging  $18.0 \pm 0.6$  ka ( $17.8 \pm 1.4$  ka) and a minimum age for deglaciation of Slieve League in south-west Donegal is provided by three consistent  $^{10}\text{Be}$  exposure ages averaging  $17.3 \pm 0.6$  ka ( $17.1 \pm 1.5$  ka) obtained for samples from rockslide run out debris (Ballantyne *et al.*, 2013b).

A  $^{14}\text{C}$  age of  $15.38 \pm 0.12$  cal ka BP from the basal organic mud of Lough Nadourcan (Watson *et al.*, 2010) provides a minimum age for deglaciation of the low ground along the eastern margin of the Derryveagh Mountains. However, this age is  $\sim 700$  years earlier than the rapid warming identified in the Greenland ice core records and INTIMATE event stratigraphy as marking the onset of the Lateglacial Interstadial at  $\sim 14.7$  ka (Rasmussen *et al.*, 2014), suggesting that the Lough Nadourcan basal  $^{14}\text{C}$  age may be compromised by the incorporation of reworked older carbon. Nevertheless, it is unlikely that ice on low ground survived much beyond the start of interstadial warming even if small glaciers persisted in the mountains.

Legacy ages from sites in north Mayo, along the south side of Donegal Bay, and BRITICE-CHRONO ages from Donegal Bay (Fig. 1) are relevant to the deglaciation chronology of south Donegal, and therefore are also considered here. McCabe *et al.* (1986, 2005) reported eight  $^{14}\text{C}$  ages obtained for marine shells and foraminifera within glacial marine sediments at Fiddauntawnanoneen and Belderg Pier on the north coast of Mayo. Seven of these ages range between  $20.38 \pm 0.31$  and  $19.16 \pm 0.21$  cal ka BP; the remaining age ( $22.09 \pm 0.28$  cal ka BP) is significantly older and may indicate the reworking of older sediment (Clark *et al.*, 2012b). Deglaciation of these adjacent sites and, by inference, the outer reaches of Donegal Bay therefore appears to have occurred around or slightly before  $\sim 20$  ka (Ballantyne and Ó Cofaigh, 2017).

To the north-east of these two sites, a distinct ice margin position is represented by the Donegal Bay Moraine (DBM), an offshore moraine that extends for 35 km north–south across outer Donegal Bay (Benetti *et al.*, 2010; Ó Cofaigh *et al.*, 2012). Deformation of stratified glacial marine deposits indicates that the moraine represents a readvance of the ice margin. Radiocarbon ages for mixed benthic foraminifera within glacial marine sediments in 76–99 m water depth on either side of the moraine (Fig. 1) constrain moraine formation to between  $20.24 \pm 0.24$  and  $17.92 \pm 0.16$  cal ka BP (Ó Cofaigh *et al.*, 2019), and moraine formation at 20–19 ka was inferred by Ó Cofaigh *et al.* (2019).

Finally, eight cosmogenic  $^{10}\text{Be}$  exposure ages from vein quartz in glacially transported boulders at three sites associated with the Tawnywaddyduff moraine system on the northern slopes of the Ox Mountains south of Sligo Bay (Fig. 1) returned ages ranging from  $20.9 \pm 1.5$  ka ( $21.1 \pm 2.3$  ka) to  $15.7 \pm 1.5$  ka ( $16.0 \pm 2.0$  ka). The overall average of these ages ( $\sim 18$  ka) was taken by Clark *et al.* (2009c) to represent the timing of a readvance of the ice sheet and construction of the moraine. However, Ballantyne and Ó Cofaigh (2017) questioned this conclusion, noting that the age range spanned  $>5$  ka and that two distinct age groupings are represented, with three older ages [mean  $20.2 \pm 1.1$  ka ( $20.3 \pm 1.9$  ka)] and five younger ages [mean  $16.6 \pm 0.6$  ka ( $16.7 \pm 1.5$  ka)]. The older sample ages are from a site on the west side of the Ox Mountains and may be compromised by nuclide inheritance as they are inconsistent with the wider dating evidence; four of the younger ages are from the east side and the other age came from the northern slopes.

**Table 1.** Terrestrial legacy ages and BRITICE-CHRONO ages pertaining to the deglaciation of Donegal, Donegal Bay and north Mayo.

Site	<sup>14</sup> C age ( $\pm 1\sigma$ , cal ka BP) <sup>1</sup>	<sup>10</sup> Be age ( $\pm 1\sigma$ , LLPR) <sup>2</sup>	<sup>10</sup> Be age ( $\pm 1\sigma$ , CRONUScal) <sup>3</sup>	<sup>36</sup> Cl age ( $\pm 1\sigma$ ) <sup>4</sup>	Material and context	Reference
<b>DONEGAL</b>						
Malin Head	17.06 $\pm$ 0.18			25.1 $\pm$ 1.1	Glacially smoothed quartzite bedrock	Bowen <i>et al.</i> (2002)
Bloody Foreland	19.03 $\pm$ 0.19			31.0 $\pm$ 17.0	Not specified, but granite bedrock or boulder	Bowen <i>et al.</i> (2002)
Corvish	18.32 $\pm$ 0.18				Marine microfauna: <i>Elphidium clavatum</i>	McCabe and Clark (2003)
	19.50 $\pm$ 0.50				Marine microfauna: <i>Elphidium clavatum</i>	
	18.24 $\pm$ 0.13				Marine microfauna: <i>Elphidium clavatum</i>	
	20.68 $\pm$ 0.16				Marine microfauna: <i>Elphidium clavatum</i>	
Bloody Foreland		21.2 $\pm$ 1.1 (1.0)	21.0 $\pm$ 2.0 (1.1)		Glacially transported granite boulder	Ballantyne <i>et al.</i> (2007)
		18.5 $\pm$ 0.9 (0.8)	18.6 $\pm$ 1.7 (0.9)		Glacially transported granite boulder	
		17.9 $\pm$ 1.7 (1.6)	18.0 $\pm$ 2.3 (1.8)		Glacially transported granite boulder	Clark <i>et al.</i> (2009b)
		33.5 $\pm$ 2.7 (2.6)	34.0 $\pm$ 4.0 (2.9)		Glacially transported granite boulder	
		21.8 $\pm$ 1.6 (1.5)	22.0 $\pm$ 2.4 (1.7)		Glacially transported granite boulder	
		21.2 $\pm$ 1.7 (1.6)	21.4 $\pm$ 2.5 (1.8)		Glacially transported granite boulder	
		21.2 $\pm$ 1.9 (1.9)	21.4 $\pm$ 2.7 (2.1)		Glacially transported granite boulder	
		23.6 $\pm$ 2.0 (1.9)	23.8 $\pm$ 2.8 (2.1)		Glacially transported granite boulder	
		21.7 $\pm$ 2.1 (2.0)	21.9 $\pm$ 2.8 (2.2)		Glacially transported granite boulder	
		22.1 $\pm$ 2.0 (2.0)	22.3 $\pm$ 2.8 (2.2)		Glacially transported granite boulder	
		<b>21.6 <math>\pm</math> 0.7</b>	<b>21.7 <math>\pm</math> 1.8</b>			
Aran Island		21.8 $\pm$ 0.9 (0.7)	21.6 $\pm$ 1.9 (0.7)		Glacially transported granite boulder	Cullen (2013)
		21.5 $\pm$ 0.9 (0.7)	21.3 $\pm$ 1.8 (0.7)		Granite bedrock	
		<b>21.7 <math>\pm</math> 0.8</b>	<b>21.5 <math>\pm</math> 1.8</b>			
Glencolumbkille <sup>7</sup>		17.8 $\pm$ 0.6 (0.5)	17.9 $\pm$ 1.5 (0.5)		Vein quartz in glacially transported schist boulder	Ballantyne <i>et al.</i> (2007)
		19.6 $\pm$ 0.7 (0.5)	19.8 $\pm$ 1.7 (0.6)		Vein quartz in schist roche moutonnée	
Errigal col		17.6 $\pm$ 0.8 (0.6)	17.4 $\pm$ 1.5 (0.6)		Glacially plucked quartzite bedrock	Ballantyne <i>et al.</i> (2013b)
		18.2 $\pm$ 0.7 (0.6)	18.0 $\pm$ 1.5 (0.6)		Glacially plucked quartzite bedrock	
		18.1 $\pm$ 0.8 (0.6)	17.9 $\pm$ 1.6 (0.6)		Glacially plucked quartzite bedrock	
		<b>18.0 <math>\pm</math> 0.6</b>	<b>17.8 <math>\pm</math> 1.4</b>			
Slieve League		17.1 $\pm$ 0.8 (0.7)	16.9 $\pm$ 1.5 (0.7)		Quartzite boulder from rockslope-failure debris	Ballantyne <i>et al.</i> (2013b)
		17.8 $\pm$ 1.0 (0.9)	17.6 $\pm$ 1.7 (0.9)		Quartzite boulder from rockslope-failure debris	
		17.1 $\pm$ 1.0 (0.9)	16.9 $\pm$ 1.6 (0.9)		Quartzite boulder from rockslope-failure debris	
		<b>17.3 <math>\pm</math> 0.6</b>	<b>17.1 <math>\pm</math> 1.5</b>			
Lough Nadourcan	15.38 $\pm$ 0.12				Organic mud, basal lake sediment, bulk sample	Watson <i>et al.</i> (2010)
<b>NORTH MAYO</b>						
Fiddauntawnanoneen	20.38 $\pm$ 0.31				Marine shell: <i>Macoma calcaria</i>	McCabe <i>et al.</i> (1986)
Belderg Pier	19.16 $\pm$ 0.21				Marine shell: <i>Macoma calcaria</i>	McCabe <i>et al.</i> (1986, 2005)
	19.23 $\pm$ 0.26				Marine shell: <i>Macoma calcaria</i>	
	19.51 $\pm$ 0.29				Marine microfauna: <i>Elphidium clavatum</i>	
	19.77 $\pm$ 0.35				Marine microfauna: <i>Elphidium clavatum</i>	
	19.88 $\pm$ 0.33				Marine shell: <i>Macoma calcaria</i>	
	19.92 $\pm$ 0.34				Marine shell: <i>Macoma calcaria</i>	
	22.09 $\pm$ 0.28				Marine microfauna: <i>Quinqueloculina seminulum</i>	Ó Cofaigh <i>et al.</i> (2019)
	20.24 $\pm$ 0.24				Mixed benthic foraminifera	
	17.92 $\pm$ 0.16				Vein quartz in glacially transported gneissic boulder	Clark <i>et al.</i> (2009c)
Donegal Bay		16.9 $\pm$ 1.4 (1.4)	17.0 $\pm$ 2.0 (1.5)		Vein quartz in glacially transported gneissic boulder	
Ox Mountains		15.7 $\pm$ 1.5 (1.4)	16.0 $\pm$ 2.0 (1.6)		Vein quartz in glacially transported gneissic boulder	
		16.4 $\pm$ 1.3 (1.3)	16.4 $\pm$ 1.9 (1.4)		Vein quartz in glacially transported gneissic boulder	

Table 1. (Continued)

Site	$^{14}\text{C}$ age ( $\pm 1\sigma$ , cal ka BP) <sup>1</sup>	$^{10}\text{Be}$ age ( $\pm 1\sigma$ , LLPR) <sup>2</sup>	$^{10}\text{Be}$ age ( $\pm 1\sigma$ , CRONUScalc) <sup>3</sup>	$^{36}\text{Cl}$ age ( $\pm 1\sigma$ ) <sup>4</sup>	Material and context	Reference
	16.9 $\pm$ 1.3 (1.2)	16.9 $\pm$ 1.3 (1.2)	16.9 $\pm$ 1.9 (1.4)		Vein quartz in glacially transported gneissic boulder	
	17.0 $\pm$ 1.7 (1.7)	17.0 $\pm$ 1.7 (1.7)	17.0 $\pm$ 2.3 (1.8)		Vein quartz in glacially transported gneissic boulder	
<b>Mean<sup>5</sup></b>	<b>16.6 <math>\pm</math> 0.6</b>	<b>16.6 <math>\pm</math> 0.6</b>	<b>16.7 <math>\pm</math> 1.5</b>		Vein quartz in glacially transported gneissic boulder	Clark <i>et al.</i> (2009c)
	19.1 $\pm$ 1.6 (1.5)	19.1 $\pm$ 1.6 (1.5)	19.1 $\pm$ 2.3 (1.7)		Vein quartz in glacially transported gneissic boulder	
	20.9 $\pm$ 1.5 (1.4)	20.9 $\pm$ 1.5 (1.4)	21.1 $\pm$ 2.3 (1.6)		Vein quartz in glacially transported gneissic boulder	
	20.5 $\pm$ 1.9 (1.8)	20.5 $\pm$ 1.9 (1.8)	20.7 $\pm$ 2.6 (2.0)		Vein quartz in glacially transported gneissic boulder	
<b>Mean<sup>6</sup></b>	<b>20.2 <math>\pm</math> 1.1</b>	<b>20.2 <math>\pm</math> 1.1</b>	<b>20.3 <math>\pm</math> 1.9</b>			

<sup>1</sup>Ages calculated using OxCal 4.2; Marine-13 dataset used for Corvish and Donegal Bay samples (Bronk Ramsey, 2009; Reimer *et al.*, 2013).

<sup>2</sup>Exposure age based on v 3.0 of the online exposure age calculator formerly known as the CRONUS-Earth online exposure age calculators (wrapper: 3.0-dev; get\_age: 3.0.2-dev; muons: 1A, alpha = 1; conists: 3.0.3-dev) with the Loch Lomond production rate (LLPR), LM scaling, and assuming 1 mm ka<sup>-1</sup> erosion. Uncertainties are total (external) uncertainties. Analytical uncertainties are given in parentheses.

<sup>3</sup>Exposure age based on v 2.0 of CRONUScalc with LM scaling and assuming 1 mm ka<sup>-1</sup> surface erosion rate. Uncertainties are total (external) uncertainties. Analytical uncertainties are given in parentheses.

<sup>4</sup>Chlorine-36 ages cited as published by original authors.

<sup>5</sup>Mean derived from six consistent values reported by Clark *et al.* (2009b) and one age reported by Ballantyne *et al.* (2007).

<sup>6</sup>Uncertainty-weighted mean.

<sup>7</sup>Site is called Malin Beg in Ballantyne *et al.* (2007).

## Field sites and methods

The six sites sampled for TCN surface exposure dating were selected to provide deglaciation-age transects along the north and south coasts of Donegal and an additional deglaciation age for the northern (Derryveagh) mountains (Fig. 1). For the north coast transect we sampled on the headlands of Rosguill and Malin Head, respectively 29 and 63 km north-east of the Bloody Foreland site dated by Ballantyne *et al.* (2007) and Clark *et al.* (2009b). For the southern transect we sampled at Glencolumbkille, close to the western extremity of the Slieve League peninsula, at Kilcar on the south-west coast of Donegal, and on the lower southern slopes of the Blue Stack Mountains. The latter two sites are, respectively, 14 km south-east and 41 km east of Glencolumbkille. In the northern mountains we obtained samples from a prominent valley-floor boulder limit in the Poisoned Glen, Derryveagh Mountains.

Samples were collected from the upper surface of large, glacially deposited boulders or ice-scoured bedrock using a hammer and chisel. Twelve boulder samples comprised whole rock (granite, conglomerate sandstone or dolerite), four were from protruding quartz veins in quartzite or schist boulders, two consisted of quartz pebbles embedded in conglomerate boulders, and two samples were from quartzite bedrock (Fig. 2; Table 2). A compass and clinometer were used to record the geometry of the sampled surfaces and the skyline topography. Locations and altitudes were determined with a handheld GPS unit cross-referenced to a 1: 50 000 topographic map. Sample thickness was measured using callipers, density was determined by the displacement of sub-samples in water, and topographic shielding was calculated using the online calculators formerly known as the CRONUS-Earth online calculators (Table 2; Balco *et al.*, 2008: <http://hess.ess.washington.edu/math/>).

Samples were processed for cosmogenic  $^{10}\text{Be}$  and  $^{36}\text{Cl}$  analysis at the NERC Cosmogenic Isotope Analysis Facility (CIAF). For  $^{10}\text{Be}$ , samples were crushed and sieved to 250–500  $\mu\text{m}$  and quartz was separated in a Frantz isodynamic magnetic mineral separator, before being repeatedly etched with HF (Kohl and Nishiizumi, 1992). Purified quartz was spiked with  $\sim 0.22$  mg of  $^9\text{Be}$  and dissolved. Be was extracted and isolated following the methodology described in Child *et al.* (2000) before being precipitated as  $\text{Be}(\text{OH})_2$  and baked to  $\text{BeO}$  in a quartz crucible.  $\text{BeO}$  was mixed with Nb and pressed into a copper cathode. For  $^{36}\text{Cl}$ , samples were crushed and sieved to  $< 500 \mu\text{m}$ , leached in hot  $\text{HNO}_3$  (trace metal analysis grade) and then washed thoroughly with ultrapure water to remove meteoric  $^{36}\text{Cl}$  contamination from grain surfaces. Each sample was then split into two fractions: about 2 g for elemental analysis by inductively coupled plasma (ICP) optical emission spectrometry and ICP mass spectrometry, and about 20 g for analysis of  $^{36}\text{Cl}$  by accelerator mass spectrometry (AMS). Chlorine was extracted and purified from the 125–250  $\mu\text{m}$  fraction of leached samples and precipitated as  $\text{AgCl}$  using a modified version of procedures developed by Stone *et al.* (1996). Samples were spiked with  $\sim 1.26$  mg of Cl and sample Cl concentrations were determined by AMS isotope dilution (Di Nicola *et al.*, 2009). Samples were processed together with full chemistry blanks.

$^{10}\text{Be}/^9\text{Be}$ ,  $^{36}\text{Cl}/^{35}\text{Cl}$  and  $^{36}\text{Cl}/^{37}\text{Cl}$  ratios were measured using the 5 MW pelletron at SUERC (Xu *et al.*, 2010; Wilcken *et al.*, 2013) and normalized to NIST SRM4325 with a  $^{10}\text{Be}/^9\text{Be}$  ratio of  $2.79 \times 10^{-11}$  (Nishiizumi *et al.*,



**Figure 2.** (a) Some of the granite boulders of the moraine at Bloody Foreland; (b) part of the spread of glacially transported granite boulders at Rosguill; (c) Rosguill boulder ROS-01. The survey pole is divided at intervals of 0.2 m; (d) sampled bedrock at Malin Head, site MH-02; (e) Glencolumbkille boulder GC-02 undergoing sampling of projecting quartz vein; (f) Blue Stack Mountains boulder BS-01, showing projecting quartz pebbles. Scale bar is divided at intervals of 0.06 m.

2007), and Z93-0005 (PRIME Lab, Purdue) with a  $^{36}\text{Cl}/\text{Cl}$  ratio of  $1.2 \times 10^{-12}$ .

Cosmogenic nuclide concentrations include a blank correction of 3–14% for  $^{10}\text{Be}$  and 1–5% for  $^{36}\text{Cl}$  (Table 3). The uncertainties in the cosmogenic nuclide concentrations include the AMS counting statistics and scatter uncertainties from sample, procedural blank and standards measurements.

### Age calculation and filtering

The cosmogenic  $^{10}\text{Be}$  exposure ages were calculated using two methods. First, ages were determined using version 3.0 of the online calculators formerly known as the CRONUS-Earth exposure age calculators (Balco *et al.*, 2008; <http://hess.ess.washington.edu/math/>) using the independently-constrained LLPR (Fabel *et al.*, 2012) with time-dependent LM scaling (Lal, 1991; Stone, 2000) and assuming  $1 \text{ mm ka}^{-1}$  of post-depositional surface erosion (cf. André, 2002; Nicholson, 2009; Larsen *et al.*, 2012). The value for LLPR generated by version 3.0 of the online calculators is  $3.953 \pm 0.093 \text{ atoms g}^{-1} \text{ a}^{-1}$ . The uncertainty in this value (2.4%) represents the standard deviation of the measurements in the calibration data set and is unlikely to capture the real scaling uncertainty in the production rate estimate, thus leading to a likely underestimate of the computed external uncertainty in exposure ages (for further details see <http://sites.google.com/a/bgc.org/v3docs/home/4-ancillar-calculations-and-plots>). Second, we used CRONUScalc (<http://cronus.cosmogenicnuclides.rocks/2.0/>; Marrero *et al.*, 2016a) with the default global  $^{10}\text{Be}$  production rate of  $3.92 \text{ atoms g}^{-1} \text{ a}^{-1}$  for LM scaling (Borchers *et al.*, 2016), again assuming an erosion rate of  $1 \text{ mm ka}^{-1}$ . Both production rates agree within  $\pm 1\sigma$

uncertainties with the range of production rates determined for other high-latitude sites in the northern hemisphere (Phillips *et al.*, 2016).  $^{36}\text{Cl}$  ages were determined using CRONUScalc with LM scaling, an erosion rate of  $1 \text{ mm ka}^{-1}$  and production rates of  $56 \pm 4.1 \text{ at } ^{36}\text{Cl} (\text{g Ca})^{-1} \text{ a}^{-1}$  for Ca spallation,  $155 \pm 11 \text{ at } ^{36}\text{Cl} (\text{g K})^{-1} \text{ a}^{-1}$  for K spallation and  $759 \pm 180 \text{ neutrons } (\text{g air})^{-1} \text{ a}^{-1}$  (Marrero *et al.*, 2016b). Table 3 presents the  $^{10}\text{Be}$  and  $^{36}\text{Cl}$  data and exposure ages with associated uncertainties ( $\pm 1\sigma$ ). The data files for the online calculators are provided as supplementary data (Table S1). The effect of varying the assumed erosion rates between 0 and  $2 \text{ mm ka}^{-1}$  for both  $^{10}\text{Be}$  and  $^{36}\text{Cl}$  calculations results in <2% change for ages up to 18 ka, 3% change for ages 19–25 ka, and up to 13% for  $^{36}\text{Cl}$  ages between 32 and 43 ka. None of these variations affected our conclusions.

Within-site consistency of ages was tested using the reduced chi-square statistic ( $\chi^2_{\text{R}}$ ) (Bevington and Robinson, 2003). Where the  $\chi^2_{\text{R}}$  value for a sample of ages from a single site exceeds the critical value at the 95% level, it was inferred that geological uncertainty contributed to the observed age scatter. In such cases outlier ages were manually removed until a  $\chi^2_{\text{R}}$  value less than the critical value was obtained; the remaining ages were regarded as consistent with and representative of a single age population, with age scatter being due to measurement error alone (Balco, 2011; Applegate *et al.*, 2012; Small and Fabel, 2016; Small *et al.*, 2017a). For sites having two or more internally consistent ages, the uncertainty-weighted mean was determined and is regarded as providing the best estimate exposure age for the site. As with the legacy TCN ages discussed above, we cite the  $^{10}\text{Be}$ -weighted mean ages determined with the LLPR first, followed by the equivalent ages calculated with CRONUScalc in parentheses.

**Table 2.** Details of samples for TCN dating from Donegal.

Sample code	Grid reference	Latitude (°N)	Longitude (°W)	Altitude (m OD)	Thickness (cm)	Density (g cm <sup>-3</sup> )	Topographic shielding	Material and context
<b>North Donegal</b>								
<i>Rosguill</i>								
ROS-01	C 0999 4222	55.22690	7.84304	65	4.0	2.66	0.9938	Glacially transported granite boulder
ROS-02	C 1014 4203	55.22520	7.84062	105	5.0	2.67	0.9997	Glacially transported granite boulder
ROS-04	C 1015 4191	55.22412	7.84055	105	3.0	2.66	0.9967	Glacially transported granite boulder
<i>Malin Head</i>								
MH-02	C 3977 5955	55.38112	7.37255	65	5.0	2.59	0.9939	Ice-scoured quartzite bedrock
MH-03	C 3947 5960	55.38156	7.37716	30	3.0	2.65	0.9996	Vein quartz in quartzite bedrock
MH-04	C 3964 5946	55.38033	7.37458	55	2.5	2.58	0.6380	Ice-scoured quartzite bedrock
<i>Poisoned Glen</i>								
PG-01	B 9317 1863	55.01505	8.10675	73	2.0	2.61	0.9962	Glacially transported granite boulder
PG-04	B 9319 1862	55.01495	8.10653	73	3.0	2.62	0.9891	Glacially transported granite boulder
PG-05	B 9324 1862	55.01498	8.10572	75	2.5	2.56	0.9969	Glacially transported granite boulder
<b>South Donegal</b>								
<i>Glencolumbkille</i>								
GCS-02	G 5100 8474	54.70850	8.76100	25	1.5	2.65	0.9974	Vein quartz in glacially transported schist boulder
GCS-03	G 5107 8468	54.70790	8.75990	40	2.0	2.65	0.9983	Vein quartz in glacially transported schist boulder
GCS-04	G 5114 8464	54.70760	8.75890	35	1.0	2.65	0.9982	Vein quartz in glacially transported schist boulder
<i>Kilcar</i>								
KC-01	G 6063 7468	54.61870	8.60960	50	5.0	3.05	0.9980	Glacially transported dolerite boulder
KC-02	G 6063 7468	54.61870	8.60960	50	4.0	3.04	0.9987	Glacially transported dolerite boulder
KC-03	G 6063 7468	54.61870	8.60960	50	3.5	3.04	0.9665	Glacially transported dolerite boulder
KC-04	G 6066 7470	54.61890	8.60900	45	6.0	3.03	0.9838	Glacially transported dolerite boulder
<i>Blue Stacks</i>								
BS-01	G 9276 8608	54.72280	8.11320	150	1.0	2.65	0.9988	Quartz pebbles in glacially transported conglomerate boulder
BS-02	G 9270 8604	54.72250	8.11400	148	3.5	2.40	0.9967	Glacially transported conglomerate sandstone boulder
BS-03	G 9254 8611	54.72310	8.11650	150	2.0	2.65	0.9993	Quartz pebbles in glacially transported conglomerate boulder
BS-04	G 9242 8620	54.72390	8.11840	163	5.0	2.26	0.9993	Glacially transported conglomerate sandstone boulder

## Results

The 20 new TCN surface exposure ages and uncertainty-weighted mean values for internally consistent ages for each site are given in Table 3 and Fig. 3. These ages are assessed below in relation to published ages for the region (Table 1).

The ages for Rosguill and Malin Head, on the north coast of Donegal, complement published deglacial age estimates for the northern sites of Aran Island, Bloody Foreland and Corvish. A mean value was not calculated for Malin Head because the three samples failed to yield an acceptable  $\chi^2$  R-value, due to their wide age scatter (~25.7–20.9 and 25.8–20.7 ka). However, sample MH-03 yielded an age of  $20.9 \pm 0.9$  ka ( $20.7 \pm 1.8$  ka), reasonably consistent with the TCN mean ages of  $21.7 \pm 0.8$  ka ( $21.5 \pm 1.8$  ka) for Aran Island,  $21.6 \pm 0.7$  ka ( $21.7 \pm 1.8$  ka) for Bloody Foreland and  $19.0 \pm 0.7$  ka ( $18.8 \pm 1.6$  ka) for Rosguill; in addition, the MH-03 age is consistent with the minimum deglaciation  $^{14}\text{C}$  age of  $20.68 \pm 0.16$  cal ka BP for Corvish, and it therefore provides the best fit age of the three Malin Head ages.

Furthermore, Malin Head is the most easterly of our sites and is unlikely to have been deglaciated before the more westerly sites. The other two Malin Head samples are considered compromised by nuclide inheritance.

The mean age of  $16.9 \pm 0.7$  ka ( $16.7 \pm 1.5$  ka) for the Poisoned Glen boulder limit in north Donegal is statistically indistinguishable from the mean age of  $18.0 \pm 0.6$  ka ( $17.8 \pm 1.4$  ka) obtained from ice-plucked bedrock on Errigal, 2.2 km north and 350 m higher. Together these two sites indicate that the northern mountains were largely deglaciated by ~18–17 ka.

In south Donegal, the consistent exposure ages obtained from three boulders from Glencolumbkille yield a mean age of  $16.7 \pm 0.6$  ka ( $16.6 \pm 1.4$  ka). Two legacy samples from this location had returned ages of  $17.8 \pm 0.6$  ka ( $17.9 \pm 1.5$  ka) and  $19.6 \pm 0.7$  ka ( $19.8 \pm 1.7$  ka). The former age is statistically indistinguishable from the three new ages, and collectively all four ages produce an uncertainty-weighted mean age of  $17.2 \pm 0.6$  ka ( $17.0 \pm 1.4$  ka) ( $\chi^2_{\text{R}} = 1.24$  and 1.77, respectively). The latter age possibly reflects the

**Table 3.** Cosmogenic ( $^{10}\text{Be}$  and  $^{36}\text{Cl}$ ) data and surface exposure ages with total uncertainties at  $1\sigma$  for the Donegal samples. Analytical uncertainties ( $1\sigma$ ) are given in parentheses.

Sample code	AMS ID	$^{10}\text{Be}$ ( $10^4$ atoms $\text{g}^{-1}$ )	Blank $^{10}\text{Be}$ ( $10^4$ atoms)	$^{36}\text{Cl}$ ( $10^4$ atoms $\text{g}^{-1}$ )	Blank $^{36}\text{Cl}$ ( $10^4$ atoms)	$^{10}\text{Be}$ exposure age <sup>1</sup> (LLPR, LM)	$^{10}\text{Be}$ exposure age <sup>2</sup> (CRONUScalc LM)	$^{36}\text{Cl}$ exposure age <sup>3</sup>
<b>North Donegal</b>								
<i>Rosguill</i>								
ROS-01	b10320	$7.828 \pm 0.253$	$0.609 \pm 0.102$			$18.9 \pm 0.8$ (0.6)	$18.7 \pm 1.6$ (0.6)	
ROS-02	b10322	$9.257 \pm 0.418$	$0.609 \pm 0.102$			$21.5 \pm 1.1$ (1.0)	$21.0 \pm 2.0$ (1.0)	
ROS-04	b10323	$8.331 \pm 0.254$	$0.609 \pm 0.102$			$19.1 \pm 0.8$ (0.6)	$18.9 \pm 1.6$ (0.6)	
<b>Mean<sup>4,5</sup></b>						<b><math>19.0 \pm 0.7</math></b>	<b><math>18.8 \pm 1.6</math></b>	
<i>Malin Head</i>								
MH-02	b10324	$9.602 \pm 0.363$	$0.609 \pm 0.102$			$23.4 \pm 1.1$ (0.9)	$23.2 \pm 2.1$ (0.9)	
MH-03	b10286	$8.433 \pm 0.272$	$0.609 \pm 0.102$			$20.9 \pm 0.9$ (0.7)	$20.7 \pm 1.8$ (0.7)	
MH-04	b10426	$6.881 \pm 0.472$	$0.827 \pm 0.142$			$25.7 \pm 1.9$ (1.8)	$25.8 \pm 2.8$ (1.8)	
<i>Poisoned Glen</i>								
PG-01	b10628	$7.413 \pm 0.328$	$0.785 \pm 0.108$			$17.4 \pm 0.9$ (0.8)	$17.2 \pm 1.6$ (0.8)	
PG-04	b10629	$6.876 \pm 0.315$	$0.785 \pm 0.108$			$16.4 \pm 0.9$ (0.8)	$16.2 \pm 1.5$ (0.8)	
PG-05	b10630	$5.605 \pm 0.276$	$0.785 \pm 0.108$			$13.1 \pm 0.7$ (0.7)	$13.0 \pm 1.2$ (0.6)	
<b>Mean<sup>4,6</sup></b>						<b><math>16.9 \pm 0.7</math></b>	<b><math>16.7 \pm 1.5</math></b>	
<b>South Donegal</b>								
<i>Glencolumbkille</i>								
GCS-02	b8570	$6.672 \pm 0.291$	$0.894 \pm 0.159$			$16.4 \pm 0.8$ (0.7)	$16.2 \pm 1.5$ (0.7)	
GCS-03	b8572	$7.192 \pm 0.306$	$0.894 \pm 0.159$			$17.4 \pm 0.9$ (0.8)	$17.3 \pm 1.6$ (0.7)	
GCS-04	b8573	$6.873 \pm 0.293$	$0.894 \pm 0.159$			$16.6 \pm 0.8$ (0.7)	$16.5 \pm 1.5$ (0.7)	
<b>Mean<sup>4</sup></b>						<b><math>16.7 \pm 0.6</math></b>	<b><math>16.6 \pm 1.4</math></b>	
<b>Mean<sup>4,7</sup></b>						<b><math>17.2 \pm 0.6</math></b>	<b><math>17.0 \pm 1.4</math></b>	
<i>Kilcar</i>								
KC-01	c4058			$11.767 \pm 0.451$	$6.872 \pm 1.722$			$18.0 \pm 1.7$ (0.7)
KC-02	c4059			$12.378 \pm 0.598$	$6.872 \pm 1.722$			$34.3 \pm 5.6$ (1.7)
KC-03	c4060			$11.796 \pm 0.473$	$6.872 \pm 1.722$			$39.4 \pm 5.7$ (1.7)
KC-04	c4061			$13.951 \pm 0.558$	$6.872 \pm 1.722$			$34.9 \pm 5.1$ (1.5)
<i>Blue Stacks</i>								
BS-01	b9962	$6.163 \pm 0.331$	$1.683 \pm 0.177$			$13.2 \pm 0.8$ (0.7)	$13.1 \pm 1.3$ (0.7)	
BS-02	b8568	$7.097 \pm 0.301$	$0.894 \pm 0.159$			$15.5 \pm 0.8$ (0.7)	$15.4 \pm 1.4$ (0.7)	
BS-03	b8569	$6.984 \pm 0.303$	$0.894 \pm 0.159$			$15.1 \pm 0.8$ (0.7)	$14.9 \pm 1.4$ (0.7)	
BS-04	b10285	$6.729 \pm 0.206$	$0.814 \pm 0.103$			$14.6 \pm 0.6$ (0.5)	$14.4 \pm 1.2$ (0.4)	
<b>Mean<sup>4,8</sup></b>						<b><math>15.0 \pm 0.5</math></b>	<b><math>14.8 \pm 1.2</math></b>	

<sup>1</sup>Exposure age based on v 3.0 of the online exposure age calculator formerly known as the CRONUS-Earth online exposure age calculators (wrapper: 3.0-dev; get\_age: 3.0.2-dev; muons: 1A, alpha=1; consts: 3.0.3-dev) with the Loch Lomond production rate (LLPR), LM scaling, and assuming 1 mm  $\text{ka}^{-1}$  erosion.

<sup>2</sup>Exposure age based on CRONUScalc v2.0 with LM scaling and 1 mm  $\text{ka}^{-1}$  erosion.

<sup>3</sup>Exposure age based on CRONUScalc v2.0 with LM scaling and 1 mm  $\text{ka}^{-1}$  erosion.

<sup>4</sup>Uncertainty-weighted mean value.

<sup>5</sup>Mean value based on ROS-01 and -04 only.

<sup>6</sup>Mean value based on PG-01 and -04 only.

<sup>7</sup>Mean value includes an additional boulder sample age from Table 1 reported by Ballantyne *et al.* (2007).

<sup>8</sup>Mean value based on BS-02, BS-03 and BS-04 only.

influence of nuclide inheritance. The Glencolumbkille ages are also consistent with three legacy samples from rockslide run out debris on Slieve League (7 km SE of Glencolumbkille) that yielded a mean minimum age for deglaciation of  $17.3 \pm 0.6$  ka ( $17.1 \pm 1.5$  ka).

Four samples from dolerite boulders at Kilcar gave  $^{36}\text{Cl}$  exposure ages ranging from  $18.0 \pm 1.7$  to  $39.4 \pm 5.7$  ka. Samples KC-02, -03 and -04 returned ages that pre-date the LGM and are probably compromised by nuclide inheritance. The other sample (KC-01:  $18.0 \pm 1.7$  ka) is consistent with the wider geochronological evidence for the timing of deglaciation in south Donegal and north Mayo.

Three of the four samples obtained from boulders on low ground ( $\sim 150$  m) at the foot of the Blue Stack Mountains yielded consistent ( $\chi^2_{\text{R}} < 1.0$ ) ages ranging from  $15.5 \pm 0.8$  ka ( $15.4 \pm 1.4$  ka) to  $14.6 \pm 0.6$  ka ( $14.4 \pm 1.2$  ka), and an uncertainty-weighted mean age of  $15.0 \pm 0.5$  ka ( $14.8 \pm 1.2$  ka). The fourth sample (BS-01) is younger than the other three. BS-01 was collected from a boulder showing significant differential weathering (Fig. 2f). The assumed erosion rate

necessary to bring the age in line with the youngest of the other three Blue Stack Mountain samples is an unlikely  $10 \text{ mm ka}^{-1}$ . Alternatively, the weathering may indicate that the sampled surface was covered by peat or soil in the past, resulting in lower  $^{10}\text{Be}$  production. Either scenario decreases confidence in this sample and we do not include it in further discussions.


The results from three Blue Stack Mountain boulders imply deglaciation of all low ground at the head of Donegal Bay before  $\sim 14.7$  ka. They also suggest that ice persisted much later in the Blue Stack Mountains than in the Derryveagh Mountains of northern Donegal, where the available dating evidence suggests deglaciation of the Errigal col at  $\sim 18.0$  ka and ice withdrawal from the Poisoned Glen boulder limit at  $\sim 16.9$  ka.

## Discussion

In conjunction with the published legacy ages discussed earlier, the new ages presented here provide spatially consistent constraints on the timing of deglaciation in Donegal



**Figure 3.** Location map of Donegal and parts of Sligo and Mayo with all ages (legacy and new) of relevance to deglaciation. The new TCN ages are given in the same format as the legacy ages in Fig. 1. Note that the Malin Head age is a single age and the Glencolumbkille age is the mean of three new ages and the legacy age given in Fig. 1. Inferred ice margins (isochrones) are shown as solid lines for ~26–25 ka (ILGM), ~22, ~21, ~18, ~16 and ~15 ka, and as broken lines for readvance limits at ~19.5 ka (Donegal Bay Moraine) and ~18 ka (Corvish).



(Fig. 3). Below we discuss deglaciation of northern and southern Donegal separately, because during the ILGM northern ice fed the Hebrides/Malin Sea Ice Stream that drained ice from western Scotland across the Malin Shelf, whereas southern ice contributed to the Donegal Bay Ice Lobe that drained ice from the Irish Midlands to the shelf edge through Donegal Bay.

### Deglacial chronology of north Donegal

The TCN ages relating to deglaciation of Aran Island [mean =  $21.7 \pm 0.8$  ka ( $21.5 \pm 1.8$  ka)], Bloody Foreland [mean =  $21.6 \pm 0.7$  ka ( $21.7 \pm 1.8$  ka)], and Malin Head [a single TCN age of  $20.9 \pm 0.9$  ka ( $20.7 \pm 1.8$  ka)], together with the oldest  $^{14}\text{C}$  age from Corvish ( $20.68 \pm 0.16$  cal ka BP) indicate progressive eastward retreat of the ice margin along the northern coast of Donegal between ~21.7 and ~20.7 ka. The Bloody Foreland and Aran Island ages imply that decoupling of ice sourced in Donegal from the Scottish-sourced Hebridean Ice Stream commenced within the interval ~22–21 ka; this is slightly earlier than previous estimates, which have placed initial disengagement of these two ice masses after ~21 ka (Small *et al.*, 2017b). The single Malin Head TCN age and the oldest  $^{14}\text{C}$  age at Corvish indicate that separation of Scottish-sourced ice and Donegal-sourced ice was complete by ~20.7 ka, implying that by this time a marine embayment extended eastward along the north coast of Donegal, separating ice flowing north and north-east from the Donegal Ice Centre from the retreating Hebrides/Malin Sea Ice Stream.

The timing of ice retreat inland towards the Derryveagh Mountains of northern Donegal is provided by the TCN ages obtained for the two sites in the heart of this range, at ~420 m OD on Errigal col [mean =  $18.0 \pm 0.6$  ka ( $17.8 \pm 1.4$  ka)] and, 2.2 km to the south, a low-level site (~74 m OD) at the mouth of the Poisoned Glen [mean =  $16.9 \pm 0.7$  ka ( $16.7 \pm 1.5$  ka)]. Although these two ages are statistically indistinguishable within uncertainties (Fig. 4), the difference between them may imply exposure of Errigal col by downwasting ice several centuries before retreat of ice in the Poisoned Glen. Irrespective of whether this was the case, the deglaciation ages for both sites imply

that ~3000–5000 years elapsed between deglaciation of Aran Island and Bloody Foreland and deglaciation of the Derryveagh Mountains (Figs 3 and 4). A further implication is that net ice margin retreat rates were extremely slow. The Errigal col and Poisoned Glen sites lie, respectively, 15 and 17 km south-east of the Bloody Foreland site; if the mean deglaciation ages for these sites are representative, then the net ice-margin retreat rate from Bloody Foreland to both sites was  $\sim 4 \text{ m a}^{-1}$ ; taking the associated uncertainties into account suggests that net retreat rate is unlikely to have exceeded  $5 \text{ m a}^{-1}$ , and may have been as low as  $3 \text{ m a}^{-1}$ . By contrast, assuming that the ice margin began to retreat from the shelf edge within the interval 26.3–24.8 ka (Ó Cofaigh *et al.*, 2019), the implied net rate of offshore ice-margin retreat from the shelf break to Bloody Foreland falls within the range  $\sim 19.2\text{--}33.3 \text{ m a}^{-1}$ . Ó Cofaigh *et al.* (2019) inferred that ice-sheet retreat from the shelf edge was initiated by calving associated with high sea levels induced by glacio-isostatic depression rather than changing climate, and the marked slowing of retreat after the ice margin had become land-based in northern Donegal appears consistent with this interpretation: the inferred slow net retreat rates of Donegal ice in this sector over the period ~21–~17 ka suggest that the retreating ice was close to equilibrium with prevailing climate, and experienced only a slight net negative mass balance during this period.


Averaged net retreat rates, however, may obscure oscillations of the ice margin, with periods of retreat alternating with limited readvances. At present, there is dated stratigraphic evidence for only one such readvance, at Corvish, near the head of Trawbreaga Bay (Fig. 3). Readvance occurred over a distance of at least 5 km according to McCabe and Clark (2003). At this site, the youngest radiocarbon age obtained for *Elphidium clavatum* tests in deformed marine silts ( $18.32 \pm 0.18$  cal ka BP) and a single age for *E. clavatum* tests in overlying undeformed silts ( $17.06 \pm 0.18$  cal ka BP) have been interpreted by McCabe and Clark (2003) as bracketing the timing of readvance of the ice margin on the northern coast of Donegal. They placed the timing of this readvance at ~18 ka, although the dating evidence appears consistent with readvance of the

ice margin at any time within the interval  $\sim 18.5$ – $16.9$  ka. On the assumption that the Corvish readvance occurred at  $\sim 18$  ka, McCabe *et al.* (2007) suggested that it correlates with the Clogher Head Readvance (CHR) in north-east Ireland, although reinterpretation of the stratigraphic and dating evidence indicates that the CHR was a short-lived event that peaked rather earlier, at  $\sim 18.4$  ka (Ballantyne and Ó Cofaigh, 2017). Thus, although the two readvances may be coeval and represent a regional-scale event that occurred in response to climatic forcing (McCabe *et al.*, 2007; Clark *et al.*, 2012b) it is equally feasible that they occurred at different times and represent localized oscillations of the ice margin. Issues associated with recognition and correlation of readvances are discussed by Clark *et al.* (2012a). The TCN mean age of  $19.0 \pm 0.7$  ka ( $18.8 \pm 1.6$  ka) indicative of the timing of deglaciation at Rosguill pre-dates the bracketing ages for the readvance at Corvish ( $\sim 18.5$ – $16.9$  ka), but because of the uncertainties associated with the TCN age we cannot preclude the possibility that the Rosguill site was reoccupied by glacier ice during the same readvance event.

The TCN mean age for the Poisoned Glen [ $16.9 \pm 0.7$  ka ( $16.7 \pm 1.5$  ka)] implies that ice persisted in the Derryveagh Mountains until  $\sim 17$ – $16$  ka, but the  $^{14}\text{C}$  age of  $15.38 \pm 0.12$  cal ka BP from Lough Nadourcan (Watson *et al.*, 2010) suggests that ice had disappeared from low ground surrounding these mountains before the rapid warming associated with the onset of the Lateglacial Interstadial at  $\sim 14.7$  ka. Valley heads, cirques and plateaus in the Derryveagh Mountains may, however, have retained ice until early in the interstadial, as appears to have been the case for the Blue Stack Mountains of south Donegal (see below).

### Deglacial chronology of south Donegal and Donegal Bay

As noted earlier, the DBM that crosses outer Donegal Bay (Fig. 3) represents the limit of a readvance of ice fed from Donegal Bay. Radiocarbon ages of  $20.24 \pm 0.24$  and  $17.92 \pm 0.16$  cal ka BP obtained for foraminifera retrieved, respectively, from the distal and proximal sides of the moraine constrain its age (Ó Cofaigh *et al.*, 2019). This broad interval encompasses the timing of both the



**Figure 4.** Equal-area Gaussian probability distributions representing the uncertainty-weighted means and associated uncertainties for the cosmogenic  $^{10}\text{Be}$  exposure ages obtained for samples from the 'coastal' sites on Aran Island ( $n=2$ ) and Bloody Foreland ( $n=7$ ), and the 'inland mountain' sites of Errigal Col ( $n=3$ ) and Poisoned Glen ( $n=2$ ). These distributions illustrate the overlap in the ages obtained for the two 'coastal' sites and for the two 'inland mountain' sites, and also the temporal interval of  $\geq 4$  ka that separates the two sets of ages.

readvance at Corvish in northern Donegal and that of the CHR in north-east Ireland (McCabe and Clark, 2003; McCabe *et al.*, 2007; Clark *et al.*, 2012b; Ballantyne and Ó Cofaigh, 2017), but the resolution of the dating evidence is inadequate to establish contemporaneity. The position and alignment of the DBM suggests that the sites at Belderg Pier and Fiddauntawnoneen on the south coast of outer Donegal Bay lay outside the readvance, and the radiocarbon ages of  $\sim 20$ – $19$  cal ka BP obtained by McCabe *et al.* (1986, 2005) for *in situ* marine fauna within glaci-marine sediments at these sites (Table 1) are consistent with this interpretation (Fig. 3). Conversely, the aggregated TCN mean age for Glencolumbkille in south-west Donegal [ $17.2 \pm 0.6$  ka ( $17.0 \pm 1.4$  ka)], the single TCN age for Kilcar ( $18.0 \pm 1.7$  ka) and the minimum deglaciation age represented by postglacial rockslide debris at nearby Slieve League [ $17.3 \pm 0.6$  ka ( $17.1 \pm 1.5$  ka)] suggest that southern Donegal lay within the limits of the readvance that produced the DBM. Similarly, the five 'younger' TCN ages [mean =  $16.6 \pm 0.6$  ka ( $16.7 \pm 1.5$  ka)] reported by Clark *et al.* (2009c) for the Twanywaddyduff moraine system of the northern Ox Mountains (Fig. 3; Table 1) indicate persistence of ice cover along the inner part of Donegal Bay after  $\sim 17$  ka. Collectively, these two sets of ages suggest that much or all of Donegal Bay continued to support ice cover as late as  $\sim 17$  ka, although it is possible that a calving margin along the axis of the bay led to development of an ice-free marine corridor between its northern and southern shores.

The TCN mean age of  $15.0 \pm 0.5$  ka ( $14.8 \pm 1.2$  ka) from the southern flanks of the Blue Stack Mountains suggests that Donegal Bay had become ice free by  $\sim 15$  ka but that ice still occupied mountain valleys and cirques near the head of the bay (Fig. 3). The implication of this age is that mountain ice probably persisted for some time following the onset of the Lateglacial Interstadial at  $\sim 14.7$  ka. The Blue Stack ages are the youngest ages for deglaciation hitherto reported for Ireland (cf. Ballantyne and Ó Cofaigh, 2017), suggesting that these and possibly other mountains in north-west Ireland supported the last remnants of the last Irish Ice Sheet before complete disappearance of glacier ice under the warmer conditions of the Lateglacial Interstadial. The Blue Stack Mountains, along with other mountain areas in Ireland, hosted glaciers during the Younger Dryas Stadial ( $\sim 12.9$ – $11.7$  ka) (Barr *et al.*, 2017; Barth *et al.*, 2018; Tomkins *et al.*, 2018), but it has not yet been demonstrated that these glaciers had persisted throughout the Lateglacial Interstadial.

The sampling sites at Glencolumbkille and the Blue Stack Mountains are separated by a distance of 41 km. The mean TCN ages for these two sites imply that net ice margin retreat between these two sites occurred over  $\sim 2200$  years, implying a net retreat rate of  $\sim 19$  m  $\text{a}^{-1}$ ; taking the associated age uncertainties into account implies that net retreat rate of the ice margin along the northern shore of Donegal Bay fell within the range  $12$ – $24$  m  $\text{a}^{-1}$ , markedly faster than the net rate inferred above ( $3$ – $5$  m  $\text{a}^{-1}$ ) for ice retreat inland from Bloody Foreland to the Derryveagh Mountains. Although subject to the same caveat (that retreat may have been interrupted by one or more ice margin readvances), there is neither morphological nor seismostratigraphic evidence for later readvances of the ice margin as it retreated eastward from the DBM. Ó Cofaigh *et al.* (2019) noted that the sediment cover east of the moraine comprises undeformed, acoustically stratified conformable glaci-marine sediments overlain by postglacial marine deposits. For comparison, the average net rate calculated by Ó Cofaigh *et al.* (2019) for

ice-margin retreat from the shelf edge to the DBM is  $11.2\text{--}14.0\text{ m a}^{-1}$ , although they considered that this probably incorporated moderately rapid retreat at a minimum rate of  $35.7\text{ m a}^{-1}$  from the shelf edge to mid-shelf, followed by much slower oscillatory retreat at a net rate of  $5.5\text{ m a}^{-1}$  between the mid-shelf and the DBM.

### Wider implications

Collectively, the chronological data reported above indicate a marked contrast in both the timing and the rate of net ice margin recession of land-based ice in northern Donegal ( $\sim 21\text{--}17\text{ ka}$ ) and retreat of the ice margin in southern Donegal adjacent to Donegal Bay ( $\sim 17\text{--}15\text{ ka}$ ). This contrast suggests that the timing of ice-margin retreat was at least partly conditioned by the relationship between ice fed from the Donegal Ice Centre and adjacent ice streams and lobes. Northern Donegal lay in an inter-ice-stream/lobe location, between the Hebrides/Malin Sea Ice Stream to the north and the Donegal Bay Ice Lobe to the south, and here the early ( $\sim 22\text{--}21\text{ ka}$ ) decoupling of Donegal ice from the extended Hebrides/Malin Sea Ice Stream appears to have created an ice-free marine embayment along the north coast of Donegal, so that ice flowing from the Donegal Ice Centre was effectively unconstrained, and subsequently retreated gradually in response to a slight net negative mass balance. In contrast, the south coast of Donegal lay near the axis of the Donegal Bay Ice Lobe, which was fed not only by ice from the Donegal Ice Centre, but also by ice from the Irish Midlands. Following initial rapid retreat, the oscillating margin of the Donegal Bay Ice Lobe retreated slowly from mid-shelf to the DBM (Ó Cofaigh *et al.*, 2019), so that ice cover persisted over south Donegal until  $\sim 17\text{ ka}$ , after which it retreated to the footslopes of the Blue Stack Mountains. This contrast in behaviour implies that different dynamics apply to extended marine-based ice streams and lobes, which are sensitive to changes in sea level, confinement and bed slope (Smedley *et al.*, 2017; Ó Cofaigh *et al.*, 2019; Small *et al.*, 2018) and land-based ice masses in inter-ice-stream/lobe locations, which respond mainly to changes in climate inputs.

Evidence for marked slowing of ice-margin retreat as the shrinking BISS stabilized at or near the present coastline is not limited to northern Donegal. TCN ages reported by Small *et al.* (2017b) for the Sea of the Hebrides to the west of Scotland suggest that termination of ice streaming after  $\sim 20.6\text{ ka}$  was succeeded by a  $\sim 3000\text{--}4000\text{-year}$  interval during which the ice margin experienced oscillatory net retreat of only  $50\text{--}70\text{ km}$  as it became progressively land-based among the islands of the Inner Hebrides. The slowing of ice margin retreat in this area coincides closely with the period of very gradual ice-margin recession in northern Donegal.

The mean exposure age of the samples from low ground ( $\sim 150\text{ m}$ ) at the foot the Blue Stack Mountains [ $15.0 \pm 0.5\text{ ka}$  ( $14.8 \pm 1.2\text{ ka}$ )] represents the youngest age for the timing of ice-sheet deglaciation hitherto reported for Ireland and implies that by  $\sim 15.0\text{ ka}$  the Donegal Ice Centre had shrunk to a small ice cap or transection complex centred on high ground. An analogous situation occurred in south-west Scotland,  $240\text{ km}$  to the east, where seven (recalibrated) TCN ages indicate that only fragmented upland remnants of the Galloway Hills Ice Centre remained by  $\sim 15.1\text{ ka}$  (Ballantyne *et al.*, 2013a). The Galloway Hills TCN ages are statistically indistinguishable from the Blue Stack ages, and both confirm that almost all ice retreat occurred under stadial conditions before  $\sim 15.0\text{ ka}$ . In both areas it is unlikely that remnant

glacier ice survived subsequent rapid warming, when mean July temperatures inferred from subfossil chironomid assemblages rose rapidly by  $5\text{--}6\text{ }^{\circ}\text{C}$  (Brooks and Birks, 2000; Lang *et al.*, 2010; Watson *et al.*, 2010; Van Asch *et al.*, 2012). A more general implication is that all of Ireland and southern Scotland were probably completely deglaciated early in the Lateglacial Interstadial. For the British Isles as a whole, present evidence suggests that remnants of the BISS survived the interstadial (if at all) only in the Highlands of Scotland (Finlayson *et al.*, 2011; Ballantyne and Small, 2018).

### Conclusions

1. Twenty new TCN ages obtained for sites in northern and southern Donegal complement (and are broadly consistent with) previously published TCN and radiocarbon ages, and reveal marked contrasts in the timing and rate of deglaciation in northern and southern Donegal.
2. The TCN ages for northern Donegal indicate decoupling of ice fed from the Donegal Ice Centre from the Hebrides/Malin Shelf Ice Stream and associated development of a marine embayment north of Donegal by  $\sim 22\text{--}21\text{ ka}$ . Conversely, the new TCN ages for south Donegal confirm that ice persisted in much or all Donegal Bay and covered south-west Donegal as late as  $\sim 17\text{ ka}$ .
3. In northern Donegal our TCN data imply very gradual ice margin retreat inland towards mountain source areas at a net rate of  $4 \pm 1\text{ m a}^{-1}$ ; by comparison, the inferred net rate of ice margin retreat from south-west Donegal to the foothills of the Blue Stack Mountains near the head of Donegal Bay averaged  $18 \pm 6\text{ m a}^{-1}$ .
4. We suggest that the above contrast in timing and rate of ice retreat reflects differences in location relative to those of major ice streams/lobes. Northern Donegal occupied an inter-ice-stream location, and after early decoupling of Donegal-sourced ice from the Hebrides/Malin Sea Ice Stream the former was unconstrained and retreated mainly in response to changes in climatic inputs. Conversely, southern Donegal lay near to the axis of the Donegal Bay Ice Lobe, which occupied (or reoccupied) much of Donegal Bay as late as  $\sim 18\text{ ka}$ , delaying deglaciation along the southern coast of Donegal until after  $\sim 17\text{ ka}$ .
5. A mean TCN age of  $\sim 15.0 \pm 0.5\text{ ka}$  ( $14.8 \pm 1.2\text{ ka}$ ) obtained for the footslopes of the Blue Stack Mountains in southern Donegal is the youngest deglacial age hitherto reported for Ireland, and implies that shortly before the onset of rapid warming at the beginning of the Lateglacial Interstadial ( $\sim 14.7\text{ ka}$ ) the Donegal Ice Centre had shrunk to a small ice cap or ice field of very limited extent, and probably disappeared completely during the early part of the interstadial. This date also confirms that virtually all the retreat of the Irish Ice Sheet occurred under stadial conditions before the onset of interstadial warming.

### Supporting information

Additional supporting information can be found in the online version of this article.

**Acknowledgements.** This work was supported by the UK Natural Environment Research Council consortium grant: BRITICE-CHRONO NE/J009768/1. The TCN analyses were undertaken at the NERC

Cosmogenic Isotope Analysis Facility (allocation 9155/1014). Staff at the Scottish Universities Environmental Research Centre AMS Laboratory, East Kilbride, are thanked for  $^{10}\text{Be}$  and  $^{36}\text{Cl}$  isotope measurements, as are two anonymous referees for their constructive comments on the initial draft of the paper.

**Abbreviations.** AMS, accelerator mass spectrometry; BIIIS, British–Irish Ice Sheet; CHR, Clogher Head Readvance; CIAF, Cosmogenic Isotope Analysis Facility; DBM, Donegal Bay Moraine; gLGM, global Last Glacial Maximum; ILGM, local Last Glacial Maximum; LLPR, Loch Lomond production rate; TCN, terrestrial cosmogenic nuclide.

## References

- André M-F. 2002. Rates of postglacial rock weathering on glacially scoured outcrops (Abisko-Riksgränsen area, 68°N). *Geografiska Annaler: Series A, Physical Geography* **84**: 139–150.
- Applegate PJ, Urban NM, Keller K *et al.* 2012. Improved moraine age interpretations through explicit matching of geomorphic process models to cosmogenic nuclide measurements from single landforms. *Quaternary Research* **77**: 293–304.
- Balco G. 2011. Contributions and unrealized potential contributions of cosmogenic-nuclide exposure dating to glacier chronology, 1990–2010. *Quaternary Science Reviews* **30**: 3–27.
- Balco G, Stone JO, Lifton NA *et al.* 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from  $^{10}\text{Be}$  and  $^{26}\text{Al}$  measurements. *Quaternary Geochronology* **3**: 174–195.
- Ballantyne CK, McCarroll D, Stone JO. 2007. The Donegal ice dome, northwest Ireland: dimensions and chronology. *Journal of Quaternary Science* **22**: 773–783.
- Ballantyne CK, McCarroll D, Stone JO. 2011. Periglacial trimlines and the extent of the Kerry–Cork Ice Cap, SW Ireland. *Quaternary Science Reviews* **30**: 3834–3845.
- Ballantyne CK, Ó Cofaigh C. 2017. The last Irish Ice Sheet: extent and chronology. In *Advances in Irish Quaternary Studies*, Coxon P, Mitchell F, McCarron S (eds). Atlantic Press: Paris; 101–149.
- Ballantyne CK, Rinterknecht V, Gheorghiu DM. 2013a. Deglaciation chronology of the Galloway Hills Ice Centre, southwest Scotland. *Journal of Quaternary Science* **28**: 412–420.
- Ballantyne CK, Small D. 2018. The last Scottish Ice Sheet. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*: DOI: 10.1007/S1755691018000038.
- Ballantyne CK, Wilson P, Schnabel C *et al.* 2013b. Lateglacial rock slope failures in north-west Ireland: age, causes and implications. *Journal of Quaternary Science* **28**: 789–802.
- Barr ID, Roberson S, Flood R *et al.* 2017. Younger Dryas glaciers and climate in the Mourne Mountains, Northern Ireland. *Journal of Quaternary Science* **32**: 104–115.
- Barth AM, Clark PU, Clark J *et al.* 2018. Persistent millennial-scale glacier fluctuations in Ireland between 24 ka and 10 ka. *Geology* **46**: 151–154.
- Benetti S, Dunlop P, Cofaigh C. 2010. Glacial and glacially related features on the continental margin of northwest Ireland mapped from marine geophysical data. *Journal of Maps* **6**: 14–29.
- Bevington PR, Robinson DK. 2003. *Data Reduction and Error Analysis for the Physical Sciences*. McGraw-Hill: New York.
- Borchers B, Marrero S, Balco G *et al.* 2016. Geological calibration of spallation production rates in the CRONUS-Earth project. *Quaternary Geochronology* **31**: 188–198.
- Bowen DQ, Rose J, McCabe AM *et al.* 1986. Correlation of Quaternary glaciations in England, Ireland, Scotland and Wales. *Quaternary Science Reviews* **5**: 299–340.
- Bowen DQ, Phillips FM, McCabe AM *et al.* 2002. New data for the Last Glacial Maximum in Great Britain and Ireland. *Quaternary Science Reviews* **21**: 89–101.
- Bronk Ramsey C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**: 337–360.
- Brooks SJ, Birks HJB. 2000. Chironomid-inferred Late-glacial air temperatures at Whitrig Bog, Southeast Scotland. *Journal of Quaternary Science* **15**: 759–764.
- Charlesworth JK. 1924. The glacial geology of north-west Ireland. *Proceedings of the Royal Irish Academy* **36B**: 174–314.
- Child D, Elliott G, Mifsud C *et al.* 2000. Sample processing for earth science studies at ANTARES. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **172**: 856–860.
- Clark CD, Hughes ALC, Greenwood SL *et al.* 2012a. Pattern and timing of retreat of the last British-Irish Ice Sheet. *Quaternary Science Reviews* **44**: 112–146.
- Clark J, McCabe AM, Bowen DQ *et al.* 2012b. Response of the Irish Ice Sheet to abrupt climate change during the last deglaciation. *Quaternary Science Reviews* **35**: 100–115.
- Clark J, McCabe AM, Schnabel C *et al.* 2009b.  $^{10}\text{Be}$  chronology of the last deglaciation of County Donegal, northwestern Ireland. *Boreas* **38**: 111–118.
- Clark J, McCabe AM, Schnabel C *et al.* 2009c. Cosmogenic  $^{10}\text{Be}$  chronology of the last deglaciation of western Ireland, and implications for sensitivity of the Irish Ice Sheet to climate change. *Geological Society of America Bulletin* **121**: 3–16.
- Clark PU, Dyke AS, Shakun JD *et al.* 2009a. The Last Glacial Maximum. *Science* **325**: 710–714.
- Colhoun EA. 1973. Two Pleistocene sections in south-eastern Donegal and their relation to the last glaciation of the Glengesh plateau. *Irish Geography* **6**: 594–609.
- Cullen C. 2013. *Deciphering the geomorphic and sedimentary record of the last Irish Ice Sheet in NW Donegal, Ireland: implications for glacial dynamics and decay configurations*. PhD Thesis, National University of Ireland, Galway.
- Di Nicola L, Schnabel C, Wilcken KM *et al.* 2009. Determination of chlorine concentrations in whole rock: comparison between prompt-gamma activation and isotope-dilution AMS analysis. *Quaternary Geochronology* **4**: 501–507.
- Dunlop P, Shannon R, McCabe M *et al.* 2010. Marine geophysical evidence for ice sheet extension and recession on the Malin Shelf: new evidence for the western limits of the British Irish Ice Sheet. *Marine Geology* **276**: 86–99.
- Dury GH. 1957. A glacially breached watershed in Donegal. *Irish Geography* **3**: 171–180.
- Dury GH. 1958. Glacial morphology of the Blue Stack area, Donegal. *Irish Geography* **3**: 242–253.
- Dury GH. 1964. Aspects of the geomorphology of Slieve League Peninsula, Donegal. *Proceedings of the Geologists' Association* **75**: 445–459.
- Fabel D, Ballantyne CK, Xu S. 2012. Trimlines, blockfields, mountain-top erratics and the vertical dimensions of the last British-Irish Ice Sheet in NW Scotland. *Quaternary Science Reviews* **55**: 91–102.
- Finlayson A, Golledge N, Bradwell T *et al.* 2011. Evolution of a Lateglacial mountain icecap in northern Scotland. *Boreas* **40**: 536–554.
- Greenwood SL, Clark CD. 2009a. Reconstructing the last Irish Ice Sheet 1: changing flow geometries and ice flow dynamics deciphered from the glacial landform record. *Quaternary Science Reviews* **28**: 3085–3100.
- Greenwood SL, Clark CD. 2009b. Reconstructing the last Irish Ice Sheet 2: a geomorphologically-driven model of ice sheet growth, retreat and dynamics. *Quaternary Science Reviews* **28**: 3101–3123.
- Hubbard A, Bradwell T, Golledge N *et al.* 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and deglaciation of the British-Irish ice sheet. *Quaternary Science Reviews* **28**: 758–776.
- Knight J. 2003. Geomorphic and sedimentary evidence for patterns of late Midlandian ice retreat in the Tempo valley, north-central Ireland. *Irish Geography* **36**: 127–144.
- Knight J. 2012. The last glaciation of Aran Island and Cruit Island, Co. Donegal, north-west Ireland. *Irish Journal of Earth Sciences* **30**: 49–58.
- Knight J, McCabe AM. 1997. Drumlin evolution and ice sheet oscillations along the NE Atlantic margin, Donegal Bay, western Ireland. *Sedimentary Geology* **111**: 57–72.
- Kohl CP, Nishiizumi K. 1992. Chemical isolation of quartz for measurement of in-situ  $\alpha$ -produced cosmogenic nuclides. *Geochimica et Cosmochimica Acta* **56**: 3583–3587.
- Lal D. 1991. Cosmic ray labeling of erosion surfaces: *in situ* nuclide production rates and erosion models. *Earth and Planetary Science Letters* **104**: 424–439.

- Lang B, Brooks SJ, Bedford A *et al.* 2010. Regional consistency in Lateglacial chironomid-inferred temperatures from five sites in north-west England. *Quaternary Science Reviews* **29**: 1528–1538.
- Larsen NK, Linge H, Håkansson L *et al.* 2012. Investigating the last deglaciation of the Scandinavian Ice Sheet in southwest Sweden with  $^{10}\text{Be}$  exposure dating. *Journal of Quaternary Science* **27**: 211–220.
- Long CB, McConnell BJ. 1997. Geology of North Donegal. *Geological Survey of Ireland: Dublin*.
- Long CB, McConnell BJ. 1999. Geology of South Donegal. *Geological Survey of Ireland: Dublin*.
- Marrero SM, Phillips FM, Borchers B *et al.* 2016a. Cosmogenic nuclide systematics and the CRONUScalc program. *Quaternary Geochronology* **31**: 160–187.
- Marrero SM, Phillips FM, Caffee MW *et al.* 2016b. CRONUS-Earth cosmogenic  $^{36}\text{Cl}$  calibration. *Quaternary Geochronology* **31**: 199–219.
- McCabe AM. 1985. Glacial geomorphology. In *The Quaternary History of Ireland*, Edwards KJ, Warren WP (eds). Academic Press: London; 67–93.
- McCabe AM. 1995. Quaternary geology. In *North-West Donegal*. Field Guide No. 19, Wilson P (ed.). Irish Association for Quaternary Studies: Dublin; 15–20.
- McCabe AM, Bowen DQ, Penney DN. 1993. Glaciomarine facies from the western sector of the last British Ice Sheet, Malin Beg, County Donegal, Ireland. *Quaternary Science Reviews* **12**: 35–45.
- McCabe AM, Clark PU. 2003. Deglacial chronology from County Donegal, Ireland: implications for deglaciation of the British-Irish ice sheet. *Journal of the Geological Society* **160**: 847–855.
- McCabe AM, Clark PU, Clark J. 2005. AMS  $^{14}\text{C}$  dating of deglacial events in the Irish Sea Basin and other sectors of the British-Irish ice sheet. *Quaternary Science Reviews* **24**: 1673–1690.
- McCabe AM, Clark PU, Clark J *et al.* 2007. Radiocarbon constraints on readvances of the British-Irish Ice Sheet in the northern Irish Sea Basin during the last deglaciation. *Quaternary Science Reviews* **26**: 1204–1211.
- McCabe AM, Haynes JR, Macmillan NF. 1986. Late-Pleistocene tidewater glaciers and glaciomarine sequences from north County Mayo, Republic of Ireland. *Journal of Quaternary Science* **1**: 73–84.
- Nicholson DT. 2009. Holocene microweathering rates and processes on ice-eroded bedrock, Rødal area, Hardangervidda, southern Norway. In *Periglacial and Paraglacial Processes and Environments, Special Publications* Vol. 320, Knight J, Harrison S (eds). Geological Society: London; 29–49.
- Nishiizumi K, Imamura M, Caffee MW *et al.* 2007. Absolute calibration of  $^{10}\text{Be}$  AMS standards. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **258**: 403–413.
- Ó Cofaigh C, Dunlop P, Benetti S. 2012. Marine geophysical evidence for Late Pleistocene ice sheet extent and recession off northwest Ireland. *Quaternary Science Reviews* **44**: 147–159.
- Ó Cofaigh C, Weillbach K, Lloyd JM *et al.* 2019. Early deglaciation of the British-Irish Ice Sheet on the Atlantic shelf northwest of Ireland driven by glacioisostatic depression and high relative sea level. *Quaternary Science Reviews*.
- Phillips FM, Argento DC, Balco G *et al.* 2016. The CRONUS-Earth Project: a synthesis. *Quaternary Geochronology* **31**: 119–154.
- Rasmussen SO, Bigler M, Blockley SP *et al.* 2014. A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary Science Reviews* **106**: 14–28.
- Reimer PJ, Bard E, Bayliss A *et al.* 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radio-carbon* **55**: 1869–1887.
- Sejrup HP, Hjelstuen BO, Torbjørn Dahlgren KIT *et al.* 2005. Pleistocene glacial history of the NW European continental margin. *Marine and Petroleum Geology* **22**: 1111–1129.
- Sellier D. 1995. Derryveagh Mountains – glacial evidence. In *North-West Donegal*. Field Guide No. 19, Wilson P (ed.). Irish Association for Quaternary Studies: Dublin; 57–58.
- Small D, Benetti S, Dove D *et al.* 2017b. Cosmogenic exposure age constraints on deglaciation and flow behaviour of a marine-based ice stream in western Scotland, 21–16 ka. *Quaternary Science Reviews* **167**: 30–46.
- Small D, Clark CD, Chiverrell RC *et al.* 2017a. Devising quality assurance procedures for assessment of legacy geochronological data relating to deglaciation of the last British-Irish Ice Sheet. *Earth-Science Reviews* **164**: 232–250.
- Small D, Fabel D. 2016. Response to Bromley, *et al.* “Comment on ‘Was Scotland deglaciated during the Younger Dryas?’” *Quaternary Science Reviews* **152**: 206–208.
- Small D, Smedley RK, Chiverrell RC *et al.* 2018. Trough geometry was a greater influence than climate-ocean forcing in regulating retreat of the marine-based Irish-Sea Ice Stream. *GSA Bulletin* **130**: 1981–1999.
- Smedley RK, Chiverrell RC, Ballantyne CK *et al.* 2017. Internal dynamics condition centennial-scale oscillations in marine-based ice-stream retreat. *Geology* **45**: 787–790.
- Smith MJ, Knight J. 2011. Palaeoglaciology of the last Irish ice sheet reconstructed from striae evidence. *Quaternary Science Reviews* **30**: 147–160.
- Stephen N, Synge FM. 1965. Late-Pleistocene shorelines and drift limits in north Donegal. *Proceedings of the Royal Irish Academy* **64B**: 131–153.
- Stone JO. 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical Research: Solid Earth* **105**: 23753–23759.
- Stone JO, Allan GL, Fifield LK *et al.* 1996. Cosmogenic chlorine-36 from calcium spallation. *Geochimica et Cosmochimica Acta* **60**: 679–692.
- Tomkins MD, Dortch JM, Hughes PD *et al.* 2018. Timing of glacial retreat in the Wicklow Mountains, Ireland, conditioned by glacier size and topography. *Journal of Quaternary Science* **33**: 611–623.
- Van Asch N, Lutz AF, Duijkers MCH *et al.* 2012. Rapid climate change during the Weichselian Lateglacial in Ireland: Chironomid-inferred summer temperatures from Fiddaun, Co. Galway. *Palaeogeography, Palaeoclimatology, Palaeoecology* **315–316**: 1–11.
- Watson JE, Brooks SJ, Whitehouse NJ *et al.* 2010. Chironomid-inferred late-glacial summer air temperatures from Lough Nadourcan, Co. Donegal, Ireland. *Journal of Quaternary Science* **25**: 1200–1210.
- Wilcken KM, Freeman SPHT, Schnabel C *et al.* 2013.  $^{36}\text{Cl}$  accelerator mass spectrometry with a bespoke instrument. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **294**: 107–114.
- Xu S, Dougans AB, Freeman SPHT *et al.* 2010. Improved  $^{10}\text{Be}$  and  $^{26}\text{Al}$  AMS with a 5 MV spectrometer. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **268**: 736–738.