

Accepted Manuscript

Research paper

Reactivity of hemilabile pyridyl- and methyl-substituted pyrimidylselenolates with $[MCl_2(dppf)]$ ($M = Pd, Pt$; $dppf = \text{bis}(\text{diphenylphosphino})\text{ferrocene}$)

Rohit Singh Chauhan, David B. Cordes, Alexandra M.Z. Slawin, Seema Yadav, Chandrakanta Dash


PII: S0020-1693(18)30176-2
DOI: <https://doi.org/10.1016/j.ica.2018.03.034>
Reference: ICA 18179

To appear in: *Inorganica Chimica Acta*

Received Date: 31 January 2018
Revised Date: 20 March 2018
Accepted Date: 21 March 2018

Please cite this article as: R.S. Chauhan, D.B. Cordes, A.M.Z. Slawin, S. Yadav, C. Dash, Reactivity of hemilabile pyridyl- and methyl-substituted pyrimidylselenolates with $[MCl_2(dppf)]$ ($M = Pd, Pt$; $dppf = \text{bis}(\text{diphenylphosphino})\text{ferrocene}$), *Inorganica Chimica Acta* (2018), doi: <https://doi.org/10.1016/j.ica.2018.03.034>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Reactivity of hemilabile pyridyl- and methyl-substituted pyrimidylselenolates with $[MCl_2(dppf)]$ ($M = Pd, Pt$; $dppf = \text{bis}(\text{diphenylphosphino})\text{ferrocene}$)

Rohit Singh Chauhan^{*a}, David B. Cordes^b, Alexandra M. Z. Slawin^b, Seema Yadav^c, Chandrakanta Dash^c

^aDepartment of Chemistry, K. J. Somaiya College of Science and Commerce, Mumbai- 400 078, Email: rohit.chuhan@somaiya.edu

^bEastCHEM School of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST Email: amzs@st-andrews.ac.uk

^cDepartment of Chemistry, Central University of Rajasthan, Ajmer-305817, Rajasthan, Email: ckdash@curaj.ac.in

Abstract:

The bis(diphenylphosphino)ferrocene (dppf) derived palladium analogue of $[PdCl_2(dppf)]$ on reaction with the sodium salt of pyridyl/pyrimidyl selenolate yielded mononuclear *cis* configured complex $[Pd(SeAr)_2(dppf)]$ ($Ar = C_5H_4N, C_4H(4,6-Me)_2N_2$), as did the platinum precursor $[PtCl_2(dppf)]$ with methyl substituted pyrimidyl salt giving $[Pt\{SeC_4H(4,6-Me)_2N_2\}_2(dppf)]$. In contrast, the reaction of platinum precursor with the sodium salt of pyridylselenolate gave two different complexes $[Pt_2(\mu-SeC_5H_4N)_2(dppf)_2]^{2+}$ and $[Pt(Cl)(SeC_5H_4N)(dppf)]$ depending in the reaction solvent. These complexes were characterized by elemental analysis and NMR ($^1H, ^{31}P$) spectroscopy. The molecular structure of $[Pt\{SeC_4H(4,6-Me)_2N_2\}_2(dppf)]$ was established by single crystal X-ray diffraction analysis.

Keywords: Selenolate, NMR, X-ray, dppf.

1. Introduction

The ferrocenylphosphine ligand has been extensively used as an efficient auxiliary ligand mainly due to its rich coordination chemistry without tormenting the intrinsic property of derived complex [1-3]. This inherent characteristic makes it an efficient materials and catalyst for the alkene hydroformylation, alkoxyacylation and heck coupling reactions [1, 4]. Even though ferrocenyl phosphine ligand was reported more than three decades ago but its chemical reactivity and application were not been unearthed until recently [1, 5-7].

The reaction of $[MCl_2(dppf)]$ ($M = Pd, Pt$) with Na_2S has afforded a binuclear product $[M_2(\mu-S)_2(dppf)_2]$ which transformed to $[Pt_2(\mu-S)(\mu-SCH_2Cl)(dppf)_2]$ and terminal thio alkyl complex $[Pt(SCH_2Cl)_2(dppf)]$ in chlorinated solvents [8]. The latter mononuclear product has also been reported by an oxidative reaction between $[Pt(dppf)_2]$ and bis(chloromethyl)disulfide [9-11]. The sulfurized dppf on reaction with $[PdCl_2(PhCN)_2]$ yielded a mononuclear complex of composition $[PdCl_2(L)]$ ($L = Fe(C_5H_4P(S)Ph_2)(C_5H_4PPh_2)$) [4]. However the reaction of other chelating phosphine precursors $[M(P\cap P)_2]$ ($M = Pd, Pt$; $P\cap P = dppe, dppp$) with Ar_2E_2 ($Ar = C_5H_4N, C_5H_3N_2$; $E = Se, Te$) yielded a mononuclear compound $[M(EAr)_2(P\cap P)]$ as well as the binuclear product $[M_2(\mu-TeAr)_2(P\cap P)_2]$ ($M = Pd, Pt$; $P\cap P = dppe$; $Ar = C_5H_4N$) [12-14] which decomposes to high nuclearity complex along with unidentified product in chlorinated solvents [15-17]. The facile transformation of bridging chalcogenolates to terminal one is predominantly due to high nucleophilicity of the complex which makes them highly susceptible towards attack on chlorinated solvent [8, 16, 18, 19].

In this report, we describe the synthesis and characterization of various ferrocenyl phosphine based palladium and platinum(II) selenolate complexes derived from pyridyl and methyl substituted pyrimidyl selenolate ligand and studies their reactivity in chlorinated solvent.

2. Experimental:

All manipulations were carried out under a nitrogen atmosphere by using standard Schlenk technique. Solvents used in the reactions were degassed with nitrogen and distilled using standard procedures. The precursor complex $[\text{PtCl}_2(\text{PhCN})_2]$ [20] and the ligands $(\text{SeC}_5\text{H}_4\text{N})_2$, [21] and $\{\text{SeC}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2$ [17] were prepared by literature methods. The metal precursors PdCl_2 , and dppf were procured from Strem chemicals. The ^1H , and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra were recorded on a Bruker Avance-II spectrometer operating at 500, 202.40 MHz, respectively and referenced for the solvent peak of chloroform at δ 7.26 for ^1H and externally to 85% H_3PO_4 (0.00 ppm) for ^{31}P NMR. Elemental analyses were carried out on a Thermo Fischer Flash EA1112 CHNS analyzer.

$[\text{Pt}\{\text{SeC}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2(\text{dppf})].0.5\text{C}_6\text{H}_6$ (**1a**) were collected at 173K on using a Rigaku FR-XUltrahigh Brilliance Microfocus RA generator/confocal optics and XtaLAB P200 diffractometer, Mo- K_α radiation ($\lambda = 0.71075 \text{ \AA}$). Intensity data were collected using ω steps accumulating area detector images spanning at least a hemisphere of reciprocal space. All data were corrected for Lorentz polarization effects. A multi-scan absorption correction was applied by using Crystal Clear [22]. The structures were solved by heavy-atom Patterson methods [23], expanded using Fourier techniques and least-squares refinement [23] was on F^2 using data converged with unweighted and weighted agreement factors. The non-hydrogen atoms were refined anisotropically and riding a model was used for hydrogen atoms. Molecular structures were drawn using ORTEP [24]. Crystallographic and structural determination data are listed in Table 1.

2.1 Synthesis of complexes

2.1.1 Preparation of $[Pd\{2-Se-C_4H(4,6-Me)_2N_2\}_2(dppf)]$ (**1a**):

To a benzene-methanolic solution (10 cm^3) of $NaSeC_4H(4,6-Me)_2N_2$ [freshly prepared from 45 mg, 0.12 mmol of $(SeC_4H(4,6-Me)_2N_2)_2$, methanol (5 cm^3) and $NaBH_4$ (10 mg, 0.26 mmol)] a solution (10 cm^3) of $[PdCl_2(dppf)]$ (98 mg, 0.12 mmol) was added in the same solvents with stirring, which continued for 4 hours at room temperature. The wine red solution was concentrated to 5 cm^3 under reduced pressure. A reddish brown precipitate was formed which was washed thoroughly with hexane and diethyl ether. The residue was recrystallized from dichloromethane and few drops of hexane to afford reddish brown powder. (yield: 54 mg, 48%; m.p.: $151\text{ }^\circ\text{C}$). Anal. Calcd for $C_{46}H_{42}FeN_4P_2PdSe_2$: C, 53.49; H, 4.10; N, 5.42%. Found: C, 53.88; H, 3.94; N, 5.39%. UV-Vis: 270 nm (br), 405 nm (sh). 1H NMR ($CDCl_3$) δ : 1.25 (s, 6H, Me), 1.74 (s, 6H, Me), 4.31 (br, 4H, C_5H_4), 4.69 (br, 4H, C_5H_4), 6.86 (d, 4.8Hz, 2H, Ph), 7.49-7.56 (m, 8H, Ph), 7.60-7.64 (td, 2.6Hz, 4H, Ph), 7.66-7.72 (m, 6H, Ph), 7.80 (d, 4.8Hz, 2H, $C_4H(4,6-Me)_2N_2$); $^{31}P\{^1H\}$ NMR ($CDCl_3$) δ : 28.6 ppm.

2.1.2 Preparation of $[Pd(SeC_5H_4N)_2(dppf)]$ (**1b**):

Prepared in a similar manner to complex **1a** using $[PdCl_2(dppf)]$ (65 mg, 0.08 mmol) and $(SeC_5H_4N)_2$ (25 mg, 0.08 mmol) as the ligand precursor and recrystallized from dichloromethane-hexane to give a brown powder. (yield 67 mg, 92%; m.p.: $110\text{ }^\circ\text{C}$). Anal. Calcd. for $C_{44}H_{36}N_2P_2FePdSe_2$: C, 54.21; H, 3.72; N, 2.87%. Found: C, 54.61; H, 3.67; N, 2.47%. UV-Vis (λ_{max}): 278 nm (sh), 328 nm (br). 1H NMR ($CDCl_3$) δ : 4.26 (t, 1.5 Hz, 4H, C_5H_4), 4.70 (t, 1.8 Hz, 4H, C_5H_4), 6.41 (d, 3.2 Hz, 2H, C_5H_4N), 7.34-7.41 (m, 8H, Ph), 7.58-7.64 (m, 4H, Ph),

7.68-7.73(m, 10H, o-Ph+ C₅H₄N) 7.80 (d, 4.5 Hz, 2H, C₅H₄N), 8.2(d, 4.8 Hz, 2H, C₅H₄N);
³¹P{¹H}NMR (CDCl₃) δ : 28.3 ppm.

2.1.3 Preparation of [Pt{2-Se-C₄H(4,6-Me)₂N₂}₂(dppf)] (2a):

Prepared in a similar manner to complex **1a**, using [PtCl₂(dppf)] (99 mg, 0.12 mmol) and Na{2-Se-C₄H(4,6-Me)₂N₂} [freshly prepared from {2-Se-C₄H(4,6-Me)₂N₂}₂ (45 mg, 0.12 mmol) in benzene and NaBH₄ (10 mg, 0.26 mmol) in methanol] and recrystallized from benzene-acetone to afford yellow crystals in 72% yield (80 mg), m.p.: 176 °C (dec.). Anal. Calcd for C₄₆H₄₂FeN₄P₂PtSe₂: C, 49.26; H, 3.77; N, 4.99%. Found: C, 49.68; H, 3.52; N, 4.74%. UV-Vis (λ_{max}) : 348 nm (br), 435 nm (sh). ¹H NMR (CDCl₃) δ : 1.26 (s, 6H, Me), 1.74 (s, 6H, Me), 4.19 (br, 4H, C₅H₄), 4.38 (br, 4H, C₅H₄), 7.28-7.35 (m, 4H, Ph), 7.38-7.48 (m, 8H, Ph), 7.67 (d, 4.5Hz, 2H, C₅HN₂), 7.82-7.87 (m, 8H, Ph) ; ³¹P{¹H}NMR (CDCl₃) δ : 12.6 [¹J(Pt-P) = 3183 Hz] ppm.

2.1.4 Preparation of [Pt₂(μ-SeC₅H₄N)₂(dppf)₂]Cl₂ (3):

To a benzene solution (15 cm³) of [PtCl₂(dppf)] (110 mg, 0.14 mmol), a methanolic solution (8 cm³) of Na(SeC₅H₄N) [freshly prepared from (C₅H₄N)₂Se₂ (42 mg, 0.14 mmol) and NaBH₄ (11 mg, 0.29 mmol)] was added and the mixture was stirred for 4 hours. The solvents were evaporated under vacuum and the residue was extracted with benzene and passed through celite. The ensuing solution on slow evaporation afforded a yellow powder (Yield 54 mg, 43%) m.p.: 118°C (dec.). Anal. Calcd. for C₇₈H₆₄N₂Cl₂P₄Fe₂Pt₂Se₂: C, 55.47; H, 3.82; N, 1.66%. Found: C, 55.78; H, 4.14; N, 1.88%. UV-Vis (λ_{max}) : 435 nm (br), 509 nm (br). ¹H NMR (CD₃OH) δ: 4.32 (br, 4H, C₅H₄), 4.67 (br, 4H, C₅H₄), 6.38 (d, 2.8 Hz, 2H, C₅H₄N), 7.32-7.42 (m, 8H, Ph), 7.56-7.66 (m, 4H, Ph+C₅H₄N), 7.81-7.88 (m, 8H, Ph+C₅H₄N), 8.48 (d, 5.0Hz, 2H,

C_5H_4N); $^{31}P\{^1H\}$ NMR (CD_3OH) δ : 12.3 [$^1J(Pt-P) = 3169Hz$] ppm. On recording the NMR of titled complex **3** in $CDCl_3$ Solution; 1H NMR ($CDCl_3$) δ : 4.14 (br, 4H, C_5H_4), 4.32 (br, 4H, C_5H_4), 7.08-7.13 (m, 4H, Ph), 7.33-7.39 (m, 9H, Ph+ C_5H_4N), 7.66-7.73 (m, 9H, Ph+ C_5H_4N), 8.48 (d, 4.1Hz, 1H, C_5H_4N); $^{31}P\{^1H\}$ NMR ($CDCl_3$) δ : 9.9 [$^1J(Pt-P) = 3762Hz$], 14.9 [$^1J(Pt-P) = 3149Hz$] ppm.

2.1.5 Preparation of $[Pt(Cl)(SeC_5H_4N)(dppf)]$ (**4**):

To a dichloromethane solution (15 cm^3) of $[PtCl_2(dppf)]$ (165 mg, 0.21 mmol) a methanolic solution (8 cm^3) of $Na(SeC_5H_4N)$ [freshly prepared from $(C_5H_4N)_2Se_2$ (65 mg, 0.21 mmol) and $NaBH_4$ (16 mg, 0.42 mmol)] was added and the mixture was stirred for 4 hour. The solvents were reduced *in vacuo* and the residue was extracted with dichloromethane which on slow evaporation afforded a yellow powder (Yield 139 mg, 73%) m.p. $161\text{ }^\circ C$ (dec.). Anal. Calcd. for $C_{39}H_{32}NCIP_2FePtSe$: C, 49.73; H, 3.42; N, 1.49%. Found: C, 49.48; H, 3.59; N, 1.58%. UV-Vis (λ_{max}) : 277 nm (sh), 307 nm (br). 1H NMR ($CDCl_3$) δ : 4.21 (br, 4H, C_5H_4), 4.34 (br, 4H, C_5H_4), 7.11-7.16 (m, 4H, Ph), 7.31-7.36 (m, 9H, Ph+ C_5H_4N), 7.68-7.73 (m, Ph+ C_5H_4N), 8.41 (d, 4.4Hz, 4H, C_5H_4N); $^{31}P\{^1H\}$ NMR ($CDCl_3$) δ : 10.1 [$^1J(Pt-P) = 3766Hz$], 14.6 [$^1J(Pt-P) = 3143Hz$] ppm.

3. Results and Discussion:

Treatment of $[PdCl_2(dppf)]$ with two equivalents of sodium aryl selenolate $NaSeAr$ ($Ar = C_5H_4N, C_4H(4,6-Me)_2N_2$) prepared by the reductive cleavage of the Se-Se bond of diselenide by $NaBH_4$ in methanol, afforded an orange mononuclear product of composition $[Pd(SeAr)_2(dppf)]$ ($Ar = C_4H(4,6-Me)_2N_2$: **1a**, C_5H_4N : **1b**, scheme 1). The ^{31}P NMR of both of the complexes showed one resonance at ~ 28 ppm. It is noteworthy that an oxidative reaction of $[Pd(dppe)_2]$

with dipyridyldiselenides resulted the similar mononuclear product $[\text{Pd}\{\text{SeC}_5\text{H}_3(3\text{-R})\text{N}\}_2(\text{dppe})]$ ($\text{R} = \text{H}, \text{Me}$) [16]. Similarly, reaction of platinum precursor with the sodium salt of methyl substituted pyrimidylselenolate in a benzene-methanol mixture gave rise to a yellow powder of composition $[\text{Pt}\{2\text{-Se-C}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2(\text{dppf})]$ (**2a**, Scheme 1). The observed coupling constant 3183 Hz is in good agreement with reported value for the mononuclear complexes [17, 25, 26].

The molecular structures of $[\text{Pt}\{2\text{-Se-C}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2(\text{dppf})] \cdot 0.5\text{C}_6\text{H}_6$ (**1a**) has been established by X-ray diffraction (Figures 2). Selected interatomic parameters for **1a**·0.5C₆H₆ are summarized in Tables 2. The complex **1a**, adopts the expected distorted square planar geometry around the platinum atom, defined by the coordination of two phosphorus and two selenium atoms. The Pt–Se distances [2.4578(10) and 2.4595(11) Å] fall in the middle of the range of reported Pt–Se distance for both complexes of the type $[\text{Pt}(\text{SeR})_2(\text{dppf})]$ (Pt–Se range 2.453–2.477 Å) [27–30] and $[\text{Pt}(\text{SeAr})_2(\text{P}\cap\text{P})]$ (Pt–Se range 2.418–2.511 Å) [27, 31–34]. The Pt–P distances [2.295(2) and 2.301(2) Å] are slightly longer than other reported Pt–P bonds for compounds of the type values for $[\text{Pt}(\text{SeAr})_2(\text{P}\cap\text{P})]$ (2.238–2.274 Å) [31–34]; however, when comparing against Pt(dppf) complexes with any selenolate ligands, $[\text{Pt}(\text{SeR})_2(\text{dppf})]$, the Pt–Se bond distances in **3a** fall into the known range (2.279–2.306 Å) [27, 32–34]. The Se–Pt–Se angle is acute [84.38(3)°] as a result of the bond angle imposed by the wide bite-angle of dppf [P–Pt–P 101.83(8)°]. These angles are similar to those found in $[\text{Pt}(\text{SeC}\equiv\text{C}(\text{CH}_2)_4\text{Me})_2(\text{dppf})] \cdot \text{C}_6\text{H}_6$ (Se–Pt–Se 85.08°, P–Pt–P 97.98°) [29].

The reaction of $[\text{PtCl}_2(\text{dppf})]$ with two equivalents of sodium 2-pyridylselenolate in a benzene-methanol mixture yielded a binuclear product of composition $[\text{Pt}_2(\mu\text{-SeC}_5\text{H}_4\text{N})_2(\text{dppf})_2]\text{Cl}_2$ (**3**). In contrast, a similar reaction in a dichloromethane-methanol mixture afforded exclusively a product of the type $[\text{Pt}(\text{Cl})(\text{SeC}_5\text{H}_4\text{N})(\text{dppf})]$ (**4**) (scheme 2). The ³¹P

NMR of complex **4** displayed two resonances at 14.6 and 10.1 ppm with $^1J(\text{Pt-P})$ values of 3143 and 3766 Hz, respectively. These two resonances, with different Pt-P coupling constants arise, due to the differing chemical environments of each phosphorus; one being *trans* to the selenolate, the other *trans* to the chloride [15, 35]. In an attempt to substitute both the chloride, the reaction has been performed between $[\text{PtCl}_2(\text{dppf})]$ and $[\text{Pb}(\text{SeC}_5\text{H}_4\text{N})]$ which also has resulted similar composition of complex **4**. It is noteworthy that similarly an oxidative addition reaction of $[\text{Pd}(\text{PPh}_3)_4]$ with dipyrindyl ditellurides gave an expected oxidative addition product $[\text{Pd}\{2\text{-TeC}_5\text{H}_3(3\text{-R})\text{N}\}_2(\text{PPh}_3)_2]$ (R = H, Me) which on keeping in CDCl_3 solution resulted derivative $[\text{PdCl}\{2\text{-Te}(\text{Cl})_2\text{C}_5\text{H}_3(3\text{-Me})(\text{PPh}_3)\}]$ [16]. Even the reaction of $[\text{Pd}_2\text{Cl}_2(\mu\text{-Cl}_2)(\text{PPh}_3)_2]$ with sodium salt of pyridyl telluroate afforded complex $[\text{PdCl}\{2\text{-TeC}_5\text{H}_3(3\text{-Me})(\text{PPh}_3)\}]$ [35]. The complex $[\text{Pt}_2(\mu\text{-SeC}_5\text{H}_4\text{N})_2(\text{dppf})_2]^{2+}$ (**3**) in CDCl_3 solution also exhibited similarly two phosphorus peaks corroborated with platinum satellites as discussed above for the complex **4** (scheme 2). The ^{31}P NMR of complex **3** in CD_3OH exhibited a single resonance at 12.6 ppm with a $^1J(\text{Pt-P})$ coupling of 3169 Hz, corresponding to a symmetrical phosphorus environment, with a selenolate ligand *trans* to the phosphine ligand [15, 26, 35, 36]. In contrast, in CDCl_3 solution, complex **3** exhibited two phosphorus peaks with almost identical chemical shifts and coupling to platinum as seen for complex **4**. This suggests, that in CDCl_3 solution, the cationic binuclear complex **3** converts to the mononuclear complex **4** (Scheme 2). The four membered ring of $\{\text{Pt}_2\text{Se}_2\}$ in complex **3** has been stabilized due to π bonding effect [8, 37]. The chlorination of seleno bridge, has been anticipated due to perturbation of π cloud in presence of electron-withdrawing group to weaken the Pt-Se bond and open up the four membered ring [38]. An ionic nature of complex **3** has also been monitored by molar conductivity measurement where complex $[\text{Pt}_2(\mu\text{-SeC}_5\text{H}_4\text{N})_2(\text{dppf})_2]^{2+}$ (**3**) showed a non electrolyte nature in CHCl_3 (1.8 μS

$\text{cm}^2 \text{mole}^{-1}$) however, it is 1:2 electrolyte in methanol ($133.2 \mu\text{S cm}^2 \text{mole}^{-1}$). This indicates the existence of neutral species in dichloromethane, while in polar solvent like CH_3OH complex adopts ionic structure. A similar reactivity pattern has been reported for the complex like $[\text{Pt}_2(\mu\text{-S})_2(\text{dppf})_2]^{2+}$ [8], $[\text{Pt}_2(\mu\text{-SeCH}_2\text{CH}_2\text{NMe}_2)_2(\text{dppe})_2]^{2+}$ [35, 36], and $[\text{Pt}_2(\mu\text{-TeC}_5\text{H}_3(3\text{-Me})_2(\text{dppe})_2]^{2+}$ (dppe=1,2-bis(diphenylphosphino)ethane) [15]. The rapid conversion of binuclear complex to mononuclear compound is well documented in literatures [8, 15, 18, 19]. The possibility of the formation of a mononuclear species is expected to be due to the strong trans effect of phosphine ligand which weakens the bridging chalcogenolate linkage and results in the formation of the mononuclear mixed halide/chalcogenolates complexes.

All complexes are colored and show absorption bands in the visible range of electronic spectra. The bands are red-shifted with respect to the diselenide ligands. The highest occupied molecular orbital (HOMO) is located on the selenolate ligands, whereas the lowest unoccupied molecular orbital (LUMO) is probably delocalised over the phosphine ligand [35].

4. Conclusion:

The substitution reaction of pyridyl/pyrimidyl selenolate with palladium and platinum(II) metal precursors $[\text{MCl}_2(\text{dppf})]$ ($\text{M} = \text{Pd}, \text{Pt}$) afforded mono or binuclear products. The pyrimidyl derived selenolate gave rise solely to mononuclear complexes with both platinum and palladium, and the pyridylselenolate gave rise to equivalent complexes with palladium. In contrast, on reaction with platinum, the pyridyl derived selenolate yielded either binuclear, $[\text{Pt}_2(\mu\text{-SeC}_5\text{H}_4\text{N})_2(\text{dppf})_2]\text{Cl}_2$, or mononuclear $[\text{Pt}(\text{Cl})(\text{SeC}_5\text{H}_4\text{N})(\text{dppf})]$ products. The isolation of these mononuclear $[\text{M}(\text{SeAr})_2(\text{dppf})]$ ($\text{M} = \text{Pd}, \text{Pt}$; $\text{Ar} = \text{C}_5\text{H}_4\text{N}, \text{C}_4\text{H}(2,6\text{-Me})\text{N}_2$) and ionic compounds depend upon the substituent present on the phenyl ring and the solvent involved during the course of the reaction. The instant conversion of binuclear to mono-nuclear species is mainly

aroused due to nucleophilic attack of chlorinated solvent on selenolate bridged binuclear complex.

Acknowledgement:

One of the authors (RSC) is grateful to DST for the financial support under the DST young scientist scheme YSS/2014/000797. We thank to Drs. V. K. Jain and Vijay Joshi for encouragement of this work and valuable suggestions for the preparation of this article.

Supporting Information

CCDC-Nos. 1574352-1574354 for $[\text{PdCl}_2(\text{dppf})]\cdot\text{CH}_2\text{Cl}_2$, $[\text{PtCl}_2(\text{dppf})]\cdot 0.5\text{CH}_2\text{Cl}_2$ and $[\text{Pt}\{2\text{-Se-C}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2(\text{dppf})]\cdot 0.5\text{C}_6\text{H}_6$, respectively contain the supplementary crystallographic data for this paper. These data can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html or from the Cambridge Crystallographic Data Center, 12 Union Road, Cambridge CB2 1EZ, UK [Fax: + 44-1223/336-033; E-mail: deposit@ccdc.cam.ac.uk].

References:

- 1) S. W. Chien, T. S. A. Hor in *Ferrocenes: Ligands, Materials and Biomolecules* (Ed.: P. Štěpnička), Wiley, Chichester, 2008, chapter 2, pp. 33–116; b) K.-S. Gan, T. S. A. Hor in *Ferrocenes: Homogeneous Catalysis, Organic Synthesis Materials Science* (Eds.: A. Togni, T. Hayashi), Wiley-VCH, Weinheim, 1995, chapter 1, pp. 3–104; c) G. Bandoli, A. Dolmella, *Coord. Chem. Rev.* 209 (2000) 161–196.

- 2) A) Y. K. Yan, H.S.O. Chan, T.S.A. Hor, K.-L. Tan, L.-K. Liu and Y.S. Wen, *J. Chem. Soc. Dalton Trans.* (1992) 423–426; b) L.-T. Phang, S.C.F. Au-Yeung, T.S.A. Hor, S.B. Khoo, Z.-Y. Zhou, T.C.W. Mak, *J. Chem. Soc. Dalton Trans.* (1993) 165; c) P.M.N. Low, Y.K. Yan, H.S.O. Chan, T.S.A. Hor, *J. Organomet. Chem.* 454 (1993) 205–209; d) P.F. Kelly, A.M.Z. Slawin, D.J. Williams, J.D. Woollins, *Polyhedron* 7 (1988) 1925–1930;
- 3) B.S. Haggerty, C.E. Housecroft, A. L. Rheingold, B.A.M. Shaykh, *J. Chem. Soc. Dalton Trans.* (1991) 2175–2184.
- 4) R. Broussier, E. Bentabet, M. Laly, P. Richard, L.G. Kuz'mina, P. Serp, N. Wheatley, P. Kalck, B. Gautheron, *J. Organomet. Chem.* 613 (2000) 77–85.
- 5) P. Štěpnička, I. Cisařova, J. Schulz, *Organometallics* 30 (2011) 4393–4403; b) P. Štěpnička, I. Cisařova, *J. Organomet. Chem.* 716 (2012) 110–119.
- 6) P. Štěpnička, I. Cisařova, *Dalton Trans.* 2013, 42, 3373–3389.
- 7) M. Zabransky, A. Machara, I. Cisařova and P. Štěpnička, *Eur. J. Inorg. Chem.* (2017) 4850–4860.
- 8) M.-C. F. Lam, K.F. Mok, P.-H. Leung, T.S. A. Hor, *J. Organomet. Chem.* 476 (1994) C 32–34.
- 9) P.R. Schleyer, E.D. Jemmis and G.W. Spitznagel, *J. Am. Chem. Soc.*, 107 (1985) 6393–6394.
- 10) (a) J.S. Grossert and R.F. Langler, *Can. J. Chem.*, 55 (1977) 407; (b) I.W.J. Still and G.W. Kutney, *Tetrahedron Lett.*, 22 (1981) 1939–1940.
- 11) Z.-G. Fang, S.-C. Ng and T.S.A. Hor, *J. Organomet. Chem.* 484 (1994) 17.
- 12) R.S. Chauhan, N. Shivran, *RSC Advances* 7 (2017) 55175–55198.

- 13) V.K. Jain, R.S. Chauhan, *Coord. Chem. Rev.* 306 (2016) 270–301.
- 14) G. Kedarnath, V. K. Jain, *Coord. Chem. Rev.* 257 (2013) 1409–1435.
- 15) R.S. Chauhan, G. Kedarnath, A. Wadawale, A.L. Rheingold, A. Munoz-Castro, R. Arratia-Perez and V. K. Jain, *Organometallics*, 31 (2012) 1743–1750.
- 16) R. S. Chauhan, G. Kedarnath, A. Wadawale, A.M.Z. Slawin and V.K. Jain, *Dalton Trans.* 42 (2013) 259–269.
- 17) R. S. Chauhan, R. K. Sharma, G. Kedarnath, D. B. Cordes, A. M. Z. Slawin and V. K. Jain, *J. Organomet. Chem.* 17 (2012) 180–186.
- 18) (a) T.S.A. Hor, DPhil Thesis, University of Oxford, UK, 1983; (b) C.E. Briant, C.J. Gardner, T.S.A. Hor, N.D. Howells and D.M.P. Mingos, *J. Chem. Soc., Dalton Trans.*, (1984) 2645; (c) R.R. Gukathasan, R.H. Morris and A. Walker, *Can. J. Chem.*, 61 (1983) 2490.
- 19) R. Ugo, G. La Monica, S. Cenini, A. Segre and F. Conti, *J. Chem. Soc. A*, (1971) 522–528.
- 20) M.S. Inkpen, A.J.P. White, T. Albrecht, N.J. Long, *Chem. Commun.* 49 (2013), 5663–5665.
- 21) K. K. Bhasin, V. Arora, S. K. Sharma, P. Venugopalan, *Applied Organomet. Chem.* 119 (2005) 161–166.
- 22) P. T. Beurskens, G. Beurskens, R. de Gelder, S. Garcia-Granda, R. O. Gould, R. Israel, J. M. M. Smits, DIRDIF-99, Crystallography Laboratory, University of Nijmegen, The Netherlands, 1999.
- 23) Crystal Clear-SM Expert v2.1. Rigaku Americas, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan, 2015.

- 24) C.K. Johnson, ORTEP II, Report ORNL-5136, Oak Ridge National Laboratory, Oak Ridge TN, 1976.
- 25) R. Oilunkaniemi, R.S. Laitinen and M. Ahlgren, *J. Organomet. Chem.* 595 (2000) 232–240.
- 26) S. Narayan, V.K. Jain, B. Varghese, *J. Chem. Soc., Dalton Trans.* (1998) 2359–2366.
- 27) R. S. Chauhan, G. Kedarnath, A. Wadawale, D. K. Maity, J. A. Golen, A. L. Rheingold, V. K. Jain, *J. Organomet Chem.*, 737 (2013) 40–46.
- 28) T. Niksch, H. Görls, M. Friedrich, R. Oilunkaniemi, R. Laitinen and W. Weigand, *Eur. J. of Inorg. Chem.*, (2010) 74–94.
- 29) S. Schäfer, C. Moser, J. J. Tirré, M. Nieger and R. Pietschnig, *Inorganic Chemistry* 44 (2005) 2798-2802.
- 30) Minna M. Karjalainen, Wolfgang Weigand, Raija Oilunkaniemi, Risto S. Laitinen, *Polyhedron* 101 (2015) 244-250.
- 31) V.K. Jain, S. Kannan, R.J. Butcher, J.P. Jasinski, *J. Chem. Soc. Dalton Trans.* (1993) 1509–1513.
- 32) K.V. Vivekananda, S. Dey, A. Wadawale, N. Bhuvanesh and V.K. Jain, *Dalton Trans.*, 42 (2013) 14158-14167.
- 33) N.K. Lokanath, H.C. Devarajegowda, S.M. Anadalwar, J.S. Prasad, S. Narayan and V.K. Jain, *Anal. Sci.* 17 (2001) 565–566.
- 34) M. Risto, E.M. Jahr, M.S. Hannu-Kurre, R. Oilunkaniemi and R.S. Laitinen, *J. Organomet. Chem.* 692 (2007) 2193–2204.
- 35) S. Dey, V.K. Jain, J. Singh, V. Trehan, K.K. Bhasin, B. Varghese, *Eur. J. Inorg. Chem.* (2003) 744–750

- 36) S. Dey, V. K. Jain, A. Knoedler, A. Klein, W. Kaim, S. Zallis, *Eur. J. Inorg. Chem.* (2001) 2965–2973.
- 37) (a) J. Chatt and F.A. Hart, *J. Chem. Soc.*, 12 (1960) 807; (b) T.S.A.Hor and A.L.C. Tan, *Inorg. Chim. Acta*, 142 (1988) 173.
- 38) E.M. Padilla, J.A. Golen, P.N. Richmann and C.M. Jensen, *Polyhedron*, 10 (1991) 1343.

ACCEPTED MANUSCRIPT

Table 1. Crystallographic and structural determination data for [Pt{2-Se-C₄H(4,6-Me)₂N₂}₂(dppf)]. 0.5C₆H₆ (**1a**)

Complex	1a
Chemical formula	C ₄₉ H ₄₅ FeN ₄ P ₂ PtSe ₂
Formula wt.	1160.73
Crystal size (mm ³)	0.09 x 0.03 x 0.01
Crystal system	Triclinic
Space group	<i>P</i> $\bar{1}$ ₁
Unit cell dimensions	
a (Å)	11.3463(18)
b (Å)	15.7103(15)
c (Å)	15.896(2)
α (°)	117.826(7)
β (°)	101.96(2)
γ (°)	93.179(15)
Volume (Å ³)	2413.3(7)
ρ_{cacl} , g cm ⁻³	1.597
Z	2
μ (mm ⁻¹)/F(000)	4.795/1138
Limiting indices	-13 ≤ h ≤ 13 -18 ≤ k ≤ 18 -17 ≤ l ≤ 19
θ for data collection(°)	1.864-25.388
No of reflections collected	29339
No of independent reflection (R_{int})	8733 (0.0749)
Data/restraints/parameters	8733/36/551
Final R_1 , wR_2 indices ($I > 2\sigma I$)	0.0487, 0.1081
R_1 , wR_2 (all data)	0.0881, 0.1200
Goodness of fit on F^2	0.933

Table 2: Selected bond lengths (Å) and angles (°) of [Pt{2-Se-C₄H(4,6-Me)₂N₂}₂(dppf)]
1a·0.5C₆H₆

Pt(1)-P(1)	2.301(2)	Pt(1)-P(2)	2.295(2)
Pt(1)-Se(1)	2.4595(11)	Pt(1)-Se(2)	2.4578(10)
Se(1)-C(35)	1.901(10)	Se(2)-C(43)	1.922(10)
P(2)-Pt(1)-P(1)	101.83(8)	Se(1)-Pt(1)-Se(2)	84.38(3)
P(1)-Pt(1)-Se(1)	85.52(6)	P(2)-Pt(1)-Se(2)	88.84(6)
P(1)-Pt(1)-Se(2)	167.23(6)	P(2)-Pt(1)-Se(1)	171.66(6)

ACCEPTED MANUSCRIPT

Figure captions

Figure 1: ORTEP diagram of $[\text{Pt}\{2\text{-Se-C}_4\text{H}(\text{4,6-Me})_2\text{N}_2\}_2(\text{dppf})]\cdot 0.5\text{C}_6\text{H}_6$ (**1a** $\cdot 0.5\text{C}_6\text{H}_6$) with atomic numbering scheme. The ellipsoids were drawn at the 50% probability level. Hydrogen atoms and solvent molecules are omitted for clarity.

ACCEPTED MANUSCRIPT






Figure 1: ORTEP diagram of [Pt{2-Se-C₄H(4,6-Me)₂N₂}₂(dppf)].0.5C₆H₆ (**1a**) with atomic numbering scheme. The ellipsoids were drawn at the 50% probability level. Hydrogen atoms and solvent molecules are omitted for clarity.




M	X	R	
Pd	N	Me	1a
Pd	C	H	1b
Pt	N	Me	2a

Scheme 1



Scheme 2



The bis(diphenylphosphino)ferrocene (dppf) derived palladium analogue of $[\text{PdCl}_2(\text{dppf})]$ on reaction with the sodium salt of pyridyl/pyrimidyl selenolate yielded mononuclear *cis* configured complex $[\text{Pd}(\text{SeAr})_2(\text{dppf})]$ ($\text{Ar} = \text{C}_5\text{H}_4\text{N}$, $\text{C}_4\text{H}(4,6\text{-Me})_2\text{N}_2$), as did the platinum precursor $[\text{PtCl}_2(\text{dppf})]$ with methyl substituted pyrimidyl salt giving $[\text{Pt}\{\text{SeC}_4\text{H}(4,6\text{-Me})_2\text{N}_2\}_2(\text{dppf})]$. In contrast, the reaction of platinum precursor with the sodium salt of pyridylselenolate gave two different complexes $[\text{Pt}_2(\mu\text{-SeC}_5\text{H}_4\text{N})_2(\text{dppf})_2]^{2+}$ and $[\text{Pt}(\text{Cl})(\text{SeC}_5\text{H}_4\text{N})(\text{dppf})]$ depending in the reaction solvent.

Highlights

- Studied the reaction of $[\text{PdCl}_2(\text{dppf})]$ with the sodium salt of pyridyl/pyrimidyl selenolate.
- Reaction of $[\text{PtCl}_2(\text{dppf})]$ with sodioim salt of pyridyl selenolate ligands.
- Isolation of binuclear and mononuclear products depending upon the solvent.

ACCEPTED MANUSCRIPT