

Enumeration of idempotents in planar diagram monoids

Igor Dolinka,¹ James East,² Athanasios Evangelou,³ Des FitzGerald,³
Nicholas Ham,³ James Hyde,⁴ Nicholas Loughlin,⁵ James D. Mitchell,⁴

August 10, 2018

Abstract

We classify and enumerate the idempotents in several planar diagram monoids: namely, the Motzkin, Jones (a.k.a. Temperley-Lieb) and Kauffman monoids. The classification is in terms of certain vertex- and edge-coloured graphs associated to Motzkin diagrams. The enumeration is necessarily algorithmic in nature, and is based on parameters associated to cycle components of these graphs. We compare our algorithms to existing algorithms for enumerating idempotents in arbitrary (regular $*$ -) semigroups, and give several tables of calculated values.

Keywords: Diagram monoids, partition monoids, Motzkin monoids, Jones monoids, Temperley-Lieb monoids, Kauffman monoids, idempotents, enumeration.

MSC: 05E15, 20M20, 20M17, 05A18.

1 Introduction

Diagram monoids arise in numerous branches of mathematics and science, including representation theory, statistical mechanics and knot theory [5, 18, 22–24, 26, 28, 39]. Many studies of diagram monoids have been combinatorial in nature [8–10, 12, 14, 15, 27], and idempotents have played a large role in several of these works. Historically, it is interesting to note that products of idempotents [12, 13, 27] were understood several years before the idempotents themselves were [9]; this is largely due to the fact that diagram monoids have natural anti-involutions that give them regular $*$ -semigroup structures [13, 34], meaning that arbitrary idempotents are products of simpler idempotents known as projections (see Section 2 for definitions).

In [9], classifications and enumerations were given for the idempotents in the partition, Brauer and partial Brauer monoids, and also for the idempotent basis elements in the corresponding diagram algebras. The current paper continues in this direction, focusing this time on planar diagram monoids, such as the Motzkin, Jones (a.k.a. Temperley-Lieb) and Kauffman monoids. However, the methods employed here are necessarily different to those of [9], as the planarity constraint means that the set symmetries used to study the monoids in [9] are no longer available. In fact, we believe that enumeration of the planar idempotents is inimical to closed-form solution; for one thing, the highly complex meandric numbers [6, 7] occur during the enumeration of Kauffman idempotents, as noted below in Section 5. We seek to fill this gap by presenting methods for computing the numbers of idempotents in the Motzkin, Jones and Kauffman monoids, with attention given to efficiency of the algorithms involved.

The article is organised as follows. In Section 2, we describe two existing algorithms for enumerating the idempotents of arbitrary finite (regular $*$ -) semigroups, commenting on their respective complexities in general and in the context of the Jones and Motzkin monoids in particular. Section 3 develops a theory of idempotents in the Motzkin, Jones and Kauffman monoids. A key role is played by certain graphs, called interface graphs, associated to arbitrary Motzkin elements. These graphs are used to classify the idempotents in Proposition 3.4 and Corollaries 3.5 and 3.6, and then to enumerate them in Theorems 3.16, 3.17 and 3.19; see also Proposition 3.15. Section 4 presents a number of algorithms, based on the theory developed in

¹Department of Mathematics and Informatics, University of Novi Sad, Trg Dositeja Obradovića 4, 21101 Novi Sad, Serbia. *Email:* dockie@dmi.uns.ac.rs

²Centre for Research in Mathematics, School of Computing, Engineering and Mathematics, University of Western Sydney, Locked Bag 1797, Penrith NSW 2751, Australia. *Email:* J.East@uws.edu.au

³School of Mathematics and Physics, University of Tasmania, Private Bag 37, Hobart 7001, Australia. *Emails:* aoost@utas.edu.au, D.FitzGerald@utas.edu.au, nicholas.ham@utas.edu.au

⁴School of Mathematics and Statistics, University of St Andrews, St Andrews, UK. *Emails:* jth4@st-andrews.ac.uk, jdm3@st-andrews.ac.uk

⁵School of Mathematics and Statistics, Newcastle University, Newcastle NE1 7RU, UK. *Email:* n.j.loughlin@newcastle.ac.uk

Section 3, that calculate the number of idempotents in the Jones, Kauffman and Motzkin monoids; C++ implementations of these algorithms can be found at [30]. Finally, Section 5 gives several tables of calculated values, including comparative run-times of the various algorithms.

The reader is referred to the monographs of Higgins [19] and Howie [20] for background on semigroups in general, and to the introduction of [9] and references therein—in particular to foundational articles of Jones [23], Martin [28] and Halverson and Ram [18]—for background and relevant detail on the partition, Brauer and partial Brauer monoids.

2 Existing algorithms

In this section, we describe two existing approaches to counting idempotents in semigroups such as those we study in this article. The first applies to any finite semigroup, while the second applies to any finite regular $*$ -semigroup; see below for the definitions. In Section 5, we will discuss the performance of these two approaches when applied to the diagram monoids we are concerned with, and we will compare them with the new algorithms presented in Section 4.

The first approach to counting idempotents in an arbitrary finite semigroup S is simple: create the elements of S , and then check if $x^2 = x$ for each $x \in S$; see Algorithm 1. If the semigroup S is generated by $A \subseteq S$, then S can be enumerated using the Froidure-Pin Algorithm [16]. If the complexity of multiplying elements in S is assumed to be constant, then the complexity of the Froidure-Pin Algorithm is $O(|S||A|)$, so Algorithm 1 has complexity $O(|S||A| + |S|) = O(|S||A|)$.

Algorithm 1 Count idempotents in a semigroup S

```

1:  $n := 0$ 
2: for  $x \in S$  do
3:   if  $x^2 = x$  then
4:      $n \leftarrow n + 1$ 
5: return  $n$ 

```

The approach just described requires the creation of each element of S in order to check which elements are idempotents. When S is very large, this can be impractical, in terms of both space and time. The second approach improves on the first in the case that S is a regular $*$ -semigroup. To describe it, we must first recall some background.

Recall from [34] that a semigroup S is a *regular $*$ -semigroup* if there is a unary operation $*$: $S \rightarrow S$ such that $x^{**} = x$, $(xy)^* = y^*x^*$ and $xx^*x = x$ for all $x, y \in S$. For the remainder of this section, we fix a finite regular $*$ -semigroup S . Recall that *Green's relations* \mathcal{R} , \mathcal{L} , \mathcal{J} , \mathcal{H} and \mathcal{D} are defined on S as follows. Let $x, y \in S$. We say that $x \mathcal{R} y$ if $xS = yS$, that $x \mathcal{L} y$ if $Sx = Sy$, and that $x \mathcal{J} y$ if $SxS = SyS$. The relation \mathcal{H} is defined to be the intersection of \mathcal{R} and \mathcal{L} , while \mathcal{D} is defined to be the join of \mathcal{R} and \mathcal{L} ; that is, \mathcal{D} is the least equivalence on S containing both \mathcal{R} and \mathcal{L} . It is well known that $\mathcal{D} = \mathcal{R} \circ \mathcal{L} = \mathcal{L} \circ \mathcal{R}$, and that $\mathcal{D} = \mathcal{J}$ since S is finite; see [20, Chapter 2] for more background on Green's relations. We write $E(S) = \{x \in S : x^2 = x\}$ for the set of all idempotents of S . An element $x \in S$ is called a *projection* if $x^2 = x = x^*$. The set of all projections of S is denoted by $\text{Proj}(S)$. The next result is true of any regular $*$ -semigroup, whether finite or infinite; proofs of the various parts may be found in [21, 34, 35].

Lemma 2.1. *Let S be a regular $*$ -semigroup. Then*

- (i) $\text{Proj}(S) = \{xx^* : x \in S\} = \{x^*x : x \in S\}$,
- (ii) $E(S) = \{xy : x, y \in \text{Proj}(S)\}$,
- (iii) every element of S is \mathcal{R} -related to a unique projection,
- (iv) every element of S is \mathcal{L} -related to a unique projection,
- (v) for any $x \in \text{Proj}(S)$ and any $a \in S$, $a^*xa \in \text{Proj}(S)$,
- (vi) for any $x, y \in S$, $x \mathcal{R} y$ if and only if $xx^* = yy^*$,
- (vii) for any $x, y \in S$, $x \mathcal{L} y$ if and only if $x^*x = y^*y$. □

It follows from Lemma 2.1(v), and the identity $(ab)^* = b^*a^*$, that S has a right action on $\text{Proj}(S)$, defined by

$$x \cdot a = a^*xa \quad \text{for } x \in \text{Proj}(S) \text{ and } a \in S. \quad (2.2)$$

For $x \in \text{Proj}(S)$, we will write

$$[x] = \{y \in \text{Proj}(S) : x = y \cdot a \text{ and } y = x \cdot b \text{ for some } a, b \in S\}$$

for the *strongly connected component* of x under the action (2.2). For any subset $A \subseteq S$, we will write $E(A) = A \cap E(S)$ and $\text{Proj}(A) = A \cap \text{Proj}(S)$ for the set of all idempotents and projections belonging to A , respectively. If \mathcal{K} is any of Green's relations, and if $x \in S$, we denote the \mathcal{K} -class of x by K_x . Since S is finite, it follows that S has the *stability* property: namely, for any $x, y \in S$, $xy \mathcal{D} x$ implies $xy \mathcal{R} x$, and $xy \mathcal{D} y$ implies $xy \mathcal{L} y$; see [36, Section A.2]. The various parts of the next lemma may be known, but we give proofs for completeness.

Lemma 2.3. *Let S be a finite regular $*$ -semigroup. Then*

- (i) *for any $x \in \text{Proj}(S)$, $[x] = \text{Proj}(D_x) = \{y \in \text{Proj}(S) : x \mathcal{D} y\}$,*
- (ii) *for any \mathcal{D} -class D of S , and for any projections $x, y \in \text{Proj}(D)$, $R_x \cap L_y$ contains an idempotent if and only if $xy \mathcal{D} x$, in which case this idempotent is xy ,*
- (iii) *for any \mathcal{D} -class D of S , the number of idempotents in D is equal to the cardinality of the set*

$$\{(x, y) : x, y \in \text{Proj}(D), xy \mathcal{D} x\}.$$

Proof. (i). Let $x \in \text{Proj}(S)$. If $y \in [x]$, then $x = y \cdot a = a^*ya$ and $y = x \cdot b = b^*xb$ for some $a, b \in S$, so that $x \mathcal{J} y$, whence $x \mathcal{D} y$ as S is finite. Conversely, suppose $y \in \text{Proj}(S)$ and $x \mathcal{D} y$. Then $x \mathcal{R} a$ and $a \mathcal{L} y$ for some $a \in S$. By Lemma 2.1(vi) and (vii), and since $x, y \in \text{Proj}(S)$, we obtain $aa^* = xx^* = x$ and $a^*a = y^*y = y$. It follows that $y = a^*a = a^*aa^*a = a^*xa = x \cdot a$, and similarly $x = y \cdot a^*$, so that $y \in [x]$.

(ii). Suppose first that $R_x \cap L_y$ contains some idempotent e . Since $e \mathcal{R} x$, Lemma 2.1(vi) gives $ee^* = xx^* = x$, and similarly $e^*e = y$. Then $e = ee^*e = e(ee^*)^*e = (ee^*)(e^*e) = xy$. Also, $xy = e \mathcal{R} x$ implies $xy \mathcal{D} x$.

Conversely, suppose $xy \mathcal{D} x$. Then stability gives $xy \mathcal{R} x$. Since $xy \mathcal{D} x \mathcal{D} y$, stability also gives $xy \mathcal{L} y$. Thus, $xy \in R_x \cap L_y$. Lemma 2.1(ii) gives $xy \in E(S)$.

(iii). First note that the number of idempotents in D is equal to the number of \mathcal{H} -classes in D containing an idempotent, since each \mathcal{H} -class contains at most one idempotent. If $a \in D$, then $H_a = R_a \cap L_a$, and by Lemma 2.1(iii) and (iv) we have $R_a = R_x$ and $L_a = L_y$ for unique projections $x, y \in \text{Proj}(D)$. That is, every \mathcal{H} -class in D is equal to $R_x \cap L_y$ for unique projections $x, y \in \text{Proj}(D)$. By part (ii), just proved, this \mathcal{H} -class contains an idempotent if and only if $xy \mathcal{D} x$. \square

Parts (i) and (iii) of Lemma 2.3 form the basis of the second approach to computing the number of idempotents in the finite regular $*$ -semigroup S , given by a generating set $A \subseteq S$; see Algorithm 2. Roughly speaking, the steps of this algorithm are:

- (1) We first create $\text{Proj}(S)$.
- (2) We then create the sets $\text{Proj}(D)$, as D runs over the set of all \mathcal{D} -classes of S .
- (3) For each \mathcal{D} -class D , we then find the cardinality of the set given in Lemma 2.3(iii), and sum over all D .

Step (1) can be achieved using the action from (2.2) in a simple orbit algorithm whose input is the generators A ; see [11, Algorithm 1]. If the complexity of determining $x \cdot a$ is assumed to be constant, then the complexity of [11, Algorithm 1], and hence the complexity of Step (1), is $O(|\text{Proj}(S)||A|)$.

By Lemma 2.3(i), the sets $\text{Proj}(D)$ correspond to the strongly connected components of the action of S on $\text{Proj}(S)$ given in (2.2). These can be found using standard algorithms from graph theory, such as Tarjan's [38] or Gabow's [17], for example; see also the monograph of Sedgewick [37]. The complexity of these algorithms, and thus the complexity of Step (2), is $O(|\text{Proj}(S)| + |A|)$, which is bounded above by $O(|\text{Proj}(S)||A|)$, the complexity of Step (1).

If the \mathcal{D} -classes of S are D_1, \dots, D_r , and if these \mathcal{D} -classes have m_1, \dots, m_r projections, respectively, then Step (3) involves $m_1^2 + \dots + m_r^2$ products and checks for \mathcal{D} -relatedness (modulo some optimisations discussed below). Thus, the total complexity of Algorithm 2 is

$$O(|\text{Proj}(S)||A| + m_1^2 + \dots + m_r^2).$$

The \mathcal{H} -classes in a single \mathcal{D} -class of a semigroup all have the same size [20, Lemma 2.2.3]. If the \mathcal{H} -classes in the \mathcal{D} -class D_i have size h_i , then $|D_i| = m_i^2 h_i$, since D_i has m_i \mathcal{R} - and \mathcal{L} -classes (Lemma 2.1(iii) and (iv)), and so

$$m_1^2 + \dots + m_r^2 \leq m_1^2 h_1 + \dots + m_r^2 h_r = |D_1| + \dots + |D_r| = |S|.$$

This upper bound is realised if and only if $h_i = 1$ for all i : i.e., if S is \mathcal{H} -trivial. In this worst case, the total complexity of Algorithm 2 is $O(|\text{Proj}(S)||A| + |S|)$. When we compare this to the complexity of Algorithm 1, which we noted above was $O(|S||A|)$, we see that Algorithm 2 has a significant advantage if $|\text{Proj}(S)|$ is small relative to $|S|$. Note that $|\text{Proj}(S)| = m_1 + \dots + m_r$.

We note that Algorithm 2, presented below, contains a number of simple optimisations. First, the \mathcal{H} -class $H_x = R_x \cap L_x$ of a projection $x \in \text{Proj}(D)$ always contains an idempotent: namely, x itself (see Line 6). Secondly, if $x, y \in \text{Proj}(D)$, then $xy \in D$ if and only if $yx = y^* x^* = (xy)^* \in D$, so we only need to test one of xy or yx for membership in D (see Lines 7–10). We also note that when S is any of the diagram monoids we consider, the \mathcal{D} -relation is given by equality of the *ranks* of elements of S , and is easily checked computationally; see Section 3 for the definition of rank, and also [10, 25, 40] for more on Green's relations on diagram monoids. Finally, we note that Algorithm 2 can be derived from [11, Algorithm 10], which counts idempotents in a fixed \mathcal{R} -class.

Algorithm 2 Count idempotents in a regular $*$ -semigroup S

```

1: Find  $\text{Proj}(S)$ 
2: Find the strongly connected components  $C_1, \dots, C_r$  of the action of  $S$  on  $\text{Proj}(S)$  defined in (2.2)
3:  $n := 0$ 
4: for  $i \in \{1, \dots, r\}$  do
5:   if  $C_i = \{x_1, x_2, \dots, x_m\}$ 
6:      $n \leftarrow n + m$ 
7:   for  $j \in \{1, \dots, m\}$  do
8:     for  $k \in \{j + 1, \dots, m\}$  do
9:       if  $x_i x_j \mathcal{D} x_i$  then
10:         $n \leftarrow n + 2$ 
11: return  $n$ 

```

This paper mostly concerns the case in which S is a Jones, Motzkin or Kauffman monoid. As noted above, the definitions of these monoids are given in Section 3, but here we make some brief comments relevant to the current discussion. For each non-negative integer n , we have a regular $*$ -monoid \mathcal{J}_n (Jones) and \mathcal{M}_n (Motzkin). The sizes of these monoids, and the sizes of their sets of projections, are given (see [3, 10, 14]) by

$$|\mathcal{J}_n| = C_n, \quad |\mathcal{M}_n| = \mu(2n, 0), \quad |\text{Proj}(\mathcal{J}_n)| = \sum_{r=0}^n \frac{r+1}{n+1} \binom{n+1}{\frac{n-r}{2}}, \quad |\text{Proj}(\mathcal{M}_n)| = \sum_{r=0}^n \mu(n, r).$$

Here, $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n th Catalan number; we interpret a binomial coefficient $\binom{m}{k}$ to be 0 if k is not an integer between 0 and m ; and the *Motzkin triangle* numbers $\mu(n, r)$ are defined by the recurrence

$$\begin{aligned} \mu(0, 0) &= 1, & \mu(n, r) &= 0 & & \text{if } n < r \text{ or } r < 0, \\ \mu(n, r) &= \mu(n-1, r-1) + \mu(n-1, r) + \mu(n-1, r+1) & & & & \text{if } 0 \leq r \leq n \text{ and } n \geq 1. \end{aligned} \quad (2.4)$$

The numbers $\mu(n, r)$ are also given by the formula

$$\mu(n, r) = \sum_{j=0}^n \binom{n}{j} \left[\binom{n-j}{r+j} - \binom{n-j}{r+j+2} \right]. \quad (2.5)$$

See Sequences A000108, A001006, A026300 in [1]. Values of $|S|$ and $|\text{Proj}(S)|$ are given in Table 1 for $S = \mathcal{J}_n$ or \mathcal{M}_n with $n \leq 15$, by way of indicating the relative complexities of Algorithms 1 and 2. Run times are given in Section 5, further highlighting the advantage of Algorithm 2 over Algorithm 1 in these cases.

n	$ \mathcal{J}_n $	$ \text{Proj}(\mathcal{J}_n) $	$ \mathcal{M}_n $	$ \text{Proj}(\mathcal{M}_n) $
0	1	1	1	1
1	1	1	2	1
2	2	2	9	2
3	5	3	51	5
4	14	6	323	13
5	42	10	2188	35
6	132	20	15 511	96
7	429	35	113 634	267
8	1430	70	853 467	750
9	4862	126	6536 382	2123
10	16 796	252	50 852 019	6046
11	58 786	462	400 763 223	17 303
12	208 012	924	3192 727 797	49 721
13	742 900	1716	25 669 818 476	143 365
14	2674 440	3432	208 023 278 209	414 584
15	9694 845	6435	1697 385 471 211	1201 917

Table 1: The sizes of the Jones and Motzkin monoids, \mathcal{J}_n and \mathcal{M}_n , and of their sets of projections, $\text{Proj}(\mathcal{J}_n)$ and $\text{Proj}(\mathcal{M}_n)$.

Algorithms 1 and 2, as presented above, are both *embarrassingly parallel*. Parallel versions of the algorithms are implemented in the Semigroups package for GAP [31].

Finally, we note that the Kauffman monoid \mathcal{K}_n (also defined in Section 3) is infinite, and also not a regular $*$ -semigroup, so neither of the algorithms discussed in this section apply to it. It is possible to define a finite quotient of \mathcal{K}_n that has only one more idempotent than \mathcal{K}_n [25], but this quotient is still not a regular $*$ -semigroup, so only Algorithm 1 would apply.

3 Idempotents of planar diagram monoids

In this section, we define the diagram monoids we will be concerned with, before describing methods to classify and enumerate the idempotents of these monoids. The classification involves certain graphs, called *interface graphs*, associated to arbitrary Motzkin elements. The enumeration is based on natural parameters associated to certain cycle components of the interface graphs, as well as a map that sends Motzkin idempotents to lower-rank idempotents.

3.1 Definitions and preliminaries

Let n be a positive integer, and write $\mathbf{n} = \{1, \dots, n\}$ and $\mathbf{n}' = \{1', \dots, n'\}$. The *partition monoid of degree n* , denoted \mathcal{P}_n , is the monoid of all set partitions of $\mathbf{n} \cup \mathbf{n}'$ under a product described below. Thus, an element of \mathcal{P}_n is a set $\alpha = \{A_1, \dots, A_k\}$, for some k , where the A_i are pairwise disjoint non-empty subsets of $\mathbf{n} \cup \mathbf{n}'$ whose union is all of $\mathbf{n} \cup \mathbf{n}'$; the A_i are called the *blocks* of α . By convention, \mathcal{P}_0 contains a single element, the empty partition, but we will assume $n \geq 1$ since all results concerning \mathcal{P}_0 are trivial.

A partition $\alpha \in \mathcal{P}_n$ may be pictured (non-uniquely) as a graph with vertex set $\mathbf{n} \cup \mathbf{n}'$, and with any edge set having the property that the connected components of the graph correspond to the blocks of the partition; the vertices of such a graph are always drawn with $1, \dots, n$ on an upper row, increasing from left to right, and vertices $1', \dots, n'$ directly below. For example, the partitions

$$\alpha = \{\{1, 4\}, \{2, 3, 4', 5'\}, \{5, 6\}, \{1', 2', 6'\}, \{3'\}\} \quad \text{and} \quad \beta = \{\{1, 2\}, \{3, 4, 1'\}, \{5, 4', 5', 6'\}, \{6\}, \{2'\}, \{3'\}\}$$

from \mathcal{P}_6 are pictured in Figure 1. As usual, we will generally identify a partition with any graph representing it.

A block A of a partition α is referred to as a *transversal* if $A \cap \mathbf{n} \neq \emptyset$ and $A \cap \mathbf{n}' \neq \emptyset$, or a *non-transversal* otherwise. For example, $\alpha \in \mathcal{P}_6$ defined above has $\{2, 3, 4', 5'\}$ as its only transversal, and has upper non-transversals $\{1, 4\}$ and $\{5, 6\}$, and lower non-transversals $\{1', 2', 6'\}$ and $\{3'\}$.

The *domain* and *codomain* of $\alpha \in \mathcal{P}_n$ are the subsets of \mathbf{n} defined by

$$\begin{aligned} \text{dom}(\alpha) &= \{i \in \mathbf{n} : i \text{ belongs to a transversal of } \alpha\}, \\ \text{codom}(\alpha) &= \{i \in \mathbf{n} : i' \text{ belongs to a transversal of } \alpha\}. \end{aligned}$$

The *rank* of $\alpha \in \mathcal{P}_n$, denoted $\text{rank}(\alpha)$, is defined to be the number of transversals of α . For example, with $\alpha \in \mathcal{P}_6$ as defined above, $\text{rank}(\alpha) = 1$, $\text{dom}(\alpha) = \{2, 3\}$ and $\text{codom}(\alpha) = \{4, 5\}$.

The product of two partitions $\alpha, \beta \in \mathcal{P}_n$ is defined as follows. Write $\mathbf{n}'' = \{1'', \dots, n''\}$. Let α^\vee be the graph obtained from α by changing the label of each lower vertex i' to i'' , and let β^\wedge be the graph obtained from β by changing the label of each upper vertex i to i'' . Consider now the graph $\Pi(\alpha, \beta)$ on the vertex set $\mathbf{n} \cup \mathbf{n}' \cup \mathbf{n}''$ obtained by joining α^\vee and β^\wedge together so that each lower vertex i'' of α^\vee is identified with the corresponding upper vertex i'' of β^\wedge . Note that $\Pi(\alpha, \beta)$, which we call the *product graph*, may contain pairs of parallel edges. We define $\alpha\beta \in \mathcal{P}_n$ to be the partition satisfying the property that $x, y \in \mathbf{n} \cup \mathbf{n}'$ belong to the same block of $\alpha\beta$ if and only if x and y are connected by a path in $\Pi(\alpha, \beta)$. This process is illustrated in Figure 1, with $\alpha, \beta \in \mathcal{P}_6$ defined above. The operation is associative, so \mathcal{P}_n is a semigroup: in fact, a monoid, with identity element $\{\{1, 1'\}, \dots, \{n, n'\}\}$.




Figure 1: Two partitions $\alpha, \beta \in \mathcal{P}_6$ (left), their product $\alpha\beta \in \mathcal{P}_6$ (right), and the product graph $\Pi(\alpha, \beta)$ (centre).

Note that the product graph $\Pi(\alpha, \beta)$ may contain connected components that only involve vertices from \mathbf{n}'' ; these are called *floating components* of $\Pi(\alpha, \beta)$. These play a crucial role in the definition of the *partition algebras*, and also the *twisted partition monoids*, which we now describe. Specifically, if $\tau(\alpha, \beta)$ denotes the number of floating components in the product graph $\Pi(\alpha, \beta)$, then one easily checks that

$$\tau(\alpha, \beta) + \tau(\alpha\beta, \gamma) = \tau(\alpha, \beta\gamma) + \tau(\beta, \gamma) \quad \text{for all } \alpha, \beta, \gamma \in \mathcal{P}_n.$$

It then follows that the product \star defined on the set $\mathcal{P}_n^\tau = \mathbb{N} \times \mathcal{P}_n = \{(i, \alpha) : i \in \mathbb{N}, \alpha \in \mathcal{P}_n\}$ by

$$(i, \alpha) \star (j, \beta) = (i + j + \tau(\alpha, \beta), \alpha\beta) \quad (3.1)$$

is associative. (Here, $\mathbb{N} = \{0, 1, 2, \dots\}$ denotes the set of natural numbers.) We call \mathcal{P}_n^τ with this product the *twisted partition monoid*.

A partition $\alpha \in \mathcal{P}_n$ is *planar* if there is a graphical representation of α in which:

- (i) all the edges are drawn within the rectangle determined by the vertices; and
- (ii) there are no crossings within the interior of the rectangle.

For example, of the two partitions $\alpha, \beta \in \mathcal{P}_6$ defined above, α is not planar, but β is. Since the product of two planar partitions is clearly planar, it follows that the set of all such planar partitions forms a submonoid of \mathcal{P}_n , and we denote this planar submonoid by \mathcal{PP}_n .

The *partial Brauer monoid* and the *Brauer monoid* are the submonoids of \mathcal{P}_n defined by

$$\mathcal{PB}_n = \{\alpha \in \mathcal{P}_n : \text{all blocks of } \alpha \text{ have size 1 or 2}\} \quad \text{and} \quad \mathcal{B}_n = \{\alpha \in \mathcal{P}_n : \text{all blocks of } \alpha \text{ have size 2}\}.$$

The *Motzkin monoid* and the *Jones monoid* are the planar submonoids of \mathcal{P}_n defined by

$$\mathcal{M}_n = \mathcal{PB}_n \cap \mathcal{PP}_n \quad \text{and} \quad \mathcal{J}_n = \mathcal{B}_n \cap \mathcal{PP}_n.$$

A Motzkin element $\gamma \in \mathcal{M}_{20}$ is pictured in Figure 2; the reader is invited to verify that γ is in fact an idempotent. It is well known that \mathcal{PP}_n is isomorphic to \mathcal{J}_{2n} [18].

The *twisted* versions of all the above monoids, \mathcal{PP}_n^τ , \mathcal{PB}_n^τ , \mathcal{B}_n^τ , \mathcal{M}_n^τ and \mathcal{J}_n^τ , are the corresponding submonoids of \mathcal{P}_n^τ : thus, for example, the twisted Brauer monoid \mathcal{B}_n^τ has underlying set $\mathbb{N} \times \mathcal{B}_n$ and product \star given by (3.1). In particular, the twisted Jones monoid \mathcal{J}_n^τ is known in the literature as the *Kauffman monoid* and is denoted \mathcal{K}_n [4, 25]. Despite the above-mentioned isomorphism of \mathcal{PP}_n and \mathcal{J}_{2n} , there is no such isomorphism between the twisted monoids \mathcal{PP}_n^τ and $\mathcal{J}_{2n}^\tau = \mathcal{K}_{2n}$. Indeed, \mathcal{PP}_n^τ and \mathcal{K}_{2n} do not have the same number of idempotents; see Tables 2 and 7 in Section 5.

The idempotents of the monoids $\mathcal{P}_n, \mathcal{B}_n, \mathcal{PB}_n$ (and their associated algebras and twisted versions) were classified and enumerated in [9], and the purpose of the current article is to undertake the same program for their planar counterparts. We conclude this subsection with a simple lemma.

Lemma 3.2. *For $\alpha \in \mathcal{M}_n$, the following are equivalent:*

- (i) $\alpha = \alpha^2$,
- (ii) $\text{dom}(\alpha) = \text{dom}(\alpha^2)$,
- (iii) $\text{codom}(\alpha) = \text{codom}(\alpha^2)$,
- (iv) $\text{rank}(\alpha) = \text{rank}(\alpha^2)$.

Proof. (i) \Rightarrow (ii) and (i) \Rightarrow (iii). These are obvious.

(ii) \Rightarrow (iv) and (iii) \Rightarrow (iv). If (ii) holds, then $\text{rank}(\alpha) = |\text{dom}(\alpha)| = |\text{dom}(\alpha^2)| = \text{rank}(\alpha^2)$, so that (iv) holds. The other implication is dual.

(iv) \Rightarrow (i). Suppose $\text{rank}(\alpha) = \text{rank}(\alpha^2)$. Every non-transversal of α is trivially a block of α^2 , so it remains to show that every transversal of α is a block of α^2 . Let the transversals of α be $\{i_1, j'_1\}, \dots, \{i_r, j'_r\}$ where $r = \text{rank}(\alpha)$ and $i_1 < \dots < i_r$, noting that this implies $\text{dom}(\alpha) = \{i_1, \dots, i_r\}$, $\text{codom}(\alpha) = \{j_1, \dots, j_r\}$ and $j_1 < \dots < j_r$ (the latter by planarity). Since $\text{dom}(\alpha^2) \subseteq \text{dom}(\alpha)$ and $\text{codom}(\alpha^2) \subseteq \text{codom}(\alpha)$, every transversal of α^2 is of the form $\{i_s, j'_t\}$ for some $s, t \in \{1, \dots, r\}$. Since $\text{rank}(\alpha^2) = \text{rank}(\alpha) = r$, there must be r such transversals of α^2 , and so these must be $\{i_1, j'_{1\pi}\}, \dots, \{i_r, j'_{r\pi}\}$ for some permutation π of $\{1, \dots, r\}$. Since α^2 is planar, this permutation must be the identity, so it follows that α^2 contains the transversals $\{i_1, j'_1\}, \dots, \{i_r, j'_r\}$, as required. \square

Remark 3.3. We have not referred to Green's relations or the regular \ast -semigroup structure on \mathcal{M}_n , or any of the other diagram monoids we study, since neither plays a role in the theory developed in this section (though they do in the algorithms presented in Section 4). Green's relations on $\mathcal{M}_n, \mathcal{J}_n$ and \mathcal{K}_n are characterised in [10, Theorem 2.4], [40, Theorem 18] and [25, Theorem 5.1], respectively, in terms of domains, ranks, and other parameters. We will not need to know the exact formulations of these results, so we will not state them here, but it is worth noting that two elements of \mathcal{M}_n are \mathcal{D} -related if and only if they have the same rank, and that \mathcal{M}_n is \mathcal{H} -trivial. These two facts lead to a simpler proof of Lemma 3.2, since for an element x of a finite semigroup, $x \mathcal{D} x^2 \Leftrightarrow x \mathcal{R} x^2 \Leftrightarrow x \mathcal{L} x^2 \Leftrightarrow x \mathcal{H} x^2$ (see [36, Theorems A.2.4 and A.3.4], for example); in particular, if the finite semigroup is \mathcal{H} -trivial, then these are also equivalent to $x = x^2$. The anti-involution $\ast : \mathcal{P}_n \rightarrow \mathcal{P}_n$ that gives \mathcal{P}_n , and hence all the submonoids considered in this article, a regular \ast -semigroup structure corresponds to reflecting (diagrams representing) elements of \mathcal{P}_n in a horizontal axis midway between the two rows of vertices.

3.2 Interface graphs and characterisation of idempotents

A key role in our study is played by the so-called *interface graph* of a Motzkin element. In this subsection, we define these graphs, and show how they may be used to characterise the idempotents of $\mathcal{M}_n, \mathcal{J}_n$ and \mathcal{K}_n .


A block A of a Motzkin element $\alpha \in \mathcal{M}_n$ is called an *upper hook* if $A \subseteq \mathbf{n}$ and $|A| = 2$, or an *upper singleton* if $A \subseteq \mathbf{n}$ and $|A| = 1$. *Lower hooks* and *lower singletons* are defined analogously.

Let $\alpha \in \mathcal{M}_n$. The *interface graph* Γ_α is a vertex- and edge-coloured graph defined as follows. The vertex set of Γ_α is simply \mathbf{n} , and the colour $c(v) \in \mathbb{Z}_2 \times \mathbb{Z}_2$ of a vertex $v \in \mathbf{n}$ is defined to be the column vector

$$c(v) = \begin{bmatrix} a \\ b \end{bmatrix} \quad \text{where} \quad a = \begin{cases} 1 & \text{if } v \in \text{codom}(\alpha) \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad b = \begin{cases} 1 & \text{if } v \in \text{dom}(\alpha) \\ 0 & \text{otherwise} \end{cases}$$

For each upper hook $\{i, j\}$ of α , Γ_α has an edge $\{i, j\}$ coloured -1 . For each lower hook $\{k', l'\}$ of α , Γ_α has an edge $\{k, l\}$ coloured $+1$. (Note that Γ_α may have two edges between a pair of vertices, but these edges will always have opposite colours.)

When drawing the interface graph Γ_α of a Motzkin element $\alpha \in \mathcal{M}_n$, we always draw the vertices in a horizontal row, in the order $1, \dots, n$, increasing from left to right. We draw the edges coloured $+1$ or -1 above or below the line of vertices, respectively, as line segments:



with the “height” of such line segments chosen so that the diagram is planar (we can do this since $\alpha \in \mathcal{M}_n$ is itself planar). And we indicate the colour $c(v) = \begin{bmatrix} a \\ b \end{bmatrix}$ of a vertex $v \in \mathbf{n}$ by drawing a small line segment above and/or below vertex v if $a = 1$ and/or $b = 1$, respectively. Roughly speaking, this picture of Γ_α is obtained by cutting (a diagram representing) α in half, horizontally along the middle of the diagram, and then connecting the top half to the bottom half by identifying the two rows of vertices. Figure 2 pictures a Motzkin element $\gamma \in \mathcal{M}_{20}$ and its interface graph Γ_γ .




Figure 2: A Motzkin element $\gamma \in \mathcal{M}_{20}$ (above) and its interface graph Γ_γ (below).

If $\alpha \in \mathcal{M}_n$, then every vertex in Γ_α has degree at most 2. Hence, every connected component of Γ_α is either a cycle or a path; we regard a singleton component of Γ_α as a path of length 0. A vertex cannot be the endpoint of two edges with the same colour, because blocks of α have size at most 2, so it follows that the edges along any path in Γ_α alternate in colour; in particular, all cycles have even length. It is also apparent that a vertex of degree 2 can only be coloured $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$. Now consider a path component $v_1 - v_2 - \dots - v_k$ of Γ_α . As above, $c(v_2) = \dots = c(v_{k-1}) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$. We call the path *inactive* if also $c(v_1) = c(v_k) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$. We call the path *active* if either $k = 1$ and $c(v_1) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ or else $k \geq 2$ and $c(v_1), c(v_k) \in \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Otherwise, we say the path is *mixed*.

The next result characterises the idempotents of \mathcal{M}_n in terms of interface graphs. Of crucial importance is the fact that the product graph $\Pi(\alpha, \alpha)$ contains an isomorphic copy of Γ_α in the middle layer. If C is a connected component of a graph Γ , so that C is itself a graph, we will slightly abuse notation and identify C with its set of vertices, so we sometimes write “ $u \in C$ ” to mean “ u is a vertex of C ”.

Proposition 3.4. *A Motzkin element $\alpha \in \mathcal{M}_n$ is an idempotent if and only if every connected component of the interface graph Γ_α is one of:*

- (i) a cycle,
- (ii) an inactive path, or
- (iii) an active path of even length.

Proof. (\Leftarrow). Suppose first that all components of Γ_α are of types (i)–(iii). By Lemma 3.2, and since clearly $\text{rank}(\alpha^2) \leq \text{rank}(\alpha)$, to show that $\alpha^2 = \alpha$, it is enough to show that $\text{rank}(\alpha^2) \geq \text{rank}(\alpha)$.

Suppose the components of Γ_α of type (iii) are C_1, \dots, C_r . For each i , let the endpoints of C_i be u_i and v_i . Since C_i is of even length, we may assume that the bottom coordinate of $c(u_i)$ and the top coordinate of $c(v_i)$ are both 1, even if $u_i = v_i$.

As noted above, the colour of any vertex from a component of types (i) or (ii) is $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$; this is also the case for any interior vertices of the components of type (iii). It follows that $\text{dom}(\alpha) = \{u_1, \dots, u_r\}$ and $\text{codom}(\alpha) = \{v_1, \dots, v_r\}$, so that $\text{rank}(\alpha) = r$. It also follows that there is a permutation π of $\{1, \dots, r\}$ such that $\{u_1, v'_{\pi(1)}\}, \dots, \{u_r, v'_{\pi(r)}\}$ are the transversals of α .

Now fix some $1 \leq i \leq r$. Since C_i is a path from u_i to v_i , it follows that the product graph $\Pi(\alpha, \alpha)$ has a path from u''_i to v''_i . Since $\{u_{\pi^{-1}(i)}, v'_i\}$ and $\{u_i, v'_{\pi(i)}\}$ are transversals of α , $\Pi(\alpha, \alpha)$ contains the edges $\{u_{\pi^{-1}(i)}, v''_i\}$ and $\{u''_i, v'_{\pi(i)}\}$. So $u_{\pi^{-1}(i)}$ and $v'_{\pi(i)}$ are connected by a path in $\Pi(\alpha, \alpha)$, and it follows that $\{u_{\pi^{-1}(i)}, v'_{\pi(i)}\}$ is a transversal of α^2 . Thus, $\text{dom}(\alpha^2) \supseteq \{u_{\pi^{-1}(1)}, \dots, u_{\pi^{-1}(r)}\} = \{u_1, \dots, u_r\}$, and it follows that $\text{rank}(\alpha^2) \geq r = \text{rank}(\alpha)$, as required.

(\Rightarrow). For this implication, we prove the contrapositive. Suppose Γ_α contains a component not of types (i)–(iii). Then this component must be either

(iv) an active path of odd length, or

(v) a mixed path.

Suppose first that C is a component of Γ_α of type (iv), and let u, v be the endpoints of C . Since C is of odd length, it follows that $u \neq v$, and that $c(u) = c(v) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ or $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$. We suppose the latter is the case (the proof for the former is similar). So $u, v \in \text{dom}(\alpha)$ belong to (distinct) transversals of α : $\{u, x'\}$ and $\{v, y'\}$, say. So $\{u'', x'\}$ and $\{v'', y'\}$ are edges in the product graph $\Pi(\alpha, \alpha)$. Since C gives rise to a path from u'' to v'' in $\Pi(\alpha, \alpha)$, it follows that $\{x', y'\}$ is a block of α^2 . But $\{x', y'\}$ is not a block of α , since $x, y \in \text{codom}(\alpha)$, so it follows that $\alpha^2 \neq \alpha$.

On the other hand, suppose C is a component of Γ_α of type (v), and let u, v be the endpoints of C . Suppose $c(v) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, so that $c(u) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ or $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Again, we just consider the latter case. So u belongs to a transversal $\{u, x'\}$ of α . Since $c(v) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, it follows that the connected component in the product graph $\Pi(\alpha, \alpha)$ containing x' is $\{x'\} \cup \{y'' : y \in C\}$. We deduce that $\{x'\}$ is a block of α^2 , and it again follows that $\alpha^2 \neq \alpha$, since $x \in \text{codom}(\alpha)$. This completes the proof. \square

The interface graph Γ_α of a Jones element $\alpha \in \mathcal{J}_n$ can only have cycles and active paths, since all blocks of α are of size 2. So we may immediately deduce from Proposition 3.4 the following characterisation of Jones idempotents.

Corollary 3.5. *A Jones element $\alpha \in \mathcal{J}_n$ is an idempotent if and only if every connected component of Γ_α is a cycle or an active path of even length.* \square

The characterisation of idempotents in the twisted Motzkin and Jones monoids, \mathcal{M}_n^τ and $\mathcal{J}_n^\tau = \mathcal{K}_n$, is as follows.

Corollary 3.6. *A twisted Motzkin element $(i, \alpha) \in \mathcal{M}_n^\tau$ is an idempotent if and only if $i = 0$ and every connected component of Γ_α is an active path of even length. Consequently, $E(\mathcal{M}_n^\tau) = E(\mathcal{K}_n)$.*

Proof. Note that $(i, \alpha) \star (i, \alpha) = (2i + \tau(\alpha, \alpha), \alpha^2)$, so $(i, \alpha) \in E(\mathcal{M}_n^\tau)$ if and only if $i = 0$, $\tau(\alpha, \alpha) = 0$ and $\alpha \in E(\mathcal{M}_n)$. Cycles and inactive paths in the interface graph Γ_α correspond to floating components in the product graph $\Pi(\alpha, \alpha)$, so it follows from Proposition 3.4 that $[\tau(\alpha, \alpha) = 0 \text{ and } \alpha \in E(\mathcal{M}_n)]$ is equivalent to Γ_α having only active paths of even length. The last assertion of the lemma follows quickly. \square

Remark 3.7. Corollary 3.6 also applies to the *Temperley-Lieb* and *Motzkin algebras*. If S is any of the monoids $\mathcal{P}_n, \mathcal{PB}_n, \mathcal{B}_n, \mathcal{PP}_n, \mathcal{M}_n$ or \mathcal{J}_n , and if \mathbb{F} is a field with some fixed element $\xi \in \mathbb{F}$, then we may form the *twisted semigroup algebra* $\mathbb{F}^\xi[S]$, as in [40]; these are the partition [23, 28], partial Brauer [29], Brauer [5], planar partition [23], Motzkin [3] and Temperley-Lieb [39] algebras, respectively. The algebra $\mathbb{F}^\xi[S]$ has basis S , and multiplication \circ defined on basis elements $\alpha, \beta \in S$ (and extended linearly) by $\alpha \circ \beta = \xi^{\tau(\alpha, \beta)}(\alpha\beta)$. If ξ is not a root of unity or if it is an M th root of unity where $M > n$ (the so-called *generic case*), then an element $\alpha \in S$ satisfies $\alpha = \alpha \circ \alpha$ if and only if $\tau(\alpha, \alpha) = 0$ and $\alpha = \alpha^2$ in S ; thus, in this case, Corollary 3.6 shows that an element $\alpha \in \mathcal{M}_n$ is an idempotent basis element of the Motzkin algebra if and only if $i = 0$ and every connected component of Γ_α is an active path of even length. As in [9, Section 6], if ξ is an M th root of unity with $M \leq n$, then a Motzkin element $\alpha \in \mathcal{M}_n$ is an idempotent basis element of $\mathbb{F}^\xi[\mathcal{M}_n]$ if and only every connected component of Γ_α is one of types (i)–(iii) as listed in Proposition 3.4, with the combined number of components of types (i)–(ii) being a multiple of M . On the other hand, if α is an arbitrary idempotent of S (where S is any of the above diagram monoids), and if $\xi \neq 0$, then $\xi^{-m(\alpha, \alpha)}\alpha$ is an idempotent of $\mathbb{F}^\xi[S]$; we thank Zajj Daugherty for this last observation.

3.3 A mapping on $E(\mathcal{M}_n)$ and an enumeration method

Now that we have characterised the idempotents of $\mathcal{M}_n, \mathcal{J}_n$ and \mathcal{K}_n , we wish to enumerate them. In this subsection, we describe a method for doing so. We make crucial use of a map $D : E(\mathcal{M}_n) \rightarrow E(\mathcal{M}_n)$ to be defined shortly, and the interface graphs defined in Subsection 3.2. This map also played a crucial role in the classification of *congruences* on \mathcal{J}_n and \mathcal{B}_n in [15]; a different, but closely-related, map was used for $\mathcal{M}_n, \mathcal{P}_n, \mathcal{PB}_n$ and \mathcal{PP}_n .

Lemma 3.8. *Suppose $\alpha \in E(\mathcal{M}_n)$ and $\{i, j'\}$ is a transversal of α . Then Γ_α contains an active path of even length with i and j as its endpoints. In particular, if $\text{dom}(\alpha) = \{i_1, \dots, i_r\}$ where $r = \text{rank}(\alpha)$, then i_1, \dots, i_r belong to r distinct connected components of Γ_α .*

Proof. Since $\alpha = \alpha^2$, $\{i, j'\}$ is a transversal of α^2 , so there is a path from i to j' in the product graph $\Pi(\alpha, \alpha)$. Since $\{i, j''\}$ and $\{i'', j'\}$ are both edges of $\Pi(\alpha, \alpha)$, it follows that there is a path from j'' to i'' in $\Pi(\alpha, \alpha)$ involving only vertices in the middle row; this gives rise to a path from i to j in Γ_α . Since $i \in \text{dom}(\alpha)$ and $j \in \text{codom}(\alpha)$, it follows that $c(i) \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ and $c(j) \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, so that i and j are indeed the endpoints of this path. Since this path is active, Proposition 3.4 tells us that it is of even length. This proves the first assertion of the lemma.

For the second assertion, suppose the transversals of α are $\{i_1, j'_1\}, \dots, \{i_r, j'_r\}$, and let $1 \leq k < l \leq r$. We must show that i_k and i_l belong to different connected components of Γ_α . To do so, suppose to the contrary that i_k and i_l belong to the same component. By the previous paragraph, the component of Γ_α containing i_k is a path from i_k to j_k , and the component containing i_l is a path from i_l to j_l . Thus, since we have assumed these are the same components, and since $i_k \neq i_l$, we must have $i_k = j_l$ and $i_l = j_k$. But then $\{i_k, i'_l\}$ and $\{i_l, i'_k\}$ are both transversals of α , contradicting planarity. \square

To define the mapping $D : E(\mathcal{M}_n) \rightarrow E(\mathcal{M}_n)$, we first define a map $d : E(\mathcal{M}_n) \rightarrow \mathcal{M}_n$. Before we do this, note first that the set of transversals of a Motzkin element $\alpha \in \mathcal{M}_n$ inherits an obvious total ordering from the natural ordering on \mathbf{n} . For example, the transversals of the Motzkin element $\gamma \in \mathcal{M}_{20}$ pictured in Figure 2 are ordered by $\{1, 3'\} < \{8, 8'\} < \{20, 17'\}$. So we may speak of the first and second transversals of α , and so on.

Now let $\alpha \in E(\mathcal{M}_n)$. If $\text{rank}(\alpha) \leq 1$, then we define $d(\alpha) = \alpha$. Otherwise, if $\{i, j'\}$ and $\{k, l'\}$ are the first two transversals of α , then we define $d(\alpha)$ to be the element of \mathcal{M}_n obtained from α by replacing these two transversals by the upper and lower hooks $\{i, k\}$ and $\{j', l'\}$. Note that blocks of $d(\alpha)$ trivially have size ≤ 2 , while the planarity of α ensures that of $d(\alpha)$; see Figure 3. Although it is not readily apparent that $d(\alpha)$ is necessarily an idempotent, we will soon see that it is.




Figure 3: The map $d : E(\mathcal{M}_n) \rightarrow E(\mathcal{M}_n)$. Shaded regions of α are assumed to be identical to the corresponding shaded regions of $d(\alpha)$.

The next result gathers some important properties of the d map, including the fact that d does indeed map into $E(\mathcal{M}_n)$. Let A be an upper non-transversal of a Motzkin element $\alpha \in \mathcal{M}_n$. We say that A is to the left of a transversal $\{i, j'\}$ of α if $\max(A) < i$. Similarly, we say a lower non-transversal B' is to the left of $\{i, j'\}$ if $\max(B') < j$. We say that A is nested (in α) if there exists an upper hook $\{k, l\}$ of α such that $k < \min(A)$ and $\max(A) < l$; otherwise, we say that A is unnnested. We define nested and unnnested lower non-transversals analogously. An outer hook of α is defined to be an upper or lower unnnested hook of α that is to the left of any transversal of α . For the statements of parts (ii) and (iii) of the next lemma, recall that we informally identify a connected component of a graph with its underlying vertex set.

Lemma 3.9. *Let $\alpha \in E(\mathcal{M}_n)$ with $\text{rank}(\alpha) \geq 2$, and suppose $\{i, j'\}$ and $\{k, l'\}$ are the first two transversals of α . Suppose $C_1, C_2, C_3, \dots, C_s$ are the connected components of Γ_α , where $i \in C_1$ and $k \in C_2$. Then*

- (i) $\{i, k\}$ and $\{j', l'\}$ are outer hooks of $d(\alpha)$,
- (ii) $C_1 \cup C_2, C_3, \dots, C_s$ are the connected components of $\Gamma_{d(\alpha)}$,
- (iii) i, j, k, l belong to $C_1 \cup C_2$, and this is a cycle component of $\Gamma_{d(\alpha)}$, and
- (iv) $d(\alpha) \in E(\mathcal{M}_n)$.

Proof. (i). Planarity of α , and the fact that there are no other transversals between $\{i, j'\}$ and $\{k, l'\}$ ensures that $\{i, k\}$ and $\{j', l'\}$ are unnested in $d(\alpha)$; cf. Figure 3. The fact that $\{i, j'\}$ and $\{k, l'\}$ are the *first* two transversals of α ensures that $\{i, k\}$ and $\{j', l'\}$ are to the left of any transversal of $d(\alpha)$.

(ii) and (iii). By Lemma 3.8, C_1 is a path in Γ_α from i to j , and C_2 a path from k to l . The only change from Γ_α to $\Gamma_{d(\alpha)}$ is the addition of the edges $\{i, k\}$ and $\{j, l\}$, coloured -1 and $+1$, respectively, and the recolouring of the vertices i, j, k, l (four 1's are changed to 0's, regardless of whether these are four distinct vertices). Since the addition of these edges joins C_1 and C_2 into a single cycle component of $\Gamma_{d(\alpha)}$, (iii) follows. Since no other components of Γ_α are modified, (ii) also follows.

(iv). Since $\alpha \in E(\mathcal{M}_n)$, the components C_3, \dots, C_s are all of the forms specified in Proposition 3.4. Since $C_1 \cup C_2$ is a cycle, (iv) follows. \square

For any $\alpha \in E(\mathcal{M}_n)$, the sequence $\alpha, d(\alpha), d^2(\alpha), \dots$ eventually terminates in an idempotent of rank 0 or 1, depending on the parity of $\text{rank}(\alpha)$, and we write $D(\alpha)$ for this idempotent. In fact, $D(\alpha) = d^s(\alpha)$, where $s = \lfloor \text{rank}(\alpha)/2 \rfloor$; we consider d^0 to be the identity mapping. For example, consider the Motzkin element $\gamma \in \mathcal{M}_{20}$ pictured in Figure 2. Here we have $\text{rank}(\gamma) = 3$, so that $D(\gamma) = d(\gamma)$; we have pictured $\delta = d(\gamma)$ and Γ_δ in Figure 4.




Figure 4: The Motzkin element $\delta = d(\gamma) \in \mathcal{M}_{20}$ (above) and its interface graph Γ_δ (below), where $\gamma \in \mathcal{M}_{20}$ is pictured in Figure 2. New edges are drawn in red.

From now on, we will write

$$\Delta(\mathcal{M}_n) = D(E(\mathcal{M}_n)) = \{D(\alpha) : \alpha \in E(\mathcal{M}_n)\} \quad \text{and} \quad \Delta(\mathcal{J}_n) = D(E(\mathcal{J}_n)) = \{D(\alpha) : \alpha \in E(\mathcal{J}_n)\}.$$

Since D acts as the identity on idempotents of rank ≤ 1 , and since all Jones elements of minimal rank are idempotents, it follows that

$$\Delta(\mathcal{M}_n) = \{\alpha \in E(\mathcal{M}_n) : \text{rank}(\alpha) \leq 1\} \quad \text{and} \quad \Delta(\mathcal{J}_n) = \{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) \leq 1\}.$$

Note that there are elements in \mathcal{J}_n of rank 0 or 1, depending on the parity of n , but not both.

The next result shows that enumeration of $E(\mathcal{M}_n)$ and $E(\mathcal{J}_n)$ reduces to the enumeration of preimages under the D map. This latter task is itself quite difficult; Section 4 and the remainder of Section 3 are devoted to achieving it.

Lemma 3.10. *If S is one of \mathcal{M}_n or \mathcal{J}_n , then $|E(S)| = \sum_{\alpha \in \Delta(S)} |D^{-1}(\alpha)|$.*

Proof. The statement about $|E(\mathcal{M}_n)|$ is clear. If $\alpha \in E(\mathcal{M}_n)$, then Corollary 3.5 and Lemma 3.9 imply that $\alpha \in \mathcal{J}_n$ if and only if $D(\alpha) \in \mathcal{J}_n$. The statement about $|E(\mathcal{J}_n)|$ follows. \square

For an arbitrary Motzkin element $\alpha \in \mathcal{M}_n$, we write $\Theta(\alpha)$ for the set of all cycle components of the interface graph Γ_α containing at least one edge corresponding to an upper outer hook of α and at least one edge corresponding to a lower outer hook of α ; outer hooks were defined before Lemma 3.9. For $\theta \in \Theta(\alpha)$, let

$$U_\theta(\alpha) = \{\{i, j\} : \{i, j\} \text{ is an upper outer hook of } \alpha \text{ and } i, j \in \theta\},$$

$$L_\theta(\alpha) = \{\{i', j'\} : \{i', j'\} \text{ is a lower outer hook of } \alpha \text{ and } i, j \in \theta\},$$

and write $u_\theta(\alpha) = |U_\theta(\alpha)|$ and $l_\theta(\alpha) = |L_\theta(\alpha)|$. For example, we have $\Theta(\gamma) = \emptyset$ for $\gamma \in \mathcal{M}_{20}$ from Figure 2 (note that γ has only one outer hook). However, with $\delta = d(\gamma) \in \mathcal{M}_{20}$ from Figure 4, we have $\Theta(\delta) = \{\theta_1, \theta_2\}$, where $\theta_1 = \{1, 2, 3, 4, 7, 8\}$ and $\theta_2 = \{9, 12, 15, 16\}$, and

$$U_{\theta_1}(\delta) = \{\{1, 8\}\}, \quad L_{\theta_1}(\delta) = \{\{1', 2'\}, \{3', 8'\}\}, \quad U_{\theta_2}(\delta) = \{\{9, 12\}\}, \quad L_{\theta_2}(\delta) = \{\{9', 16'\}\}.$$

So $l_{\theta_1}(\delta) = 2$, while $u_{\theta_1}(\delta) = u_{\theta_2}(\delta) = l_{\theta_2}(\delta) = 1$. The next result shows why the sets we have just defined are important.

Lemma 3.11. *Suppose $\alpha \in E(\mathcal{M}_n)$ is such that $\Theta(\alpha) \neq \emptyset$. Let $\theta \in \Theta(\alpha)$, and let $\{i, k\} \in U_\theta(\alpha)$ and $\{j', l'\} \in L_\theta(\alpha)$. Let β be obtained from α by replacing the blocks $\{i, k\}$ and $\{j', l'\}$ by $\{i, j'\}$ and $\{k, l'\}$. Then $\beta \in E(\mathcal{M}_n)$ and $d(\beta) = \alpha$. Further, every element of $d^{-1}(\alpha) \setminus \{\alpha\}$ may be constructed in this way, for some $\theta \in \Theta(\alpha)$ and some pair of edges from $U_\theta(\alpha) \times L_\theta(\alpha)$.*

Proof. Since $\{i, k\}$ and $\{j', l'\}$ are outer hooks of α , it follows that $\beta \in \mathcal{M}_n$. To show that β is an idempotent, we need to check that each component of Γ_β is of one of the forms specified in Proposition 3.4. By construction, the components of Γ_α other than θ are still components of Γ_β , and these must all be of the specified form, since $\alpha \in E(\mathcal{M}_n)$. But θ , a cycle component of Γ_α , is split into two active path components of Γ_β . To complete the proof that β is an idempotent, it remains to show that these paths are of even length. But this follows quickly from the fact that we are removing an edge coloured $+1$ and an edge coloured -1 from a cycle whose edge colours alternate between $+1$ and -1 .

Since $\{i, k\}$ and $\{j', l'\}$ are to the left of any transversals of α , as they are outer hooks of α , it follows that $\{i, j'\}$ and $\{k, l'\}$ are the first two transversals of β , and then it follows immediately that $d(\beta) = \alpha$.

Finally, suppose $\gamma \in E(\mathcal{M}_n)$ is such that $d(\gamma) = \alpha$, and let the first two transversals of γ be $\{u, v'\}$ and $\{x, y'\}$, respectively. By Lemma 3.9, $\{u, x\}$ and $\{v', y'\}$ are outer hooks of $d(\gamma) = \alpha$, and u, v, x, y belong to the same cycle component of $\Gamma_{d(\gamma)} = \Gamma_\alpha$. If we denote this cycle component by σ , then $\{u, x\} \in U_\sigma(\alpha)$ and $\{v', y'\} \in L_\sigma(\alpha)$, and we see that γ is constructed in the manner described in the lemma, with respect to σ , $\{u, x\}$ and $\{v', y'\}$. \square

Lemma 3.11 gives information about preimages under the d map. In order to extend this to preimages under the D map, we require the next two intermediate lemmas. The first, Lemma 3.12, concerns curves in the plane, and the second, Lemma 3.13, applies this to the situation in which the curves are part of the interface graph of a Motzkin element.

Lemma 3.12. *Let A, B, C, D be distinct points on the x -axis, with x -coordinates $a < b < c < d$, respectively. Suppose \mathcal{C}_1 and \mathcal{C}_2 are smooth non-self-intersecting curves in the plane such that*

- (i) \mathcal{C}_1 joins A to C , while \mathcal{C}_2 joins B to D ,
- (ii) apart from the endpoints stated above, both curves are contained in the region $a < x < d$, and
- (iii) \mathcal{C}_1 and \mathcal{C}_2 never go below the points B or C : that is, no point (x, y) on either curve satisfies $[x = b \text{ and } y < 0]$ or $[x = c \text{ and } y < 0]$.

Then \mathcal{C}_1 and \mathcal{C}_2 intersect.

Proof. Consider the curve \mathcal{C} obtained from \mathcal{C}_1 by adding the positive half \mathcal{L}_1 of the line $x = a$ and the negative half \mathcal{L}_2 of the line $x = c$, as shown in Figure 5. By the stated assumptions, \mathcal{C} has no self-intersections, and so divides the plane into two regions: one containing B and one containing D . But \mathcal{C}_2 joins B to D , so it follows that that \mathcal{C}_2 and \mathcal{C} intersect. Assumptions (ii) and (iii), respectively, tell us that \mathcal{C}_2 does not intersect \mathcal{L}_1 or \mathcal{L}_2 . So \mathcal{C}_2 must intersect \mathcal{C}_1 . \square

Recall that we are identifying a connected component C of a graph on vertex set \mathbf{n} with the underlying vertex set of C . In this way, we may also write $\min(C)$ to mean the vertex of C with minimum value in the natural ordering on \mathbf{n} .





Figure 5: The curve $\mathcal{C} = \mathcal{L}_1 \cup \mathcal{C}_1 \cup \mathcal{L}_2$ from the proof of Lemma 3.12.

Lemma 3.13. *Suppose $\alpha \in E(\mathcal{M}_n)$ and $\theta_1, \theta_2 \in \Theta(\alpha)$, where $\min(\theta_1) < \min(\theta_2)$.*

- (i) *If $\{i, j\} \in U_{\theta_1}(\alpha)$ and $\{k, l\} \in U_{\theta_2}(\alpha)$ with $i < j$ and $k < l$, then $j < k$.*
- (ii) *If $\{i', j'\} \in L_{\theta_1}(\alpha)$ and $\{k', l'\} \in L_{\theta_2}(\alpha)$ with $i < j$ and $k < l$, then $j < k$.*

Proof. We just prove (i), as (ii) is dual. Suppose to the contrary that $j > k$. Since $\{i, j\}$ and $\{k, l\}$ are unnested blocks of α , and since α is planar, it follows that $k < l < i < j$. In what follows, we consider the cycles θ_1 and θ_2 as (closed, non-self-intersecting) curves in the plane, with each vertex $v \in \mathbf{n}$ drawn at the point $(v, 0)$, and with edges drawn in the usual way to join vertices as:



We first claim that $\max(\theta_2) < \max(\theta_1)$. Indeed, suppose to the contrary that $\max(\theta_2) > \max(\theta_1)$. Put $a = \min(\theta_1)$, $b = l$, $c = i$ and $d = \max(\theta_2)$, and let $A = (a, 0)$, $B = (b, 0)$, and so on. Note that θ_1 is the union of two paths joining A and C ; let \mathcal{C}_1 be either of these paths. Similarly, θ_2 is the union of two paths joining B and D ; let \mathcal{C}_2 be either of these paths. It is easy to check that conditions (i)–(iii) of Lemma 3.12 are satisfied, using the fact that $\{i, j\}$ and $\{k, l\}$ are unnested to verify condition (iii). It follows that \mathcal{C}_1 and \mathcal{C}_2 , and hence θ_1 and θ_2 , intersect, a contradiction. This completes the proof of the claim that $\max(\theta_2) < \max(\theta_1)$. It follows that $\min(\theta_1) < \min(\theta_2) < \max(\theta_2) < \max(\theta_1)$.

Now, θ_1 is also the union of two paths joining $(\min(\theta_1), 0)$ and $(\max(\theta_1), 0)$; let \mathcal{C}_3 be either of these paths. So \mathcal{C}_3 is a non-self-intersecting curve in the plane and, apart from its endpoints, it lies in the region $\min(\theta_1) < x < \max(\theta_1)$. In particular, it divides the region $\min(\theta_1) < x < \max(\theta_1)$ into upper and lower regions. Since θ_1 and θ_2 do not intersect, θ_2 is contained wholly within one of these two regions. Since $\{k, l\}$ is unnested, it lies in the lower region and, hence, it follows that θ_2 is contained in this lower region. But then every edge of θ_2 lies under the curve $\mathcal{C}_3 \subseteq \theta_1$. It follows that every lower hook of α corresponding to an edge of θ_2 is nested in α , so that $L_{\theta_2}(\alpha) = \emptyset$, contradicting the assumption that $\theta_2 \in \Theta(\alpha)$. \square

Remark 3.14. As the last paragraph of the above proof indicates, the assumption that θ_2 has outer upper and lower hooks is necessary to prove the conclusion of Lemma 3.13(i). Indeed, consider the Jones idempotent $\alpha = \{\{1, 2\}, \{3, 4\}, \{5, 6\}, \{1', 6'\}, \{2', 5'\}, \{3', 4'\}\} \in E(\mathcal{J}_6)$. The interface graph Γ_α has two connected components: $\theta_1 = \{1, 2, 5, 6\}$ and $\theta_2 = \{3, 4\}$. Both are cycles, and $\min(\theta_1) < \min(\theta_2)$, yet $\{5, 6\}$ and $\{3, 4\}$ are upper outer hooks of θ_1 and θ_2 , respectively, and we do not have $6 < 3$. However, while θ_1 does belong to $\Theta(\alpha)$, θ_2 does not.

We are now ready to combine the preceding series of lemmas in order to to enumerate the preimages under the D map. Proposition 3.15 below (and its proof) shows that for any $\alpha \in \Delta(\mathcal{M}_n)$, the idempotents from $D^{-1}(\alpha)$ are obtained from α by selecting some collection $\theta_1, \dots, \theta_t$ of cycle components from Γ_α , each containing at least one upper outer hook and at least one lower outer hook, and then replacing $2t$ such hooks (one upper and one lower outer hook from each component) by suitable transversals. The number of idempotents in $D^{-1}(\alpha)$ corresponding to the collection $\theta_1, \dots, \theta_t$ is found by calculating the numbers of upper and lower outer hooks of these components and multiplying all $2t$ of these numbers together. The total

size of $D^{-1}(\alpha)$ is then the sum of all such products over all collections of cycle components; algebraically, this sum of products then may be simplified into a single product. For the statement of the next result, if $0 \leq r \leq n$, we will write $E_r(\mathcal{M}_n) = \{\alpha \in E(\mathcal{M}_n) : \text{rank}(\alpha) = r\}$.

Proposition 3.15. *Let $\alpha \in \Delta(\mathcal{M}_n)$, and write $q = \text{rank}(\alpha)$ and $k = |\Theta(\alpha)|$.*

- (i) *For any $\beta \in D^{-1}(\alpha)$, $\text{rank}(\beta) = q + 2t$ for some $0 \leq t \leq k$.*
- (ii) *For any $0 \leq t \leq k$, the set $D^{-1}(\alpha) \cap E_{q+2t}(\mathcal{M}_n)$ has cardinality $\sum_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi|=t}} \prod_{\theta \in \Psi} (u_\theta(\alpha)l_\theta(\alpha))$.*
- (iii) *We have $|D^{-1}(\alpha)| = \prod_{\theta \in \Theta(\alpha)} (u_\theta(\alpha)l_\theta(\alpha) + 1)$.*

Proof. (i). Suppose $\beta \in D^{-1}(\alpha)$. Let $t \geq 0$ be minimal so that $\alpha = d^t(\beta)$. In the sequence

$$\beta, d(\beta), \dots, d^t(\beta) = \alpha,$$

the rank of each term is 2 more than the rank of the next term, by definition of the d map. It follows that $\text{rank}(\beta) = \text{rank}(\alpha) + 2t = q + 2t$. We have already noted that $t \geq 0$. By Lemma 3.9,

$$\Theta(\beta) \subsetneq \Theta(d(\beta)) \subsetneq \dots \subsetneq \Theta(d^t(\beta)) = \Theta(\alpha).$$

Thus, $k = |\Theta(\alpha)| \geq |\Theta(\beta)| + t \geq t$.

(ii). Fix some $0 \leq t \leq k$, and write

$$\Sigma = D^{-1}(\alpha) \cap E_{q+2t}(\mathcal{M}_n) \quad \text{and} \quad \sigma = \sum_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi|=t}} \prod_{\theta \in \Psi} (u_\theta(\alpha)l_\theta(\alpha)).$$

If $t = 0$, then $\Sigma = \{\alpha\}$ and $\sigma = \prod_{\theta \in \Theta} (u_\theta(\alpha)l_\theta(\alpha)) = 1$, as the latter is an empty product. Now suppose $t \geq 1$. To complete the proof of (ii), it suffices to find mutually inverse maps

$$f : \Sigma \rightarrow \bigcup_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi|=t}} \prod_{\theta \in \Psi} (U_\theta(\alpha) \times L_\theta(\alpha)) \quad \text{and} \quad g : \bigcup_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi|=t}} \prod_{\theta \in \Psi} (U_\theta(\alpha) \times L_\theta(\alpha)) \rightarrow \Sigma.$$

Here, " $\prod_{\theta \in \Psi}$ " denotes the direct product.

To define f , let $\beta \in \Sigma$, and write $\text{dom}(\beta) = \{i_1, \dots, i_{q+2t}\}$ and $\text{codom}(\beta) = \{l_1, \dots, l_{q+2t}\}$, where $i_1 < \dots < i_{q+2t}$ and $l_1 < \dots < l_{q+2t}$. By t applications of Lemma 3.9, we see that for each $1 \leq h \leq t$, the pair $(\{i_{2h-1}, i_{2h}\}, \{l'_{2h-1}, l'_{2h}\})$ belongs to $U_{\theta_h}(\alpha) \times L_{\theta_h}(\alpha)$ for some $\theta_h \in \Theta(\alpha)$, and that the components $\theta_1, \dots, \theta_t \in \Theta(\alpha)$ are distinct. So we may define

$$f(\beta) = \left((\{i_1, i_2\}, \{l'_1, l'_2\}), \dots, (\{i_{2t-1}, i_{2t}\}, \{l'_{2t-1}, l'_{2t}\}) \right).$$

To define g , let $\Psi = \{\theta_1, \dots, \theta_t\} \subseteq \Theta(\alpha)$ with $\min(\theta_1) < \dots < \min(\theta_t)$ and, for each $1 \leq h \leq t$, let $\{i_{2h-1}, i_{2h}\} \in U_{\theta_h}(\alpha)$ and $\{l'_{2h-1}, l'_{2h}\} \in L_{\theta_h}(\alpha)$, where $i_{2h-1} < i_{2h}$ and $l_{2h-1} < l_{2h}$. By Lemma 3.13, it follows that $i_1 < \dots < i_{2t}$ and $l_1 < \dots < l_{2t}$. Since these vertices belong to unnested edges of α , we may define $\beta \in \mathcal{M}_n$ to be the Motzkin element obtained from α by replacing the non-transversals $\{i_1, i_2\}, \dots, \{i_{2t-1}, i_{2t}\}$ and $\{l'_1, l'_2\}, \dots, \{l'_{2t-1}, l'_{2t}\}$ by the transversals $\{i_1, l'_1\}, \dots, \{i_{2t}, l'_{2t}\}$. Note that β is obtained from α by t applications of the process described in Lemma 3.11, treating the components in the order $\theta_t, \dots, \theta_1$. In particular, $\beta \in E(\mathcal{M}_n)$ and $\alpha = D(\beta)$. By construction, $\text{rank}(\beta) = \text{rank}(\alpha) + 2t = q + 2t$. It follows that $\beta \in \Sigma$, so we may then define

$$g\left((\{i_1, i_2\}, \{l'_1, l'_2\}), \dots, (\{i_{2t-1}, i_{2t}\}, \{l'_{2t-1}, l'_{2t}\}) \right) = \beta.$$

It is easy to check that f and g are mutual inverses.

(iii). We use parts (i) and (ii), and the identity $\prod_{i \in I} (a_i + 1) = \sum_{J \subseteq I} \prod_{j \in J} a_j$, to calculate

$$|D^{-1}(\alpha)| = \sum_{t=0}^k \sum_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi|=t}} \prod_{\theta \in \Psi} (u_\theta(\alpha)l_\theta(\alpha) + 1) = \sum_{\Psi \subseteq \Theta(\alpha)} \prod_{\theta \in \Psi} (u_\theta(\alpha)l_\theta(\alpha) + 1) = \prod_{\theta \in \Theta(\alpha)} (u_\theta(\alpha)l_\theta(\alpha) + 1). \quad \square$$




Figure 6: Interface graphs of the six Motzkin idempotents $\alpha_1, \dots, \alpha_6 \in E(\mathcal{M}_{20})$ satisfying $D(\alpha) = \delta$, where $\delta \in \Delta(\mathcal{M}_{20})$ is pictured in Figure 4. Note that $d(\alpha_1) = d(\alpha_2) = d(\alpha_3) = d(\alpha_4) = \alpha_1 = \delta$, while $d(\alpha_5) = d(\alpha_6) = \alpha_4$. Note also that α_3 is the Motzkin element γ from Figure 2.

To continue the example started above, let $\delta \in \Delta(\mathcal{M}_{20})$ be as in Figure 4. Using the values calculated before the statement of Lemma 3.11, Proposition 3.15(iii) gives

$$|D^{-1}(\delta)| = (u_{\theta_1}(\delta)l_{\theta_1}(\delta) + 1)(u_{\theta_2}(\delta)l_{\theta_2}(\delta) + 1) = (1 \cdot 2 + 1)(1 \cdot 1 + 1) = 6.$$

The interface graphs of the six elements of $D^{-1}(\delta)$ are depicted in Figure 6.

Lemma 3.10 and Proposition 3.15(iii) immediately give the following.

Theorem 3.16. *If S is one of \mathcal{M}_n or \mathcal{J}_n , then $|E(S)| = \sum_{\alpha \in \Delta(S)} \prod_{\theta \in \Theta(\alpha)} (u_{\theta}(\alpha)l_{\theta}(\alpha) + 1)$. □*

To give the corresponding statement for the Kauffman monoid \mathcal{K}_n , for $\alpha \in \mathcal{M}_n$, we write $\Xi(\alpha)$ for the set of *all* cycle components of the interface graph Γ_{α} , noting that $\Theta(\alpha) \subseteq \Xi(\alpha)$. We may identify \mathcal{J}_n with a subset (but not a submonoid) of \mathcal{K}_n , by identifying $\alpha \in \mathcal{J}_n$ with $(0, \alpha) \in \mathcal{K}_n$. By Corollary 3.6, it follows that $E(\mathcal{K}_n) \subseteq E(\mathcal{J}_n)$.

Theorem 3.17. *We have $|E(\mathcal{K}_n)| = \sum_{\substack{\alpha \in \Delta(\mathcal{J}_n) \\ \Xi(\alpha) = \Theta(\alpha)}} \prod_{\theta \in \Theta(\alpha)} (u_{\theta}(\alpha)l_{\theta}(\alpha))$.*

Proof. First, note that

$$E(\mathcal{K}_n) = E(\mathcal{J}_n) \cap E(\mathcal{K}_n) = \left(\bigcup_{\alpha \in \Delta(\mathcal{J}_n)} D^{-1}(\alpha) \right) \cap E(\mathcal{K}_n) = \bigcup_{\alpha \in \Delta(\mathcal{J}_n)} (D^{-1}(\alpha) \cap E(\mathcal{K}_n)).$$

As the sets $D^{-1}(\alpha)$, $\alpha \in \Delta(\mathcal{J}_n)$, are pairwise disjoint, it follows that $|E(\mathcal{K}_n)| = \sum_{\alpha \in \Delta(\mathcal{J}_n)} |D^{-1}(\alpha) \cap E(\mathcal{K}_n)|$. So it remains to show that

$$|D^{-1}(\alpha) \cap E(\mathcal{K}_n)| = \begin{cases} \prod_{\theta \in \Theta(\alpha)} (u_{\theta}(\alpha)l_{\theta}(\alpha)) & \text{if } \Xi(\alpha) = \Theta(\alpha) \\ 0 & \text{otherwise.} \end{cases}$$

With this in mind, let $\alpha \in \Delta(\mathcal{J}_n)$, and write $q = \text{rank}(\alpha)$. If there is a cycle component $\theta \in \Xi(\alpha) \setminus \Theta(\alpha)$, then θ is a cycle component of any $\beta \in D^{-1}(\alpha)$, by Lemma 3.11, and it then follows from Corollary 3.6 that $D^{-1}(\alpha) \cap E(\mathcal{K}_n) = \emptyset$. Next, suppose $\Xi(\alpha) = \Theta(\alpha)$, and put $k = |\Theta(\alpha)|$. By Proposition 3.15(i), an element of $D^{-1}(\alpha)$ has rank $q + 2t$ for some $0 \leq t \leq k$. The interface graph of such an element contains $k - t$ cycle components, so (again using Corollary 3.6) we only obtain an element of $E(\mathcal{K}_n)$ in the case $t = k$, in which

case *all* elements of $D^{-1}(\alpha) \cap E_{q+2t}(\mathcal{M}_n)$ belong to $E(\mathcal{K}_n)$. Proposition 3.15(ii) then gives the stated value of $|D^{-1}(\alpha) \cap E(\mathcal{K}_n)| = |D^{-1}(\alpha) \cap E_{q+2k}(\mathcal{M}_n)|$, since the only term in the sum in Proposition 3.15(ii) when $t = k$ is the $\Psi = \Theta(\alpha)$ term. \square

Remark 3.18. As in Remark 3.7, Theorem 3.17 also gives the number of idempotent basis elements of the Temperley-Lieb and Motzkin algebras in the generic case. As in [9, Section 6], the formula in Theorem 3.17 could be adapted to treat the case in which the twisting parameter ξ is an M th root of unity with $M \leq n$, but we omit the details.

We may also give formulae for the number of idempotents of $\mathcal{M}_n, \mathcal{J}_n, \mathcal{K}_n$ of fixed rank. Recall that for $0 \leq r \leq n$, we write $E_r(\mathcal{M}_n) = \{\alpha \in E(\mathcal{M}_n) : \text{rank}(\alpha) = r\}$. If S is one of \mathcal{J}_n or \mathcal{K}_n , we will write $E_r(S) = S \cap E_r(\mathcal{M}_n)$. Recall that we are identifying $E(\mathcal{K}_n)$ with a subset of $E(\mathcal{J}_n)$. Note that $E_r(\mathcal{J}_n) = E_r(\mathcal{K}_n) = \emptyset$ if $r \not\equiv n \pmod{2}$.

Theorem 3.19. *Let $0 \leq r \leq n$, and write $r = q + 2t$ where $q \in \{0, 1\}$.*

$$(i) \text{ If } S \text{ is one of } \mathcal{M}_n \text{ or } \mathcal{J}_n, \text{ then } |E_r(S)| = \sum_{\substack{\alpha \in \Delta(S) \\ \text{rank}(\alpha) = q}} \sum_{\substack{\Psi \subseteq \Theta(\alpha) \\ |\Psi| = t}} \prod_{\theta \in \Psi} (u_\theta(\alpha) l_\theta(\alpha)).$$

$$(ii) \text{ We have } |E_r(\mathcal{K}_n)| = \sum_{\alpha} \prod_{\theta \in \Theta(\alpha)} (u_\theta(\alpha) l_\theta(\alpha)), \text{ where the sum is over all } \alpha \in \Delta(\mathcal{J}_n) \text{ with } \text{rank}(\alpha) = q, \\ \Xi(\alpha) = \Theta(\alpha) \text{ and } |\Theta(\alpha)| = t.$$

Proof. Part (i) follows quickly from Proposition 3.15 (and its proof), and part (ii) from the proof of Theorem 3.17. \square

4 The algorithms

In this section we describe algorithms for enumerating the idempotents in the Jones, Kauffman and Motzkin monoids, based on the theoretical results obtained in Section 3. These are presented in Algorithms 3–5, below.

In the algorithms described in this section, it is necessary to enumerate the interface graphs of the elements of the Jones monoid \mathcal{J}_n of minimal rank, and of the elements of the Motzkin monoid \mathcal{M}_n of ranks 0 and 1. Roughly speaking, in accordance with Theorems 3.16 and 3.17, the algorithms then involve finding connected components of these interface graphs and counting the number of upper and lower outer hooks involved in every cycle component. This could be achieved using standard graph theoretic algorithms, and there would be essentially nothing further to describe. In general, however, determining the connected components of a graph with v vertices and e edges has complexity $O(v + e)$; see [17, 37, 38]. Hence, in the case of the even degree Jones monoid \mathcal{J}_{2n} , for example, the complexity of this approach would be $O(4nC_n^2)$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n th Catalan number. This is because the interface graph of any element of $\Delta(\mathcal{J}_{2n})$ has $2n$ vertices and $2n$ edges, and since $\Delta(\mathcal{J}_{2n})$ has size C_n^2 [10, Proposition 2.7(iii)]. However, the interface graphs under consideration have several special properties that we can exploit to substantially reduce the run time of our algorithms. In order to explain these properties, and hence the nature of the algorithms, we must first discuss a number of concepts relevant to the enumeration of interface graphs.

4.1 Background on Dyck and Motzkin words

A *Dyck word* is a balanced string of left and right brackets. *Balanced* in this context means that the numbers of left and right brackets are equal, and that at any point, reading from left to right, the number of left brackets is at least the number of right brackets. A Dyck word necessarily has even length. For example, $u = ((\))(\))$ is a Dyck word of length 8. A recent algorithm of Neri [32, 33] allows for fast generation of Dyck words. We will write D_{2n} for the set of all Dyck words of length $2n$. So $|D_{2n}| = C_n$ is the n th Catalan number.

A *Motzkin word* is a string of left and right brackets and dots, such that the subword consisting only of the brackets is a Dyck word. A Motzkin word can have any length. For example, $v = ()(\cdot())\cdot(\cdot)(\cdot)(\cdot)$ is a Motzkin word of length 18. A Motzkin word of length n can be thought of as a pair consisting of a Dyck word of length m , for some even $m \leq n$, and a subset of $\mathbf{n} = \{1, \dots, n\}$ of size $n - m$; the subset specifies

the positions of the dots in the Motzkin word. For example, the above Motzkin word v corresponds to the Dyck word $()((()))((())) \in D_{14}$ and the subset $\{4, 8, 10, 11\}$. We will write M_n for the set of all Motzkin words of length n ; there should be no confusion with the Motzkin monoid itself, which is denoted by \mathcal{M}_n . So $|M_n| = \mu(n, 0)$, where the numbers $\mu(n, r)$ were defined in (2.4) and (2.5). It is relatively straightforward to produce the sets M_n for the values of n we are concerned with here, namely for $n \leq 20$. For instance, there are 50 852 019 Motzkin words of length 20, and these can be produced and stored in a convenient format for use in Algorithm 5 in about 35 seconds, and using about 2 GB of memory. In particular, creating and storing the Motzkin words of a given length represents a tiny fraction of the time taken by Algorithm 5. We will not describe the process for producing the Motzkin words in more detail here.

Denote by \mathcal{S}_n the *symmetric group* of degree n , which consists of all permutations of \mathbf{n} . There is a natural injective map

$$\mathbf{p} : M_n \rightarrow \mathcal{S}_n$$

taking a Motzkin word $w \in M_n$ to the permutation $\mathbf{p}(w) \in \mathcal{S}_n$ defined, for $i \in \mathbf{n}$, by

$$\mathbf{p}(w)(i) = \begin{cases} i & \text{if } w \text{ has a dot in position } i \\ j & \text{if } w \text{ has a bracket in position } i \text{ that is matched by a bracket in position } j. \end{cases}$$

So, if we suppose $w \in M_n$ has left brackets at positions $i_1 < \dots < i_m$, and that these are matched by right brackets at positions j_1, \dots, j_m , respectively, then $\mathbf{p}(w)$ can be written as a product of commuting transpositions as $\mathbf{p}(w) = (i_1 j_1) \cdots (i_m j_m)$. For example, with $u \in D_8$ and $v \in M_{18}$ as defined above,

$$\mathbf{p}(u) = (1\ 8)(2\ 5)(3\ 4)(6\ 7) \in \mathcal{S}_8 \quad \text{and} \quad \mathbf{p}(v) = (1\ 2)(3\ 7)(5\ 6)(9\ 12)(13\ 18)(14\ 15)(16\ 17) \in \mathcal{S}_{18}.$$

In general, for a Motzkin word $w \in M_n$, $\mathbf{p}(w)$ has no fixed points if and only if $w \in D_n$; in this case, n must be even. Note also that if $w \in M_n$ has at least one bracket, then $\mathbf{p}(w)$ is an involution (a permutation of order 2); if w consists only of dots, then $\mathbf{p}(w)$ is the identity element of \mathcal{S}_n .

Now consider a Motzkin element $\alpha \in \mathcal{M}_n$ with $\text{rank}(\alpha) = 0$. The upper blocks of α induce a Motzkin word $w_1 \in M_n$ in a natural way; for each upper hook $\{i, j\}$ of α with $i < j$, w_1 has a left bracket at position i and a right bracket at position j , while the upper singletons of α correspond to the dots of w_1 . Similarly, the lower blocks of α induce a second Motzkin word $w_2 \in M_n$. We will write $\mathbf{m}(\alpha) = (w_1, w_2) \in M_n \times M_n$ for the pair consisting of these two words. Conversely, given any pair $(w_1, w_2) \in M_n \times M_n$, it is easy to see that there is a Motzkin element $\alpha \in \mathcal{M}_n$ of rank 0 with $\mathbf{m}(\alpha) = (w_1, w_2)$.

To describe the Motzkin elements of rank 1 in terms of Motzkin words, we first define M'_{n+1} to be the subset of M_{n+1} consisting of all Motzkin words of length $n+1$ whose last symbol is a right bracket. Equivalently, a Motzkin word $w \in M_{n+1}$ belongs to M'_{n+1} if $\mathbf{p}(w)(n+1) \neq n+1$. Now consider a Motzkin element $\alpha \in \mathcal{M}_n$ with $\text{rank}(\alpha) = 1$, and let the unique transversal of α be $\{i, j\}$. We define the Motzkin word $w_1 \in M'_{n+1}$ to have a left bracket at position i , a right bracket at position $n+1$, and where the remaining symbols of w_1 are determined by the upper blocks of α in the same way as in the previous paragraph. We define w_2 analogously, in terms of j and the lower blocks of α . Again, we will write $\mathbf{m}(\alpha) = (w_1, w_2) \in M'_{n+1} \times M'_{n+1}$ for the pair consisting of these two words. Again, given any pair $(w_1, w_2) \in M'_{n+1} \times M'_{n+1}$, there is a Motzkin element $\alpha \in \mathcal{M}_n$ of rank 1 with $\mathbf{m}(\alpha) = (w_1, w_2)$.

The previous two paragraphs describe a bijection

$$\mathbf{m} : \{\alpha \in \mathcal{M}_n : \text{rank}(\alpha) \leq 1\} \rightarrow (M_n \times M_n) \cup (M'_{n+1} \times M'_{n+1}).$$

We denote by \mathbf{d} the restriction of \mathbf{m} to the Jones elements $\{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) \leq 1\}$ of rank at most 1. The restriction \mathbf{d} is a bijection onto its image, which is $D_{2\lceil \frac{n}{2} \rceil} \times D_{2\lfloor \frac{n}{2} \rfloor}$: that is, either $D_n \times D_n$ or $D_{n+1} \times D_{n+1}$, according to whether n is even or odd, respectively. In particular, Jones elements of minimum rank correspond to certain pairs of Dyck words of an appropriate length.

We noted above that $|D_{2n}| = C_n$ and that $|M_n| = \mu(n, 0)$ for any n . It is also known [10, Proposition 2.8] that $|M'_{n+1}| = \mu(n, 1)$. In the algorithms presented in this section we will fix (arbitrary) orderings on the sets D_{2n} , M_n and M'_{n+1} , and will denote the elements of these sets as

$$D_{2n} = \{u_i : 1 \leq i \leq C_n\}, \quad M_n = \{w_i : 1 \leq i \leq \mu(n, 0)\}, \quad M'_{n+1} = \{w'_i : 1 \leq i \leq \mu(n, 1)\}.$$

If $w \in M_n$ is a Motzkin word, then we say that a left bracket of w is an *outer bracket* if this bracket is not enclosed by any other brackets. We define $\mathbf{O}(w)$ to be the subset of \mathbf{n} for which $i \in \mathbf{O}(w)$ if and only

if w has an outer bracket at position i . For example, for $u \in D_8$ and $v \in M_{18}$ defined above, $\mathbf{O}(u) = \{1\}$ and $\mathbf{O}(v) = \{1, 3, 9, 13\}$.

Finally, recall that in Subsection 3.3 we defined and studied a map

$$D : E(\mathcal{M}_n) \rightarrow \{\alpha \in E(\mathcal{M}_n) : \text{rank}(\alpha) \leq 1\}.$$

In this section, for convenience, we will write $\hat{\alpha} = D(\alpha)$ for any $\alpha \in E(\mathcal{M}_n)$.

4.2 The algorithm for Jones idempotents

Algorithm 3 contains pseudocode for counting the number of idempotents in the Jones monoid \mathcal{J}_n . A C++ implementation of this algorithm can be found at [30]. Roughly speaking, the algorithm begins by enumerating the elements of $\Delta(\mathcal{J}_n)$ in terms of pairs (u_i, u_j) of Dyck words of length n or $n+1$, as appropriate. It then proceeds to count the outer hooks in each connected component of the interface graph of the Jones element $\mathbf{d}^{-1}(u_i, u_j) \in \Delta(\mathcal{J}_n)$; this then yields the number of idempotents $\alpha \in E(\mathcal{J}_n)$ with $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$, according to Proposition 3.15(iii). The algorithm then concludes by summing these values.

Algorithm 3 Count the number of idempotents in the Jones monoid \mathcal{J}_n

1:	$N := 0$	<i>[Number of idempotents]</i>
2:	for $i \in \{1, \dots, C_{\lceil n/2 \rceil}\}$ do	<i>[Loop over Dyck words of length n or $n+1$]</i>
3:	$N \leftarrow N + 2^{ \mathbf{O}(u_i) \setminus \{\mathbf{p}(u_i)(n+1)\} }$	<i>[\(\alpha \in E(\mathcal{J}_n)\) such that $\mathbf{d}(\hat{\alpha}) = (u_i, u_i)\)$</i>
4:	for $j \in \{i+1, \dots, C_{\lceil n/2 \rceil}\}$ do	<i>[Loop over Dyck words]</i>
5:	$M := 1$	<i>[Number of idempotents α with $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$]</i>
6:	$m := 0$	<i>[Largest value seen in any cycle of $\Gamma_{\mathbf{d}^{-1}(u_i, u_j)}$]</i>
7:	while $m < \max\{\mathbf{O}(u_j) \setminus \{\mathbf{p}(u_j)(n+1)\}\}$ do	<i>[Loop over cycles of $\Gamma_{\mathbf{d}^{-1}(u_i, u_j)}$]</i>
8:	$k, l \leftarrow \min\{x \in \mathbf{O}(u_j) : x \geq m\}$	<i>[Start of the next cycle]</i>
9:	$I, J := 0$	<i>[Count the number of outer hooks in this cycle]</i>
10:	repeat	<i>[Loop within the current cycle]</i>
11:	if $l \in \mathbf{O}(u_i)$ then	<i>[Found an outer bracket of u_i in current cycle]</i>
12:	$I \leftarrow I + 1$	
13:	if $l \in \mathbf{O}(u_j)$ then	<i>[Found an outer bracket of u_j in current cycle]</i>
14:	$J \leftarrow J + 1$	
15:	$m \leftarrow \max\{m, \mathbf{p}(u_j)(l)\}$	
16:	$l \leftarrow \mathbf{p}(u_i)\mathbf{p}(u_j)(l)$	<i>[Go to the next position in the current cycle]</i>
17:	until $l = k$	<i>[Returned to the start of the cycle]</i>
18:	$M \leftarrow M(IJ + 1)$	<i>[Multiply by number of outer brackets in current cycle]</i>
19:	$N \leftarrow N + 2M$	<i>[Add number of idempotents α with $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$]</i>
20:	return N	

Before moving on to the other algorithms, we first comment on a number of features of Algorithm 3, including some simple optimisations that have been included.

First, if $\alpha \in \Delta(\mathcal{J}_n)$ is such that $\mathbf{d}(\alpha) = (u_i, u_i)$ for some i , then every component of the interface graph Γ_α is a cycle of length 2 or an active path of length 0 (the latter only occurs when n is odd, in which case there is a unique such path). As such, Proposition 3.15(iii) tells us that $D^{-1}(\alpha)$ has size 2^k if n is even, or 2^{k-1} if n is odd, respectively, where k is the number of outer (left) brackets of u_i . The reason for subtracting 1 from k in the case n is odd is that the last outer bracket of $u_i \in D_{n+1}$ corresponds to the path component of Γ_α . See Line 3 of Algorithm 3. Lines 3 and 7 refer to $\mathbf{p}(u_i)(n+1)$, which is only defined when n is odd, and which can be ignored when n is even. In the implementation in [30], Algorithm 3 is split into two parts covering the even and odd cases separately.

If u_i and u_j are distinct Dyck words, then there are the same number of idempotents $\alpha \in E(\mathcal{J}_n)$ with $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$ as there are with $\mathbf{d}(\hat{\alpha}) = (u_j, u_i)$. This corresponds to the anti-involution $*$: $\mathcal{J}_n \rightarrow \mathcal{J}_n$, and the fact that a Jones element α is an idempotent if and only if α^* is, since $[\mathbf{d}^{-1}(u_i, u_j)]^* = \mathbf{d}^{-1}(u_j, u_i)$.

A further optimization along these lines is available in the case that n is even. Namely, for any n , whether even or odd, there is an involution \dagger : $\mathcal{J}_n \rightarrow \mathcal{J}_n$, where α^\dagger is the result of reflecting α in a vertical axis midway between points 1 and n . The involution \dagger was studied along with the anti-involution $*$ in [2]. It is again clear that $\alpha \in \mathcal{J}_n$ is an idempotent if and only if α^\dagger is. For an even value of n , and for $\alpha \in \mathcal{J}_n$ of rank 0,

if $\mathbf{d}(\alpha) = (u_i, u_j)$, then $\mathbf{d}(\alpha^\dagger) = (\text{rev}(u_i), \text{rev}(u_j))$, where $\text{rev}(w)$ is the result of writing w in reverse and interchanging left and right brackets. This means that in the case that α and α^\dagger are not equal, we only need to calculate the size of one of $D^{-1}(\alpha)$ or $D^{-1}(\alpha^\dagger)$. The implementation of this optimisation is rather technical, and only applies in the even case (since the active paths in the interface graphs of rank 1 Jones elements interfere with the \dagger map in the case of odd n), so we have not included it in the pseudocode for Algorithm 3. This optimisation is included in the implementation [30].

As a further note, it is not necessary to check if $l \in \mathbf{O}(u_i)$ and $l \in \mathbf{O}(u_j)$ in Lines 11 and 13 of Algorithm 3, since it can be shown that if $l \in \mathbf{O}(u_i)$, then $l \notin \mathbf{O}(u_j)$, and vice versa, unless l is the minimum vertex in its component. In fact, apart from the above-mentioned exception, it is only possible to have $l \in \mathbf{O}(u_j)$ before the first time that $l \in \mathbf{O}(u_i)$, and this could be separated into another loop to reduce the number of branches in the innermost loops. However, for the sake of brevity we do not include this optimization in the pseudocode in Algorithm 3, although it is included in the implementation [30].

Finally, we note that Algorithm 3 is *embarrassingly parallel*, in the sense that the number of idempotents $\alpha \in E(\mathcal{J}_n)$ such that $\mathbf{d}(\hat{\alpha}) = (u_i, u_j) \in D_{2\lceil n/2 \rceil} \times D_{2\lceil n/2 \rceil}$ can be enumerated independently for different values of i and j .

4.3 The algorithm for Kauffman idempotents

The key difference between Algorithm 3 for the Jones monoid \mathcal{J}_n and Algorithm 4 (below) for the Kauffman monoid \mathcal{K}_n is that if, for some $\alpha \in \Delta(\mathcal{J}_n)$, there is a cycle in the interface graph Γ_α containing no upper outer hooks or no lower outer hooks, then $D^{-1}(\alpha) \cap E(\mathcal{K}_n)$ is empty; see the proof of Theorem 3.17. Hence, in Algorithm 4, *every* cycle of Γ_α must be considered and not only those starting at an outer hook as was the case in Algorithm 3. Again, a number of optimisations are included in Algorithm 4, but we will not comment explicitly on these, as they are virtually identical to those in Algorithm 3.

Note that for any $1 \leq i \leq C_{\lceil n/2 \rceil}$, there are no idempotents $\alpha \in E(\mathcal{K}_n)$ with $\mathbf{d}(\hat{\alpha}) = (u_i, u_i)$ unless $u_i = ()() \dots ()$, in which case the identity element is the only such idempotent. This is why we start with $N = 1$ in Line 1 of Algorithm 4, and why we only consider pairs (u_i, u_j) with $i \neq j$; see Lines 2 and 3.

Algorithm 4 Count the number of idempotents in the Kauffman monoid \mathcal{K}_n

```

1:  $N := 1$  [Number of idempotents]
2: for  $i \in \{1, \dots, C_{\lceil n/2 \rceil}\}$  do [Loop over Dyck words of length  $n$  or  $n + 1$ ]
3:   for  $j \in \{i + 1, \dots, C_{\lceil n/2 \rceil}\}$  do [Loop over Dyck words]
4:      $M := 1$  [Number of idempotents  $\alpha \in E(\mathcal{K}_n)$  with  $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$ ]
5:      $b := 1$  [The current vertex]
6:      $B := \emptyset$  [The vertices of  $\Gamma_{\mathbf{d}^{-1}(u_i, u_j)}$  seen already]
7:     while  $M \neq 0$  and  $b < n + 1$  do [Loop over cycles of  $\Gamma_{\mathbf{d}^{-1}(u_i, u_j)}$ ]
8:        $I, J := 0$  [Count the number of outer hooks in this cycle]
9:       repeat
10:        if  $b \in \mathbf{O}(u_i)$  then [Found an outer bracket of  $u_i$  in current cycle]
11:           $I \leftarrow I + 1$ 
12:        if  $b \in \mathbf{O}(u_j)$  then [Found an outer bracket of  $u_j$  in current cycle]
13:           $J \leftarrow J + 1$ 
14:           $B \leftarrow B \cup \{b\}$ 
15:           $b \leftarrow \mathbf{p}(u_i)\mathbf{p}(u_j)(b)$  [Go to the next position in the current cycle]
16:        until  $b \in B$ 
17:         $M \leftarrow MIJ$  [Multiply by number of outer brackets in current cycle]
18:        while  $M \neq 0$  and  $b \in B$  do [Find the next vertex  $b$  not seen already]
19:           $b \leftarrow b + 1$ 
20:      $N \leftarrow N + 2M$  [Add number of idempotents  $\alpha$  with  $\mathbf{d}(\hat{\alpha}) = (u_i, u_j)$ ]
21: return  $N$ 

```

4.4 The algorithm for Motzkin idempotents

Algorithm 5 contains pseudocode for calculating the number of idempotents in the Motzkin monoid \mathcal{M}_n . The basic idea of Algorithm 5 is similar to Algorithm 3, except that separate parts are required to count

idempotents of even rank (Lines 2–20) and odd rank (Lines 21–41). Similar optimisations to Algorithms 3 and 4 have been included.

Algorithm 5 Count the number of idempotents in the Motzkin monoid \mathcal{M}_n

```

1:  $N := 0$  [Number of idempotents]
2: for  $i \in \{1, \dots, \mu(n, 0)\}$  do [Loop over Motzkin words of length  $n$ ]
3:    $N \leftarrow N + 2^{|\mathbf{O}(w_i)|}$  [ $\alpha \in E(\mathcal{M}_n)$  with  $\mathbf{m}(\hat{\alpha}) = (w_i, w_i)$ ]
4:   for  $j \in \{i + 1, \dots, \mu(n, 0)\}$  do [Loop over Motzkin words]
5:      $M := 1$  [Number of idempotents  $\alpha$  with  $\mathbf{m}(\hat{\alpha}) = (w_i, w_j)$ ]
6:      $m := 0$  [Largest value seen in any path of  $\Gamma_{\mathbf{m}^{-1}(w_i, w_j)}$ ]
7:     while  $m < \max\{\mathbf{O}(w_j)\}$  do [Loop over paths of  $\Gamma_{\mathbf{m}^{-1}(w_i, w_j)}$ ]
8:        $k, l \leftarrow \min\{x \in \mathbf{O}(w_j) : x \geq m\}$  [Start of the next path]
9:        $I, J := 0$  [Count the number of outer hooks in this cycle]
10:      repeat [Follow the current path]
11:        if  $l \in \mathbf{O}(w_i)$  then [Found an outer bracket of  $w_i$  in current path]
12:           $I \leftarrow I + 1$ 
13:        if  $l \in \mathbf{O}(w_j)$  then [Found an outer bracket of  $w_j$  in current path]
14:           $J \leftarrow J + 1$ 
15:           $m \leftarrow \max\{m, \mathbf{p}(w_j)(l)\}$ 
16:           $l \leftarrow \mathbf{p}(w_i)\mathbf{p}(w_j)(l)$  [Go to the next position in the current path]
17:        until  $l = k$  or  $\mathbf{p}(w_i)(l) = l$  or  $\mathbf{p}(w_j)\mathbf{p}(w_i)(l) = \mathbf{p}(w_i)(l)$ 
18:        if  $l = k$  then [The current path is a cycle]
19:           $M \leftarrow M(IJ + 1)$  [Multiply by number of outer brackets in current cycle]
20:       $N \leftarrow N + 2M$  [Add number of idempotents  $\alpha$  with  $\mathbf{m}(\hat{\alpha}) = (w_i, w_j)$ ]
21: for  $i \in \{1, \dots, \mu(n, 1)\}$  do [Loop over Motzkin words of length  $n + 1$  in  $\mathcal{M}'_{n+1}$ ]
22:    $N \leftarrow N + 2^{|\mathbf{O}(w_i)|}$  [ $\alpha \in E(\mathcal{M}_n)$  with  $\mathbf{m}(\hat{\alpha}) = (w'_i, w'_i)$ ]
23:   for  $j \in \{i + 1, \dots, \mu(n, 1)\}$  do [Loop over elements of  $\mathcal{M}'_{n+1}$ ]
24:      $M := 1$  [Number of idempotents  $\alpha$  with  $\mathbf{m}(\hat{\alpha}) = (w'_i, w'_j)$ ]
25:      $m := n + 1$  [Smallest value seen in any path of  $\Gamma_{\mathbf{m}^{-1}(w'_i, w'_j)}$ ]
26:     while  $m > \min\{\mathbf{O}(w'_j)\}$  and  $M \neq 0$  do [Loop over paths of  $\Gamma_{\mathbf{m}^{-1}(w'_i, w'_j)}$ ]
27:        $k, l \leftarrow \max\{\mathbf{p}(w'_j)(x) : x \in \mathbf{O}(w'_j), x \leq m\}$  [Start of the next path]
28:        $I, J := 0$  [Count the number of outer hooks in this cycle]
29:       repeat [Follow the current path]
30:         if  $\mathbf{p}(w'_j)(l) \in \mathbf{O}(w'_i)$  then [Found an outer bracket of  $w'_i$  in current path]
31:            $I \leftarrow I + 1$ 
32:         if  $\mathbf{p}(w'_j)(l) \in \mathbf{O}(w'_j)$  then [Found an outer bracket of  $w'_j$  in current path]
33:            $J \leftarrow J + 1$ 
34:            $m \leftarrow \min\{m, l\}$ 
35:            $l \leftarrow \mathbf{p}(w'_i)\mathbf{p}(w'_j)(l)$  [Go to the next position in the current path]
36:         until  $l = k$  or  $\mathbf{p}(w'_i)(l) = l$  or  $\mathbf{p}(w'_j)\mathbf{p}(w'_i)(l) = \mathbf{p}(w'_i)(l)$ 
37:         if  $l = k$  then [The current path is a cycle]
38:            $M \leftarrow M(IJ + 1)$  [Multiply by number of outer brackets in current cycle]
39:         else if  $k = n + 1$  then [The current path is not a cycle]
40:            $M \leftarrow 0$  [There are no  $\alpha$  with  $\mathbf{m}(\hat{\alpha}) = (w'_i, w'_j)$  to count]
41:        $N \leftarrow N + 2M$  [Add number of idempotents  $\alpha$  with  $\mathbf{m}(\hat{\alpha}) = (w'_i, w'_j)$ ]
42: return  $N$ 

```

5 Values and benchmarking

In this section, we give some calculated values of the various number sequences we have considered. Tables 2 and 3 give the number of idempotents in the Jones, Kauffman and Motzkin monoids, as well as comparative running times for the various algorithms described in the paper. For each algorithm, these values were computed using GAP or [30], as appropriate, on an IBM power8 (8247-22L), with 24 cores at 3,026 GHz

(giving 192 threads) running powerKVM. At the time of writing, these represent the largest known values of $|E(\mathcal{J}_n)|$, $|E(\mathcal{K}_n)|$ and $|E(\mathcal{M}_n)|$; cf. Sequences A225798, A281438 and A256672 on [1]. Note that values of $|E(\mathcal{J}_{2n})|$ can be computed faster than $|E(\mathcal{J}_{2n-1})|$ because of the \dagger map discussed in Subsection 4.2.

n	$ E(\mathcal{J}_n) $	Alg. 1	Alg. 2	Alg. 3
0	1			
1	1			
2	2			
3	5			
4	12			
5	36			
6	96			
7	311			
8	886			
9	3000			
10	8944			
11	31 192			
12	96 138			
13	342 562	2		
14	1083 028	8		
15	3923 351	32		
16	12 656 024	5901	1	
17	46 455 770	-	4	
18	152 325 850	-	16	
19	565 212 506	-	51	
20	1878 551 444	-	214	
21	7033 866 580	-	689	2
22	23 645 970 022	-	-	2
23	89 222 991 344	-	-	29
24	302 879 546 290	-	-	23
25	1150 480 017 950	-	-	522
26	3938 480 377 496	-	-	500
27	15 047 312 553 918	-	-	7260
28	51 892 071 842 570	-	-	5520
29	199 274 492 098 480	-	-	101 160
30	691 680 497 233 180	-	-	77 100

n	$ E(\mathcal{K}_n) $	Alg. 4
0	1	
1	1	
2	1	
3	3	
4	5	
5	15	
6	31	
7	93	
8	215	
9	653	
10	1619	
11	4979	
12	12 949	
13	40 293	
14	108 517	
15	341 241	
16	943 937	
17	2996 127	
18	8465 319	
19	27 092 419	
20	77 878 271	
21	251 073 791	5
22	732 129 719	5
23	2375 764 351	60
24	7012 025 277	67
25	22 886 955 207	787
26	68 254 122 669	912
27	223 946 197 065	10 740
28	673 885 100 857	12 300
29	2221 505 541 773	147 300
30	6737 598 265 009	165 720

Table 2: Left: the number of idempotents in the Jones monoid \mathcal{J}_n , and the time in seconds to calculate these numbers using Algorithms 1, 2 and 3. Right: the number of idempotents in the Kauffman monoid \mathcal{K}_n , and the time in seconds to calculate these numbers using Algorithm 4; note that Algorithms 1 and 2 do not apply to \mathcal{K}_n , as it is neither finite nor a regular $*$ -semigroup; see Section 2.

Tables 4, 5 and 6 give values of $|E_r(\mathcal{J}_n)|$, $|E_r(\mathcal{K}_n)|$ and $|E_r(\mathcal{M}_n)|$, respectively, for values of $n \leq 10$; recall that $E_r(S)$ is the set of all idempotents of S of rank r , where S is any of \mathcal{J}_n , \mathcal{K}_n or \mathcal{M}_n ; cf. Sequences A281441, A281442 and A269736 on [1]. These values were calculated using the Semigroups package for GAP [31]. Higher values of these sequences could be calculated, by modifying Algorithms 3, 4 and 5 in light of Theorem 3.19, but we have not done so. Note also that for odd n , $|E_1(\mathcal{K}_n)|$ is a *meandric number*; see Sequences A005315 and A005316 in [1], and also [6, 7].

As noted earlier, even though the monoid \mathcal{PP}_n of all planar partitions of degree n is isomorphic to the Jones monoid \mathcal{J}_{2n} of degree $2n$, this is not true of their twisted versions, \mathcal{PP}_n^τ and $\mathcal{J}_{2n}^\tau = \mathcal{K}_{2n}$. In general, \mathcal{J}_{2n} contains more idempotents than \mathcal{PP}_n^τ . The methods of this paper do not lead to algorithms for counting the idempotents of \mathcal{PP}_n^τ . However, for completeness, we used GAP [31] to calculate the number of these idempotents for $n \leq 10$. Table 7 gives the total number of these idempotents, while Table 8 gives the number of idempotents of a fixed rank; cf. Sequences A286867 and A289620 on [1].

n	$ E(\mathcal{M}_n) $	Alg. 1	Alg. 2	Alg. 5
0	1			
1	2			
2	7			
3	31			
4	153			
5	834			
6	4839			
7	29 612			
8	188 695	3		
9	1243 746	30	2	
10	8428 597	-	2	
11	58 476 481	-	12	
12	413 893 789	-	81	
13	2980 489 256	-	640	2
14	21 787 216 989	-	5424	18
15	161 374 041 945	-	46 330	212
16	1209 258 743 839	-	-	1917
17	9155 914 963 702	-	-	16 200
18	69 969 663 242 487	-	-	136 980
19	539 189 056 700 627	-	-	1096 320

Table 3: The number of idempotents in the Motzkin monoid \mathcal{M}_n , and the time in seconds to calculate these numbers using Algorithms 1, 2 and 5.

Acknowledgements

We would like to thank Dr Jan De Beule (Vrije Universiteit Brussel) and the Department of Mathematics (WE01) of Ghent University for the use of their IBM power8.

References

- [1] The on-line encyclopedia of integer sequences. Published electronically at <http://oeis.org/>, 2017.
- [2] Karl Auinger, Igor Dolinka, and Mikhail V. Volkov. Equational theories of semigroups with involution. *J. Algebra*, 369:203–225, 2012.
- [3] Georgia Benkart and Tom Halverson. Motzkin algebras. *European J. Combin.*, 36:473–502, 2014.
- [4] Mirjana Borisavljević, Kosta Došen, and Zoran Petrić. Kauffman monoids. *J. Knot Theory Ramifications*, 11(2):127–143, 2002.
- [5] Richard Brauer. On algebras which are connected with the semisimple continuous groups. *Ann. of Math. (2)*, 38(4):857–872, 1937.
- [6] Michael La Croix. *Approaches to the enumerative theory of meanders*, Available at <http://www.math.uwaterloo.ca/~malacroix/Latex/Meanders.pdf>, 2003.
- [7] P. Di Francesco, O. Golinelli, and E. Guitter. Meanders and the Temperley-Lieb algebra. *Comm. Math. Phys.*, 186(1):1–59, 1997.
- [8] Igor Dolinka and James East. The idempotent-generated subsemigroup of the Kauffman monoid. *Glasg. Math. J.*, 59(3):673–683, 2017.
- [9] Igor Dolinka, James East, Athanasios Evangelou, Des FitzGerald, Nicholas Ham, James Hyde, and Nicholas Loughlin. Enumeration of idempotents in diagram semigroups and algebras. *J. Combin. Theory Ser. A*, 131:119–152, 2015.
- [10] Igor Dolinka, James East, and Robert D. Gray. Motzkin monoids and partial Brauer monoids. *J. Algebra*, 471:251–298, 2017.
- [11] J. East, A. Egri-Nagy, J. D. Mitchell, and Y. Péresse. Computing finite semigroups. *J. Symbolic Comput.*, to appear, [arXiv:1510.01868](https://arxiv.org/abs/1510.01868).
- [12] James East. On the singular part of the partition monoid. *Internat. J. Algebra Comput.*, 21(1-2):147–178, 2011.
- [13] James East and D. G. FitzGerald. The semigroup generated by the idempotents of a partition monoid. *J. Algebra*, 372:108–133, 2012.

$n \setminus r$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1		1									
2	1		1								
3		4		1							
4	4		7		1						
5		25		10		1					
6	25		57		13		1				
7		196		98		16		1			
8	196		522		148		19		1		
9		1764		1006		207		22		1	
10	1764		5206		1673		275		25		1

Table 4: The number of rank r idempotents in the Jones monoid \mathcal{J}_n .

$n \setminus r$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1		1									
2	0		1								
3		2		1							
4	0		4		1						
5		8		6		1					
6	0		22		8		1				
7		42		40		10		1			
8	0		140		62		12		1		
9		262		288		88		14		1	
10	0		992		492		118		16		1

Table 5: The number of rank r idempotents in the Kauffman monoid \mathcal{K}_n .

- [14] James East and Robert D. Gray. Diagram monoids and Graham–Houghton graphs: Idempotents and generating sets of ideals. *J. Combin. Theory Ser. A*, 146:63–128, 2017.
- [15] James East, James D. Mitchell, Nik Ruškuc, and Michael Torpey. Congruence lattices of finite diagram monoids. *Adv. Math.*, 333:931–1003, 2018.
- [16] Véronique Froidure and Jean-Eric Pin. Algorithms for computing finite semigroups. In *Foundations of computational mathematics (Rio de Janeiro, 1997)*, pages 112–126. Springer, Berlin, 1997.
- [17] Harold N. Gabow. Path-based depth-first search for strong and biconnected components. *Inform. Process. Lett.*, 74(3-4):107–114, 2000.
- [18] Tom Halverson and Arun Ram. Partition algebras. *European J. Combin.*, 26(6):869–921, 2005.
- [19] Peter M. Higgins. *Techniques of semigroup theory*. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1992.
- [20] John M. Howie. *Fundamentals of semigroup theory*, volume 12 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 1995. Oxford Science Publications.
- [21] Teruo Imaoka. On fundamental regular $*$ -semigroups. *Mem. Fac. Sci. Shimane Univ.*, 14:19–23, 1980.
- [22] V. F. R. Jones. Hecke algebra representations of braid groups and link polynomials. *Ann. of Math. (2)*, 126(2):335–388, 1987.
- [23] V. F. R. Jones. The Potts model and the symmetric group. In *Subfactors (Kyuzeso, 1993)*, pages 259–267. World Sci. Publ., River Edge, NJ, 1994.
- [24] Louis H. Kauffman. An invariant of regular isotopy. *Trans. Amer. Math. Soc.*, 318(2):417–471, 1990.
- [25] Kwok Wai Lau and D. G. FitzGerald. Ideal structure of the Kauffman and related monoids. *Comm. Algebra*, 34(7):2617–2629, 2006.
- [26] Gustav Lehrer and Ruibin Zhang. The second fundamental theorem of invariant theory for the orthogonal group. *Ann. of Math. (2)*, 176(3):2031–2054, 2012.
- [27] Victor Maltcev and Volodymyr Mazorchuk. Presentation of the singular part of the Brauer monoid. *Math. Bohem.*, 132(3):297–323, 2007.

$n \setminus r$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1	1	1									
2	4	2	1								
3	16	11	3	1							
4	81	48	19	4	1						
5	441	266	93	28	5	1					
6	2601	1492	549	152	38	6	1				
7	16 129	9042	3211	947	226	49	7	1			
8	104 329	56 712	20 004	5784	1480	316	61	8	1		
9	697 225	369 689	127 676	37 048	9432	2169	423	74	9	1	
10	4787 344	2477 806	841 945	241 268	62 149	14 402	3036	548	88	10	1

Table 6: The number of rank r idempotents in the Motzkin monoid \mathcal{M}_n .

n	0	1	2	3	4	5	6	7	8	9	10
$ E(\mathcal{PP}_n^T) $	1	1	6	44	362	3226	30 488	301 460	3090 020	32 618 046	345 515 557

Table 7: The number of idempotents in the twisted planar partition monoid \mathcal{PP}_n^T .

- [28] Paul Martin. Temperley-Lieb algebras for nonplanar statistical mechanics—the partition algebra construction. *J. Knot Theory Ramifications*, 3(1):51–82, 1994.
- [29] Paul Martin and Volodymyr Mazorchuk. On the representation theory of partial Brauer algebras. *Q. J. Math.*, 65(1):225–247, 2014.
- [30] J. D. Mitchell. <https://github.com/james-d-mitchell/Jones>, 2016.
- [31] J. D. Mitchell et al. *Semigroups - GAP package, Version 3.0.16*, May 2018, <http://dx.doi.org/10.5281/zenodo.592893>.
- [32] C. Neri. <https://github.com/cassioneri/Dyck>, 2016.
- [33] C. Neri. A loopless and branchless $O(1)$ algorithm to generate the next Dyck word. *Preprint*, 2016, arXiv:1602.06426.
- [34] T. E. Nordahl and H. E. Scheiblich. Regular $*$ -semigroups. *Semigroup Forum*, 16(3):369–377, 1978.
- [35] Mario Petrich. Certain varieties of completely regular $*$ -semigroups. *Boll. Un. Mat. Ital. B (6)*, 4(2):343–370, 1985.
- [36] John Rhodes and Benjamin Steinberg. *The q -theory of finite semigroups*. Springer Monographs in Mathematics. Springer, New York, 2009.
- [37] Robert Sedgewick. *Algorithms*. Addison-Wesley Series in Computer Science. Addison-Wesley Publishing Company, Advanced Book Program, Reading, MA, 1983.
- [38] Robert Tarjan. Depth-first search and linear graph algorithms. *SIAM J. Comput.*, 1(2):146–160, 1972.
- [39] H. N. V. Temperley and E. H. Lieb. Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the “percolation” problem. *Proc. Roy. Soc. London Ser. A*, 322(1549):251–280, 1971.
- [40] Stewart Wilcox. Cellularity of diagram algebras as twisted semigroup algebras. *J. Algebra*, 309(1):10–31, 2007.

$n \setminus r$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1	0	1									
2	0	5	1								
3	0	33	10	1							
4	0	253	93	15	1						
5	0	2147	880	178	20	1					
6	0	19 593	8599	1982	288	25	1				
7	0	188 837	86 762	21 723	3684	423	30	1			
8	0	1899 107	900 997	238 419	44 767	6111	583	35	1		
9	0	19 761 209	9595 264	2638 114	531 656	81 606	9388	768	40	1	
10	0	211 447 863	104 447 385	29 503 900	6255 952	1044 248	136 740	13 640	978	45	1

Table 8: The number of rank r idempotents in the twisted planar partition monoid \mathcal{PP}_n^T .