

Efficient Singlet Oxygen Photogeneration by Zinc Porphyrin-Dimers Upon One- and Two-Photon Excitation

Leszek Mateusz Mazur, Thomas J. Roland, Stephanie Leroy-Lhez, Vincent Sol, Marek Samoc, Ifor D.W. Samuel, and Katarzyna Matczyszyn

J. Phys. Chem. B, **Just Accepted Manuscript** • DOI: 10.1021/acs.jpcc.8b12561 • Publication Date (Web): 05 Mar 2019

Downloaded from <http://pubs.acs.org> on March 12, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Efficient Singlet Oxygen Photogeneration by Zinc Porphyrin-Dimers Upon One- and Two-Photon Excitation.

Leszek M. Mazur^a, Thomas Roland^b, Stephanie Leroy-Lhez^c, Vincent Sol^c, Marek

*Samoc^a, Ifor D.W. Samuel^b, Katarzyna Matczyszyn^{*a}*

^a Advanced Materials Engineering and Modelling Group, Faculty of Chemistry, Wrocław

University of Science and Technology, Wybrzeże Wyspińskiego 27, 50-370 Wrocław,

Poland

^b Organic Semiconductor Centre, SUPA, School of Physics and Astronomy, University

of St Andrews, Fife, KY16 9SS, UK

^c Laboratoire PEIRENE EA 7500, Faculté des Sciences et Techniques 123, Avenue

Albert Thomas, Limoges cedex 87060, France

ABSTRACT

The development of photodynamic therapy at depth requires photosensitizers which have both sufficient quantum yield for singlet oxygen generation and strong two-photon absorption. Here we show that this can be achieved by conjugated linkage of zinc porphyrins to make dimers. We determined the quantum yield of generation of $^1\text{O}_2$, ϕ_Δ , by measuring emission at 1270 nm using a near infra-red streak camera and found it to increase from 15% for a single porphyrin unit to 27-47% for the dimers with a conjugated linker. Then, we measured the spectra of two-photon absorption cross section, σ_2 , by a focus-tunable Z-scan method, which allows for nondestructive investigation of light-sensitive materials. We observed a strong enhancement of the two-photon absorption coefficient in the dimers, especially those with an alkyne linker. These results lead to an excellent figure of merit for two-photon production of singlet oxygen (expressed by the product $\sigma_2 \times \phi_\Delta$) in the porphyrin dimers, of around 3700 GM, which is very promising for applications involving treatment of deep tumors by photodynamic therapy.

1. INTRODUCTION

1
2
3
4 Photodynamic therapy (PDT) is a widely killing *Propionibacterium acnes*),
5
6
7 used cancer treatment, particularly for the periodontology⁶, treatment of viral lesions⁷
8
9
10 treatment of superficial cancers such as and age-related macular degeneration⁸.
11
12
13 many skin cancers¹. PDT involves the use In order to efficiently photogenerate
14
15
16 of light in combination with a singlet oxygen, a material should fulfill a
17
18
19 photosensitizer to cause a photochemical few criteria. It needs to absorb excitation
20
21
22 reaction, leading to the generation of light and have a triplet energy level higher
23
24
25 reactive oxygen species (ROS)², which than 397 kJ/mol to allow for efficient
26
27
28 then kill neighboring tumor cells. Whilst energy transfer to ground state oxygen
29
30
31 most often used to kill cancer cells, the and hence the generation of singlet
32
33
34 reactive oxygen species can also be used oxygen. Moreover, it should have high
35
36
37 to kill bacteria thereby providing a triplet quantum yield and long triplet state
38
39
40
41
42 powerful alternative to antibiotics in the lifetime². Despite these requirements,
43
44
45 management of topical infections^{3, 4}. For there are many materials capable of
46
47
48 example, in addition to cancers, PDT has photogenerating singlet oxygen, they can
49
50
51
52
53 been used in the treatment of acne⁵ (by be grouped into: (1) the organic dyes and
54
55
56
57
58
59
60

1
2
3 aromatics⁹; (2) the porphyrins, chlorins, wavelengths in the biological
4
5
6 phthalocyanines, and related transparency windows of tissue at
7
8
9
10 macrocycles¹⁰; (3) semiconductors¹¹; and 650-950 nm and 1000-1250 nm¹⁴. As only
11
12
13
14 (4) metal nanoparticles^{12, 13}. Among these few photosensitizers have a linear
15
16
17 photosensitizers, porphyrin derivatives absorption band in this wavelength
18
19
20 attracted significant attention and some of region¹⁵, two-photon excitation in which
21
22
23
24 them are already in clinical use. two longer wavelength photons are
25
26
27

28 Photodynamic therapy has been widely simultaneously absorbed is a natural way
29
30
31 used to kill tumor cells, which are located to achieve the desired excitation energy
32
33
34 close to skin surface, due to the fact that with longer wavelength light that can
35
36
37 the photosensitizers are excited with penetrate further into tissue. The
38
39
40 wavelengths which are strongly absorbed photosensitizers which aim to be used in
41
42
43 and scattered by the tissue. PDT could two-photon excited PDT should be
44
45
46 treat a much wider range of conditions if characterized by high quantum yield of
47
48
49 could work at greater depth which could singlet oxygen generation, ϕ_{Δ} , and large
50
51
52 be achieved by using excitation two-photon absorption cross-sections.
53
54
55
56
57
58
59
60

1
2
3
4 Typical porphyrin derivatives¹⁶⁻¹⁸, even consider this figure of merit when
5
6
7 those accepted for clinical use like normalized by molecular weight i.e.
8
9
10 Photofrin[®], Verteporfin[®] or Foscan[®] are $\sigma_2 \times \phi_{\Delta} / M$, marked as $(\sigma_2 \times \phi_{\Delta})^*$ to facilitate
11
12
13 indeed efficient singlet oxygen generators comparison of large and small candidate
14
15
16 but suffer from low two-photon absorption photosensitizers. In the case of
17
18
19 cross-sections, on the order of 1 GM^{19, 20}. porphyrins, in order to maximize the
20
21
22 In that context, there is a need to design parameter $\sigma_2 \times \phi_{\Delta}$, insertion of a metal ion
23
24
25 and synthesize new molecules with leads to a trade-off as it causes reduction
26
27
28 excellent figure of merit for two-photon of the singlet oxygen quantum yield²¹ and
29
30
31 production of singlet oxygen (expressed increase of the 2PA cross-section.
32
33
34 by the product $\sigma_2 \times \phi_{\Delta}$). It is also relevant to
35
36
37
38
39
40
41

42 In the field of nonlinear optics, the been shown that centrosymmetric
43
44
45 general rules for molecular design to quadrupolar compounds have larger
46
47
48 maximize 2PA cross-section exist in the values of σ_2 than their dipolar
49
50
51 literature and are supported by both counterparts²². In addition, introducing
52
53
54 theoretical and experimental work. It has strong electron donating (D) or electron
55
56
57
58
59
60

1
2
3 withdrawing (A) groups to yield the
4
5
6 structures like D-A-D or A-D-A further
7
8
9 improves 2PA properties²³. Finally, the
10
11 process of simultaneous absorption of two
12
13 photons requires strong overlap of the
14
15
16 π -orbitals over a large system, making it
17
18 sensitive to molecular conformation
19
20 changes, which may alter the electronic
21
22 coupling.


23
24
25
26
27
28
29
30
31
32
33
34
35 The general structure-property rules can
36
37 be applied to porphyrin based
38
39 compounds. It has been shown that 2PA
40
41
42 properties may be tuned by metal
43
44
45 insertion, adding electron donating or
46
47
48 withdrawing groups and expanding the
49
50
51 π -conjugated system by creating dimers


or higher order oligomers^{17, 23, 24}. In the
present work we use metal containing
porphyrins and explore the effect of
increasing electron delocalization by
conjugated links between macrocycles. In
particular, we explore the photogeneration
of singlet oxygen and two-photon
absorption in a series of zinc porphyrins,
either being a monomer (compound **1**), or
a dimer, linked by ethyne (**2**) or di-ethyne
link (**3**) or by a single bond (**4**), as depicted
in Scheme 1.

There are numerous papers
investigating either the two-photon
absorption or singlet oxygen
photogeneration properties of porphyrin

1
2
3 based compounds, but not so many where
4
5
6
7 the combination of these two effects is
8
9
10 presented²⁵⁻²⁷. Our goal was to carry out
11
12
13 interdisciplinary research covering the
14
15
16 nonlinear optical properties and singlet
17
18 oxygen photogeneration, thus providing
19
20
21 strong support to the results and
22
23
24 conclusions concerning the structure-
25
26
27 property optimization. Singlet oxygen
28
29
30 generation was studied using an infra-red
31
32
33
34
35
36
37

streak camera. The two-photon absorption
properties were investigated by the focus-
tunable Z-scan method, which has the
important advantage over alternative
methods of being quick, thereby avoiding
photodegradation which is a particular
consideration in the measurement of
materials intended to cause
photochemistry.





Scheme 1. Structures of investigated compounds.

2. EXPERIMENTAL SECTION

The molecules used in this work were synthesized and purified according to the protocols described in²⁸. The molecules were dissolved in chloroform (Sigma-Aldrich) and quartz cuvettes were filled with the solutions for the photophysical measurements. UV-vis absorption spectra were measured using a Perkin Elmer Lambda 1050 UV/Vis/NIR

spectrophotometer. The concentration of the solution was adjusted to reach optical densities suitable for the PL and emission from singlet oxygen experiments. Steady-state PL spectra were recorded using an automated spectrofluorometer (Fluorolog, Horiba Jobin-Yvon), with a 450 W Xenon lamp excitation source. The solution singlet oxygen quantum yield (SOQY) (ϕ_{Δ})

values were determined using PHAROS regenerative amplifier (output tetraphenylporphyrin (H₂TPP) from Fisher Scientific in chloroform solution as a standard reference because of its well-known SOQY of 55%²⁹. The direct emission of singlet oxygen at 1270 nm was detected by a NIR streak camera from Hamamatsu Photonics. The camera was working in Photon Counting mode and the experiment was automatically stopped after 50 000 frames for each sample. In order to prevent the degradation, the samples solutions were placed in a rotating cell, placed in such a way that the exciting beam was shifted from the center.

The beam from a Light Conversion

PHAROS regenerative amplifier (output wavelength 1028 nm, pulse duration 290 fs, repetition rate 5 kHz) was directed to an optical parametric amplifier (OPA, a Light Conversion OPHEUS-F). The OPA generated 828 nm, which was further frequency doubled to obtain 414 nm used to excite the samples. The output pulse duration was less than 100 fs. The quantum yield of singlet oxygen generation may be determined using the modified method used by Tanielian and Heinrich³⁰ and using equation 1²¹:

$$\phi_{\Delta x} = \phi_{\Delta r} \frac{A_r(\lambda_r) I(\lambda_r) D_x \tau_r}{A_x(\lambda_x) I(\lambda_x) D_r \tau_s} \quad (1)$$

1
2
3 where ϕ_{Δ} stands for the quantum yield of
4
5
6
7 singlet oxygen generation, the subscripts
8
9
10 x and r refer to the compound to be
11
12
13 characterized and to the reference
14
15
16 solutions, respectively. λ is the excitation
17
18 wavelength, $A(\lambda)$ is the absorbance, $I(\lambda)$ is
19
20
21 the relative intensity of the excitation light,
22
23
24
25
26
27 τ is the lifetime of singlet oxygen emission
28
29
30
31 in the sample and reference solvent, and
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
 D is the integrated area under the
emission spectra.

As, the samples and reference solutions
are excited at the same conditions and
they are dissolved in chloroform, the
above equation can be simplified to:

$$\phi_{\Delta x} = \phi_{\Delta r} \frac{A_r(\lambda_r) D_x}{A_x(\lambda_x) D_r} \quad (2)$$

The two-photon absorption cross-sections were determined by f-scan method described in³¹. The laser system used for this study was a Quantronix Integra Ti:Sapphire regenerative amplifier (output wavelength: 800 nm; pulse duration: 130 fs; repetition rate: 1 kHz) that pumped an optical parametric amplifier (OPA, a Quantronix Palitra) to provide tunable excitation. The sample solution was prepared by dissolving the corresponding chromophore in chloroform at concentration of 0.5% (w/w) and placed in a glass cell with 1 mm optical path-

length. The measurements were carried out in a relative manner, calibrating all the data against f-scans carried out on a fused silica plate and taking into account the nonlinear signals obtained from a cell containing pure solvent.

3. RESULTS AND DISCUSSION

Steady-state electronic absorption.

The absorption and photoluminescence spectra of the investigated compounds were measured in dilute CHCl_3 solutions with 1×10^{-6} mol/L concentration. Maximum absorption wavelengths and maximum emission wavelengths are listed in Table 1. The absorption spectra are displayed in Figure 1. For the monomeric





porphyrin, compound **1**, (Figure 1a), two absorption bands are clearly visible. The Soret band is located at 414 nm and so called Q-band at 543 nm. For the dimers, the Q-bands are intensified they show bathochromic shift compared to molecule **1**. Moreover, both Soret and Q-bands are much broader with additional peaks visible in the absorption spectra for compounds **2-4**. It is worth to mention that the stronger Q-band absorption is responsible for enhanced 2PA properties in the NIR. Fluorescence occurs from the Q-band, whichever band is excited. The fluorescence bands of compounds **2** and **3** show a bathochromic shift with respect

1
2
3 to compound **1** (Figure 1a-c). The as $B_x(0-0)$ transition. There are also other
4 bathochromic shift of the absorption and B_x and B_y transitions which span a wide
5
6 fluorescence bands can be explained by region to the blue from the $B_x(0-0)$
7 the decreased HOMO-LUMO gap due to transition. In the case of compound **4** the
8 π -delocalization between the neighboring molecule is no longer planar due to steric
9 hindrance, and so the dimerization does
10 not lead to increased electron
11 delocalization. There is an almost equally
12 intense split in the Soret absorption and Q-
13 band emission of **4** (Figure 1d), which can
14 be explained by the excitonic coupling
15 between the two parallel strong dipole
16 transitions of each porphyrin ring^{28, 32}. The
17 magnitude of the absorption band split is
18 equal to 2002 cm^{-1} and the emission band
19 split is equal to 854 cm^{-1} .
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The crystallographic structure analysis performed by another group yielded the torsional angle of the porphyrin dimer linked by a single bond to be $69.1(2)^{\circ}$ ³³.

The torsional angle of the dimers linked by ethyne and di-ethyne bonds were

determined by DFT calculations to be zero, which means that the molecules are planar³⁴, which enhances both 2PA and singlet oxygen photogeneration due to faster intersystem crossing (S_1-T_1), compared to twisted conformers³⁵.



1
2
3
4 **Figure 1.** Normalized absorption (black curves) and steady-state fluorescence spectra
5
6 (red curves) of investigated porphyrins. Figures a) to d) correspond to molecules **1** to **4**,
7
8 respectively.
9
10
11

12 13 14 15 **Singlet oxygen detection.**

16
17
18 The singlet oxygen quantum yield, ϕ_{Δ} , emission. The spectra obtained for sample
19
20 was determined by recording direct **4** and the reference are shown in Figure 2.
21
22 emission from singlet oxygen centered at Finally, the spectra were integrated to
23
24 1270 nm by a NIR streak camera. The obtain the D factors necessary to calculate
25
26 transition from first excited singlet state the quantum yield of singlet oxygen
27
28 $^1\Delta_g$ to ground state being triplet, $^3\Sigma_g^-$, is photogeneration. The values of ϕ_{Δ} of all
29
30 spin forbidden². This is the reason for the investigated compounds are summarized
31
32 emission lifetime on the order of μ s, which in Table 1. The least efficient is the
33
34 strongly depends on the solvent²¹. monomeric unit (**1**). The dimerization
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 The signals were integrated from 8 to leads to enhanced properties of singlet
50
51 50 μ s to obtain the spectra of $^1\text{O}_2$ oxygen photogeneration, regardless the
52
53
54
55
56
57
58
59
60

1
2
3 type of the link. One can notice that the
4
5
6
7 value of ϕ_{Δ} for compound **4**, which is not
8
9
10 conjugated due to steric hindrance, is
11
12 approximately doubled, compared to **1**.
13
14
15 On the other hand, compounds **2** and **3**,
16
17
18 which have large π -delocalization and are
19
20
21 close to planar, are characterized with
22
23
24 much larger ϕ_{Δ} being equal to 35 and
25
26
27 47 %. These findings are in good
28
29
30 agreement with other works on porphyrins
31
32
33 dimers, e.g. with porphyrin units linked by
34
35
36 butadiyne bond linked to TIPS-protected
37
38
39 ethynyl groups³⁶. The determined ϕ_{Δ} of
40
41
42 such porphyrin dimer in DCM was
43
44
45 reported to be 37 %, which means that the
46
47
48
49 structural design of compounds **2** and **3**
50
51
52
53
54
55
56
57
58
59
60

are more efficient in terms of singlet oxygen production.




Figure 2. Singlet oxygen emission spectra of **4** and H₂TPP, obtained by integrating the streak camera images from 8 to 50 μ s.

Table 1. Steady-state photophysical properties of the investigated compounds (**1-4**) and their singlet oxygen quantum yields. The positions of the most intensive bands are given in bold.

porphyrins	λ_{abs} (nm)	λ_{em} (nm)	ϕ_{Δ} (%)
-------------------	-----------------------------	----------------------------	---------------------

	Soret band	Q-band		
1	393 414	542	554 585 637 700	15
2	420 441 450 482	566 622 673	627 681 751	35

Two-photon absorption cross-section spectra.


The spectra of two-photon absorption cross-section may be obtained by many methods, two-photon excited fluorescence (TPEF)³⁷ and fs Z-scan³⁸⁻⁴¹ being the most common nowadays. They strongly differ from each other, as in TPEF one detects fluorescence spectrum at various wavelengths and compares the integrated PL with a standard reference with known PLQY and σ_2 . In Z-scan one

3	414 431 444 481	562 681	599 654 706 785	47
4	417 455	538 558 600	567 624 658	27

measures the changes of sample transmission as a function of the sample position in Z direction and fit the data recorded on so called Open Aperture curve as described in the literature^{38, 41} to determine the value of two-photon absorption cross-section at the specific wavelength. In order to obtain reliable data by these two aforementioned techniques, the samples have to be perfectly photostable.

1
2
3
4 Upon photoexcitation with one- and the f-scan technique, instead of translating
5
6
7 two-photons the singlet oxygen which is the sample, the focal point is shifted by an
8
9
10 generated by the investigated porphyrins electrically controlled lens³¹. Combined
11
12
13
14 causes their degradation. It is important to with improved data acquisition technique,
15
16
17 mention that each photosensitizer this method reduces the time necessary
18
19
20
21 molecule can typically produce 10^3 - 10^5 for a single scan from a few minutes to
22
23
24 molecules of $^1\text{O}_2$ before being degraded several seconds, thus limiting the
25
26
27
28 through photobleaching by $^1\text{O}_2$ or by some possibility of photodegradation. While the
29
30
31 other process². Thus, it is crucial to vast majority of published papers
32
33
34
35 minimize the time of irradiation especially described only the singlet oxygen
36
37
38
39 with high power laser used to study the photogeneration process, or two-photon
40
41
42 nonlinear effects. Hence we use the so- absorption properties of the studied
43
44
45 called “f-scan” technique which is a photosensitizers, in this work both are
46
47
48
49 modification of the standard Z-scan treated with equal attention and detail,
50
51
52 method, to determine the spectra of allowing for more reliable conclusions.
53
54
55
56 two-photon absorption cross-section. In

The spectra of two-photon absorption cross-sections of investigated porphyrins obtained by f-scan technique are shown in Figure 3. They are overlaid with linear absorption spectra, shown as red curves, plotted vs. twice the wavelength. For all porphyrins, both the monomer (1) and dimers (2-4) the maximum of two-photon absorption cross-sections is within first biological window which exists at wavelengths between 650 nm and 950 nm¹⁴.



1
2
3 **Figure 3.** Spectra of two-photon absorption cross-sections of the investigated porphyrins
4
5
6
7 (1-4). The spectra were overlaid with linear absorption (red curves) plotted vs. twice the
8
9
10 wavelength. Figures a) to d) correspond to compounds 1 to 4.

16 In case of the monomer 1, the maximum means that it is favorable for the
17 value of σ_2 is around 1000 GM at 725 nm, molecules to be excited to higher vibronic
18 as shown in Figure 3. a). One can see that state upon 2PA excitation, compared to
19 the band of 2PA is split into two, even 1PA process. Despite the fact that linear
20 though the linear absorption shows only absorption of 4 is split into two equally
21 one peak at 414 nm. It is caused by intense bands, the 2PA spectrum is
22 different selection rules for one- and different. One can see that the band at
23 two-photon absorption, as described in the 725 nm is much more intensive than that at
24 literature²². It is worth noticing that the high 875 nm. The maximum value of σ_2 ,
25 energy 2PA band at 725 nm is observed measured at 725 nm is approximately
26 for all the compounds and has the highest doubled compared to those of compound
27 value of σ_2 among other 2PA bands. It 1 and is equal to 2500 GM. In the studies

1
2
3 of two-photon absorption cross-sections but its position remains the same, the
4
5
6 various normalization factors are used, wavelength of maximum σ_2 is equal to
7
8
9 among which the normalization to molar 725 nm. However, the low energy band,
10
11 mass is the most popular. If the maximum which is located at 875 nm for other
12
13 σ_2 is divided by molar mass, a similar investigated compounds, is
14
15 value is obtained, which means that such bathochromically shifted to 1050 nm.
16
17
18 structural change does not impose neither Similarly to compound 1, there is a dip in
19
20 cooperation nor deleterious effects on the the two-photon absorption cross-section
21
22 multiphoton absorption properties of the spectra at twice the wavelength of
23
24 investigated compounds 1 and 4. In case maximum 1PA band. The 2PA spectrum
25
26 of compounds 2 and 3, the spectra of of compound 3 is presented in Figure 3. c).
27
28 two-photon absorption cross-section show The higher energy band is
29
30 more complex structures, similarly to their bathochromically shifted by 50 nm to
31
32 1PA spectra. For compound 2 775 nm and the value of σ_2 is around 8000
33
34 (Figure 3. b), the higher energy 2PA band GM being slightly lower than for
35
36 is broader than for compounds 1 and 4, compound 2. For both compounds 2 and
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3, the maximum value of two-photon absorption cross-section is enhanced few times, while keeping the molar mass similar to compound **4**. The results prove that a simple modification of the linker, from single bond (**4**) to ethyne (**2**) and di-ethyne (**3**) links between the zinc porphyrins cores cause cooperation effect, enhancing the σ_{2^*} figure of merit.

The values of maximum two-photon absorption cross-section and the figure of merit are listed in Table 2. The dimers of zinc porphyrins, especially connected by ethyne and di-ethyne link, are much more effective two-photon photosensitizers than tetraphenyl porphyrin, or other porphyrin derivatives that are commercially used for PDT.

Table 2. Two-photon absorption cross-section, merit factors and two-photon excited oxygen sensitization properties of the investigated porphyrins (**1-4**).

	λ_{2PA}^{max} (nm)	σ_2^{max} (GM)	σ_2^{max*} (GM·mol/g)	$\sigma_2^{max} \cdot \phi_{\Delta}$ (GM)	$\sigma_2^{max} \cdot \phi_{\Delta}^*$ (GM·mol/g)	Two-photon excited oxygen sensitization enhancement factor ^[c]
1	725	1030	1.4	155	0.21	21
2	725	8900	6.1	3115	2.15	433
3	775	7950	5.4	3737	2.53	519
4	725	2500	1.7	675	0.45	94
H ₂ TPP	790	12 ^[a]	0.02	7.2 ^[b]	0.01	1

^[a] taken from reference ⁴²

1
2
3 [b] taken from reference ²⁷
4

5 [c] Enhancement factor: $\sigma_2^{max} \cdot \phi_{\Delta}$ of the compound normalized to that of H₂TPP.
6
7

8 It was previously shown e.g. by TPEF conditions in the three-level system and
9 measurements¹⁷, that dimerization of dramatic enhancement of the excited-
10 porphyrin units leads to enhanced 2PA state singlet-singlet transition due to linear
11 cross-sections and enhanced two-photon elongation of the π -conjugated system.
12 production of singlet oxygen. The σ_2 Insertion of side groups, being either
13 determined for the porphyrin dimer linked electron acceptors (A) or donors (D)
14 by ethyne and di-ethyne bonds were on influence the σ_2 in a way that two-photon
15 the order of 8200 and 5500 GM, which is absorption cross-section is the lowest for
16 similar to the values found in the current neutral porphyrins, higher for the
17 work for similar porphyrin derivatives. The porphyrins extended with A and the
18 cooperative effect was explained by a highest for compounds with D groups⁴³.
19 combination of several factors, including Even though the vast majority of
20 strong enhancement of the lowest one- molecules targeted to be used in 2PA PDT
21 photon Q-transition, better resonance are based on the porphyrin structure,
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 there are some alternative approaches
5
6
7 with reasonable figures of merit. One
8
9
10 example is a macromolecular
11
12
13 photosensitizer, which was obtained via
14
15
16 the covalent anchoring of several
17
18
19 molecular PS based on a
20
21
22 dibromobenzene (DBB) core onto a water-
23
24
25 soluble polymer chain based on
26
27
28 poly(N-acryloylmorpholine). It resulted in
29
30
31 large 2PA cross-section up to 2600 GM at
32
33
34
35 740 nm and was proved to be an efficient
36
37
38
39 candidate to cause melanoma cells death
40
41
42 upon photo irradiation⁴⁴.
43
44
45
46
47
48

49 4. CONCLUSIONS

50
51
52 In this work, we investigated two-photon
53
54
55 absorption and singlet oxygen
56
57
58
59
60

photogeneration properties in a series of
zinc porphyrins, either in a monomer or
dimer form. The dimers were connected
by single bond, or by ethyne, or di-ethyne
link. We observed strong enhancement of
the obtained quantum yields for
two-photon excited singlet oxygen
generation (expressed by the product
 $\sigma_2^{max} \phi_{\Delta}$) in the porphyrin dimer
connected by di-ethyne link, around
3700 GM, which is around 500 times more
efficient than tetraphenyl porphyrin. It can
be seen that in case of zinc porphyrin
derivatives, structure optimization towards
high two-photon absorption cross-section
simultaneously enhances the properties of

1
2
3
4 singlet oxygen photogeneration. Thus, in
5
6
7 order to obtain a material capable of
8
9
10 efficient production of singlet oxygen upon
11
12
13 two-photon excitation, one should
14
15
16
17 increase the conjugation length, so that
18
19
20
21 π -delocalization is maximized, while
22
23
24 maintaining planar structure, as shown for
25
26
27
28 compounds **2** and **3**. The results are very
29
30
31 promising for applications involving
32
33
34
35 treatment of deep tumors by
36
37
38 photodynamic therapy.
39
40
41

42 AUTHOR INFORMATION

43 Corresponding Author

44
45
46 katarzyna.matczyszyn@pwr.edu.pl;
47
48 idws@st-andrews.ac.uk
49

50 ORCID

51
52
53 Leszek M. Mazur: 0000-0002-4562-0743
54
55
56
57
58
59
60

Vincent Sol: 0000-0003-0175-0156

Marek Samoc: 0000-0002-5404-2455

Ifor D.W. Samuel: 0000-0001-7821-7208

Katarzyna Matczyszyn: 0000-0001-8578-8340

Conflict of interest statement

There are no conflicts to declare.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the support from National Science Centre (Poland) under Opus program DEC-2013/09/B/ST5/03417, the EPSRC of the UK through grant EP/L017008/1 and the Conseil Régional du Limousin in France for financial support. The work was financed by a statutory activity subsidy

1
2
3 from the Polish Ministry of Science and
4
5
6
7 Higher Education for the Faculty of
8
9
10 Chemistry of Wrocław University of
11
12
13 Science and Technology. IDWS
14
15
16 acknowledges a Royal Society Wolfson
17
18
19 Research Merit Award. We thank
20
21
22 the European Union as this work is a part
23
24
25 of the ITN-EJD-2017 project POLYTHEA
26
27
28 which has received funding from the
29
30
31 European Union's Horizon 2020 research
32
33
34 and innovation programme under the
35
36
37 Marie Skłodowska-Curie grant agreement
38
39
40 n°764837.
41
42
43
44
45
46
47
48
49

50 REFERENCES:

51
52 1. Brown, S. B.; Brown, E. A.; Walker, I.,
53 The Present and Future Role of Photodynamic
54 Therapy in Cancer Treatment. *Lancet Oncol.* **2004**,
55 *5*, 497-508.
56
57
58
59
60

2. DeRosa, M. C.; Crutchley, R. J.,
Photosensitized Singlet Oxygen and Its
Applications. *Coord. Chem. Rev.* **2002**, *233–234*,
351-371.
3. Yin, R.; Agrawal, T.; Khan, U.; Gupta, G.
K.; Rai, V.; Huang, Y.-Y.; Hamblin, M. R.,
Antimicrobial Photodynamic Inactivation in
Nanomedicine: Small Light Strides against Bad
Bugs. *Nanomedicine* **2015**, *10*, 2379-2404.
4. Tardivo, J. P.; Adami, F.; Correa, J. A.;
Pinhal, M. A. S.; Baptista, M. S., A Clinical Trial
Testing the Efficacy of PDT in Preventing
Amputation in Diabetic Patients. *Photodiagn.*
Photodyn. Ther. **2014**, *11*, 342-350.
5. Bissonnette, R., Treatment of Acne with
Photodynamic Therapy. *G. Ital. Dermatol.*
Venerol. **2011**, *146*, 445-456.
6. Kumar, V.; Sinha, J.; Verma, N.; Nayan,
K.; Saimbi, C.; Tripathi, A., Scope of
Photodynamic Therapy in Periodontics. *Indian J.*
Dent. Res. **2015**, *26*, 439-442.
7. Kharkwal, G. B.; Sharma, S. K.; Huang,
Y.-Y.; Dai, T.; Hamblin, M. R., Photodynamic
Therapy for Infections: Clinical Applications.
Lasers Surg. Med. **2011**, *43*, 755-767.
8. Boscia, F.; Furino, C.; Sborgia, L.;
Reibaldi, M.; Sborgia, C., Photodynamic Therapy
for Retinal Angiomatous Proliferations and
Pigment Epithelium Detachment. *Am. J.*
Ophthalmol. **2004**, *138*, 1077-1079.
9. Redmond, R. W.; Gamlin, J. N., A
Compilation of Singlet Oxygen Yields from
Biologically Relevant Molecules. *Photochem.*
Photobiol. **1999**, *70*, 391-475.
10. Murtinho, D.; Pineiro, M.; Pereira, M. M.;
Rocha Gonsalves, A. M. d. A.; Arnaut, L. G.;
Miguel, M. d. G.; Burrows, H. D., Novel
Porphyrins and a Chlorin as Efficient Singlet
Oxygen Photosensitizers for Photooxidation of
Naphthols or Phenols to Quinones. *J. Chem. Soc.,*
Perkin Trans. 2 **2000**, 2441-2447.
11. Yamamoto, Y.; Imai, N.; Mashima, R.;
Konaka, R.; Inoue, M.; Dunlap, W. C., Singlet
Oxygen from Irradiated Titanium Dioxide and
Zinc Oxide. *Methods Enzymol* **2000**, *319*, 29-37.
12. Chen, C.-W.; Hsu, C.-Y.; Lai, S.-M.; Syu,
W.-J.; Wang, T.-Y.; Lai, P.-S., Metal Nanobullets
for Multidrug Resistant Bacteria and Biofilms.
Adv. Drug Del. Rev. **2014**, *78*, 88-104.
13. Huang, C.-C.; Liu, T.-M., Controlled Au-
Polymer Nanostructures for Multiphoton Imaging,

- Prodrug Delivery, and Chemo-Photothermal Therapy Platforms. *ACS Appl. Mater. Interfaces* **2015**, *7*, 25259-25269.
14. Smith, A. M.; Mancini, M. C.; Nie, S., Bioimaging: Second Window for in vivo Imaging. *Nat. Nanotechnol.* **2009**, *4*, 710-711.
15. Habermeyer, B.; Guillard, R., Some Activities of Porphyrin Illustrated by the Applications of Porphyrinoids in PDT, PIT and PDI. *Photochem. Photobiol. Sci.* **2018**, *17*, 1675-1690.
16. Goyan, R. L.; Cramb, D. T., Near-Infrared Two-Photon Excitation of Protoporphyrin IX: Photodynamics and Photoproduct Generation. *Photochem. Photobiol.* **2000**, *72*, 821-827.
17. Drobizhev, M.; Stepanenko, Y.; Dzenis, Y.; Karotki, A.; Rebane, A.; Taylor, P. N.; Anderson, H. L., Extremely Strong near-IR Two-Photon Absorption in Conjugated Porphyrin Dimers: Quantitative Description with Three-Essential-States Model. *J. Phys. Chem. B* **2005**, *109*, 7223-7236.
18. Nowak-Król, A.; Wilson, C. J.; Drobizhev, M.; Kondratuk, D. V.; Rebane, A.; Anderson, H. L.; Gryko, D. T., Amplified Two-Photon Absorption in Trans-A2B2-Porphyrins Bearing Nitrophenylethynyl Substituents. *ChemPhysChem* **2012**, *13*, 3966-3972.
19. Balaz, M.; Collins, H. A.; Dahlstedt, E.; Anderson, H. L., Synthesis of Hydrophilic Conjugated Porphyrin Dimers for One-Photon and Two-Photon Photodynamic Therapy at NIR Wavelengths. *Org. Biomol. Chem.* **2009**, *7*, 874-888.
20. Karotki, A.; Khurana, M.; Lepock, J. R.; Wilson, B. C., Simultaneous Two-Photon Excitation of Photofrin in Relation to Photodynamic Therapy. *Photochem. Photobiol.* **2006**, *82*, 443-452.
21. Mathai, S.; Smith, T. A.; Ghiggino, K. P., Singlet Oxygen Quantum Yields of Potential Porphyrin-Based Photosensitizers for Photodynamic Therapy. *Photochem. Photobiol. Sci.* **2007**, *6*, 995-1002.
22. Pawlicki, M.; Collins, H. A.; Denning, R. G.; Anderson, H. L., Two-Photon Absorption and the Design of Two-Photon Dyes. *Angew. Chem. Int. Ed.* **2009**, *48*, 3244-3266.
23. Wilkinson, J. D.; Wicks, G.; Nowak-Król, A.; Lukasiewicz, L. G.; Wilson, C. J.; Drobizhev, M.; Rebane, A.; Gryko, D. T.; Anderson, H. L., Two-Photon Absorption in Butadiyne-Linked Porphyrin Dimers: Torsional and Substituent Effects. *J. Mater. Chem. C* **2014**, *2*, 6802-6809.
24. Kim, K. S.; Lim, J. M.; Osuka, A.; Kim, D., Various Strategies for Highly-Efficient Two-Photon Absorption in Porphyrin Arrays. *J. Photochem. Photobiol., C* **2008**, *9*, 13-28.
25. Yao, D.; Hugues, V.; Blanchard-Desce, M.; Mongin, O.; Paul-Roth, C. O.; Paul, F., Dendritic Molecular Assemblies for Singlet Oxygen Generation: Meso-Tetraphenylporphyrin-Based Biphotonic Sensitizers with Remarkable Luminescence. *New J. Chem.* **2015**, *39*, 7730-7733.
26. McIlroy, S. P.; Cló, E.; Nikolajsen, L.; Frederiksen, P. K.; Nielsen, C. B.; Mikkelsen, K. V.; Gothelf, K. V.; Ogilby, P. R., Two-Photon Photosensitized Production of Singlet Oxygen: Sensitizers with Phenylene-Ethynylene-Based Chromophores. *J. Org. Chem.* **2005**, *70*, 1134-1146.
27. Yao, D.; Zhang, X.; Triadon, A.; Richy, N.; Mongin, O.; Blanchard-Desce, M.; Paul, F.; Paul-Roth, C. O., New Conjugated Meso-Tetrafluorenylporphyrin-Cored Derivatives as Fluorescent Two-Photon Photosensitizers for Singlet Oxygen Generation. *Chem. - Eur. J.* **2017**, *23*, 2635-2647.
28. Jiblaoui, A.; Baudequin, C.; Chaleix, V.; Ducourthial, G.; Louradour, F.; Ramondenc, Y.; Sol, V.; Leroy-Lhez, S., An Easy One-Pot Desilylation/Copper-Free Sonogashira Cross-Coupling Reaction Assisted by Tetra-Butylammonium Fluoride (TBAF): Synthesis of Highly π -Conjugated Porphyrins. *Tetrahedron* **2013**, *69*, 5098-5103.
29. Poon, C. T.; Zhao, S. S.; Wong, W. K.; Kwong, D. W. J., Synthesis, Excitation Energy Transfer and Singlet Oxygen Photogeneration of Covalently Linked N-Confused Porphyrin-Porphyrin and Zn(II) Porphyrin Dyads. *Tetrahedron Lett.* **2010**, *51*, 664-668.
30. Tanielian, C.; Heinrich, G., Effect of Aggregation on the Hematoporphyrin-Sensitized Production of Singlet Molecular Oxygen. *Photochem. Photobiol.* **1995**, *61*, 131-135.
31. Kolkowski, R.; Samoc, M., Modified Z - Scan Technique Using Focus-Tunable Lens. *J. Opt.* **2014**, *16*, 125202.
32. L. Anderson, H., Building Molecular Wires from the Colours of Life: Conjugated

- 1
2
3 Porphyrin Oligomers. *Chem. Commun.* **1999**,
4 2323-2330.
- 5 33. Senge Mathias, O.; Pinteá, M.; Ryan
6 Aoife, A., Synthesis and Crystal Structure of a
7 Meso-Meso Directly Linked Bisporphyrin. *Z.*
8 *Naturforsch., B: Chem. Sci.* **2011**, *66*, 553-558.
- 9 34. Peeks, M. D.; Neuhaus, P.; Anderson, H.
10 L., Experimental and Computational Evaluation of
11 the Barrier to Torsional Rotation in a Butadiyne-
12 Linked Porphyrin Dimer. *Phys. Chem. Chem.*
13 *Phys.* **2016**, *18*, 5264-5274.
- 14 35. Kuimova, M. K.; Balaz, M.; Anderson, H.
15 L.; Ogilby, P. R., Intramolecular Rotation in a
16 Porphyrin Dimer Controls Singlet Oxygen
17 Production. *J. Am. Chem. Soc.* **2009**, *131*, 7948-
18 7949.
- 19 36. Schmitt, J.; Heitz, V.; Jenni, S.; Sour, A.;
20 Bolze, F.; Ventura, B., π -Extended Porphyrin
21 Dimers as Efficient near-Infrared Emitters and
22 Two-Photon Absorbers. *Supramol. Chem.* **2017**,
23 *29*, 769-775.
- 24 37. Deiana, M.; Mettra, B.; Mazur, L. M.;
25 Andraud, C.; Samoc, M.; Monnereau, C.;
26 Matczyszyn, K., Two-Photon Macromolecular
27 Probe Based on a Quadrupolar Anthracenyl
28 Scaffold for Sensitive Recognition of Serum
29 Proteins under Simulated Physiological
30 Conditions. *ACS Omega* **2017**, *2*, 5715-5725.
- 31 38. Van Stryland, E. W.; Sheik-Bahae, M., Z-
32 Scan Measurements of Optical Nonlinearities. In
33 *Characterization Techniques and Tabulations for*
34 *Organic Nonlinear Materials*, Kuzyk, M. G.; Dirk,
35 C., Eds. Marcel Dekker: 1998; pp 655-692.
- 36 39. Gao, B.; Mazur, L. M.; Morshedi, M.;
37 Barlow, A.; Wang, H.; Quintana, C.; Zhang, C.;
38 Samoc, M.; Cifuentes, M. P.; Humphrey, M. G.,
39 Exceptionally Large Two- and Three-Photon
40 Absorption Cross-Sections by OPV
41 Organometalation. *Chem. Commun.* **2016**, *52*,
42 8301-8304.
- 43 40. Lin, T.-C.; Chien, W.; Mazur, L. M.; Liu,
44 Y.-Y.; Jakubowski, K.; Matczyszyn, K.; Samoc,
45 M.; Amini, R. W., Two- and Three-Photon
46 Absorption Properties of Fan-Shaped Dendrons
47 Derived from 2,3,8-Trifunctionalized
48 Indenoquinoxaline Units: Synthesis and
49 Characterization. *J. Mater. Chem. C* **2017**, *5*, 8219-
50 8232.
- 51 41. Sheik-Bahae, M.; Said, A. A.; Wei, T. H.;
52 Hagan, D. J.; Stryland, E. W. V., Sensitive
53 Measurement of Optical Nonlinearities Using a
54 Single Beam. *IEEE J. Quantum Electron.* **1990**,
55 *26*, 760-769.
- 56 42. Makarov, N. S.; Drobizhev, M.; Rebane,
57 A., Two-Photon Absorption Standards in the 550–
58 1600 nm Excitation Wavelength Range. *Opt.*
59 *Express* **2008**, *16*, 4029-4047.
- 60 43. Koszelewski, D.; Nowak-Krol, A.;
Drobizhev, M.; Wilson, C. J.; Haley, J. E.; Cooper,
T. M.; Romiszewski, J.; Gorecka, E.; Anderson, H.
L.; Rebane, A.; Gryko, D. T., Synthesis and Linear
and Nonlinear Optical Properties of Low-Melting
 π -Extended Porphyrins. *J. Mater. Chem. C* **2013**,
1, 2044-2053.
44. Cepraga, C.; Marotte, S.; Ben Daoud, E.;
Favier, A.; Lanoë, P.-H.; Monnereau, C.; Baldeck,
P.; Andraud, C.; Marvel, J.; Charreyre, M.-T.;
Leverrier, Y., Two-Photon Photosensitizer–
Polymer Conjugates for Combined Cancer Cell
Death Induction and Two-Photon Fluorescence
Imaging: Structure/Photodynamic Therapy
Efficiency Relationship. *Biomacromolecules*
2017, *18*, 4022-4033.

