

# Habitat use of a coastal delphinid population investigated using passive acoustic monitoring

Palmer, K.J.<sup>1</sup>, Brookes, Kate L.<sup>2</sup>, Davies, I.M.<sup>2</sup>, Ewan Edwards<sup>2</sup>, Rendell, L.<sup>1</sup>.

<sup>1</sup> Sea Mammal Research Unit, School of Biology, University of St Andrews, Fife, KY16 9TH, UK

<sup>2</sup> Marine Scotland Science, 375 Victoria Road, Aberdeen, AB11 9DB, UK

## 1 Abstract

- 2 • The population of bottlenose dolphins in eastern Scotland has undergone significant  
3 range expansion since the 1990's, when a special area of conservation was established for  
4 the population.
- 5 • Distribution of this population is well described within areas of its range, where intensive  
6 work has been carried out, such as the inner Moray Firth, St Andrews Bay, and the Tay  
7 estuary area. However, elsewhere in their range, habitat use is less well understood.
- 8 • In this study, a large-scale and long-term passive acoustic array was used to gain a better  
9 understanding of bottlenose dolphin habitat use in eastern Scottish waters,  
10 complementing and augmenting existing visual surveys.
- 11 • Data from the array were analysed using a three-stage approach. First, acoustic  
12 occupancy results were reported; second, temporal trends were modelled; and third, a  
13 spatial-temporal-habitat model of acoustic occupancy was created.
- 14 • Results from the acoustic occupancy are in agreement with visual studies that found areas  
15 near known foraging locations were consistently occupied. Results from the trend

16 analysis were inconclusive. Habitat modelling showed that, throughout their range,  
17 bottlenose dolphins are most likely to be detected closer to shore, and, for a constant  
18 distance to shore, in deeper water.

19

20 **Keywords:** Ocean, coastal, habitat management, Marine Protected Area, protected species,  
21 mammals

22

## 23 **1. Introduction**

24 Bottlenose dolphins (*Tursiops truncatus*) are a cosmopolitan species with populations found in  
25 tropical and temperate waters worldwide (Connor, Wells, Mann, and Read, 2000). Presently, the  
26 International Union for Conservation of Nature lists the species as ‘Least concern’ indicating a  
27 low risk of extinction. However, they are also listed under Appendix 2 of the Convention on the  
28 Conservation of Migratory Species of Wild Animals indicating a need for, or benefit from,  
29 international co-operation on conservation efforts. Off the eastern coast of Scotland, there is a  
30 population of bottlenose dolphins consisting of approximately 200 individuals (Cheney et al.,  
31 2013). The population is protected by a variety of national and international regulations,  
32 including Annexes II and IV of European Union Habitats Directive (92/43/EEC), Wildlife and  
33 Countryside Act (1981), and Joint Nature Conservation Committee UK Post-2010 Biodiversity  
34 Framework. In 2005, as part of these conservation efforts, a Special Area of Conservation (SAC)  
35 was established in the Moray Firth to protect habitat important to this population. The Moray  
36 Firth SAC covers approximately 1500 km<sup>2</sup> extending west from the Beaully Firth, north to  
37 Helmsdale, and east to Lossiemouth (<http://www.jncc.defra.gov.uk>; Figure 1). The management

38 of the SAC has been implemented in such a way that the population is protected throughout its  
39 range, whereby any activity which could have an adverse effect on the integrity of the site (i.e.  
40 the protected features) is subject to a Habitats Regulations Appraisal (Arso Civil et al., 2019).

41 The range of this population extends well outside of the bounds of the SAC, with animals  
42 commonly sighted along more than 200 km of coastal habitat (Paxton, Scott-Hayward,  
43 Mackenzie, Rexstad, & Thomas, 2016). Both within and outside the SAC, animals are known to  
44 aggregate at certain locations, often associated with the mouths of rivers or estuaries (Hastie,  
45 Wilson, & Thompson, 2003, 2006; Hastie, et. al, 2004; Mendes, Turrell, Lütkebohle, &  
46 Thompson, 2002; Pirotta et al., 2014; Sargeant, Mann, Berggren, & Krutzen, 2005; Wilson,  
47 Thompson, & Hammond, 1997). Because of the high encounter rates at these locations, some  
48 have become focal areas for boat-based survey efforts, most notably the inner Moray Firth and  
49 Firth of Tay. In some of these locations, henceforth termed points of aggregation, dolphins are  
50 known to exploit tidal cycles and local bathymetry to maximise foraging efficiency (Hastie et al.,  
51 2004). This is the case at two locations within the inner Moray Firth SAC: Chanonry Point near  
52 the River Ness, and the entrance to the Cromarty Firth (Figure 1). At these locations deep  
53 channels result in higher prey density at low tides and therefore may represent increased foraging  
54 success for marine mammals (Thompson, Pierce, Hislop, Miller, & Diack, 1991). Outside of the  
55 Moray Firth SAC, points of aggregation have been observed around the mouth of the River Dee  
56 (Sini, Canning, Stockin, & Pierce, 2005), the Firth of Tay, and St Andrews Bay (Arso Civil et  
57 al., 2019). While foraging activity has been observed at some of these locations, the underlying  
58 factor(s) resulting in the higher occurrence are less clear for others. For example, dolphins are  
59 commonly sighted in and around St Andrews Bay, which is a shallow water area with a small  
60 estuary (Arso Civil et al., 2019; Quick & Janik, 2012).

61 Despite the large population range, most survey effort has focused on the Moray Firth and  
62 specifically on well-established areas of high usage (Arso Civil, 2014; Arso Civil et al., 2019;  
63 Bailey et al., 2010; Bailey & Thompson, 2006; Hastie et al., 2006; Hastie et al., 2003; Janik &  
64 Thompson, 1996; Pirotta et al., 2014; Thompson, Brookes, & Cordes, 2015; Wilson et al., 1997).  
65 While these areas clearly represent key habitat for this population (Cheney et al., 2013), effective  
66 conservation requires knowledge of habitat use throughout the population's range. Even in  
67 foraging hotspots, bottlenose dolphin sightings are often not predictable (Culloch & Robinson,  
68 2008).

69 Since 2000, there have been a handful of regional scale surveys covering a large portion of the  
70 population's habitat (Cheney et al., 2013). These include a compilation of visual and sightings  
71 data from land and boat-based surveys (Thompson et al., 2011); a series of line transect surveys  
72 between the Firth of Forth and the river Dee (Arso Civil, 2014), as well as some passive acoustic  
73 studies (Cheney et al., 2013) . Together, results from these studies suggest that bottlenose  
74 dolphins use the entirety of the coastal habitat, though less frequently outside of the Moray Firth  
75 SAC than within it, and that animals are more likely to be sighted in waters within a few  
76 kilometres of the shore. However, the relative lack of survey effort in other parts of the  
77 population's range (Paxton et al., 2016) limits understanding of how these areas are used and  
78 their relative importance to the population.

79 This lack of understanding has potential implications for the Habitats Regulations Appraisals  
80 undertaken as part of the licensing of marine activities in the region, including the development  
81 of offshore wind energy. Of particular concern is the lack of data in the regions most likely to  
82 receive noise from wind farm construction activities, along with a lack of understanding of how  
83 far offshore bottlenose dolphins range in these regions. Thompson, Brookes & Cordes (2015)

84 used a combination of fixed passive acoustic, and presence only visual survey data to model  
85 usage of offshore areas by bottlenose dolphins. While they showed that it was unlikely that the  
86 species used areas close to construction activities, the lack of data in the areas of concern  
87 reduced stakeholder confidence in the findings.

88 To address these issues, the East Coast Marine Mammal Acoustic Study (ECoMMAS) (Marine  
89 Scotland Science, 2013) was started in 2013 to improve understanding of bottlenose dolphin use  
90 of the east coast of Scotland, with effort spread more evenly throughout the region, including  
91 data collection further offshore. The study uses fixed passive acoustic monitoring to complement  
92 existing visual surveys in coastal and high-use areas. The data presented here were collected  
93 during the first three years of the study.

94 C-PODs are commercially available echolocation click train detectors widely used for  
95 monitoring cetaceans. The instruments are sold with a proprietary click train detector that  
96 discriminates between ‘noise’ and the echolocation click trains (series of echolocation clicks)  
97 produced by dolphins and porpoises. Over the last decade, studies using these devices have  
98 contributed to our understanding of the behaviour and habitat use of the Moray Firth bottlenose  
99 dolphin population (Graham et al., 2017; Pirotta, Merchant, Thompson, Barton, & Lusseau,  
100 2015; Pirotta et al., 2014).

101 Where multiple species are present however, discriminating between target (e.g. bottlenose  
102 dolphin) and non-target species constitutes a major and ongoing challenge in the field of marine  
103 passive acoustic monitoring. This is especially pertinent for studies using logging devices like C-  
104 PODS that collect few acoustic features from which to classify the detections. To account for  
105 this, users typically either deploy the instruments in habitats where only a single species is

106 expected (Jaramillo-Legorreta et al., 2017) or assume the contribution of non-target species  
107 detections to the analysis is limited (Pirodda et al., 2014; Thompson et al., 2011). Due to the scale  
108 of the ECoMMAS array, neither assumption was applicable in this study. Throughout the survey  
109 area, multiple species have been known to occur (Anderwald et al., 2010; Arso Civil, 2014;  
110 Hammond et al., 2017). There is therefore a need to incorporate both acoustic classifiers and  
111 classifier uncertainty into the analysis (Caillat, 2013).

112 In this research, a heuristic approach was taken to misclassification wherein species uncertainty  
113 is built into the model response. An acoustic classification system (Palmer, Brookes, & Rendell,  
114 2017) was applied to C-POD detections in order to group detections into one of three classes:  
115 broadband, frequency banded or unknown. The broadband category represents click trains  
116 matching bottlenose dolphin and common dolphin (*Delphinus spp.*) click characteristics and the  
117 frequency banded category represents click trains matching white-beaked (*Lagenorhynchus*  
118 *albirostris*) and Risso's (*Grampus griseus*) dolphin click characteristics (Calderan, Wittich,  
119 Harries, Gordon, & Leaper, 2013; Soldevilla et al., 2008). This analysis used the probability that  
120 each echolocation click was broadband as the predictor for bottlenose dolphin presence, thereby  
121 reducing the influence of non-target species on the model results.

122 Monitoring occupancy rates provides baseline data for future studies seeking to understand  
123 changes in distribution over long timescales. In Scottish waters, long-term acoustic studies of  
124 have been used to investigate the spatial and temporal distribution of harbour porpoises and  
125 bottlenose dolphins, as well as model the potential impacts of anthropogenic activities (Brookes,  
126 Bailey, & Thompson, 2013; Harris et al., 2017; Simon et al., 2010; Williamson et al., 2016) . In  
127 these studies, the presence of an acoustic signal characteristic of the animal (e.g. click or whistle)  
128 is used as a proxy for true occupancy (P. Thompson et al., 2011).

129 We expected to find low acoustic occupancy rates and the potential for misclassification was  
130 high, so this research took a three-stage approach. In the first stage, two acoustic occupancy rates  
131 are reported: proportion of acoustically monitored days containing acoustic encounters, and the  
132 proportion of acoustically monitored days containing one or more broadband acoustic  
133 encounters. The proportion of days with echolocation encounters is reported for the first three  
134 years of the ECoMMAS survey.

135 The second stage of the study modelled temporal trends in acoustic occupancy from the first  
136 three years of the ECoMMAS. As with baseline acoustic occupancy rates, identifying patterns in  
137 annual occupancy trends should be of interest to regulators seeking to manage the effects of  
138 offshore activities on dolphin habitat and behaviour.

139 The third stage of the analysis determined whether and to what extent it is possible to produce  
140 spatial-temporal habitat models of broadband acoustic occupancy using ECoMMAS C-POD data  
141 alone. In this portion of the analysis a model containing all available spatial and temporal  
142 covariates was fitted to acoustic detections aggregated from the first three years of the  
143 ECoMMAS study.

144

## 145 **2. Methods**

### 146 **Data Collection**

147 Data in this study were collected by 30 C-POD (version 1) echolocation click detectors  
148 (Chelonia, Ltd, UK). Deployment locations were spread across the region of interest, in ten  
149 groups of three; each group of three radiated out from the coast at approximately 5km intervals

150 to provide data at increasing distance offshore (figure 1). The 30 deployment locations are  
151 identified by the combination of the group name (based on the nearest settlement on land) and  
152 distance from shore (e.g. Cro\_05 for the Cromarty nearshore location).

153 The entire array was deployed each spring and recovered in the fall. Precise deployment and  
154 recovery times depended on ship availability and weather conditions (Table 1). With the  
155 exception of the first deployment in 2015, which was recovered prior to battery exhaustion, all  
156 C-PODS ran continuously until either storage or battery capacity was exhausted.

157

## 158 **Data Quality**

### 159 **Acoustic Data Processing**

160 C-POD data from 2013-2015 were processed with the accompanying KERNO classifier version  
161 2.042 ([www.chelonia.co.uk](http://www.chelonia.co.uk)) for the presence of high or moderate quality “other cetacean” click  
162 trains. The KERNO classifier annotates impulsive detections as narrow-band high frequency  
163 (NBHF) click trains, ‘Other cetacean’ click trains and ‘sonar’. NBHF detections are primarily  
164 produced by porpoises. ‘Other cetacean’ click trains may be indicative of a variety of dolphin  
165 species (Sarnocinska, Tougaard, Johnson, Madsen, & Wahlberg, 2016). After processing for the  
166 presence of ‘other cetacean’ clicks, click trains were grouped into acoustic encounters. Each  
167 acoustic encounter consisted of all high or moderate quality ‘other cetacean’ click trains starting  
168 within 20 minutes of the end of another click train. Acoustic encounters were subsequently  
169 processed with the categorization system described in Palmer et al. (2017). This system  
170 categorises each acoustic encounter into one of the following three categories; ‘broadband’,  
171 ‘frequency banded’, or ‘unknown’. Thus, only acoustic encounters considered by the system to

172 be at least five times more likely to be either broadband or frequency banded were categorised.  
173 Encounters that failed to meet the classification threshold for either taxonomic group were  
174 classified as unknown.

175 To incorporate classifier uncertainty into the analysis, the probability that broadband clicks were  
176 detected ( $P(\textit{Broadband})$ ) was used as the response variable in the acoustic occupancy models  
177 (Palmer et al 2017; supplementary material). Broadband click detection probability was defined  
178 as the probability that broadband clicks were actually present, given the category produced by  
179 the classification system. For days when no acoustic encounters were detected,  $P(\textit{Broadband})$   
180 was set to 0. Days when only broadband acoustic encounters (as determined by the classification  
181 system) were reported,  $P(\textit{Broadband})$  was set to 0.79, reflecting the known error rate as  
182 determined by the classification confusion matrix. Similarly, for days when only frequency-  
183 banded clicks were reported,  $P(\textit{Broadband})$  was 0.08. For days when both broadband and  
184 frequency banded click encounters were reported, complete uncertainty was assumed by setting  
185  $P(\textit{Broadband})$  to 0.5.

186

## 187 **Temporal Covariates**

188 The way temporal covariates were included in the models differed between the modelling stages.  
189 For the second stage temporal models, time of the year was measured as the Julian day (1-365)  
190 and included as a smooth continuous variable. For the third stage spatial-temporal model, there  
191 were insufficient detections to incorporate time as a smoothed variable and thus, season was  
192 included in the model as a three-level factor (Spring, Summer, or Autumn). Spring was defined  
193 as the months between April and May (March data was not available), Summer (June to August)

194 and Autumn (September to November). No data were collected over the winter season. For both  
195 analyses, year was included as a three-level categorical predictor (2013, 2014 or 2015).

## 196 **Spatial Covariates**

197 As with temporal covariates, spatial covariates were included as either continuous or factor  
198 variables. Previous studies have identified the following spatial covariates as potential predictors  
199 for the presence of bottlenose dolphins: distance to nearest point of aggregation (e.g. Cromarty  
200 Firth and River Dee), distance to shore, the gradient of the seabed (henceforth slope), and depth  
201 (Thompson et al., 2015).

202 Distance to the nearest point of aggregation was included as a continuous variable in the spatial-  
203 temporal model. Known points of aggregation have previously been shown to drive spatial and  
204 temporal distribution of animals in this population and, in some areas, have been linked to  
205 foraging (Hastie, Wilson, Wilson, Parsons, & Thompson, 2004). Given the spatial and temporal  
206 scale of this study, estuaries that may represent important habitat for animals either transiting  
207 between the established points of aggregation or contemporaneous with local and/or ephemeral  
208 prey sources were included. Known points of aggregation included the Cromarty Firth, Firth of  
209 Tay, and the rivers Ness and Dee (Cheney et al., 2013; Hastie et al., 2004; Quick et al., 2014). To  
210 the known points of aggregation, the mouths of the rivers Spey, North Esk, and Tweed were  
211 added. River estuaries were selected from the Atlantic Salmon Rivers Database  
212 (<http://www.nasco.int/RiversDatabase.aspx>). Distance to nearest point of aggregation was  
213 reported as a continuous variable and was measured by calculating the distance between each C-  
214 POD and the nearest point of aggregation.

215 Distance to shore was measured as either a three-level factor corresponding to whether each C-  
216 POD was deployed in nearshore (05), midshore (10), or offshore (15) habitat, or as a continuous  
217 predictor. For the spatial-temporal model of acoustic occupancy, distance to shore was reported  
218 as the continuous range between the deployment location and the distance to the nearest 0 m  
219 isobath (Pante & Simon-Bouhet, 2013).

220 Deployment depth (in meters) was recorded from the ship at the time of deployment. Additional  
221 spatial covariate data were obtained from the NOAA ETOPO1 database (Amante, 2009), with 1  
222 arc-second resolution (~30m) and processed using the ‘marmap’ R package (Pante & Simon-  
223 Bouhet, 2013). Slope was calculated in radians using the Fleming and Hoffer algorithm through  
224 the ‘raster’ R package (Fleming & Hoffer, 1979; Hijmans & van Etten, 2014). Depth and slope  
225 were modelled as continuous predictors (see supplemental information for covariate details).

### 226 **Site-specific temporal trends**

227 Generalized estimating equations with splines (GEE-GAMs) were fitted to each of the ten  
228 deployment groups based on *a priori* knowledge that bottlenose dolphin behaviour changes  
229 throughout their range, depending on whether they are or are not near foraging areas (Hastie et  
230 al., 2004; Pirodda et al., 2014; Thompson et al., 2013). GEE-GAMs were chosen for their flexible  
231 modelling structures capable of handling binary data. Only data from C-PODs that returned at  
232 least two days with ‘other cetacean’ detections were included in the temporal models. Temporal  
233 autocorrelation in detections across consecutive days was accounted for by including in the  
234 models an autoregressive correlation structure (*ar1*) to detections from each individual C-POD  
235 deployment (Box, Jenkins, Reinsel, & Ljung, 2015).

236 For this analysis, model selection focused on estimating the form of the relationship between the  
237 probability of detecting a broadband acoustic encounter and the Julian day of the year. For each  
238 deployment, four models were investigated. Predictor variables for all models included  
239 ShoreDist, a three level factor for distance from shore of the deployment location (05, 10, 15), a  
240 three level factor for survey year (2013, 2014 or 2015) and an integer for Julian day of year.

241 The first model (Equation 1) assumed an interaction between the shore distance and Julian day of  
242 year, and that the pattern in detections throughout the year could be modelled by a cubic B-  
243 spline. The second model (Equation 2) assumed an interaction between the cubic B-spline and  
244 the survey year. The third model (Equation 3) had no interactions between the cubic B-spline and  
245 the shore distance or survey year, and the fourth model (Equation 4) assumed a parametric  
246 linearrelationship between the daily probability of detecting a broadband echolocation click train,  
247  $P(\textit{Broadband})$ , and the Julian day of year. In accordance with previous studies using cubic spline  
248 models a single knot was set at the median of each C-POD record (Pirotta, Matthiopoulos,  
249 MacKenzie, Scott-Hayward, & Rendell, 2011). It was not possible to include more than one knot  
250 in the spatial models, as the lost degrees of freedom prevented model convergence. All models  
251 were fitted in R v.3.3.2 using the ‘geepack’ package (Halekoh, Højsgaard, & Yan, 2006). B-  
252 splines were added to the models using the ‘splines’ package (R Core Team, 2016).

253

$$P(\textit{Broadband}) \sim \textit{Year} + \textit{ShoreDist} * \textit{bs}(\textit{JulianDay}, \textit{knots} = \textit{median}(\textit{JulianDay}))$$

Equation 1

$$P(\text{Broadband}) \sim \text{ShoreDist} + \text{Year} *$$

Equation 2

$$bs(\text{JulianDay}, \text{knots} = \text{median}(\text{JulianDay}))$$

$$P(\text{Broadband}) \sim \text{ShoreDist} + \text{Year} +$$

Equation 3

$$bs(\text{JulianDay}, \text{knots} = \text{median}(\text{JulianDay}))$$

$$P(\text{Broadband}) \sim \text{ShoreDist} + \text{Year} + \text{JulianDay}$$

Equation 4

254

255 Akaike's Information Criterion (AIC) scores are commonly used to select between candidate  
256 GAM or GLM models (Akaike, 1974). However, because GEE's are not likelihood-based  
257 models, AIC scores cannot be calculated. Instead a quasi-likelihood criterion (QIC; Pan 2001)  
258 was used to select between the four temporal acoustic occupancy models. Quasi-likelihood  
259 criterion model selection mirrors AIC-based selection in application, but is appropriate for  
260 selecting between GEE models.

261 Assessing how well the selected model fitted the data followed previous methods (Pirodda et al.,  
262 2011; Thompson et al., 2013). For each deployment group, the model with the lowest QIC was  
263 used to predict the probability of detecting a broadband echolocation click across the range of the  
264 predictors. Receiver operating curves (ROCs; Fawcett, 2006) were then created to determine the  
265 relationship between the detection threshold, and the false positive and false negative rates for  
266 each model. ROC curves show the relationship between the proportions of true positive  
267 detections, here the proportion of days with broadband echolocation click trains accurately  
268 predicted, and the proportion of false positive detections or the proportion of days the model

269 inaccurately predicted the presence of broadband echolocation click presence. True and false  
270 positive rates are then plotted for each threshold. The threshold at which the trade-off between  
271 true and false positive rates is approximately equal is referred to as the optimum threshold. Using  
272 the ROC, an optimal detection threshold was selected above which broadband echolocation  
273 clicks were assumed to be detected and below which they were not. Using optimum threshold,  
274 confusion matrices were then created to measure the proportion of detection-positive and  
275 detection-negative days correctly identified by the model. The area under the ROC curve (AUC)  
276 was used to describe the model goodness-of-fit. AUC scores represent a measure of how well the  
277 model predicts the data. AUC values of 0.5 indicate that the model correctly predicted 50% of  
278 the observations and therefore, for a binomial model, values of 0.5 represent models that  
279 performed as well as would be expected by chance alone. Considering the variation in the data, it  
280 was relevant to determine how well each model fit all locations in the group. Thus, in addition to  
281 assessing how well the selected model fit each deployment group, how well the winning model  
282 fit the data from each C-POD deployment location was also investigated. Through this process  
283 AUC scores were calculated for each model for each deployment group, as well as for all 30  
284 individual deployment locations (Figure 1). These analyses were done in R using the ‘ROCR’  
285 v1.0-7 and ‘PresenceAbsence’ v1.19 packages (Freeman, 2007; Sing, Sander, Beerenwinkel, &  
286 Lengauer, 2005). The relationship between  $P(\text{Broadband})$  and Julian day was then plotted for  
287 each of the deployed C-PODs and years (Figures 3-5).

## 288 **Spatial-Temporal Habitat Modelling**

289 Bottlenose dolphins are known to move along the east coast of Scotland for foraging and other  
290 purposes (Cheney et al., 2013; Thompson et al., 2013). The full model presented in this study

291 included independent factors for slope, distance to point of aggregation, depth, and distance to  
292 shore. Temporal covariates included only season as a factor (spring, summer, and autumn).

293 For this analysis, a generalized additive mixed model (GAMM; Wood 2006) that incorporated  
294 both spatial and temporal variables was fitted to the data. Because smooth terms are centred  
295 using the MGCV package, smooth terms were also added as a main effect, as per package  
296 recommendations. As with the temporal models, an autoregressive correlation structure with  
297 detections grouped by deployment site was included (Box et al., 2015). Only the ‘ar1’  
298 autocorrelation structure was investigated, based on biological understanding that acoustic  
299 encounters spanning several days were unlikely to be driven by the same underlying factor.

300 The limited degrees of freedom in the data precluded fitting multiple models. Rather, a full  
301 model was fitted that included at least one form of all spatial and temporal covariates. Model  
302 covariates were investigated for collinearity using variance of inflation factors (VIF), and any  
303 covariates with VIF scores greater than two were considered collinear (Craney & Surles, 2002).

304 As the goal of this analysis was to produce a comprehensive model for habitat use, model  
305 selection was limited to excluding variables with estimated degrees of freedom less than 1.

306 Adjusted r-squared and AUC scores were used to describe model fit.

307 The resulting model was used to predict the presence of broadband acoustic encounters in the  
308 Scottish North Sea. A grid size of 1 km<sup>2</sup>, the approximate detection range of the C-PODs  
309 (Nuuttila, Thomas, et al., 2013) was used. The prediction space was restricted to habitats that fell  
310 within the parameters covered by the C-POD deployments including depth (103.0 - 9.3 m),  
311 distance to nearest point of aggregation (2.3 - 67.17 km), and distance to shore (0.35 - 17.9 km).

312

313 **3. Results**

314 **Acoustic Occupancy Rates**

315 Throughout the three years of survey reported in this study, 11,663 days of C-POD recordings  
316 were collected. At only 16 deployment sites, devices were retrieved in all three years (Figure 2  
317 and Supplemental Material). C-PODs deployed at the Fraserburgh 10 site were not recovered in  
318 2013 and 2014 and did not detect any acoustic encounters in 2015. C-PODs at the Spey Bay 10  
319 and Helmsdale 10 locations returned data for two of the three years but failed to document two  
320 or more days with dolphin echolocation click trains. The C-PODs deployed at the St Andrews 10  
321 location were successfully recovered in all three years but failed to detect dolphin echolocation  
322 click trains on two or more days.

323 The C-POD deployed nearest to Cromarty Firth showed the highest acoustic occupancy rate,  
324 with 78% of the days containing at least one broadband detection in 2013, and 83% in 2015  
325 (Table 2). There was wide variation in the acoustic occupancy rate and broadband occupancy  
326 rate across the array. C-PODs deployed at the northern and southern ends of the survey area  
327 (Latheron and St Abbs ) had very low (<5%) broadband occupancy rates for all survey years.  
328 Broadband occupancy rates at the nearshore (05) deployment locations were typically greater  
329 than the more offshore (10 or 15) locations. The mean broadband occupancy rates for the 05, 10  
330 and 15 locations were 0.12, 0.03 and 0.02 detections/day respectively. Excluding the Cromarty  
331 05 C-POD, the occupancy rate for the nearshore deployments was 0.06 detections/day, nearly  
332 twice that of the mid or offshore locations (Table 2).

333 C-PODs in the Stonehaven deployment group were notable for having the second highest  
334 acoustic occupancy rates behind the Cromarty group. Both broadband and frequency branded

335 acoustic encounters were documented at these sites with similar frequency (Figure 2). The C-  
336 PODs in this group detected echolocation click encounters on more than 15% of the survey days  
337 and broadband encounters on at least 10% of the survey days.

338

### 339 **Site-Specific Temporal Trends**

340 Deployments at the Helmsdale 15, St Andrews 10, Fraserburgh 10, and Spey Bay 10 sites failed  
341 to detect broadband clicks on at least two days and were removed from the temporal analysis.

342 Delta-QIC scores for temporal model selection were less than 3.5 for half of the deployment  
343 groups indicating some uncertainty in model selection. Furthermore, AUC scores at some  
344 individual deployment sites less than 0.5 (Table 3): equal to what would be expected by chance  
345 alone. Even at sites with high AUC scores, the ability to predict days with broadband acoustic  
346 encounters was 0.53, indicating that nearly half the detections could not be explained by the  
347 model. The lowest AUC score among the ten deployment groups was at the St Abbs group (AUC  
348 = 0.62, Table 3), indicating it performed only slightly better than would be expected by chance  
349 alone. The highest AUC was 0.93 determined for both the Helmsdale and Cromarty groups.

350 When model fit was investigated at each of the 30 deployment sites, AUC ranged from 0.2 at the  
351 Cruden Bay 10 location to 0.99 at the Latheron 10 location (Table 2).

352 Low acoustic occupancy rates across most sites meant that the temporal models generally did  
353 well at predicting periods without detections, but were poor at predicting detection-positive days.  
354 Across the dataset, 43% of the days without broadband detections were accurately predicted,  
355 with the exception of Cruden Bay, where 30% were correctly classified. Apart from the

356 Cromarty group, no model was able to predict more than 20% of the broadband detection-  
357 positive days.

358 Large ( $>3$ )  $\Delta$ QIC and high ( $>0.75$ ) AUC scores indicated a more confident model selection and  
359 better model fit at the Latheron 10, St Andrews 15, Stonehaven 15, Spey Bay 10, and Helmsdale  
360 15 sites. Of these, only the Stonehaven 15 location contained broadband echolocation click  
361 trains on greater than 1 % of the days. Thus, high AUC scores at the other locations were  
362 influenced by the correct prediction of days without dolphin detections.

363 For sites with the highest acoustic occupancy of broadband click trains, e.g. Cromarty 05 and  
364 Stonehaven 15, GEE-GLM models suggested peaks in the probability of detecting broadband  
365 echolocation encounters in August and July, respectively. At other locations, including  
366 deployment sites in the Fraserburgh, Arbroath and St Andrews groups, temporal trends in  
367 acoustic occupancy were highly stochastic. Poor model fits ( $AUC < 0.50$ ) at the deployment sites  
368 within these groups make it difficult to identify the presence and/or persistence of patterns in  
369 daily acoustic occupancy (Figures 3-5).

### 370 **Spatial-Temporal Habitat Modelling**

371 VIF scores for spatial covariates were less than three and subsequently all spatial variables were  
372 retained. In the full model the estimated degrees of freedom (EDF) for slope were less than one,  
373 and the predictor was removed. In the final model, all terms were significant except season  
374 (Table 4). The AUC score of the final model was 0.86. Modelling results suggested that the  
375 probability of detecting broadband echolocation click train encounters decreased with increasing  
376 distance to shore and increasing distance to the nearest point of aggregation. However, across the

377 extent of the array, the probability of detecting broadband echolocation encounters increased  
378 with increasing depth (Figure 6).

379 When the GAMM was projected over the available habitat, higher broadband occupancy was  
380 predicted near the Inner Moray Firth and Dee river estuaries. The GAMM also predicted that C-  
381 PODs deployed in nearshore areas were more likely to detect broadband encounters than those  
382 deployed further offshore. Finally, deeper (>60m) offshore areas were projected to have a higher  
383 probability of broadband occupancy than shallow areas (Figure 7; see supplemental material for  
384 projections of the confidence intervals as well as projections for Spring and Autumn).

385

#### 386 4. Discussion

387 The primary goal of this study was to improve understanding of the patterns of habitat use by  
388 this coastal bottlenose dolphin population throughout its range. A three-step approach was taken  
389 to the analysis. First, daily acoustic occupancy rates were reported for all sites and for both  
390 unfiltered acoustic encounters and echolocation click encounters identified as ‘broadband’ by the  
391 classification system (Palmer et al., 2017). Second, models investigating temporal trends were  
392 fitted to the available data to investigate seasonal occurrence patterns. Third, a spatial-temporal  
393 model was fitted to the data to predict the animals’ habitat use.

394 The study faced two main challenges: low acoustic occupancy rates and species classification  
395 uncertainty. Low acoustic occupancy rates limited detection sample size. The autocorrelation  
396 structure in the temporal model accounted for correlation within acoustic encounters, but further  
397 limited the remaining degrees of freedom to model spatial and temporal trends in acoustic  
398 occupancy. For half of the deployment groups, model selection techniques ( $\Delta QIC$ ) did not

399 strongly favour one temporal model over another. This is indicative of variation not accounted  
400 for by any of the models in the set. Larger amounts of acoustic data will be needed to produce  
401 robust model estimates in future studies. For example, Pirodda et al. (2014) used data from eight  
402 years of continuous surveys to produce estimates for dolphin foraging rates within the Moray  
403 Firth SAC. In addition to having a longer sample period, echolocation detectors in Pirodda et al.  
404 (2014) were deployed in areas of high use and therefore registered a higher rate of detections.  
405 Future deployments of the ECoMMAS will address some of the temporal modelling issues this  
406 work encountered, as gaps in data coverage are reduced through multiple annual deployments, as  
407 was done in 2015.

408 Despite their lack of species resolution, C-PODs remain widely used instruments for passive  
409 acoustic monitoring (Cox et al., 2017; Jaramillo-Legorreta et al., 2017; Nuuttila, Meier, et al.,  
410 2013; Williamson et al., 2016; Wilson, Benjamins, & Elliott, 2013). Nearly all studies that use  
411 the instruments simply assume that the target species, here bottlenose dolphins, are responsible  
412 for the preponderance of the detections. This study improves to some degree on that assumption  
413 by applying a secondary classification system to the detections. This system is unable to  
414 discriminate between some species of dolphins and enhanced taxonomic resolution is unlikely to  
415 be achievable with these devices. Even with full spectrum recordings, it is exceptionally difficult  
416 to discriminate between echolocation clicks of common and bottlenose dolphins (Soldevilla et  
417 al., 2008).

418 Species discrimination was improved over the Chelonia classifier, but perfect dolphin  
419 classification is impossible for any passive acoustic study, regardless of the recording device or  
420 sample frequency (Caillat, 2013; Roch et al., 2011). As such, these findings emphasise the need  
421 to combine long term data from visual and acoustic surveys. In doing so, researchers will be able

422 to provide robust data on long-term trends in dolphin occurrence throughout the habitat and for  
423 areas of ecological or commercial interest (Thompson et al., 2011).

424 This is the first acoustic study that approximates the entire geographic range of this population.  
425 The ECoMMAS, in combination with Arso Civil et al. (2019), provides critical information  
426 about baseline habitat use. Such information is needed to monitor change in habitat use through  
427 time (Bailey et al., 2010). A novel classification algorithm on the ‘other cetacean’ detections  
428 reported by the C-POD software was used. The additional information produced by the  
429 classification algorithm enabled both the temporal and spatial-temporal model to more closely  
430 focus on the species of interest. Thus, this research required fewer assumptions about the impact  
431 of non-target species detections on the resulting models.

432 Patterns in broadband acoustic occupancy rates were generally consistent with previous research  
433 suggesting the bottlenose dolphins are more likely to be observed in coastal waters, within 5 km  
434 of shore (Arso Civil, 2014). While most instruments were deployed in less than 30m of water,  
435 broadband acoustic occupancy rates throughout the survey were generally higher for C-PODs  
436 closer to the shoreline (Table 3; Figure 2). This supports the work of Thompson, Brookes &  
437 Cordes (2015) and increases the confidence that bottlenose dolphins are unlikely to be present in  
438 areas that may be exposed to significant construction noise from offshore wind farms.

439 Acoustic occupancy rates and habitat modelling highlight the waters between Stonehaven and  
440 Aberdeen as a potential area of high occupancy. Instruments deployed in the Stonehaven group  
441 showed the second highest acoustic occupancy rates behind the Cromarty group. In 2013 and  
442 2015, the Stonehaven 15 and 05 (respectively) C-PODs documented dolphin presence on at least  
443 30% of the monitored days (Table 3). Moreover, both broadband and frequency banded click

444 trains were documented at these sites at nearly equal rates, suggesting a potential hotspot  
445 important for multiple species. Previous studies have shown that dolphins are present in coastal  
446 waters north and south of Stonehaven year-round (Thompson et al., 2011). Historically, white-  
447 beaked and bottlenose dolphin sightings have been common in visual surveys (Anderwald et al.,  
448 2010; Arso Civil, 2014; Weir, Stockin, & Pierce, 2007). Thus, further research to determine  
449 whether the area constitutes a biological hotspot is warranted.

450 Modelling efforts for temporal trends across the spatial and temporal extent of the array were  
451 challenged by few detections and gaps in data coverage. As such, the inference that can be made  
452 from the models is highly limited. Despite these challenges, the model for the Cromarty group  
453 did fit well and indicated a peak in broadband detections consistent with earlier visual surveys  
454 (Thompson et al., 2011). The novel approach to classification uncertainty reduced the number of  
455 days with echolocation encounters in the dataset. While this conservative approach hindered  
456 modelling efforts in this research, it will provide more robust estimates of dolphin species  
457 distributions as the survey matures (Pirota et al., 2014).

458 Spatial-temporal habitat selection modelling was more successful and generally agreed with  
459 previous studies linking smaller distances to shore with increased probability of detecting  
460 bottlenose dolphins (Arso Civil, 2014; Pirota et al., 2014; Quick et al., 2014). The spatial  
461 modelling suggested that broadband acoustic encounters were more likely to be detected in  
462 deeper water and predicted a slight increase in detections >15 km from shore (Figures 5-6).  
463 Without concurrent visual confirmation residual uncertainty remains regarding whether and to  
464 what extent echolocation encounters detected at offshore locations represented common  
465 dolphins. The spatial-temporal model indicated that distance to the nearest selected point of  
466 aggregation and depth were also important predictors of broadband occupancy. Unfortunately,

467 there were not enough detection data to model the spatial and temporal covariates together (e.g.  
468 Julian day of year and depth).

469 Bottlenose dolphins are commonly sighted in St Andrews Bay (Arso Civil et al., 2019; Quick et  
470 al., 2014), so the low number of detections at the St Andrews survey location nearest the bay (St  
471 Andrews 05) was somewhat unexpected. There are several possible reasons for this. One  
472 possibility is that the area may represent habitat associated with rest or socializing rather than  
473 foraging, so there are fewer clicks to detect. Previous studies have found lower detection rates  
474 for groups of animals, travelling and socializing animals, than single animals or foraging animals  
475 (Nuuttila, Thomas, et al., 2013). If animals near the Fife Ness survey sites were primarily  
476 travelling or socializing, they may not have been detected at rates comparable to foraging  
477 animals. These results reinforce the need to integrate visual and acoustic surveys when managing  
478 highly mobile species.

479 Unfortunately, the limited taxonomic resolution of the acoustic data means that it is not possible  
480 to say with a high degree of certainty which of the broadband or frequency banded species were  
481 present at these locations. Delphinid species classification is an issue that other studies using C-  
482 PODs have not typically had the tools to address. This study uses improved classification  
483 measures to more reliably discriminate between the various species present in the area.  
484 Furthermore, the maximum acoustic occupancy probability of 0.79 for broadband acoustic  
485 encounters is not a direct representation of true bottlenose dolphin occupancy. Thus,  
486 conservative interpretation of these results, including relative occupancy between the survey  
487 locations, is prudent.

488 In situations where species classification remains an outstanding problem it is appropriate to  
489 combine inferences from multiple survey methodologies (Cheney et al., 2013; Thompson et al.,  
490 2015). In this survey region, visual surveys provide evidence that the majority of the broadband  
491 echolocation encounters detected at the near-shore deployments originated from bottlenose  
492 dolphins (Anderwald et al., 2010; Arso Civil, 2014; Arso Civil et al., 2019; Thompson et al.,  
493 2013). Considerable uncertainty remains regarding broadband detections from offshore areas that  
494 lack consistent visual survey effort. Where there are increased broadband detections at the  
495 offshore locations, the data warrant further investigation, but classification is not possible. These  
496 areas would benefit from either increased visual survey effort or more advanced acoustic  
497 techniques that have recently shown promise in discriminating between common and bottlenose  
498 dolphins (Frasier et al., 2017).

499 Data presented here also represent a small spatial sample, and acoustic data are lacking from  
500 many important sites such as the River Dee and Tay estuary. In these, shipping activity has  
501 restricted the use of acoustic moorings which may present a potential navigational hazard. Thus,  
502 it has not been possible to deploy acoustic recorders in some known points of aggregation.

503 Appropriate sampling methods for investigating temporal and spatial trends are diametrically  
504 opposed. If the continued goal of the ECoMMAS array is to relate habitat data to acoustic  
505 occupancy, managers should consider changing deployment locations at each recovery and re-  
506 deployment. However, if the goal is to maintain a historical record of the trends in acoustic  
507 occupancy at these locations it is important that the deployment locations remain consistent.

508 From a conservation and management perspective, knowledge of where animals are is equally as  
509 valuable as knowledge of where they are not. The ECoMMAS provides continuous survey  
510 coverage for areas where consistent visual surveys are untenable. The first ecological results of

511 ECoMMAS are consistent with visual sightings highlighting the importance of particular high-  
512 usage areas to the population (Arso Civil et al., 2019). Similarly, daily acoustic occupancy rates  
513 in areas between established points of aggregation were an order of magnitude lower. In sites  
514 other than Cromarty 05, there was no clear trend in temporal detections. Thus, by themselves,  
515 these results do not suggest the need to change any of the existing regulatory framework for this  
516 population of bottlenose dolphins.

517 Bottlenose dolphins are highly mobile and adaptable generalists, capable of exploiting changing  
518 environments (Santos et al., 2001). The areas currently considered to be critical habitat for this  
519 population (e.g. the SAC) may shift with changing climate or other anthropogenic impacts. For  
520 example, the point of aggregation near the Cromarty Firth has conclusively been linked with  
521 foraging (Hastie et al. 2003). If the area no longer provides optimal foraging habitat, dolphins  
522 will likely move elsewhere. Under such dynamic systems fixed protected areas may not provide  
523 optimal conservation solutions for either protected species or human users. Dynamic ocean  
524 management plans represent a flexible conservation approach that mirror the spatial and  
525 temporal variability present in marine systems (Maxwell et al., 2015). Such management plans  
526 have been implemented in North America where vessel speed restrictions may be triggered when  
527 critically endangered North Atlantic right whales (*Eubalaena glacialis*) are visually or  
528 acoustically detected near shipping lanes (Spaulding et al., 2009; Van Parijs et al., 2009). These  
529 dynamic management areas are designed to provide maximum protection from anthropogenic  
530 mortality while limiting additional regulatory burden on users. In a changing regulatory  
531 landscape, there may be opportunities to rethink the implementation of conservation measures  
532 for highly mobile species. Since the establishment of the Moray Firth SAC, the population has  
533 grown and is now observed using the entire coastline (Cheney et al., 2013; Arso Civil et al.

534 2019). This range expansion over a relatively short period might be reflected in a dynamic  
535 management plan that considers variation in animal presence and the timing of ecological  
536 features (e.g. diadromous fish runs or seasonal patterns in habitat use). Under dynamic  
537 management plans, surveys like the ECoMMAS would be invaluable in providing detailed  
538 information about habitat over longer periods than can be provided by visual surveys alone.

### 539 **Acknowledgments**

540 We are especially grateful to the deployment and recovery crews from Marine Scotland Science,  
541 Moray First Marine, and the crew of the Northern Lighthouse Board vessel, Polestar. Sophie  
542 Smout and Dave Miller provided useful feedback on statistical analysis for which we are much  
543 indebted. LR and KJP were supported by Marine Scotland Science and the Marine Alliance for  
544 Science and Technology for Scotland (MASTS) pooling initiative, and their support is gratefully  
545 acknowledged. MASTS is funded by the Scottish Funding Council (grant reference HR09011)  
546 and contributing institutions. We are also grateful to Paul Thompson, John Baxter, and two  
547 anonymous reviewers for providing helpful comments on the manuscript.

548

549

550

551

552 **Literature Cited**

- 553 Akaike, H. (1974). A new look at the statistical model identification. *IEEE transactions on*  
554 *automatic control*, 19(6), 716-723.
- 555 Amante, C. (2009). ETOPO1 1 arc-minute global relief model: procedures, data sources and  
556 analysis: US Department of Commerce, National Oceanic and Atmospheric  
557 Administration, National Environmental Satellite, Data, and Information Service,  
558 National Geophysical Data Center, Marine Geology and Geophysics Division Colorado.  
559 <http://www.ngdc.noaa.gov/mgg/global/global.html>. doi:10.7289/V5C8276M
- 560 Anderwald, P., Evans, P. G., Canning, S., Hepworth, K., Innes, M., McDonald, P., Sim, I.,  
561 Stokin, K., Weir, C. (2010). Cetaceans of the East Grampian Region. *Sea Watch*  
562 *Foundation, Aberdeen,*  
563 *UK*. Retrieved from [https://www.seawatchfoundation.org.uk/wp-content/uploads/2012/08/East-](https://www.seawatchfoundation.org.uk/wp-content/uploads/2012/08/East-gramp-cet-rev.pdf)  
564 [gramp-cet-rev.pdf](https://www.seawatchfoundation.org.uk/wp-content/uploads/2012/08/East-gramp-cet-rev.pdf). Last Accessed: 30/04/2019.
- 565 Arso Civil, M. (2014). PhD Thesis: Population ecology of bottlenose dolphins (*Tursiops*  
566 *truncatus*) of the East Coast of Scotland. <http://hdl.handle.net/10023/6543>.
- 567 Arso Civil, M., Quick, N., Cheney, B., Pirota, E., Thompson, P., & Hammond, P. (2019).  
568 Changing distribution of the east coast of Scotland bottlenose dolphin population and the  
569 challenges of areabased management. *Aquatic Conservation: Marine and Freshwater*  
570 *Ecosystems, This Volume*.
- 571 Bailey, H., Clay, G., Coates, E. A., Lusseau, D., Senior, B., & Thompson, P. M. (2010). Using  
572 T-PODs to assess variations in the occurrence of coastal bottlenose dolphins and harbour  
573 porpoises. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 20(2), 150-158.
- 574 Bailey, H., & Thompson, P. (2006). Quantitative analysis of bottlenose dolphin movement  
575 patterns and their relationship with foraging. *Journal of Animal Ecology*, 75(2), 456-465.
- 576 Box, G. E., Jenkins, G. M., Reinsel, G. C., & Ljung, G. M. (2015). *Time series analysis:*  
577 *forecasting and control*: John Wiley & Sons.
- 578 Brookes, K. L., Bailey, H., & Thompson, P. M. (2013). Predictions from harbor porpoise habitat  
579 association models are confirmed by long-term passive acoustic monitoring. *The Journal*  
580 *of the Acoustical Society of America*, 134(3), 2523-2533.

581 Caillat, M. (2013). Assessing and correcting for the effects of species misclassification during  
582 passive acoustic surveys of cetaceans. (PhD), University of St Andrews,

583 Calderan, S., Wittich, A., Harries, O., Gordon, J., & Leaper, R. (2013). White-beaked dolphin  
584 and Risso's dolphin click characteristics and the potential for classification and species  
585 identification. *Scottish Natural Heritage Commissioned Report No. 624* Retrieved from  
586 [http://www.snh.org.uk/pdfs/publications/commissioned\\_reports/624.pdf](http://www.snh.org.uk/pdfs/publications/commissioned_reports/624.pdf). Last Accessed:  
587 21/05/2018.

588 Cheney, B., Thompson, P. M., Ingram, S. N., Hammond, P. S., Stevick, P. T., Durban, J. W.,  
589 Culloch, R.M., Elwen, S.H., Mandleberg, L., Janik, V.M. Quick, N.J. (2013). Integrating  
590 multiple data sources to assess the distribution and abundance of bottlenose dolphins  
591 *Tursiops truncatus* in Scottish waters. *Mammal Review*, 43(1), 71-88.

592 Cox, S. L., Witt, M. J., Embling, C. B., Godley, B. J., Hosegood, P. J., Miller, P. I., . . . Ingram,  
593 S. N. (2017). Temporal patterns in habitat use by small cetaceans at an oceanographically  
594 dynamic marine renewable energy test site in the Celtic Sea. *Deep-Sea Research Part Ii-*  
595 *Topical Studies in Oceanography*, 141, 178-190.

596 Craney, T. A., & Surles, J. G. (2002). Model-dependent variance inflation factor cutoff values.  
597 *Quality Engineering*, 14(3), 391-403.

598 Culloch, R. M., & Robinson, K. P. (2008). Bottlenose dolphins using coastal regions adjacent to  
599 a Special Area of Conservation in north-east Scotland. *Journal of the Marine Biological*  
600 *Association of the United Kingdom*, 88(6), 1237-1243.

601 Fleming, M. D., & Hoffer, R. M. (1979). Machine processing of Landsat MSS data and DMA  
602 topographic data for forest cover type mapping. *LARS Technical Reports*, 80.  
603 <https://pdfs.semanticscholar.org/f77d/047528cc029ac1354aeb15d2a43444211478.pdf>  
604 Last Accessed: 21/05/2018.

605 Frasier, K. E., Roch, M. A., Soldevilla, M. S., Wiggins, S. M., Garrison, L. P., & Hildebrand, J.  
606 A. (2017). Automated classification of dolphin echolocation click types from the Gulf of  
607 Mexico. *PLoS computational biology*, 13(12), e1005823.

608 Freeman, E. (2007). PresenceAbsence: an R package for presence-absence model evaluation.  
609 *USDA Forest Service, Rocky Mountain Research Station*, 507.

610 Graham, I. M., Pirodda, E., Merchant, N. D., Farcas, A., Barton, T. R., Cheney, B., . . .  
611 Thompson, P. M. (2017). Responses of bottlenose dolphins and harbor porpoises to  
612 impact and vibration piling noise during harbor construction. *Ecosphere*, 8(5), e01793.

613 Halekoh, U., Højsgaard, S., & Yan, J. (2006). The R package geepack for generalized estimating  
614 equations. *Journal of Statistical Software*, 15(2), 1-11.

615 Hammond, P., Lacey, C., Gilles, A., Viquerat, S., Boerjesson, P., Herr, H., Macleod, K., Ridoux,  
616 V., Santos, M.b., Scheidat, M., Teilmann, J., Vingada, J., Øien, N. (2017). Estimates of  
617 cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III  
618 aerial and shipboard surveys. Retrieved from: [https://synergy.st-](https://synergy.st-andrews.ac.uk/scans3/files/2017/04/SCANS-III-design-based-estimates-2017-04-28-final.pdf)  
619 [andrews.ac.uk/scans3/files/2017/04/SCANS-III-design-based-estimates-2017-04-28-](https://synergy.st-andrews.ac.uk/scans3/files/2017/04/SCANS-III-design-based-estimates-2017-04-28-final.pdf)  
620 [final.pdf](https://synergy.st-andrews.ac.uk/scans3/files/2017/04/SCANS-III-design-based-estimates-2017-04-28-final.pdf). Last Accessed: 21/05/2018.

621 Harris, C. M., Thomas, L., Falcone, E. A., Hildebrand, J., Houser, D., Kvadsheim, P. H., Lam,  
622 F.P.A., Miller, P.J., Moretti, Read, A. J. (2017). Marine mammals and sonar: dose-  
623 response studies, the risk-disturbance hypothesis and the role of exposure context.  
624 *Journal of Applied Ecology*, 55(1), 369-404.

625 Hastie, G., Wilson, B., & Thompson, P. (2003). Fine-scale habitat selection by coastal bottlenose  
626 dolphins: application of a new land-based video-montage technique. *Canadian Journal of*  
627 *Zoology-Revue Canadienne De Zoologie*, 81(3), 469-478.

628 Hastie, G., Wilson, B., & Thompson, P. (2006). Diving deep in a foraging hotspot: acoustic  
629 insights into bottlenose dolphin dive depths and feeding behaviour. *Marine Biology*,  
630 148(5), 1181-1188.

631 Hastie, G. D., Barton, T. R., Grellier, K., Hammond, P. S., Swift, R. J., Thompson, P. M., &  
632 Wilson, B. (2003). Distribution of small cetaceans within a candidate Special Area of  
633 Conservation; implications for management. *Journal of Cetacean Research and*  
634 *Management*, 5(3), 261-266.

635 Hastie, G. D., Wilson, B., Wilson, L., Parsons, K., & Thompson, P. (2004). Functional  
636 mechanisms underlying cetacean distribution patterns: hotspots for bottlenose dolphins  
637 are linked to foraging. *Marine Biology*, 144(2), 397-403.

638 Hijmans, R. J., & van Etten, J. (2014). raster: Geographic data analysis and modeling. *R package*  
639 *version*, 2, 15.

640 Janik, V. M., & Thompson, P. M. (1996). Changes in surfacing patterns of bottlenose dolphins in  
641 response to boat traffic. *Marine Mammal Science*, 12(4), 597-602.

642 Jaramillo-Legorreta, A., Cardenas-Hinojosa, G., Nieto-Garcia, E., Rojas-Bracho, L., Ver Hoef,  
643 J., Moore, J., Tregenza, N., Barlow, J., Gerrodette, T., Thomas, L., Taylor, B. (2017).  
644 Passive acoustic monitoring of the decline of Mexico's critically endangered vaquita.  
645 *Conservation Biology*, 31(1), 183-191. Retrieved from  
646 <https://www.ncbi.nlm.nih.gov/pubmed/27338145>. doi:10.1111/cobi.12789

647 Maxwell, S. M., Hazen, E. L., Lewison, R. L., Dunn, D. C., Bailey, H., Bograd, Briscoe, S. J.,  
648 Fossette, D.K., Hobday, S., Bennett, A.J., Benson, M. (2015). Dynamic ocean  
649 management: Defining and conceptualizing real-time management of the ocean. *Marine*  
650 *Policy*, 58, 42-50.

651 Mendes, S., Turrell, W., Lütkebohle, T., & Thompson, P. (2002). Influence of the tidal cycle and  
652 a tidal intrusion front on the spatio-temporal distribution of coastal bottlenose dolphins.  
653 *Marine Ecology Progress Series*, 239, 221-229.

654 Nuuttila, H. K., Meier, R., Evans, P. G. H., Turner, J. R., Bennell, J. D., & Hiddink, J. G. (2013).  
655 Identifying Foraging Behaviour of Wild Bottlenose Dolphins (*Tursiops truncatus*) and  
656 Harbour Porpoises (*Phocoena phocoena*) with Static Acoustic Dataloggers. *Aquatic*  
657 *Mammals*, 39(2), 147-161.

658 Nuuttila, H. K., Thomas, L., Hiddink, J. G., Meier, R., Turner, J. R., Bennell, J. D., Bennell, J.D.,  
659 Tregenza, N.C., Evans, P. G. (2013). Acoustic detection probability of bottlenose  
660 dolphins, *Tursiops truncatus*, with static acoustic dataloggers in Cardigan Bay, Wales.  
661 *The Journal of the Acoustical Society of America*, 134(3), 2596-2609.

662 Palmer, K. J., Brookes, K., & Rendell, L. (2017). Categorizing click trains to increase taxonomic  
663 precision in echolocation click loggers. *The Journal of the Acoustical Society of America*,  
664 142(2), 863.

665 Pante, E., & Simon-Bouhet, B. (2013). marmap: a package for importing, plotting and analyzing  
666 bathymetric and topographic data in R. *PloS one*, 8(9), e73051.

667 Paxton, C. G. M., Scott-Hayward, L. A. S., Mackenzie, M., Rexstad, E., & Thomas, L. (2016).  
668 Revised Phase III Data Analysis of Joint Cetacean Protocol Data Resources with  
669 Advisory Note, JNCC Report 517, ISSN 0963-8091. Retrieved from  
670 <http://jncc.defra.gov.uk/page-7201>. Last Accessed: 30/04/2018.

671 Perrin, W. F., Würsig, B., & Thewissen, J. (2009). *Encyclopedia of marine mammals* (Second  
672 ed.): Academic Press.

673 Pirotta, E., Matthiopoulos, J., MacKenzie, M., Scott-Hayward, L., & Rendell, L. (2011).  
674 Modelling sperm whale habitat preference: a novel approach combining transect and  
675 follow data. *Marine Ecology Progress Series*, 436, 257-272.

676 Pirotta, E., Merchant, N. D., Thompson, P. M., Barton, T. R., & Lusseau, D. (2015). Quantifying  
677 the effect of boat disturbance on bottlenose dolphin foraging activity. *Biological  
678 Conservation*, 181, 82-89.

679 Pirotta, E., Thompson, P. M., Miller, P. I., Brookes, K. L., Cheney, B., Barton, T. R., Graham, I.,  
680 Lusseau, D. (2014). Scale-dependent foraging ecology of a marine top predator modelled  
681 using passive acoustic data. *Functional Ecology*, 28(1), 206-217.

682 Quick, N. J., Arso, M., Cheney, B., Valentina, I., Janik, V., Thompson, P., & Hammond, P. S.  
683 (2014). *The east coast of Scotland bottlenose dolphin population: Improving  
684 understanding of ecology outside the Moray Firth SAC*. Retrieved from  
685 [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/346326/O](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/346326/OESEA2_east_coast_of_Scotland_bottlenose_dolphin_population.pdf)  
686 [ESEA2\\_east\\_coast\\_of\\_Scotland\\_bottlenose\\_dolphin\\_population.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/346326/OESEA2_east_coast_of_Scotland_bottlenose_dolphin_population.pdf). Last Accessed:  
687 30/04/2018.

688 Quick, N. J., & Janik, V. M. (2012). Bottlenose dolphins exchange signature whistles when  
689 meeting at sea. *Proceedings of the Royal Society B-Biological Sciences*, 279(1738), 2539-  
690 2545

691 R Core Team. (2016). R Foundation for Statistical Computing, Vienna, Austria (Version 3.2.4).  
692 Vienna, Austria: R Foundation for Statistical Computing. Retrieved from [https://www.R-](https://www.R-project.org/)  
693 [project.org/](https://www.R-project.org/)

694 Roch, M. A., Klinck, H., Baumann-Pickering, S., Mellinger, D. K., Qui, S., Soldevilla, M. S., &  
695 Hildebrand, J. A. (2011). Classification of echolocation clicks from odontocetes in the  
696 Southern California Bight. *The Journal of the Acoustical Society of America*, 129(1),  
697 467-475.

698 Santos, M., Pierce, G. J., Reid, R., Patterson, I., Ross, H., & Mente, E. (2001). Stomach contents  
699 of bottlenose dolphins (*Tursiops truncatus*) in Scottish waters. *Journal of the Marine  
700 Biological Association of the United Kingdom*, 81(5), 873-878.

701 Sargeant, B. L., Mann, J., Berggren, P., & Krutzen, M. (2005). Specialization and development  
702 of beach hunting, a rare foraging behavior, by wild bottlenose dolphins (*Tursiops sp.*).  
703 *Canadian Journal of Zoology-Revue Canadienne De Zoologie*, 83(11), 1400-1410.

704 Sarnocinska, J., Tougaard, J., Johnson, M., Madsen, P. T., & Wahlberg, M. (2016). Comparing  
705 the performance of C-PODs and SoundTrap/PAMGUARD in detecting the acoustic  
706 activity of harbor porpoises (*Phocoena phocoena*). Paper presented at the Proceedings of  
707 Meetings on Acoustics, Fourth International Conference on the Effects of Noise on  
708 Aquatic Life. Dublin, Ireland. July 10-16

709 Science, M. S. (2013). The East Coast Marine Mammal Acoustic Study (ECOMMAS). Topic  
710 Sheet NO 126 V1. . Retrieved From: <http://www.gov.scot/Resource/0042/00426891.pdf>  
711 Last Accessed: 30/04/2018.

712 Simon, M., Nuuttila, H., Reyes-Zamudio, M. M., Ugarte, F., Verfub, U., & Evans, P. G. H.  
713 (2010). Passive acoustic monitoring of bottlenose dolphin and harbour porpoise, in  
714 Cardigan Bay, Wales, with implications for habitat use and partitioning. *Journal of the*  
715 *Marine Biological Association of the United Kingdom*, 90(8), 1539-1545.

716 Sing, T., Sander, O., Beerenwinkel, N., & Lengauer, T. (2005). ROCr: visualizing classifier  
717 performance in R. *Bioinformatics*, 21(20), 3940-3941.

718 Soldevilla, M. S., Henderson, E. E., Campbell, G. S., Wiggins, S. M., Hildebrand, J. A., & Roch,  
719 M. A. (2008). Classification of Risso's and Pacific white-sided dolphins using spectral  
720 properties of echolocation clicks. *The Journal of the Acoustical Society of America*,  
721 124(1), 609-624.

722 Spaulding, E., Robbins, M., Calupca, T., Clark, C. W., Tremblay, C., Waack, A., Ward, A.,  
723 Kemp, J., Newhall, K. (2009). *An autonomous, near-real-time buoy system for automatic*  
724 *detection of North Atlantic right whale calls*. Paper presented at the Proceedings of  
725 Meetings on Acoustics (6). doi: 10.1121/1.3340128

726 Thompson, P., Brookes, K. L., Cordes, L. S., Barton, T., Cheney, B., & Graham, I. M. (2013).  
727 *Assessing the Potential Impact of Oil and Gas Exploration Operations on Cetaceans in*  
728 *the Moray Firth*. Third year Report for DECC, Scottish Government, COWRIE, and Oil  
729 & Gas UK. Retrieved from  
730 [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/50021/mf-](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/50021/mf-)  
731 [results2.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/50021/mf-results2.pdf). Last Accessed: 30/04/2018.

732  
733 Thompson, P. M., Cheney, B., Ingram, S., Stevick, P. T., Wilson, B., & Hammond, P. S. (2011).  
734 Distribution, abundance and population structure of bottlenose dolphins in Scottish  
735 waters. Scottish Government and Scottish Natural Heritage funded report. Scottish  
736 Natural Heritage Commissioned Report No. 354. Retrieved from  
737 [https://www.abdn.ac.uk/lighthouse/documents/SNH\\_Report\\_354\\_Distribution\\_abundanc](https://www.abdn.ac.uk/lighthouse/documents/SNH_Report_354_Distribution_abundance.pdf)  
738 [e.pdf](https://www.abdn.ac.uk/lighthouse/documents/SNH_Report_354_Distribution_abundance.pdf). Last Accessed: 30/04/2018.

739 Thompson, P., Pierce, G., Hislop, J., Miller, D., & Diack, J. (1991). Winter foraging by common  
740 seals (*Phoca vitulina*) in relation to food availability in the inner Moray Firth, NE  
741 Scotland. *The Journal of Animal Ecology*, 283-294.

742 Thompson, P. M., Brookes, K. L., & Cordes, L. S. (2015). Integrating passive acoustic and  
743 visual data to model spatial patterns of occurrence in coastal dolphins. *ICES Journal of*  
744 *Marine Science*, 72(2), 651-660.

745 Van Parijs, S. M., Clark, C. W., Sousa-Lima, R. S., Parks, S. E., Rankin, S., Risch, D., & van  
746 Opzeeland, I. (2009). Management and research applications of real-time and archival  
747 passive acoustic sensors over varying temporal and spatial scales. *Marine Ecology*  
748 *Progress Series*, 395, 21-36.

749 Weir, C. R., Stockin, K. A., & Pierce, G. J. (2007). Spatial and temporal trends in the distribution  
750 of harbour porpoises, white-beaked dolphins and minke whales off Aberdeenshire (UK),  
751 north-western North Sea. *Journal of the Marine Biological Association of the United*  
752 *Kingdom*, 87(1), 327-338.

753 Williamson, L. D., Brookes, K. L., Scott, B. E., Graham, I. M., Bradbury, G., Hammond, P. S.,  
754 & Thompson, P. M. (2016). Echolocation detections and digital video surveys provide  
755 reliable estimates of the relative density of harbour porpoises. *Methods in Ecology and*  
756 *Evolution*, 7(7), 762-769.

757 Wilson, B., Benjamins, S., & Elliott, J. (2013). Using drifting passive echolocation loggers to  
758 study harbour porpoises in tidal-stream habitats. *Endangered Species Research*, 22(2),  
759 125-U152.

760 Wilson, B., Thompson, P. M., & Hammond, P. S. (1997). Habitat use by bottlenose dolphins:  
761 seasonal distribution and stratified movement patterns in the Moray Firth, Scotland.  
762 *Journal of Applied Ecology*, 34(6), 1365-1374.

763 Wood, S.N. (2006) *Generalized Additive Models: An Introduction with R*. Boca Raton, FL:  
764 Chapman & Hall/CRC.

765 **Tables**

766 Table 1. Deployment and recovery months of the ECoMMAS array in the three years of data  
767 collection used in this study. In 2015 two consecutive deployments were undertaken.

Year	Deployment	Recovery
2013	June and July	October
2014	May	November
2015	April July	July November

768

769

770

771

772

773

774 Table 2. Daily acoustic occupancy rates (number of days with detections/days with acoustic coverage) for unprocessed C-POD data  
 775 (All) and detections classified as “broadband” by the classification system. Ninety-five percent binomial confidence intervals in  
 776 parenthesis. Black areas indicated C-PODs that were not recovered or failed to record data.

	2013		2014		2015	
	Occ. Rate (All)	Occ. Rate (Broadband)	Occ. Rate (All)	Occ. Rate (Broadband)	Occ. Rate (All)	Occ. Rate (Broadband)
Lat_05	0.19 (0.12 - 0.28)	0.00 (0.00- 0.04)	0.00 (0.00 - 0.04)	0.00 (0.00- 0.04)	0.20 (0.15 - 0.26)	0.00 (0.00- 0.03)
Lat_10	0.03 (0.01 - 0.08)	0.01 (0.00- 0.05)			0.04 (0.02 - 0.08)	0.01 (0.00- 0.04)
Lat_15	0.04 (0.01 - 0.09)	0.01 (0.00- 0.05)			0.02 (0.01 - 0.05)	0.00 (0.00- 0.02)
Hel_05	0.05 (0.02 - 0.12)	0.00 (0.00- 0.04)	0.12 (0.08 - 0.17)	0.03 (0.01 - 0.06)	0.14 (0.09 - 0.20)	0.07 (0.04 - 0.12)
Hel_10	0.00 (0.00 - 0.03)	0.00 (0.00- 0.03)			0.02 (0.01 - 0.06)	0.00 (0.00- 0.03)
Hel_15	0.01 (0.00 - 0.05)	0.00 (0.00- 0.03)	0.01 (0.00 - 0.06)	0.00 (0.00- 0.04)	0.00 (0.00 - 0.02)	0.00 (0.00- 0.02)
Cro_05	0.89 (0.80 - 0.94)	0.78 (0.68 - 0.86)			0.95 (0.91 - 0.97)	0.83 (0.77 - 0.87)
Cro_10	0.32 (0.17 - 0.52)	0.12 (0.04 - 0.30)	0.35 (0.26 - 0.46)	0.25 (0.17 - 0.35)	0.37 (0.27 - 0.48)	0.28 (0.19 - 0.39)
Cro_15	0.02 (0.01 - 0.08)	0.02 (0.01 - 0.08)	0.00 (0.00 - 0.04)	0.00 (0.00- 0.04)	0.04 (0.02 - 0.08)	0.03 (0.01 - 0.06)
SpB_05	0.22 (0.15 - 0.32)	0.13 (0.08 - 0.22)	0.21 (0.11 - 0.38)	0.09 (0.03 - 0.24)	0.14 (0.10 - 0.19)	0.08 (0.05 - 0.13)
SpB_10	0.00 (0.00 - 0.03)	0.00 (0.00- 0.03)			0.00 (0.00 - 0.05)	0.00 (0.00- 0.05)
SpB_15			0.01 (0.00 - 0.05)	0.01 (0.00- 0.05)	0.03 (0.01 - 0.06)	0.02 (0.01 - 0.05)
Fra_05	0.13 (0.08 - 0.21)	0.00 (0.00- 0.04)	0.21 (0.13 - 0.33)	0.20 (0.12 - 0.31)	0.11 (0.07 - 0.16)	0.06 (0.03 - 0.10)

Fra_10					0.00 (0.00 - 0.03)	0.00 (0.00- 0.03)
Fra_15			0.04 (0.02 - 0.10)	0.03 (0.01 - 0.08)	0.08 (0.05 - 0.13)	0.05 (0.03 - 0.09)
Cru_05	0.19 (0.13 - 0.26)	0.02 (0.00- 0.06)	0.04 (0.02 - 0.10)	0.01 (0.00- 0.05)	0.13 (0.07 - 0.22)	0.01 (0.00- 0.07)
Cru_10			0.15 (0.09 - 0.23)	0.04 (0.02 - 0.10)	0.00 (0.00 - 0.43)	0.00 (0.00- 0.43)
Cru_15	0.16 (0.11 - 0.23)	0.06 (0.03 - 0.10)	0.15 (0.09 - 0.23)	0.03 (0.01 - 0.09)		
Sto_05	0.17 (0.11 - 0.25)	0.10 (0.06 - 0.16)			0.36 (0.30 - 0.44)	0.27 (0.21 - 0.34)
Sto_10			0.12 (0.06 - 0.21)	0.05 (0.02 - 0.13)	0.07 (0.04 - 0.12)	0.04 (0.02 - 0.07)
Sto_15	0.30 (0.23 - 0.37)	0.11 (0.07 - 0.16)	0.10 (0.06 - 0.19)	0.01 (0.00- 0.06)	0.12 (0.06 - 0.20)	0.06 (0.03 - 0.14)
Abr_05	0.17 (0.11 - 0.26)	0.07 (0.03 - 0.14)	0.11 (0.06 - 0.18)	0.05 (0.02 - 0.12)	0.27 (0.18 - 0.38)	0.09 (0.04 - 0.17)
Abr_10	0.02 (0.01 - 0.08)	0.00 (0.00- 0.04)	0.02 (0.01 - 0.09)	0.00 (0.00- 0.05)	0.04 (0.02 - 0.08)	0.02 (0.01 - 0.06)
Abr_15	0.18 (0.13 - 0.25)	0.05 (0.03 - 0.10)			0.03 (0.01 - 0.06)	0.02 (0.01 - 0.05)
StA_05	0.18 (0.12 - 0.27)	0.09 (0.04 - 0.16)	0.07 (0.03 - 0.16)	0.03 (0.01 - 0.10)	0.07 (0.04 - 0.11)	0.03 (0.02 - 0.07)
StA_10	0.00 (0.00 - 0.04)	0.00 (0.00- 0.04)	0.01 (0.00 - 0.06)	0.01 (0.00- 0.06)	0.02 (0.01 - 0.09)	0.01 (0.00- 0.07)
StA_15	0.03 (0.01 - 0.08)	0.01 (0.00- 0.05)	0.02 (0.01 - 0.07)	0.01 (0.00- 0.06)	0.00 (0.00 - 0.05)	0.00 (0.00- 0.05)
Stb_05	0.05 (0.02 - 0.10 )	0.02 (0.01 - 0.07)	0.06 (0.03 - 0.12)	0.02 (0.01 - 0.07)	0.04 (0.01 - 0.10)	0.04 (0.01 - 0.1)
Stb_10	0.03 (0.01 - 0.09)	0.01 (0.00- 0.06)	0.02 (0.01 - 0.07)	0.02 (0.01 - 0.07)	0.02 (0.01 - 0.05)	0.01 (0.00- 0.04)
Stb_15	0.04 (0.02 - 0.08)	0.02 (0.01 - 0.06)			0.01 (0.00 - 0.07)	0.00 (0.00- 0.05)

777

778

779 Table 3 Temporal model selection results for the 10 deployment groups (Group). Formula  
 780 indicates the final model form selected via QIC for each deployment group (Equations 1-4).  
 781 Response for all models was P(Broadband). ShoreDist represents the three-level factor for  
 782 the near (05), mid (10), and offshore (15) deployment locations. Year is the two or three  
 783 level factor each survey year (2013, 2014 or 2015), Delta QIC is the difference in QIC  
 784 scores between selected model and the next best performing model. Group AUC is the area  
 785 under the ROC curve for each model applied to deployment groups. Group presence (Pres.)  
 786 and absence (Abs.) are the proportion of presences and absences correctly identified by the  
 787 model for each group. Unit is the location of each C-POD within the group and each  
 788 individual deployment location (Dep) AUC, Pres. and Abs. are the area under the curve and  
 789 proportion of presences and absences correctly predicted by the model for each of the C-  
 790 POD locations. Dashes indicated locations where modelling was not possible due to either  
 791 low numbers of detections or failure to recover the C-PODs deployed at that location.

792

Group Name	Formula Selected	Delta QIC	Group AUC	Pres.	Abs.	Dep. Name	Dep. AUC	Dep. Pres.	Dep. Abs.
Latheron	$P(\text{Broadband}) \sim \text{ShoreDist} + \text{Year} * \text{bs}(\text{JulianDay}, \text{knots} = \text{mean}(\text{JulianDay}))$	13.83	0.85	0	0.88	Lat_05	0.99	0.00	0.99
						Lat_10	0.82	0.01	0.59
						Lat_15	0.98	0.00	0.97
Helmsdale	$P(\text{Broadband}) \sim \text{ShoreDist} +$	3.51	0.93	0.01	0.77	Hel_05	0.82	0.02	0.78

	Year + JulianDay								
						Hel_10	0.94	0.00	0.93
						Hel_15			
	<i>P(Broadband)</i> ~ ShoreDist +								
	Year + bs(JulianDay, knots =	1.22	0.93	0.28	0.6	Cro_05	0.62	0.53	0.10
Cromarty	mean(JulianDay))					Cro_10	0.61	0.19	0.30
						Cro_15	0.78	0.01	0.80
	<i>P(Broadband)</i> ~ Year +								
	ShoreDist * bs(JulianDay,	5.34	0.81	0.04	0.63	SpB_05	0.63	0.05	0.63
Spey Bay	knots = mean(JulianDay))					SpB_10			
						SpB_15	0.75	0.01	0.55
	<i>P(Broadband)</i> ~ ShoreDist +								
	Year + JulianDay	2.52	0.75	0.04	0.48	Fra_05	0.70	0.03	0.81
Fraserburgh						Fra_10	0.99	0.00	0.98
						Fra_15	0.54	0.04	0.40
	<i>P(Broadband)</i> ~ ShoreDist +								
	Year + JulianDay	1.85	0.63	0.03	0.33	Cru_05	0.64	0.01	0.72
Cruden Bay						Cru_10	0.22	0.03	0.13
						Cru_15	0.61	0.02	0.82
	<i>P(Broadband)</i> ~ Year +								
	ShoreDist * bs(JulianDay,	5.73	0.79	0.06	0.77	Sto_05	0.71	0.13	0.63
Stonehaven									

	knots = mean(JulianDay))								
						Sto_10	0.63	0.02	0.67
						Sto_15	0.81	0.05	0.82
	P(Broadband)~ Year +								
	ShoreDist * bs(JulianDay,	1.39	0.82	0.03	0.51	Abr_05	0.61	0.04	0.58
Arborath	knots = mean(JulianDay))								
						Abr_10	0.98	0.01	0.96
						Abr_15	0.76	0.03	0.49
	P(Broadband)~ ShoreDist +								
	Year * bs(JulianDay, knots =	8.46	0.85	0.02	0.82	StA_05	0.83	0.04	0.72
St Andrews	mean(JulianDay))								
						StA_10	0.81	0.00	0.72
						StA_15	0.63	0.00	0.51
	P(Broadband)~ ShoreDist +								
	Year + JulianDay	3.06	0.62	0.01	0.63	Stb_05	0.51	0.02	0.18
St Abbs									
						Stb_10	0.63	0.00	0.92
						Stb_15	0.60	0.01	0.36

793

794 Table 4 GAMM summary for the parametric and smooth coefficient estimates, standard errors, estimated degrees of  
 795 freedom (EDF), reference degrees of freedom (Ref.df), F, t and p-values for the final habitat model. Smooth factors  
 796 (Distance to nearest Point Of Aggregation and Distance to Shore) are added as a main effect.

---

Model Formula

$P(\text{Broadband}) \sim s(\text{DistToPOA}, \text{bs} = \text{"ts"}, k = 3) + s(\text{Depth}, \text{bs} = \text{"ts"}) + s(\text{DistToShore}, \text{bs} = \text{"ts"}) + \text{POIName}$   
 $+ \text{Season} + \text{DistToPOA} + \text{DistToShore}$

---

Parametric coefficients

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	-2.70586	0.17708	-15.281	<.001
Dee	-0.63122	0.34879	-1.81	0.070
Esk	-1.02055	0.31654	-3.224	0.001
Spey	-1.24386	0.27673	-4.495	<.001
Tay Firth	-0.64207	0.35049	-1.832	0.067
Tweed	-2.41157	0.44156	-5.461	<.001
SeasonSpring	-0.06665	0.16495	-0.404	0.686
SeasonSummer	0.03319	0.12101	0.274	0.784

---

Approximate significance of smooth terms

	EDF	Ref.df	F	p-value
s(DistToPOA)	1.917	2	55.264	<.001
s(Depth_m)	4.686	9	6.233	<.001
s(DistToShore)	4.961	9	9.094	<.001

---

R-sq.(adj) = 0.322 , Scale est. = 1 , n = 9181


---

797

798


799 Figures

800



801 Figure 1. Study area including the Moray Firth Special Area of Conservation (yellow) and  
802 deployment locations of the East Coast Marine Mammal Acoustic Study (red points) and  
803 associated deployment group names.


804



805

806 Figure 2. Observed (points) and 95% confidence interval of the modelled occupancy (shaded) for  
 807 the 2013 near (05) mid (10) and offshore (15) deployment sites (panels). Colour indicates  
 808 distance from shore as a factor.

809




810

811 Figure 3. Observed (points) and 95% confidence interval of the modelled occupancy (shaded) for


812 the 2014 near (05) mid (10) and offshore (15) deployment sites (panels). Colour indicates

813 distance from shore as a factor.



815

816 Figure 4. Observed (points) and 95% confidence interval of the modelled occupancy (shaded) for  
 817 the 2015 near (05) mid (10) and offshore (15) deployment sites (panels). Colour indicates  
 818 distance from shore as a factor.



820

821 Figure 5. Two dimensional representations of the binomial smooths for the habitat GAMM.


822 Shaded area represents the probability of detecting a broadband echolocation click train on a

823 given C-POD as a function of the CPOD's distance to the nearest point of aggregation (top left),

824 deployment depth (top right) and distance to shore (bottom). Shaded areas represent 95%

825 confidence intervals and dashes on X-axis are rug plot of deployment variables.

826



827

828 Figure 6. Predicted broadband occupancy throughout the east coast habitat. Predictions based on  
 829 GAMM analysis of CPOD acoustic records from 2013-2015. Data are standardized to year 2015

830 and season is set to summer

Summer Acoustic Occupancy Model Predictions  
Model Average

