



<http://ijgt.ui.ac.ir/>

International Journal of Group Theory
ISSN (print): 2251-7650, ISSN (on-line): 2251-7669
Vol. x No. x (202x), pp. xx-xx.
© 202x University of Isfahan



www.ui.ac.ir

GRAPHS DEFINED ON GROUPS

PETER J. CAMERON

ABSTRACT. This paper concerns aspects of various graphs whose vertex set is a group G and whose edges reflect group structure in some way (so that, in particular, they are invariant under the action of the automorphism group of G). The particular graphs I will chiefly discuss are the power graph, enhanced power graph, deep commuting graph, commuting graph, and non-generating graph.

My main concern is not with properties of these graphs individually, but rather with comparisons between them. The graphs mentioned, together with the null and complete graphs, form a hierarchy (as long as G is non-abelian), in the sense that the edge set of any one is contained in that of the next; interesting questions involve when two graphs in the hierarchy are equal, or what properties the difference between them has. I also consider various properties such as universality and forbidden subgraphs, comparing how these properties play out in the different graphs.

I have also included some results on intersection graphs of subgroups of various types, which are often in a “dual” relation to one of the other graphs considered. Another actor is the Gruenberg–Kegel graph, or prime graph, of a group: this very small graph has a surprising influence over various graphs defined on the group.

Other graphs which have been proposed, such as the nilpotence, solvability, and Engel graphs, will be touched on rather more briefly. My emphasis is on finite groups but there is a short section on results for infinite groups. There are briefer discussions of general $\text{Aut}(G)$ -invariant graphs, and structures other than groups (such as semigroups and rings).

Proofs, or proof sketches, of known results have been included where possible. Also, many open questions are stated, in the hope of stimulating further investigation.

Communicated by Alireza Abdollahi.

MSC(2020): Primary: 20D60; Secondary: 05C25.

Keywords: power graph, commuting graph, cograph, generating graph.

Received: 04 March 2021, Accepted: 15 April 2021.

Article Type: Research Paper.

<http://dx.doi.org/110.22108/ijgt.2021.127679.1681>

1. Introduction

There are a number of graphs whose vertex set is a group G and whose edges reflect the structure of G in some way, so that the automorphism group of G acts as automorphisms of the graph. These include the commuting graph (first studied in 1955), the generating graph (from 1996), the power graph (from 2000), and the enhanced power graph (from 2007), all of which have a considerable and growing literature. A relative newcomer, not published yet, is the deep commuting graph.

This paper does not aim to be a survey of all these areas, which would be far too ambitious a task. Rather, I am interested in comparisons among the different graphs. In particular, there is a hierarchy containing the null graph, power graph, enhanced power graph, deep commuting graph, commuting graph, non-generating graph (if the group is non-abelian), and complete graph: the edge set of each is contained in that of the next.

These graphs have some similarities: for example, the enhanced power graphs, commuting graphs, deep commuting graphs, and generating graphs of finite groups all form universal families (that is, every finite graph is embeddable in one of these graphs for some group G). However, the proofs of this require rather different techniques for the different graphs.

Another question, about which relatively little is currently known, concerns the differences between graphs in the hierarchy. Even rather basic questions such as connectedness are unstudied for most of these, although Saul Freedman and coauthors have results on the difference between the non-generating graph and the commuting graph (and, at top and bottom, the difference between the complete graph and the non-generating graph is the generating graph, while the difference between the power graph and the null graph is the power graph, both of which have an extensive literature).

For some of these graphs, either the complementary graph was defined first, or the graph and its complement were studied independently. For example, the generating graph of a finite group preceded the non-generating graph; and Neumann's theorem, that if the non-commuting graph of an infinite group contains no infinite clique then it has finite clique number, clearly has an equivalent formulation in terms of the coclique number of the commuting graph. Also, several of the questions I consider have easy translations for the complement graph; for example, the universality results just mentioned immediately show that the complementary graphs are universal too. Twin reduction and the related concept of cograph do not distinguish between a graph and its complement. I have chosen to focus on those graphs which form a hierarchy; if the complements were preferred, the hierarchy would reverse.

A curious feature is the appearance of the Gruenberg–Kegel graph, which determines (or almost determines) various features of the commuting graph and the power graph. The vertex set of this graph is not the group, but the much smaller set of prime divisors of the group order. For example, if G has trivial centre, its reduced commuting graph (with the identity removed) is connected if and only if its Gruenberg–Kegel graph is. Conversely, for all graph types in the hierarchy except possibly the non-generating graph, the corresponding graph on G determines the Gruenberg–Kegel graph of G .

Authors who have studied these have used a variety of notations for them. I have tried to use a consistent and helpful notation, for example, $\text{Pow}(G)$ and $\text{Com}(G)$ for the power graph and commuting graph, respectively, of G .

The final, brief sections concern more general graphs defined on groups and invariant under group automorphisms; the graphs of the hierarchy on infinite groups; and extensions to other algebraic structures such as semigroups and rings.

Since much is not known, I have tried to emphasise open problems throughout.

Computations reported here were performed using GAP [55], with the packages GRAPE [104] for handling graphs, HAP [51] for computing Schur and Bogomolov multipliers, and LOOPS [89] for Moufang loops. Generators for specific groups were taken from the On-Line Atlas of Finite Groups [113].

I'm grateful to several people, especially Alireza Abdollahi, Saul Freedman, Michael Giudici, Michael Kinyon, Bojan Kuzma and Natalia Maslova, for helpful comments on a previous version.

1.1. Notation. I will denote a typical graph by Γ , with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. If A is a subset of $V(\Gamma)$, then the *induced subgraph* of Γ on A is the graph with vertex set A whose edges are those of Γ contained in A . A *complete graph* is one in which all pairs of vertices are joined; a *null graph* is one with no edges. A *clique* (resp. *coclique*) is a set of vertices on which the induced subgraph is complete (resp. null).

For a graph Γ ,

- the *clique number* $\omega(\Gamma)$ is the size of the largest clique;
- the *clique cover number* $\theta(\Gamma)$ is the minimum number of cliques whose union is the vertex set of Γ ;
- the *independence number* or *coclique number* $\alpha(\Gamma)$ is the size of the largest coclique;
- the *chromatic number* $\chi(\Gamma)$ is the smallest number of independent sets whose union is the vertex set of Γ (so-called because it is the smallest number of colours needed to colour the vertices so that adjacent vertices get different colours).

Note that the independence number and clique cover number of Γ are equal to the clique number and chromatic number of the complement of Γ (the graph with the same vertex set, whose edges are the non-edges of Γ).

It is clear that $\omega(\Gamma) \leq \chi(\Gamma)$ and $\alpha(\Gamma) \leq \theta(\Gamma)$. The graph Γ is called *perfect* if every induced subgraph has clique number equal to chromatic number. The *Weak Perfect Graph Theorem* of Lovász [83] asserts that, if Γ is perfect, then so is its complement; the *Strong Perfect Graph Theorem* of Chudnovsky *et al.* [43] asserts that Γ is perfect if and only if it does not contain a cycle of odd length greater than 3 or the complement of one as an induced subgraph.

The *comparability graph* of a partial order $P = (A, \leq)$ is the graph with vertex set A , in which a and b are joined if either $a \leq b$ or $b \leq a$. *Dilworth's Theorem* [49] asserts that the comparability graph of a partial order and its complement are both perfect. (The first part is easy, the second less so but follows from the first using the Weak Perfect Graph Theorem.)

Groups will almost always be finite here; my notation for finite groups is standard. (The reason for calling a graph Γ is to avoid conflict of notation, since G will be a typical group.)

1.2. Cayley graphs. One topic I will not consider, except in this section, concerns Cayley graphs. A *Cayley graph* for the group G is a graph on the vertex set G which is invariant under right translation by elements of G . (Some authors use left translation; the two concepts are equivalent, and the inversion map on G converts one into the other.) Equivalently, if S is an inverse-closed subset of $G \setminus \{1\}$, then the Cayley graph $\text{Cay}(G, S)$ is the graph with vertex set G in which g and h are adjacent whenever $gh^{-1} \in S$.

One reason for not considering these is that they have a huge literature, far more than I can survey here. I have heard the view expressed that algebraic graph theory is the study of Cayley graphs of finite groups (in fact, it is broader than this, but Cayley graphs are an important topic); while it is certainly arguable that geometric group theory is the study of Cayley graphs of finitely generated infinite groups.

The other is that Cayley graphs are not in general preserved by the automorphism group of G . I will say a few words about this.

Suppose that the set S is a *normal subset* of G , that is, closed under conjugation. Then $\text{Cay}(G, S)$ is invariant under both left and right translation. Such a graph is sometimes called a *normal Cayley graph*, see for example [71, 80]. However, the reader is warned that more recently this term has been used in a completely different sense: a Cayley graph $\Gamma = \text{Cay}(G, S)$ is normal if the group of right translations of Γ is a normal subgroup of $\text{Aut}(\Gamma)$, see for example [115].

To avoid confusion, I propose to call a normal Cayley graph in the first sense above an *inner-automorphic* Cayley graph. Note that this condition is equivalent to saying that the graph is invariant under both left and right translation. (For the composition of right translation by g and left translation by g^{-1} is conjugation by g , and so a graph invariant under two of these maps is invariant under the third also.)

Proposition 1.1. *The Cayley graph $\text{Cay}(G, S)$ is inner-automorphic if and only if S is a union of conjugacy classes in G .*

Note also that the minimal (non-null) inner-automorphic Cayley graphs for G are the relations of the *conjugacy class association scheme* on G , see [32, 58, 114] (the last of these references calls this structure the *group scheme* of G).

I shall call the Cayley graph $\text{Cay}(G, S)$ *automorphic* if it is invariant under the whole of $\text{Aut}(G)$, that is, if S is a union of orbits of $\text{Aut}(G)$ acting on G . (Thus, if $\text{Cay}(G, S)$ is automorphic, then its automorphism group contains the *holomorph* of G .) These graphs would fall under the rubric considered here, although I shall not be discussing them further.

2. Dramatis personae

This section introduces the specific graphs on a group that I will be mainly concerned with.

2.1. The commuting graph. Let G be a finite group. The *commuting graph* of G is the graph with vertex set G in which two vertices x and y are joined if $xy = yx$. This graph was introduced by Brauer and Fowler in their seminal paper [28] showing that only finitely many groups of even order can have a prescribed centraliser. (Brauer and Fowler do not use the word “graph”, but define the graph metric in the induced subgraph of the commuting graph on $G \setminus \{1\}$ and use this. The argument begins by showing that, if there are at least two conjugacy classes of involutions, then any two involutions have distance at most 3 by a path in the commuting graph which avoids the identity.)

The commuting graph has had further applications in group theory. Vertices in $Z(G)$ are joined to everything, and for investigating questions such as connectedness these are often removed; this makes no difference here.

Also the definition puts a loop at every vertex. There is good reason for doing this. It follows from results of Jerrum [73] on the “Burnside process” that the limiting distribution of the random walk on the commuting graph with loops is uniform on conjugacy classes – that is, the limiting probability of being at a vertex is inversely proportional to the size of its conjugacy class. This is useful in finding representatives of very small conjugacy classes in large groups.

Closely related is the fact that the *commuting ratio* of a group G , the probability that two randomly-chosen elements commute, is the ratio of the number of ordered edges of the commuting graph (including loops) to $|G|^2$: see for example [63, 50].

But for my purposes here, I will imagine that the loops have been silently removed.

As with all of these graphs, we can ask: Which groups are characterised by their commuting graphs? For example, abelian groups of the same order have isomorphic commuting graphs, as do the dihedral and quaternion groups of order 8. Figure 1 the commuting graphs of the two groups $D_8 = \langle a, b : a^4 = 1, b^2 = 1, b^{-1}ab = a^{-1} \rangle$ and $Q_8 = \langle a, b : a^4 = 1, b^2 = a^2, b^{-1}ab = a^{-1} \rangle$.




FIGURE 1. Commuting graph of D_8 or Q_8

It was conjectured in [4] and proved in [7, 65, 105] that any non-abelian finite simple group is characterised by its commuting graph.

I note in passing an application of the commuting graphs of finite groups to the structure of finite quotients of the multiplicative group of a finite-dimensional division algebra by Segev [100].

To conclude this section, a simple observation about the commuting graph:

Proposition 2.1. *A maximal clique in the commuting graph of G is a maximal abelian subgroup of G .*

2.2. The deep commuting graph. The *deep commuting graph* of a finite group G was introduced very recently [38]. Two elements of G are joined in the deep commuting graph if and only if their preimages in every central extension of G (that is, every group H with a central subgroup Z such that $H/Z \cong G$) commute. More specifically, take the commuting graph of a *Schur cover* [99] of G (this is a central extension H of largest order such that Z is contained in the derived group of H), and take the induced subgraph of the commuting graph of H on a transversal to Z . It can be shown that the resulting graph is independent of the choice of Schur cover.

For example, D_8 and Q_8 are Schur covers of the Klein group V_4 ; Figure 1 shows that the deep commuting graph of the Klein group is the star $K_{1,3}$, though its commuting graph is the complete graph K_4 .

We note in particular that the deep commuting graph is equal to the commuting graph if the *Schur multiplier* of G (the central subgroup Z in a Schur cover) is trivial. The converse is false, as we will see in Proposition 3.3.

Question 1. Is it true that a non-abelian finite simple group is characterised by its deep commuting graph?

2.3. The power graph. The *directed power graph* of G is the directed graph with vertex set G , with an arc $x \rightarrow y$ if $y = x^m$ for some integer m . The *power graph* of G is the graph obtained by ignoring directions and double arcs; in other words, x is joined to y if one of x and y is a power of the other. It is clearly a spanning subgraph of the commuting graph. The power graph was introduced by Kelarev and Quinn [76].

The directed power graph is a *partial preorder*, that is, a reflexive and transitive relation on G ; and the power graph is its *comparability graph* (two vertices joined if and only if they are related in the preorder). Comparability graphs of partial preorders and partial orders form the same class. For clearly every partial order is a partial preorder. For the converse, if (A, \leq) is a partial preorder, then the relation \equiv defined by $a \equiv b$ if $a \leq b \leq a$ is an equivalence relation; putting a total order on each equivalence class gives a partial order with the same comparability graph as (A, \leq) . It follows from Dilworth's theorem that the power graph of a finite group is perfect. This was first proved by Feng *et al.* [52]; see also [1].

The power graph does not uniquely determine the directed power graph; for example, if G is the cyclic group of order 6, then the identity and the two generators are indistinguishable in the power graph (they are joined to all other vertices), but one is a sink and the other two are sources in the directed power graph. However, the following is shown in [33]:

Theorem 2.2. *If two finite groups have isomorphic power graphs, then they have isomorphic directed power graphs.*

This is false for infinite groups; see [35].

For a more extensive survey of power graphs, I refer to [2].

2.4. The enhanced power graph. In 2007, Abdollahi and Hassanabadi [5, 6] studied a graph they called the *noncyclic graph* of a group G , in which two vertices x and y are joined if $\langle x, y \rangle$ is not cyclic. (For technical reasons they excluded the set of isolated vertices, the so-called *cyclicizer* of G .) Later in this paper, some of their results will be discussed.

For comparison with the other graphs considered in this paper, I will take the complement of their graph, which was independently defined in the paper [1] under the name *enhanced power graph* of G . Also, I will not initially assume that vertices joined to all others are excluded; we will examine such vertices later.

Thus, the enhanced power graph of a group G has vertex set G , with x and y joined if and only if $\langle x, y \rangle$ is cyclic. Equivalently, x and y are joined if there is an element $z \in G$ such that each of x and y is a power of z . This graph was introduced to interpolate between the power graph and the commuting graph, but has now been studied in its own right, especially by Samir Zahirović and coauthors [116, 117], and is in some respects easier to handle than the power graph.

The enhanced power graph can be obtained from the directed power graph by joining two vertices if both lie in the closed out-neighbourhood of some vertex. Thus, if two groups have isomorphic power graphs, then they have isomorphic enhanced power graphs. The converse is also true, see [117]:

Theorem 2.3. *For a pair of finite groups, the following are equivalent:*

- (a) *the power graphs are isomorphic;*
- (b) *the directed power graphs are isomorphic;*
- (c) *the enhanced power graphs are isomorphic.*

Here is a brief sketch. We are given the power graph of G . In the first step, we identify the maximal cyclic subgroups, as maximal cliques in the graph. In the next step, we identify further cyclic subgroups, as intersections of maximal cyclic subgroups. Given a maximal cyclic subgroup C , suppose that we obtain all its subgroups in this way, so that we can identify the sets of elements of each possible order in C . If $|C| = m$, then every cyclic subgroup has order a divisor r of m . To obtain the power graph, we remove edges between elements of orders r and s if neither divides the other; for the directed power graph, we direct the remaining edges from larger to smaller order. If not all the subgroups are determined, we are free to choose sets of the appropriate sizes, so long as our choices are consistent.

It is, however, not true that the power graph, directed power graph, and enhanced power graph of G have the same automorphism group. For $G = C_6$, all three automorphism groups are different.

Question 2. Is there a simple algorithm for constructing the directed power graph or the enhanced power graph from the power graph, or the directed power graph from the enhanced power graph?

Note that the enhanced power graph of G is a union of complete subgraphs on the maximal cyclic subgroups of G . Similarly, the commuting graph is a union of complete subgraphs on the maximal abelian subgroups. In fact, more is true:

Proposition 2.4. *A maximal clique in the enhanced power graph of G is a maximal cyclic subgroup of G .*

This follows from the fact that if a set of elements of G have the property that any two generate a cyclic subgroup, then the whole set generates a cyclic subgroup: see [1, Lemma 32].

At this point I mention another graph which I will not discuss in so much detail. It has been studied under the name *intersection graph* (see for example [42, 67]), but I wish to reserve this term for a different concept. Since, as we will see, it is dual (in a certain vague sense) to the enhanced power graph, I will call it the *dual enhanced power graph* of G , and denote it $\text{DEP}(G)$. Since the identity is always isolated in this graph (unlike the other graphs discussed so far), it is natural to remove it and define the vertex set of $\text{DEP}(G)$ to be $G \setminus \{1\}$.

Two non-identity elements x and y are joined in the dual enhanced power graph of G if $\langle x \rangle \cap \langle y \rangle$ is not the identity. Recall that two vertices x, y are joined in the enhanced power graph if they have a common in-neighbour in the directed power graph; we see that two vertices x and y are joined in $\text{DEP}(G)$ if they have a common out-neighbour different from the identity.

In particular, we see that the directed power graph determines the dual enhanced power graph. But, unlike the case with the power graph and enhanced power graph, it does not work in the reverse direction. If G and H are the cyclic group and quaternion group of order 8, then $\text{DEP}(G)$ and $\text{DEP}(H)$ are complete graphs on 7 vertices, but the directed power graphs of these two groups are not isomorphic.

2.5. The generating graph. The *generating graph* of a finite group G has vertex set G , with x and y joined if and only if $\langle x, y \rangle = G$. If the minimum number of generators of G is greater than 2, then the generating graph is the null graph. If G is cyclic, then its generating graph has loops; we will not be too much interested in this case. Note that, by the Classification of Finite Simple Groups, every non-abelian finite simple group is 2-generated.

The generating graph was introduced in [81], and studied further in [29]. The results are often phrased in terms of the *spread*, a graph-theoretic parameter defined as follows: Γ has spread (at least) k if every set of k vertices has a common neighbour. Thus a graph has spread 1 if it has no isolated vertex, while spread 2 is stronger than having diameter at most 2.

In [29] it was shown that the generating graph of a non-abelian finite simple group has positive *spread*. A substantial strengthening has recently been proved by Burness *et al.* [31]:

Theorem 2.5. *For a finite group G with reduced generating graph Γ , the following three conditions are equivalent:*

- (a) *any non-identity vertex has a neighbour in the generating graph (that is, Γ has spread 1);*
- (b) *any two non-identity vertices have a common neighbour in the generating graph (that is, Γ has spread 2);*
- (c) *any proper quotient of G is cyclic.*

In particular, the conditions hold for a non-abelian finite simple group. However, it is false for infinite groups: even (a) can fail for 2-generator infinite groups with all proper quotients cyclic [46].

In the sequel, I will often consider the *non-generating graph*, the complement of the generating graph.

2.6. The hierarchy. These graphs are given *ad hoc* names in the literature, but since I will be talking about all of them here, I prefer to give them names which help to distinguish them. Thus, the commuting graph of G will be $\text{Com}(G)$; the deep commuting graph $\text{DCom}(G)$; the power graph $\text{Pow}(G)$; the directed power graph $\text{DPow}(G)$; the enhanced power graph $\text{EPow}(G)$; the generating graph $\text{Gen}(G)$; and the non-generating graph $\text{NGen}(G)$. In each case, the vertex set is the group G . Sometimes I refer to the *reduced graph* of one of the above types, and denote it by a superscript $-$; this usually means that the identity element is deleted from the vertex set.

There are inclusions between these graphs, as follows. Here $E(\Gamma)$ denotes the edge set of a graph Γ ; thus $E(\Gamma_1) \subseteq E(\Gamma_2)$ means that Γ_1 is a *spanning subgraph* of Γ_2 (a subgraph using all of the vertices and some of the edges).

Proposition 2.6. *Let G be a finite group.*

- (a) $E(\text{Pow}(G)) \subseteq E(\text{EPow}(G)) \subseteq E(\text{DCom}(G)) \subseteq E(\text{Com}(G))$.
- (b) *If G is non-abelian or not 2-generated, then $E(\text{Com}(G)) \subseteq E(\text{NGen}(G))$.*

Proof. (a) All is obvious except possibly the inclusion of $E(\text{EPow}(G))$ in $E(\text{DCom}(G))$. So suppose that $\langle x, y \rangle$ is a cyclic subgroup of G , and let H be a central extension of G , with $H/Z \cong G$. The lift of $\langle x, y \rangle$ is the extension of a central subgroup Z by a cyclic group, and hence is abelian; so the lifts of x and y commute in H .

(b) If G is not 2-generated, then $\text{NGen}(G)$ is the complete graph, and the result is clear. If G is non-abelian, it cannot be generated by two commuting elements. □

Because of this, I will refer to the null graph, power graph, enhanced power graph, deep commuting graph, commuting graph, non-generating graph (in the case that G is non-abelian) and complete graph on G as the *graph hierarchy*, or just *hierarchy*, of G .

To conclude this section, I note that the dual enhanced power graph does not fit very well into the hierarchy. Recall that a superscript $-$ means that the identity is removed; the identity is isolated in $\text{DEP}(G)$ so it is natural to remove it from this graph also.

Proposition 2.7. (a) *For any finite group G , $E(\text{Pow}^-(G)) \subseteq E(\text{DEP}(G))$.*

- (b) *If $Z(G) = 1$, then $E(\text{DEP}(G)) \subseteq E(\text{NGen}^-(G))$.*
- (c) *In general $\text{DEP}(G)$ is incomparable with the other graphs in the hierarchy.*

Proof. (a) Suppose that $\{x, y\}$ is an edge of $\text{Pow}(G)$ with $x, y \neq 1$. Without loss of generality, x is a power of y ; then $\langle x \rangle \cap \langle y \rangle = \langle x \rangle$.

(b) Suppose that $Z(G) = 1$, and that $\{x, y\}$ is an edge of $\text{DEP}(G)$. Let $\langle x \rangle \cap \langle y \rangle = \langle z \rangle$, with $z \neq 1$. Then $x, y \in C_G(z) < G$, so $\langle x, y \rangle \neq G$.

(c) If $G = C_6$, then $\text{EPow}^-(G)$, and all higher graphs in the hierarchy, are complete, but $\text{DEP}(G)$ is not. If $G = Q_8$, then $\text{Com}^-(G)$, and all lower graphs in the hierarchy, are incomplete, but $\text{DEP}(G)$ is complete. \square

Question 3. For which groups G , and for which types X of graph in the hierarchy, does $\text{DEP}(G) = X^-(G)$ hold?

2.7. The Gruenberg–Kegel graph. A related graph will play a role in the investigation in several places. The *Gruenberg–Kegel graph*, also known as the *prime graph*, of a finite group G has vertex set the set of prime divisors of the order of G ; vertices p and q are joined by an edge if and only if G contains an element of order pq .

The graph was introduced in an unpublished manuscript by Gruenberg and Kegel to study the integral group ring of a finite group, and in particular the decomposability of the augmentation ideal: see [61]. The main structural result was published by Williams (a student of Gruenberg) in [112]. It asserts that groups whose Gruenberg–Kegel graph is disconnected have a very restricted structure.

Theorem 2.8. *Let G be a finite group whose Gruenberg–Kegel graph is disconnected. Then one of the following holds:*

- (a) G is a Frobenius or 2-Frobenius group;
- (b) G is an extension of a nilpotent π -group by a simple group by a π -group, where π is the set of primes in the connected component containing 2.

A 2-Frobenius group is a group G with normal subgroups H and K with $H \leq K$ such that

- K is a Frobenius group with Frobenius kernel H ;
- G/H is a Frobenius group with Frobenius kernel K/H .

A typical example is the group $G = S_4$, with $K = A_4$, $H = V_4$ (the Klein group), and $G/K \cong S_3$.

Williams went on to examine the known finite simple groups to determine which ones could occur in conclusion (b) of the Theorem. He could not handle the groups of Lie type in characteristic 2; this was completed by Kondrat'ev in 1989, and some errors corrected by Kondrat'ev and Mazurov in 2000.

The next result indicates that the Gruenberg–Kegel graph is closely connected with our hierarchy of graphs.

Theorem 2.9. *Let G_1 and G_2 be groups whose power graphs, or enhanced power graphs, or deep commuting graphs, or commuting graphs, are isomorphic. Then the Gruenberg–Kegel graphs of G_1 and G_2 are equal.*

Proof. The four possible hypotheses each imply that G_1 and G_2 have the same order, so their GK graphs have the same set of vertices.

We show that in all cases except the power graph, primes p and q are adjacent in the GK graph of G if and only if there is a maximal clique in the graph on G with size divisible by pq . This is clear

in the cases of the enhanced power graph and the commuting graph; for, as we observed earlier, the maximal cliques in these are maximal cyclic subgroups or maximal abelian subgroups of G respectively (Propositions 2.1 and 2.4), and if their order is divisible by pq then they contain elements of order pq . Conversely an element of order pq is contained in a maximal cyclic (or abelian) subgroup.

Consider the deep commuting graph of a group G . Let H be a Schur cover of G , with $H/Z \cong G$. A maximal clique has the form $A = B/Z$, where B is a maximal abelian subgroup of H (containing Z). So A is an abelian subgroup of G , and if pq divides $|A|$ then A contains an element of order pq . Conversely, suppose that p and q are joined in the GK graph, and let x and y be commuting elements of orders p and q in G , and a and b their lifts in H . Then a and b are contained in $\langle Z, ab \rangle$, which is an extension of a central subgroup by a cyclic group and hence is abelian; so a and b commute. Choosing a maximal abelian subgroup of H containing a and b and projecting onto G gives a maximal clique in $\text{DCom}(G)$ with order divisible by pq .

This fails for the power graph. Instead we use the fact that groups with isomorphic power graphs also have isomorphic enhanced power graphs, and so have equal GK graphs, by what has already been proved. \square

The proof gives a little more. Suppose, for example, that G and H are groups for which $\text{Com}(G)$ is isomorphic to $\text{EPow}(H)$. Then the Gruenberg–Kegel graphs of G and H are equal.

I do not know whether the analogous result holds for the non-generating graph of a non-abelian 2-generated group.

The Gruenberg–Kegel graph is an active topic of research; see [41] for a survey and some recent results. Some of the research concerns the question of whether a group is determined (perhaps up to finitely many possibilities) by its GK graph. There are two versions of this: two GK graphs could be equal (as graphs whose vertex set is a finite set of primes) or merely isomorphic as graphs but with possibly different labels for the vertices. For an interesting example, the GK graphs of the groups A_{10} and $\text{Aut}(J_2)$ are isomorphic, and both have vertex sets $\{2, 3, 5, 7\}$, but are not equal: the labels 2 and 3 are swapped.

2.8. Intersection graphs. Let G be a finite group, not trivial and not a cyclic group of prime order. The *intersection graph* of G is the graph whose vertices are the non-trivial proper subgroups of G , with two vertices H_1 and H_2 adjacent if $H_1 \cap H_2 \neq \{1\}$.

There are various other intersection graphs: we can restrict to subgroups in a particular class, or to maximal subgroups.

We will see a connection between some intersection graphs and some of the graphs in our hierarchy.

3. Equality and differences

For a non-abelian finite group G , there are seven graphs in the hierarchy, and a natural question is: When can two of them be equal? If they are not equal, what can be said about their difference?

3.1. **Equality.** At the two ends, things are easy:

Proposition 3.1. (a) $\text{Pow}(G)$ is equal to the null graph if and only if G is the trivial group.

(b) $\text{NGen}(G)$ is equal to the complete graph if and only if G is not 2-generated.

(c) $\text{NGen}(G) = \text{Com}(G)$ if and only if G is either abelian and not 2-generated, or a minimal non-abelian group.

Proof. Parts (a) and (b) are clear. So suppose that $\text{NGen}(G) = \text{Com}(G)$ and this is not the complete graph. Then G is non-abelian, but if x and y do not generate G then they commute; so every proper subgroup of G is abelian. Thus G is minimal non-abelian. \square

The minimal non-abelian groups were determined by Miller and Moreno [87] in 1903. There are two types: the first consists of groups of prime power order; the second are extensions of an elementary abelian p -group by a cyclic q -group, where p and q are primes.

Leaving aside the deep commuting graph from the present, the following was shown in [1]:

Proposition 3.2. Let G be a finite group.

(a) The power graph of G is equal to the enhanced power graph if and only if G contains no subgroup isomorphic to $C_p \times C_q$, where p and q are distinct primes; equivalently, the Gruenberg–Kegel graph of G is a null graph.

(b) The enhanced power graph of G is equal to the commuting graph if and only if G contains no subgroup isomorphic to $C_p \times C_p$, where p is prime; equivalently, the Sylow p -subgroups of G are cyclic or generalized quaternion groups.

(c) The power graph of G is equal to the commuting graph if and only if G contains no subgroup isomorphic to $C_p \times C_q$, where p and q are primes (equal or distinct).

Proof. (a) If G contains commuting elements of orders p and q , they are adjacent in $\text{EPow}(G)$ but not in $\text{Pow}(G)$. Conversely, suppose that x and y are adjacent in $\text{EPow}(G)$ but not in $\text{Pow}(G)$. Then x and y are contained in a cyclic group C but neither is a power of each other; C must then have order divisible by two distinct primes.

(b) If G contains commuting elements of the same prime order p but not in a cyclic subgroup of order p , they are joined in the commuting graph but not in the enhanced power graph. Conversely, suppose that x and y are adjacent in $\text{Com}(G)$ but not in $\text{EPow}(G)$. The orders of x and y must have a common factor (otherwise they generate a cyclic group); so some powers of them have prime order p and generate $C_p \times C_p$.

Now a theorem of Burnside (see [64, Theorem 12.5.2]) shows that a p -group containing no subgroup $C_p \times C_p$ is cyclic or generalized quaternion.

(c) The third part is immediate from the first two. \square

Using these results it is possible to classify the groups involved.

- (a) Any group of prime power order has Gruenberg–Kegel graph consisting of a single vertex, so has power graph equal to enhanced power graph. Any other group with this property has disconnected Gruenberg–Kegel graph, and so satisfies the conclusion of Theorem 2.8. These groups are the ones with the property that every element has prime power order; they are sometimes called *EPPO groups*. They, and some generalisations, were studied intensively in the 1960s and 1970s by Graham Higman and his students in Oxford (see for example [69]). Recently, Cameron and Maslova [41] have given a complete list of EPPO groups.
- (b) A group with all Sylow subgroups cyclic is metacyclic; indeed, if the primes dividing its order are p_1, p_2, \dots, p_r in increasing order, then it has a normal Hall subgroup corresponding to the last i primes in this list, for $1 \leq i \leq r - 1$.

By Glauberman's Z^* -theorem [57], if G has generalized quaternion Sylow 2-subgroup, and $O(G)$ is the largest normal subgroup of odd order in G , then $G/O(G)$ has a unique involution; the quotient \bar{G} by the subgroup generated by this involution has dihedral Sylow 2-subgroup, so falls into the classification by Gorenstein and Walter [59]. Of the groups in their theorem, we retain only those with cyclic Sylow subgroups for odd primes, that is, \bar{G} is isomorphic to $\text{PSL}(2, p)$ or $\text{PGL}(2, p)$ or to a dihedral 2-group. Conversely, each such group can be lifted to a unique group with a unique involution. The normal subgroup $O(G)$ has all its subgroups cyclic, so is metacyclic, as above.

Finally, the deep commuting graph lies between the enhanced power graph and the commuting graph. In order to investigate equality here, we need another construction. Recall that the Schur multiplier of G is the largest kernel Z in a stem extension H of G (with $Z \leq Z(H) \cap H'$ and $H/Z \cong G$). An extension is said to be *commutation-preserving*, or CP, if whenever two elements $x, y \in G$ commute, their preimages in H also commute. Now there is a well-defined largest kernel of a CP stem extension of G ; this is the *Bogomolov multiplier* of G , see [25, 74].

The Bogomolov multiplier first arose in connection with the work of Artin and Mumford on obstructions to Noethers conjecture on the pure transcendence of the field of invariants; it is also connected with other topics in number theory such as the Tate–Shafarevich set, and in group theory such as the coclass. However, we only need the definition given above.

Proposition 3.3. *Let G be a finite group.*

- (a) $\text{DCom}(G) = \text{EPow}(G)$ if and only if G has the following property: let H be a Schur cover of G , with $H/Z = G$. Then for any subgroup A of G , with B the corresponding subgroup of H (so $Z \leq B$ and $B/Z = A$), if B is abelian, then A is cyclic.
- (b) $\text{DCom}(G) = \text{Com}(G)$ if and only if the Bogomolov multiplier of G is equal to the Schur multiplier.

I refer to [38] for the proofs.

A precise characterisation of the groups attaining either equality is not known; but examples exist where one bound but not the other is met, or where neither bound is met (see [38]):

- If G is the symmetric or alternating group of degree at least 8, then $E(\text{EPow}(G)) \subset E(\text{DCom}(G)) \subset E(\text{Com}(G))$.
- If G is a dihedral group of order 2^n with $n \geq 3$, then $E(\text{EPow}(G)) = E(\text{DCom}(G)) \subset E(\text{Com}(G))$.
- If G is a certain group of order 64 (number 182 in the GAP library), then $E(\text{EPow}(G)) \subset E(\text{DCom}(G)) = E(\text{Com}(G))$.

Note that

- if the Schur multiplier of G is trivial, then $\text{DCom}(G) = \text{Com}(G)$;
- in general, the Bogomolov multiplier is much smaller than the Schur multiplier; for example, if G is a non-abelian finite simple group, then its Bogomolov multiplier is trivial [79].

Question 4. (a) What can be said about groups G for which $\text{DCom}(G) = \text{EPow}(G)$?

(b) What can be said about groups G for which $\text{DCom}(G) = \text{Com}(G)$?

3.2. Differences. For any pair of graphs in the hierarchy, if G is a group such that these two graphs are unequal, we could ask about the graph whose edge set is the difference. We could denote these by using, for example, $(\text{Com} - \text{Pow})(G)$ for the graph whose edges are those belonging to the commuting graph but not the power graph, with similar notation in other cases.

At the top, the difference between the complete graph and the non-generating graph is just the generating graph, which has been extensively studied. At the next level, the difference between the generating graph and the commuting graph (the graph $(\text{NGen} - \text{Com})(G)$) has been studied by Saul Freedman; the results will appear in his thesis. The most complete results are for nilpotent groups, and are reported in [34]. In particular, if G is nilpotent and the non-commuting non-generating graph is not null, then after deletion of all isolated vertices it is connected, with diameter 2 or 3.

Other differences (apart from the difference between the power graph and the null graph) have not been studied.

Question 5. For each pair of graph types in the hierarchy, what can be said about groups for which the difference is connected (after removing isolated vertices and vertices joined to all others)?

In section 9, I give a very weak partial result on the graph $(\text{Com} - \text{Pow})(G)$.

3.3. Further problems. We saw the result of [117] that two groups have isomorphic power graphs if and only if they have isomorphic enhanced power graphs.

Question 6. Are there any other implications of this kind between pairs of graphs in the hierarchy?

For a simple negative example, the groups C_{p^2} and $C_p \times C_p$ have isomorphic commuting graphs but nonisomorphic power graphs, while the group $C_p \times C_p \times C_p$ and the non-abelian group of order p^3 and exponent p have isomorphic power graphs but nonisomorphic commuting graphs.

Do there exist groups G_1 and G_2 such that, for example, $\text{Pow}(G_1)$ is isomorphic to $\text{Com}(G_2)$? This will be true if $\text{Pow}(G_1) = \text{Com}(G_1)$ and G_1 and G_2 have isomorphic commuting graphs, or if $\text{Pow}(G_2) = \text{Com}(G_2)$ and G_1 and G_2 have isomorphic power graphs.

Question 7. Can $\text{Pow}(G_1)$ and $\text{Com}(G_2)$ be isomorphic for groups G_1 and G_2 which both have power graph not equal to commuting graph? Similar questions for other pairs of graphs in the hierarchy.

4. Cliques and cocliques

It is clear that, if Γ_1 and Γ_2 share a vertex set and $E(\Gamma_1) \subseteq E(\Gamma_2)$, then $\omega(\Gamma_1) \leq \omega(\Gamma_2)$, $\chi(\Gamma_1) \leq \chi(\Gamma_2)$, $\alpha(\Gamma_1) \geq \alpha(\Gamma_2)$, and $\theta(\Gamma_1) \geq \theta(\Gamma_2)$. So these four parameters are monotonic for the graphs in the hierarchy on a given group (non-decreasing for ω and χ , non-increasing for α and θ).

Proposition 4.1. (a) $\omega(\text{EPow}(G))$ is the order of the largest cyclic subgroup of G ;
 (b) $\omega(\text{Com}(G))$ is the order of the largest abelian subgroup of G ;
 (c) $\omega(\text{DCom}(G))$ is the order of the largest subgroup of G whose inverse image in any central extension of G is abelian.

Proof. (a) and (b) follow from Propositions 2.1 and 2.4. For (c), apply (b) to a Schur cover of G . \square

The power graph of a group G is perfect, and so has equal clique number and chromatic number. These numbers do not exceed the clique number of the enhanced power graph, which is the largest order of an element of G ; but they may be smaller. For example, $\omega(\text{Pow}(C_6)) = 5$, but $\text{EPow}(C_6)$ is the complete graph K_6 .

As noted, $\omega(\text{Pow}(G)) \leq \omega(\text{EPow}(G))$. There is an inequality in the other direction:

Theorem 4.2. There is a function f on the natural numbers such that, for any finite group G , $\omega(\text{EPow}(G)) \leq f(\omega(\text{Pow}(G)))$.

Proof. Let m be the largest prime power which is the order of an element of G . The power graph of an m -cycle is complete; so $\omega(\text{Pow}(G)) \geq m$. On the other hand, the largest order of an element of G is not greater than the least common multiple of $\{1, \dots, m\}$, say $f(m)$; and so $\omega(\text{EPow}(G)) \leq f(m)$. \square

No such result holds for the commuting graph. If G is an elementary abelian group of order 2^n , then $\omega(\text{EPow}(G)) = 2$ but $\omega(\text{Com}(G)) = 2^n$.

The function f in the theorem is exponential. For there are $\pi(m) \leq (1 + o(1))m/\log m$ primes up to m , and $f(m)$ is the product of the largest power of each prime not exceeding m ; so $f(m) \leq m^{(1+o(1))m/\log m} = e^{(1+o(1))m}$. But the true value is probably much smaller.

Question 8. Find the best possible function f in Theorem 4.2.

An upper bound for the chromatic number of the enhanced power graph is given in [1, Theorem 12]:

Theorem 4.3. *Let G be a finite group, and S the set of orders of elements of G . Then*

$$\chi(\text{EPow}(G)) \leq \sum_{n \in S} \phi(n),$$

where ϕ is Euler's function.

Proof. For each $n \in S$, the set of elements of order n in G is the disjoint union of complete graphs of size $\phi(n)$, and can be coloured with $\phi(n)$ colours. If we use disjoint sets of colours for different orders, no further clashes will occur. \square

The bound is met for abelian groups. For if G is abelian, then S is the set of divisors of the exponent of G , and so the sum on the right is the exponent, which is the largest element order in G .

Question 9. Find a formula for the clique number of the power graph, or the chromatic number of the enhanced power graph, of a finite group.

Results about the independence number and clique cover number are less well developed. Since the power graph is perfect, the Weak Perfect Graph Theorem of Lovász [83] asserts that its complement is also perfect, so

$$\alpha(\text{Pow}(G)) = \theta(\text{Pow}(G))$$

for any finite group G . (Alternatively this follows from Dilworth's Theorem.)

The independence number of the non-generating graph of a finite group has been investigated by Lucchini and Maróti [84].

5. Induced subgraphs

In this section I will consider the question, for each of the graphs in our hierarchy: For which finite graphs Γ does there exist a finite group G such that Γ is isomorphic to an induced subgraph of the graph of that type defined on G ? (An *induced subgraph* of Γ on a subset A of the vertex set consists of the vertices of A and all edges of Γ which are contained in A .)

To summarise the results:

- A finite graph Γ is isomorphic to an induced subgraph of the power graph of some finite group G if and only if Γ is the comparability graph of a partial order.
- For each of the other graphs in the hierarchy, every finite graph is isomorphic to an induced subgraph of that graph defined on some finite group.

Three related questions are:

- Question 10.**
- (a) What is the smallest group for which a given graph is embeddable in the enhanced power graph/deep commuting graph/commuting graph/non-generating graph?
 - (b) What is the smallest group in which every graph on n vertices can be embedded in one of these graphs?

- (c) Which graphs occur if we restrict the group to have a particular property such as nilpotence or simplicity?

Here is a very rough lower bound for the smallest N such that every n -vertex graph can be embedded in the enhanced power graph, deep commuting graph, commuting graph, or non-generating graph of some group of order at most N . For our rough calculation, we need only consider groups of order N . It is known that there are at most $2^{c(\log N)^3}$ such groups (see [23]); each has at most N^n subsets of size n . But there are at least $2^{n(n-1)/2}/n!$ graphs on n vertices up to isomorphism. So we require

$$2^{c(\log N)^3} \cdot N^n \geq 2^{n(n-1)/2}/n!,$$

which implies that $N \geq 2^{n^{2/3-\epsilon}}$. So the exponential bound we find in some cases is not too far from the truth.

Note also that every n -vertex graph is embeddable in a Paley graph of order q , where q is a prime power congruent to 1 (mod 4) and $q > n^2 2^{2n-2}$ (see [24, 26]); so, to find a group whose commuting graph, etc., embeds all graphs of order n , we only need to embed this Paley graph.

5.1. The commuting graph.

Theorem 5.1. *Every finite graph is isomorphic to an induced subgraph of the commuting graph of a finite group. This group can be taken to be nilpotent of class 2 and exponent 4.*

Proof. Let F be the two-element field, V a vector space over F , and B a bilinear form on V . Define an operation \circ on $V \times F$ by the rule

$$(v_1, a_1) \circ (v_2, a_2) = (v_1 + v_2, a_1 + a_2 + B(v_1, v_2)).$$

It is a straightforward exercise to show that this operation makes $V \times F$ a group. This group is nilpotent of class 2 and exponent (dividing) 4, since $\{0\} \times F$ is a central subgroup with elementary abelian quotient. Moreover, (v_1, a_1) and (v_2, a_2) commute if and only if $B(v_1, v_2) = B(v_2, v_1)$.

Now a bilinear form is uniquely determined by its values on pairs of vectors taken from a basis for V ; these values can be assigned arbitrarily. So let Γ be a graph with vertex set $\{1, \dots, n\}$, and let v_1, \dots, v_n be a basis. Assign the values $B(v_i, v_j) = 0$ if $i \leq j$; for $i > j$, put $B(v_i, v_j) = 0$ if vertices i and j are adjacent, 1 if not. Then it is clear that the induced subgraph of the commuting graph on the set $\{v_1, \dots, v_n\}$ is isomorphic to Γ . □

The construction above shows that the smallest group whose commuting graph contains a given n -vertex graph has order at most 2^{n+1} if Γ has n vertices. However, it may be very much smaller; for the complete graph K_n , the answer is clearly n .

Using the remark before this subsection, we can build a group whose commuting graph embeds every n -vertex graph, by applying this construction to a sufficiently large Paley graph.

However, there is a more economical way to proceed, using a group from a familiar class, the *extraspecial p -groups*. The quaternion group Q_8 is generated by elements a, b, z satisfying $a^2 = b^2 = [a, b] = z$

and $z^2 = 1$; its centre is generated by z . Let G_n denote the central product of n copies of Q_8 . Then G_n is generated by elements a_i, b_i (for $1 \leq i \leq n$ and z , such that, for each i , a_i, b_i, z generate Q_8 , while a_i and b_i commute with a_j and b_j for $j \neq i$).

The group G_n has order 2^{2n+1} .

Theorem 5.2. *Every n -vertex graph is isomorphic to an induced subgraph of the commuting graph of G_n .*

Proof. The proof is by induction on n ; starting the induction is trivial. So suppose that the theorem is true for n , and let Γ be an $(n+1)$ -vertex graph. We can suppose that the induced subgraph on $\{v_1, \dots, v_n\}$ is already embedded as an induced subgraph of the commuting graph of the group G_n .

We modify this embedding as follows. We embed G_n in G_{n+1} in the obvious way, and replace the vertex v_i by $v_i a_{n+1}$ if v_i is not joined to v_{n+1} , leaving it as it is if these vertices are joined. Since a_{n+1} commutes with all v_i , this does not change the induced subgraph. Now we embed v_{n+1} as b_{n+1} . Clearly this gives an embedding of Γ . \square

This argument is based on a hint from Persi Diaconis, who has used similar ideas to study conjugacy classes in the Heisenberg group over a finite field and generalisations. (Recall that the random walk on the commuting graph has a limit which is uniform on conjugacy classes.) I do not have further details at present. Similar ideas can be found in the beautiful paper [48].

5.2. The deep commuting graph.

Theorem 5.3. *Every finite graph is isomorphic to an induced subgraph of the deep commuting graph of a finite group.*

Proof. Let Γ be a finite graph. As we have seen, Γ is isomorphic to an induced subgraph of the commuting graph of some group. So it is enough to show that this group can be chosen to have trivial Schur multiplier. Since the induced subgraph on a subgroup H of the commuting graph of G is the commuting graph of H , it suffices to show that every finite group can be embedded in a finite group with trivial Schur multiplier.

By Cayley's Theorem, every finite group of order n can be embedded in the symmetric group S_n . Unfortunately the symmetric group has Schur multiplier C_2 if $n \geq 8$. So we embed S_n into the general linear group $GL(n, 2)$ by permutation matrices of order n . Now the Schur multiplier of $GL(n, 2)$ is trivial except for $n = 3$ or $n = 4$ [60]. \square

5.3. The power graph. We saw that the power graph of G is the comparability graph of a partial order; hence any induced subgraph is also a comparability graph. The converse is also true:

Theorem 5.4. *A finite graph is isomorphic to an induced subgraph of the power graph of a finite group if and only if it is the comparability graph of a partial order. The group can be taken to be cyclic of squarefree order.*

Proof. One way round follows from our preliminary remarks: the power graph is the comparability graph of a partial order, and the class of such graphs is closed under taking induced subgraphs.

So suppose that we have a partial order \leq on X . For each $x \in X$, let $[x] = \{y \in X : y \leq x\}$. A routine check shows that

- $[y] \subseteq [x]$ if and only if $y \leq x$;
- $[x] = [y]$ if and only if $x = y$.

So the given partial order is isomorphic to the set of subsets of X of the form $[x]$, ordered by inclusion.

Now choose distinct prime numbers p_x for $x \in X$. Let G be the direct product of cyclic groups $C_{p_x} = \langle a_x \rangle$ of order p_x for $x \in X$. Map the subset Y of X to the element $g_Y = (g_x : x \in X)$ of the direct product, where

$$g_x = \begin{cases} a_x & \text{if } x \in Y, \\ 1 & \text{otherwise.} \end{cases}$$

It is readily checked that g_X and g_Y are adjacent in the power graph if and only if X and Y are adjacent in the comparability graph of the inclusion order on X .

To conclude, we note that G is a cyclic group of squarefree order. □

Note that a graph is the comparability graph of a partial order if and only if there is a transitive orientation of the edges. There is a list of forbidden induced subgraphs for comparability graphs [54], but it is not straightforward to state.

5.4. The enhanced power graph.

Theorem 5.5. *Every finite graph is isomorphic to an induced subgraph of the enhanced power graph of some group (which can be taken to be abelian).*

Proof. The proof is by induction. For a graph with a single vertex, there is no problem. So let Γ be a graph with vertex set $\{1, \dots, n\}$, and suppose that $i \mapsto x_i$ (for $i = 1, \dots, n - 1$) is an isomorphism to an induced subgraph of $\text{EPow}(G)$.

Choose a prime p not dividing the order of G , and let $H = \langle a, b \rangle$ be an elementary abelian group of order p^2 . Now in the group $G \times H$, replace x_i by $x_i a$ if i is not joined to n in Γ , and leave it as is if i is joined to n . Then map n to $x_n = b$.

Since $p \nmid |G|$, for any $z \in G$ we have $\langle z, a \rangle = \langle z \rangle \times \langle a \rangle$, which is cyclic. So the embedding of $\{1, \dots, n - 1\}$ is still an isomorphism to an induced subgraph. Moreover, $\langle x_i, b \rangle$ is cyclic while $\langle x_i a, b \rangle$ is not, so we have the correct edges from b to the other vertices, and the result is proved.

The resulting group is the product of n copies of $C_p \times C_p$ for distinct primes p . □

The dual enhanced power graph is even easier:

Theorem 5.6. *Every finite graph is isomorphic to an induced subgraph of the dual enhanced power graph of some group (which can be taken to be cyclic).*

Proof. Let E be the edge set of Γ ; choose distinct primes p_e for each edge $e \in E$, and let $C_{p_e} = \langle a_e \rangle$ be the cyclic group of order p_e , and G the direct product of all these cyclic groups. Now represent a vertex v by the element $b_v = \prod_{v \in e} a_e$. Then

$$\langle b_v \rangle \cap \langle b_w \rangle = \begin{cases} \langle a_e \rangle & \text{if } e = \{v, w\}, \\ \{1\} & \text{otherwise.} \end{cases}$$

So Γ is an induced subgraph of $\text{DEP}(G)$. □

5.5. The generating graph.

Theorem 5.7. *Every finite graph is isomorphic to an induced subgraph of the generating graph of a finite group.*

Proof. Let Γ be a finite graph. We proceed in a number of steps.

Step 1. Replace Γ by its complement.

Step 2. Every graph can be represented as the intersection graph of a *linear hypergraph*, a family of sets which intersect in at most one point (where intersection 1 corresponds to adjacency). The ground set E is the set of edges of the graph; the vertex v is represented by the set $S(v)$ of edges incident with v . Then for distinct vertices v, w ,

$$S(v) \cap S(w) = \begin{cases} e, & \text{if } \{v, w\} \text{ is an edge } e, \\ \emptyset & \text{if } v \text{ and } w \text{ are nonadjacent.} \end{cases}$$

Step 3. Add some dummy points, each lying in just one of the sets, so that they all have the same cardinality k , with $k \geq 3$. Now add some dummy points in none of the sets so that the cardinality n of the set Ω of points satisfies the conditions that $n > 2k$ and $n - k$ is prime.

Step 4. Now replace each set by its complement. The complements of two subsets of Ω have union Ω if and only if the two sets are disjoint. Thus, each original vertex is now represented by an $(n - k)$ -set where two such sets have union Ω if and only if the corresponding vertices are adjacent in Γ .

Step 5. Replace each set by a cyclic permutation on that set, fixing the remaining points. Each of these cycles has odd prime length, so each is an even permutation, and so lies in the alternating group A_n . Let g_v be the permutation corresponding to the vertex v of Γ .

- If v and w are nonadjacent, then the supports of g_v and g_w have union strictly smaller than Ω , so $\langle g_v, g_w \rangle \neq A_n$.
- Suppose v and w are adjacent. Then the supports of g_v and g_w have union Ω , so $H = \langle g_v, g_w \rangle$ is transitive on Ω . It is primitive: for each of g_v and g_w is a cycle of prime length $n - k > n/2$, and a block of imprimitivity either contains the cycle (and so has length greater than $n/2$, hence n) or meets it in one point (and so there are more than $n/2$ blocks, hence n blocks). Hence H is a primitive group of degree n containing a cycle of prime length p with $n/2 < p < n - 2$. By Jordan's theorem [111, Theorem 13.9], H contains the alternating group A_n . Since it is generated by even permutations, $H = A_n$.

Thus we have embedded Γ as an induced subgraph in the generating graph of A_n , as required. \square

5.6. Differences. We can also ask which graphs can be embedded in the graph whose edge set is the difference of the edge sets of two graphs in the hierarchy.

The proof that enhanced power graphs are universal uses abelian groups for the embedding. So, by embedding the complement, it shows:

Corollary 5.8. *Let Γ be a finite graph. Then there is a group G such that Γ is isomorphic to an induced subgraph of $\text{Com} - \text{EPow}(G)$.*

However, a much stronger result is true:

Theorem 5.9. *Let Γ be a finite complete graph, whose edges are coloured red, green and blue in any manner. Then there is an embedding of Γ into a finite group G so that*

- (a) *vertices joined by red edges are adjacent in the enhanced power graph;*
- (b) *vertices joined by green edges are adjacent in the commuting graph but not in the enhanced power graph;*
- (c) *vertices joined by blue edges are non-adjacent in the commuting graph.*

Proof. We begin with two observations. First, the direct product of cyclic (resp. abelian) groups of coprime orders is cyclic (resp. abelian).

Second, consider the non-abelian group of order p^3 and exponent p^2 , where p is an odd prime:

$$P = \langle a, b \mid a^{p^2} = b^p = 1, [a, b] = a^p \rangle.$$

Any two elements of $\langle a \rangle$ generate a cyclic group; and the group generated by b and x is cyclic if $x = 1$, abelian but not cyclic if $x = a^p$, and non-abelian if $x = a$.

The proof is by induction on the number n of vertices. The result is clearly true if $n = 1$. So let $\{v_1, \dots, v_n\}$ be the vertex set of Γ , and suppose that we have an embedding of $\{v_1, \dots, v_{n-1}\}$ into a group G satisfying (a)–(c).

Choose an odd prime p not dividing $|G|$, and consider the group $P \times G$, where P is as above. Modify the embedding of the first $n - 1$ vertices by replacing v_i by $(1, v_i)$ if $\{v_i, v_n\}$ is red, by (a^p, v_i) if $\{v_i, v_n\}$ is green, and by (a, v_i) if $\{v_i, v_n\}$ is blue. It is easily checked that we still have an embedding of $\{v_1, \dots, v_{n-1}\}$ satisfying (a)–(c). Moreover, if we now embed v_n as $(b, 1)$, we find that the conditions hold for the remaining pairs as well. \square

Clearly there are plenty of problems along similar lines to investigate here.

6. Products

There are a number of graph products. Here I will be chiefly concerned with the strong product, defined as follows.

Let Γ and Δ be graphs with vertex sets V and W respectively. The *strong product* $\Gamma \boxtimes \Delta$ has vertex set the Cartesian product $V \times W$; vertices (v_1, w_1) and (v_2, w_2) are joined whenever v_1 is equal or

adjacent to v_2 and w_1 is equal or adjacent to w_2 , but not equality in both places. (All of the graphs in the hierarchy naturally have loops at each vertex, which we have discarded; the strong product is the natural categorical product in the category of graphs with a loop at each vertex.)

I note that the strong product, along with the Cartesian and categorical products, is denoted by a symbol representing the corresponding product of two edges: the Cartesian product is $\Gamma \square \Delta$, while the categorical product is $\Gamma \times \Delta$.

The question of the perfectness of strong products of graphs has been studied by Ravindra and Parthasarathy [95].

The only group product that concerns us here is the direct product.

Proposition 6.1. *Let G and H be finite groups.*

- (a) $\text{Com}(G \times H) = \text{Com}(G) \boxtimes \text{Com}(H)$.
- (b) *If G and H have coprime orders, then $\text{EPow}(G \times H) = \text{EPow}(G) \boxtimes \text{EPow}(H)$.*
- (c) *If G/G' and H/H' have coprime orders, and in particular if G and H are perfect groups, then $\text{DCom}(G \times H) = \text{DCom}(G) \boxtimes \text{DCom}(H)$.*

Proof. (a) Distinct elements (g_1, h_1) and (g_2, h_2) in $G \times H$ commute if and only if g_1 and g_2 are equal or commute, and h_1, h_2 are equal or commute.

(b) Suppose that $|G|$ and $|H|$ are coprime. If $\langle g_1, g_2 \rangle$ and $\langle h_1, h_2 \rangle$ are cyclic, then (as their orders are coprime) their direct product is also cyclic and contains (g_1, h_1) and (g_2, h_2) . Conversely, again using coprimeness, if $\langle (g_1, h_1), (g_2, h_2) \rangle$ is cyclic, then it contains $(g_1, 1)$, $(g_2, 1)$, $(1, h_1)$ and $(1, h_2)$.

(c) A formula of Schur gives the Schur multiplier of G and H to be $M(G) \times M(H) \times (G \otimes H)$, where $M(G)$ is the Schur multiplier of G . If $|G/G'|$ and $|H/H'|$ are coprime, the third term is absent. It follows that a Schur cover of $G \times H$ is the direct product of Schur covers of G and H . The result follows. \square

Thus, questions about the commuting graph or enhanced power graph of a nilpotent group can be reduced to questions about the corresponding graphs for their Sylow subgroups.

The corresponding result fails for the power graph and the non-generating graph. The power graphs of C_2 and C_3 are complete but the power graph of $C_2 \times C_3$ are not. For the non-generating graph, we note that for any non-abelian finite simple group G , there is an integer m such that G^m fails to be 2-generated if $n > m$.

7. Cographs and twin reduction

Two vertices in a graph are called twins if they have the same neighbours (possibly excluding one another). Equivalently, v and w are twins if the transposition (v, w) (fixing the other vertices) is an automorphism of Γ . If G is a non-trivial group, then any of the graphs in our hierarchy based on G will contain many pairs of twins. Thus, twins will play an important part when we come to look at automorphism groups.

If a graph has twins, then we can make a new graph by merging the twins to a single vertex. The process can be continued until no pairs of twins remain. If the resulting graph has just a single vertex, the original graph is called a cograph.

Cographs also play an important part in the story, and make another link with the Gruenberg–Kegel graph. So we make a detour to look at twin reduction and cographs.

7.1. Cographs. A graph Γ is a *cograph* if either of the following equivalent conditions holds for it:

- Γ does not contain the four-vertex path P_4 as an induced subgraph;
- Γ can be constructed from the 1-vertex graph by the operations of complement and disjoint union.

In particular, a cograph is connected if and only if its complement is disconnected. This leads to a tree representation of cographs and to very efficient algorithms for determining their properties.

I remark here that a connected component of a cograph has diameter at most 2, since two vertices at distance 3 would be joined by an induced 4-vertex path.

Here is a curious fact which may (or may not) have a connection with the following material. The P_4 -structure of a graph Γ is the hypergraph whose vertex set is the same as that of Γ , the hyperedges being the 4-element sets which induce a copy of P_4 in Γ . We have the following easy observations:

- a graph and its complement have the same P_4 -structure, since the graph P_4 is self-complementary;
- a graph is a cograph if and only if its P_4 -structure is the null hypergraph.

Bruce Reed [96] proved the *semi-strong perfect graph theorem*, which had been conjectured by Vasek Chvátal, asserting that if two graphs have isomorphic P_4 -structures and one is perfect, then so is the other.

Cographs have been rediscovered a number of times, and as a result appear in the literature with very different names, such as “complement-reducible graphs”, “hereditary Dacey graphs” and “N-free graphs”. See [101, 107, 75] for information about cographs.

7.2. Twins and twin reduction. In a graph Γ , we can define two kinds of “twin relations” on vertices. The *open neighbourhood* $\Gamma(v)$ of v in Γ is the set of vertices in Γ joined to v ; the *closed neighbourhood* is $\Gamma(v) \cup \{v\}$. Two vertices v, w are *open twins* if they have the same open neighbourhoods; they are *closed twins* if they have the same closed neighbourhoods. Both of these relations are obviously equivalence relations. Two vertices are open twins in Γ if and only if they are closed twins in the complement of Γ . Note that open twins are not joined, while closed twins are joined.

If x and y are twins, then we may collapse them to a single vertex; this process is called *twin reduction*. Some, though clearly not all, properties of a graph are preserved by twin reduction. One such property is that of being a cograph, since twin reduction can neither create nor destroy an induced 4-vertex path (since no two of its vertices can be twins). We will see a more general result shortly.

7.3. The cokernel of a graph.

Theorem 7.1. *Given a graph Γ , the result of performing a sequence of twin reductions until the graph is twin-free is unique up to isomorphism, independent of the chosen sequence of reductions.*

Proof. Open and closed twin classes of sizes greater than 1 are disjoint. For suppose that x and y are open twins and y and z are closed twins. Then xy is a non-edge while yz is an edge. Since y and z are twins, x is not joined to z ; but since y and x are twins, z is joined to x . These conclusions are contradictory. (Alternatively, since y and z are twins, (y, z) is an automorphism, and so x and z are open twins; and similarly using the automorphism (x, y) , x and z are closed twins.)

We are going to prove the theorem by induction on the number of vertices. There is nothing to do for graphs with a single vertex, so let Γ have n vertices, with $n > 1$, and assume that the result is true for any graph with fewer than n vertices. Take two twin reduction sequences on Γ . Suppose that the first begins by identifying x and y , and the second by identifying u and v .

If $\{x, y\} = \{u, v\}$, then the two sequences result in the same graph, and induction finishes the job.

If $|\{x, y\} \cap \{u, v\}| = 1$, then our initial remark shows that the two pairs of twins have the same type, so the graphs obtained after one step are isomorphic, and again induction finishes the job.

Suppose that $\{x, y\} \cap \{u, v\} = \emptyset$. Then the two reductions commute. Let Δ be the graph obtained by applying the two reductions. Then Δ occurs after two steps in reduction sequences for Γ beginning by identifying x and y , or by identifying u and v . By induction the end result of either given sequence is the same as the result of reducing Δ (up to isomorphism).

The theorem is proved. □

The next result gives the connection between cographs and twin reduction.

Proposition 7.2. *A graph Γ is a cograph if and only if the cokernel of Γ is the graph with a single vertex.*

Proof. For the necessity, we show by induction that a cograph with more than one vertex contains twins. Let Γ be a cograph with more than one vertex, and suppose that any smaller cograph with more than one vertex contains twins. If Γ is disconnected, then if it has a component with more than one vertex, then this component contains twins; otherwise Γ is a null graph and all pairs of vertices are open twins. If Γ is connected, then its complement is disconnected, and we argue in the complement instead.

Now the result of twin reduction is an induced subgraph of Γ , and so also a cograph; so so the reduction continues until only one vertex remains.

Conversely, suppose that Γ is not a cograph. Then Γ contains a 4-vertex path, say (w, x, y, z) . Then any pair of these vertices are not twins, and so are not identified in any twin reduction; so the result of the reduction still contains a 4-vertex path. So no sequence of reductions can terminate in a single vertex. □

This gives another test for a cograph: apply twin reductions until the process terminates, and see whether just one vertex remains.

7.4. Cographs as comparability graphs. A partially ordered set is *N-free* if it does not contain as an induced substructure the poset whose Hasse diagram is shown in Figure 2.




FIGURE 2. The poset N

N-free posets form the smallest class of posets containing the 1-element poset and closed under disjoint union and ordered sum. (In the disjoint union of two posets, elements of different posets are incomparable; in the ordered sum, every element of the first is smaller than every element of the second.)

N-free posets arise in statistical design, where the operations of *crossing* and *nesting* correspond to disjoint union and ordered sum. An orthogonal block structure is *special* if it can be built by crossing and nesting. These constructions also apply to association schemes. Crossing and nesting were introduced by Nelder [90] and the more general poset constructions by Speed and Bailey [106]; see [16] for more detail and historical comments.

Proposition 7.3. (a) *The comparability graph of an N-free poset is a cograph.*

(b) *Conversely, any cograph is the comparability graph of an N-free poset.*

Proof. The first statement is immediate, and the second is easily proved by induction. □

7.5. Twin reduction in the hierarchy. The relevance of twin reduction to our problem is:

Proposition 7.4. *Let Γ be the power graph, enhanced power graph, deep commuting graph, commuting graph, or non-generating graph of a non-trivial group G . Then the twin relation on Γ is not the relation of equality.*

Proof. Suppose that G contains an element g of order greater than 2. Let h be an element such that $g \neq h$ but $\langle g \rangle = \langle h \rangle$. Then any element joined to one of g and h in one of the graphs listed is also joined to the other. The arguments are all easy; let us look at the least trivial, the deep commuting graph. Let H be a Schur cover of G with kernel Z , and x and y elements of H covering g and h respectively. Then $\langle Z, x \rangle$ is abelian and contains y , so x and y commute.

The groups not covered by this are elementary abelian 2-groups. In these cases, everything is clear: the power graph, enhanced power graph and deep commuting graph are stars; the commuting graph is complete; the non-generating graph is complete if the group has order greater than 4. All these graphs are cographs. □

So, for any of our classes of graphs, say X , the question “What is the cokernel of $X(G)$?” is a generalisation of “Is $X(G)$ a cograph?”

Finally for this section, I note that the class of cographs is not preserved by strong product. Figure 3 shows P_4 as an induced subgraph of $P_3 \boxtimes P_3$.




FIGURE 3. Cographs are not closed under \boxtimes

7.6. Forbidden subgraphs and twin reduction. For a graph F , let $\text{Forb } F$ denote the class of graphs containing no induced subgraph isomorphic to F .

If F is twin-free, then it is clear that identifying a pair of twins in a graph Γ can neither create nor destroy an induced copy of F . So the class $\text{Forb}(F)$ and its complement are both preserved by twin reduction, and in particular:

Theorem 7.5. *If F is twin-free, then a graph belongs to $\text{Forb } F$ if and only if its cokernel belongs to $\text{Forb } F$.*

The theorem immediately extends from a single graph F to an arbitrary class \mathcal{F} of graphs.

The Strong Perfect Graph Theorem shows that perfect graphs form the class $\text{Forb}(\mathcal{F})$, where \mathcal{F} consists of all cycles of odd length greater than 3 and their complements. All these graphs are twin-free. So as a corollary of the theorem, we see:

Corollary 7.6. *A graph is perfect if and only if its cokernel is perfect.*

This has the immediate consequence that cographs are perfect, although this is easily proved directly.

However, there are interesting graph classes defined by forbidden subgraphs which are not twin-free, including chordal graphs, split graphs, and threshold graphs. We now turn to the general question.

8. Forbidden subgraphs

In this section we consider various classes of graphs defined by forbidden induced subgraphs, and ask: for which finite groups G can one of the graphs in the hierarchy defined on G belong to this class? Not much is known.

We saw that a graph is perfect if and only if it has no induced subgraph isomorphic to an odd cycle of length greater than 3 or the complement of one. In particular, a cograph is perfect.

We have seen that power graphs are comparability graphs of partial orders, and noted that these graphs are perfect. For our other types of graph, no proper subclass defined by forbidden induced subgraphs contains them all; so we have to ask a different question: which groups have the property that one of these graphs belongs to a graph class defined in this way?

8.1. Power graphs. We begin with the question: When is the power graph of the group G a cograph? This question was considered in the paper [40]; here the Gruenberg–Kegel graph makes another appearance. The question is answered completely for nilpotent groups, but a necessary and a sufficient

condition are known for general groups in terms of the GK graph; these conditions do not coincide, and we will see that no condition just in terms of the GK graph can be both necessary and sufficient.

I begin with a simple observation.

Proposition 8.1. *Let G be a group of prime power order. Then $\text{Pow}(G)$ is a cograph.*

Proof. We use the fact that, if G is cyclic of prime power order, then $\text{Pow}(G)$ is the complete graph. Suppose that (a, b, c) is a 3-vertex induced path. In $\text{DPow}(G)$, we cannot have $a \rightarrow b \rightarrow c$, since transitivity would imply $a \rightarrow c$, contrary to assumption; similarly, $c \rightarrow b \rightarrow a$ is impossible. If $b \rightarrow a$ and $b \rightarrow c$, then $a, c \in \langle b \rangle$, a cyclic p -group, and so $a \sim b$ in the power graph, also a contradiction. So the only possibility is $a \rightarrow b$ and $c \rightarrow b$.

Now suppose that (a, b, c, d) is a 4-vertex path. Then considering the subpath (b, c, d) shows that $b \rightarrow c$ and $d \rightarrow c$. But $a \rightarrow b \rightarrow c$ is impossible; so no such path can exist. \square

Theorem 8.2. *Let G be a finite nilpotent group. Then $\text{Pow}(G)$ is a cograph if and only if either*

- (a) G has prime power order; or
- (b) G is cyclic of order pq , where p and q are distinct primes.

Theorem 8.3. (a) *Let G be a finite group whose Gruenberg–Kegel graph is a null graph. Then the power graph of G is a cograph.*

- (b) *Let G be a finite group whose power graph is a cograph. Then, with possibly one exception, a connected component of the Gruenberg–Kegel graph has at most two vertices, the exception being the component containing the prime 2. If $\{p, q\}$ is a connected component of the GK graph, with p and q odd primes, then p and q divide $|G|$ to the first power only.*

For the final claim, note that the groups $\text{PSL}(2, 11)$ and M_{11} have the same GK-graph (an edge $\{2, 3\}$ and isolated vertices $\{5\}$ and $\{11\}$), but the power graph of the first is a cograph, that of the second is not.

Question 11. Classify the groups whose power graph is a cograph.

The next question is maybe not so important in itself but might be an indication of how well we understand groups whose power graph is a cograph.

Question 12. Is the following true? Let Γ be a cograph. Then there is a finite group G such that $\text{Pow}(G)$ is a cograph and contains Γ as an induced subgraph.

There are two reasons why this might be tricky:

- The inductive scheme for cographs requires complementation. But this does not work nicely for power graphs.
- Theorem 8.3 shows that there will be no simple construction using direct products of groups of coprime order.

Probably this question will involve groups of prime power order.

The remainder of this subsection is based on the paper [40].

We saw that the power graph of a finite group is a comparability graph of a partial order, and so in particular is a perfect graph. Various interesting subclasses of the perfect graphs are defined by forbidding certain induced subgraphs. Cogroups form an example: they forbid the 4-vertex path P_4 . Here are some other graph classes.

- A graph is *chordal* if it contains no induced cycles of length greater than 3 (that is, every cycle of length greater than 3 has a chord).
- A graph is *split* if the vertex set is the disjoint union of two subsets, one inducing a complete graph and the other a null graph (with possibly some edges between them). A graph is split if it contains no induced subgraph isomorphic to C_4 , C_5 or $2K_2$.
- A graph is *threshold* if it can be constructed from the 1-vertex graph by adding vertices joined either to all or to no existing vertices. A graph is threshold if and only if it contains no induced subgraph isomorphic to P_4 , C_4 , or $2K_2$.

Theorem 8.4. *For a finite nilpotent group G , the power graph of G is chordal if and only if one of the following conditions holds:*

- (a) G has prime power order;
- (b) G is the direct product of a cyclic group of p -power order and a group of exponent q , where p and q are distinct primes.

Question 13. Which non-nilpotent groups have the property that the power graph is chordal?

Theorem 8.5. *The following conditions for a finite group are equivalent:*

- (a) $\text{Pow}(G)$ is a threshold graph;
- (b) $\text{Pow}(G)$ is a split graph;
- (c) $\text{Pow}(G)$ contains no induced subgraph isomorphic to $2K_2$;
- (d) G is cyclic of prime power order, or an elementary abelian or dihedral 2-group, or cyclic of order $2p$, or dihedral of order $2p^n$ or $4p$, where p is an odd prime.

Note that this theorem does not assume that G is nilpotent.

8.2. Other graphs. For our other classes of graphs, the problem of deciding for which groups the graph in question forbids a certain induced subgraph has not been much worked on, and a number of interesting questions are open. One difference is that, as we have seen, every finite graph occurs as an induced subgraph of each type of graph on some finite group.

In line with the preceding subsection we could ask a multipart question:

Question 14. For which finite groups is the enhanced power graph/deep commuting graph/commuting graph/nongenerating graph a perfect graph, or a cogroup, or a chordal graph, or a split graph, or a threshold graph?

We can give a partial answer for groups of prime power order.

Theorem 8.6. *Let G be a group of prime power order.*

- (a) *The power graph of G is equal to the enhanced power graph, and contains no induced P_4 or C_4 .*
- (b) *If G is 2-generated of order p^n , then the non-generating graph of G consists of $p + 1$ complete subgraphs of order p^{n-1} , any two intersecting in the same subset of size p^{n-2} .*

Proof. (a) In a group of prime power order, if two elements generate a cyclic subgroup, then one is a power of the other; this shows that the power graph and enhanced power graph coincide. Suppose that (x, y, z) is an induced path of length 2. If $x, z \in \langle y \rangle$, or if $y \in \langle x \rangle$ and $z \in \langle y \rangle$, or if $x \in \langle y \rangle$, $y \in \langle z \rangle$, then x and z are joined, a contradiction. So $y \in \langle x \rangle \cap \langle z \rangle$. Now if w is joined to x but not to y , then (w, x, y) is an induced path of length 2, so $x \in \langle y \rangle$ by the same argument. But then $y \in \langle z \rangle$, so $x \in \langle z \rangle$, a contradiction.

(b) Let $\Phi(G)$ be the Frattini subgroup of G . By the Burnside Basis Theorem, $\Phi(G)$ consists of vertices lying in no generating pair and has index p^2 in G ; moreover, the generating pairs are all pairs of elements lying in distinct non-trivial cosets of $\Phi(G)$. \square

No such result holds for the commuting graph; Theorem 5.1 shows that the commuting graphs of 2-groups form a universal class. Computation shows that the smallest group whose commuting graph is not a cograph is the symmetric group S_4 : the elements $(1, 2, 3, 4)$, $(1, 3)(2, 4)$, $(1, 2)(3, 4)$ and $(1, 3, 2, 4)$ induce a 4-vertex path. In fact, seven groups of order 32 have commuting graphs which are not cographs.

Question 14 has been considered for the commuting graph by Britnell and Gill [30], who obtained a partial description of groups for which the commuting graph is a perfect graph. Assuming that G has a component (a subnormal quasisimple subgroup), they determine all possible components of such groups.

Also, the paper [117] classifies nilpotent groups whose enhanced power graphs are perfect (but I have not been able to access this paper).

Question 15. What about other graph classes, for example planar graphs?

For planarity, this may not be too hard for graphs in the hierarchy. Since the complete graph on 5 vertices is not planar, it follows that G has no elements of order greater than 4. (For all cases except the power graph, a cyclic subgroup induces a complete graph, so the claim is clear. In the power graph, the power graph of a cyclic group of prime power order is complete, and the power graph of C_6 contains a K_5 .) So G is solvable.

Indeed, bounding the genus of a graph bounds its clique number, and so (for graphs in the hierarchy) bounds the orders of elements.

See [9] for the commuting graph and its complement.

8.3. Results for simple groups.

Proposition 8.7. *Let $G = \text{PSL}(2, q)$, with q a prime power and $q \geq 4$.*

- (a) *If q is even, then $\text{EPow}(G)$, $\text{DCom}(G)$ and $\text{Com}(G)$ are cographs; $\text{Pow}(G)$ is a cograph if and only if $q - 1$ and $q + 1$ are either prime powers or products of two distinct primes.*
- (b) *If q is odd, then $\text{EPow}(G)$ and $\text{DCom}(G)$ are cographs; $\text{Pow}(G)$ is a cograph if and only if $(q - 1)/2$ and $(q + 1)/2$ are either prime powers or products of two distinct primes.*

Proof. We consider the graphs obtained by removing the identity. We note that a graph Γ is a cograph if and only if the graph obtained by adding a vertex joined to all others is a cograph.

We begin with the case when q is a power of 2, noting that in this case $\text{PSL}(2, q)$ has trivial Schur multiplier except when $q = 4$. The elements of the group have order 2 or prime divisors of $q - 1$ or $q + 1$. The centralisers of involutions are elementary abelian of order q , while the centralisers of other elements are cyclic of order $q - 1$ and $q + 1$. In other words, centralisers are abelian and intersect only in the identity. So the commuting graph (which is the deep commuting graph if $q \neq 4$) is a disjoint union of complete graphs; the enhanced power graph is a disjoint union of complete graphs and isolated vertices (corresponding to the involutions); and the power graph is a disjoint union of the power graphs of cyclic groups of orders $q \pm 1$ and isolated vertices. Using the fact that the power graph of the cyclic group C_m is a cograph if and only if m is either a prime power or the product of two primes [40, Theorem 3.2], the result follows. The result for the deep commuting graph for $\text{PSL}(2, 4)$ can be proved by computation, or by identifying this group with $\text{PSL}(2, 5)$ (which is dealt with in the next paragraph).

Now consider the case when q is a power of an odd prime p . The centralisers of elements of order p are elementary abelian of order q ; centralisers of other elements are cyclic of orders $(q \pm 1)/2$ except for involutions, which are centralised by dihedral groups of order $q \pm 1$, whichever is divisible by 4. So, although centralisers may not be disjoint, the maximal cyclic subgroups are, so the statements about the enhanced power graph and the commuting graph are true.

If q is odd and $q \neq 9$, the Schur multiplier of $\text{PSL}(2, q)$ is cyclic of order 2, and so a Schur cover of this group is $\text{SL}(2, q)$. The unique involution in this group is $-I$, the kernel of the extension; so involutions in $\text{PSL}(2, q)$ lift to elements of order 4, which lie in unique maximal abelian subgroups.

The group $\text{PSL}(2, 9)$ has Schur multiplier of order 6. The deep commuting graph of $\text{PSL}(2, 9)$ was analysed using GAP, with generators for the Schur cover from the on-line Atlas of Finite Group Representations [113]. □

Question 16. Are there infinitely many prime powers q for which the power graph of $\text{PSL}(2, q)$ is a cograph?

Here is a preliminary analysis of this question.

Case q even. Then $q = 2^d$, say, and each of $q + 1$ and $q - 1$ is either a prime power or the product of two primes. By Catalan's conjecture (now Mihăilescu's Theorem: see [44, Section 6.11]), the only two proper powers differing by 1 are 8 and 9. So, unless $q = 8$, we conclude that each of $q + 1$ and $q - 1$ is

either prime or the product of two primes. Moreover, one of these numbers is divisible by 3, so we can say further that one of $q + 1$ and $q - 1$ is three times a prime, while the other is a prime or the product of two primes. For example, $2^{11} - 1 = 23 \cdot 89$ while $2^{11} + 1 = 3 \cdot 683$. The values of d up to 200 for which $q = 2^d$ satisfies the condition are 1, 2, 3, 4, 5, 7, 11, 13, 17, 19, 23, 31, 61, 101, 127, 167, 199.

Case q odd. If q is congruent to $\pm 1 \pmod{8}$, then one of $(q + 1)/2$ and $(q - 1)/2$ is divisible by 4, and so must be a power of 2; so either $q = 9$, or q is a Fermat or Mersenne prime. So, apart from this case, q is congruent to $\pm 3 \pmod{8}$. Now either q is an odd power of 3, or one of $(q + 1)/2$ and $(q - 1)/2$ is twice a prime, while the other is three times a prime or a power of 3 (unless $q = 11$ or $q = 13$). The odd prime powers up to 500 satisfying the condition are 3, 5, 7, 9, 11, 13, 17, 19, 27, 29, 31, 43, 53, 67, 163, 173, 243, 257, 283, 317.

Table 1 gives the numbers of vertices in cokernels for small finite simple groups. Note that a graph is a cograph if and only if its cokernel has one vertex. The last column of the table gives the number of cyclic subgroups of G , that is, equivalence classes under the relation \equiv where $x \equiv y$ if $\langle x \rangle = \langle y \rangle$. Equivalent vertices are closed twins in all our graphs, so this number is an upper bound for the number of vertices in each cokernel. (I have replaced PSL and PSU by L and U in the table to save space.)

G	$ G $	Pow(G)	EPow(G)	DCom(G)	Com(G)	NGen(G)	Cyc
A_5	60	1	1	1	1	32	32
$L_2(7)$	168	1	1	1	44	79	79
A_6	360	1	1	1	92	167	167
$L_2(8)$	504	1	1	1	1	128	156
$L_2(11)$	660	1	1	1	112	244	244
$L_2(13)$	1092	1	1	1	184	366	366
$L_2(17)$	2448	1	1	1	308	750	750
A_7	2520	352	352	352	352	842	947
$L_2(19)$	3420	1	1	1	344	914	914
$L_2(16)$	4080	1	1	1	1	784	784
$L_3(3)$	5616	756	756	808	808	1562	1796
$U_3(3)$	6048	786	534	499	499	1346	1850
$L_2(23)$	6072	1267	1	1	508	1313	1566
$L_2(25)$	7800	1627	1	1	652	1757	2082
M_{11}	7920	1212	1212	1212	1212	2444	2576

TABLE 1. Sizes of cokernels of graphs on small simple groups

The table suggests various conjectures, some of which can be proved. For example:

Theorem 8.8. *Let G be a non-abelian finite simple group. Then $\text{NGen}(G)$ is not a cograph.*

Proof. We consider the reduced graph obtained by deleting the identity vertex. The reduced non-generating graph of a simple group is connected, and has diameter at most 5 (this follows from the results of Ma, Herzog *et al.*, Shen and Freedman discussed in Section 12 below). Also, the generating graph is connected (see [29]), and indeed has diameter 2 (see [31]). But the complement of a connected cograph is disconnected. \square

Question 17. Find, or estimate, the number of vertices in the cokernel of the non-generating graph of a finite simple group. In particular, classify simple groups for which the number of vertices in the cokernel of the non-generating graph is equal to the number of cyclic subgroups.

The hypothesis of simplicity is essential here. For example, the non-generating graph of any 2-generator p -group is a cograph (it consists of a star $K_{1,p+1}$ with the central vertex blown up to a clique of size p^{n-2} and the remaining vertices to cliques of size $p^{n-2}(p-1)$, where the group has order p^n). This graph is a cograph: its complement is complete multipartite with some isolated vertices.

9. Connectedness

In this section, we examine the question of connectedness of these graphs. All those in the hierarchy (except the null graph) are connected, since the identity is joined to all other vertices. The question is non-trivial, however, if we remove vertices joined to all others. The first job is to characterise these vertices.

9.1. Centres. In the commuting graph of G , the set of vertices joined to all others is simply the centre $Z(G)$ of G . So we adapt the terminology by defining analogues of the centre for other graphs in the list. So, if X denotes Pow, EPow, DCom, Com or NGen, we define the X -centre of G , denoted $Z_X(G)$, to be the set of vertices joined to all others in $X(G)$. It turns out that (aside from the non-generating graph), in almost all cases, $Z_X(G)$ is a normal subgroup of G ; the only exception is for the power graph of a cyclic group of non-prime-power order. We note that $Z_{\text{EPow}}(G)$ has been studied by Patrick and Wepsic [92], who called it the *cyclicizer* of G : it is the set

$$\{x \in G : (\forall y \in G) \langle x, y \rangle \text{ is cyclic}\}.$$

The main results on the cyclicizer are also given in [5].

Theorem 9.1. (a) $Z_{\text{Pow}}(G)$ is equal to G if G is cyclic of prime power order; or the set consisting of the identity and the generators if G is cyclic of non-prime-power order; or $Z(G)$ if G is a generalized quaternion group; or $\{1\}$ otherwise.

(b) $Z_{\text{EPow}}(G)$ is the product of the Sylow p -subgroups of $Z(G)$ for $p \in \pi$, where π is the set of primes p for which the Sylow p -subgroup of G is cyclic or generalized quaternion; in particular, $Z_{\text{EPow}}(G)$ is cyclic.

(c) $Z_{\text{DCom}}(G)$ is the projection into G of $Z(H)$, where H is a Schur cover of G .

(d) $Z_{\text{Com}}(G) = Z(G)$.

Proof. (a) This is [33, Proposition 4].

(b) Suppose that x is joined to all other vertices in $\text{EPow}(G)$. Then $\langle x, y \rangle$ is cyclic for all $y \in G$; so certainly $x \in Z(G)$.

If three elements of a group have the property that any two of them generate a cyclic group, then all three generate a cyclic group: see [1, Lemma 32]. So $Z_{\text{EPow}}(G)$ is a subgroup, since if $x, y \in Z_{\text{EPow}}(G)$ then, for all $w \in G$, $\langle x, w \rangle$ and $\langle y, w \rangle$ are cyclic, and so if $\langle x, y \rangle = \langle z \rangle$ then $\langle z, w \rangle$ is cyclic for all $w \in G$, so that $z \in Z_{\text{EPow}}(G)$.

Now let x be an element of prime order p in $Z_{\text{EPow}}(G)$. If G contains a subgroup $C_p \times C_p$ then there is an element of order p not in $\langle x \rangle$, so not adjacent to x , a contradiction. So the Sylow p subgroup of G is cyclic or generalised quaternion, by Burnside’s theorem. But now x lies in every Sylow p -subgroup of G , so is joined to every element of p -power order, and hence to every element of G .

(c) and (d) Part (d) is clear, and (c) follows since $\text{DCom}(G)$ is a projection of the commuting graph of a Schur cover of G . □

By contrast, $Z_{\text{NGen}}(G)$ is not necessarily a subgroup of the 2-generated group G . If G is non-abelian, then $Z_{\text{NGen}}(G)$ must contain $Z(G)$, since the non-generating graph contains the commuting graph. Also, it contains the Frattini subgroup $\Phi(G)$ of G , since 2-element generating sets are minimal and so their elements do not lie in the Frattini subgroup (which consists of the elements which can be dropped from any generating set).

If the order of G is a prime power, then the Burnside basis theorem shows that $Z_{\text{NGen}}(G) = \Phi(G)$, since a set of elements generates G if and only if its projection onto $G/\Phi(G)$ generates this quotient.

Also, by the result of [29] (see Theorem 2.5), if all proper quotients of G are cyclic, then $Z_{\text{NGen}}(G) = \{1\}$.

In general, however, $Z_{\text{NGen}}(G)$ may be a subgroup different from both $Z(G)$ and $\Phi(G)$. For example, let G be the symmetric group S_4 . Then both $Z(G)$ and $\Phi(G)$ are trivial, but $Z_{\text{NGen}}(G)$ is the Klein group V_4 (the minimal normal subgroup of G).

Moreover, it may not be a subgroup at all. For example, if $G = C_6 \times C_6$, then $Z_{\text{NGen}}(G)$ consists of the elements not of order 6, since both elements in any generating pair must have order 6.

Question 18. Characterise the 2-generated groups in which $Z_{\text{NGen}}(G)$ is a subgroup of G .

9.2. Connectedness. Each of our types of graph is connected, since the corresponding “centre” is non-empty and its vertices are joined to all others. So the question becomes interesting if we ask whether the induced subgraph on the elements outside this centre is connected.

The situation for the commuting graph is well-understood, thanks to the results of [56, 88]. But first I mention another link with the Gruenberg–Kegel graph. This has been known for some time, but the first mention I know in the literature is [88, Section 3].

Theorem 9.2. *Let G be a group with trivial centre. Then the induced subgraph of the commuting graph on $G \setminus \{1\}$ is connected if and only if the Gruenberg–Kegel graph is connected.*

Proof. Suppose first that $Z(G) = 1$ and the commuting graph is connected. Let p and q be primes dividing $|G|$. Choose elements g and h of orders p and q respectively, and suppose their distance in the commuting graph is d . We show by induction on d that there is a path from p to q in the GK graph.

If $d = 1$, then g and h commute, so gh has order pq , and p is joined to q . So assume the result for distances less than d , and let $g = g_0, \dots, g_d = h$ be a path from g to h .

Let i be minimal such that p does not divide the order of g_i (so $i > 0$). Now some power of g_{i-1} , say g_{i-1}^a , has order p , while a power g_i^b of g_i has prime order $r \neq p$.

The distance from g_i^b to g_d is at most $d - i < d$, so there is a path from r to q in the GK graph. But g_{i-1}^a and g_i^b commute, so p is joined to r .

For the converse, assume that the GK graph is connected.

Note first that for every non-identity element g , some power of g has prime order, so it suffices to show that all elements of prime order lie in the same connected component of the commuting graph. Also, since a non-trivial p -group has non-trivial centre, the non-identity elements of any Sylow subgroup lie in a single connected component.

Let C be a connected component. Connectedness of the GK graph shows that C contains a Sylow p -subgroup for every prime p dividing $|G|$. Also, every element of C , acting by conjugation, fixes C . It follows that the normaliser of C is G , and hence that C contains every Sylow subgroup of G , and thus contains all elements of prime order, as required. \square

After a lot of preliminary work, much of it on specific groups, summarised in the introduction to [56], Iranmanesh and Jafarzadeh [72] conjectured that there is an absolute upper bound on the diameter of any connected component of the induced subgraph on $G \setminus Z(G)$ of the commuting graph of G . This conjecture was refuted by Giudici and Parker [56]. However, it was proved for groups with trivial centre by Morgan and Parker [88]. In these results, I use the term “reduced commuting graph” to mean the induced subgraph of the commuting graph of G on $G \setminus Z(G)$, in which no vertex is joined to all others.

Theorem 9.3. *There is no upper bound for the diameter of the reduced commuting graph of a finite group; for any given d there is a 2-group whose reduced commuting graph is connected with diameter greater than d .*

On the other hand:

Theorem 9.4. *Suppose that the finite group G has trivial centre. Then every connected component of its reduced commuting graph has diameter at most 10.*

For the power graph and enhanced power graph, we note that, if the group G is not cyclic or generalized quaternion, then the corresponding “centre” is just the identity. So the natural question is: if G is not cyclic or generalized quaternion, is the induced subgraph of the power graph on non-identity elements connected? This question has been considered in several papers, for example [37, 117].

The next result shows that we have only one rather than two problems to consider.

Proposition 9.5. *Let G be a group with $Z(G) = \{1\}$. Then the reduced power graph of G is connected if and only if the reduced enhanced power graph of G is connected.*

Proof. If g and h are joined in the power graph, they are joined in the enhanced power graph; if they are joined in the enhanced power graph, then they lie at distance at most 2 in the power graph; both g and h are powers of the intermediate vertex, which is thus not the identity. \square

The argument shows that, if these graphs are connected, the diameter of the power graph is at least as great as, and at most twice, the diameter of the enhanced power graph. Can these bounds be improved?

I have already quoted the result of Burness *et al.* on the generating graph. For the non-generating graph, the results of Freedman and others on the difference between the non-generating graph and the commuting graph (that is, the graph $(\text{NGen} - \text{Com})(G)$) have been mentioned also.

As promised, here is a weak result on the graph $(\text{Com} - \text{Pow})(G)$.

Theorem 9.6. *Suppose that the finite group G satisfies the following conditions:*

- (a) *The Gruenberg–Kegel graph of G is connected.*
- (b) *If P is any Sylow subgroup of G , then $Z(P)$ is non-cyclic.*

Then the induced subgraph of $(\text{Com} - \text{Pow})(G)$ on $G \setminus \{1\}$ either has an isolated vertex or is connected.

Proof. Let $\Gamma(G)$ denote the induced subgraph of $(\text{Com} - \text{Pow})(G)$ on $G \setminus \{1\}$. Note that, if H is a subgroup of G , then the induced subgraph of $\Gamma(G)$ on $H \setminus \{1\}$ is $\Gamma(H)$.

First we show that, if P is a p -group, then $\Gamma(P)$ is connected. Let $Q \leq Z(P)$ with $Q \cong C_p \times C_p$. Then the induced subgraph on $Q \setminus \{1\}$ is complete multipartite with $p + 1$ blocks of size $p - 1$, corresponding to the cyclic subgroups of Q . So it suffices to show that any element z of $P \setminus \{1\}$ has a neighbour in $Q \setminus \{1\}$. We see that z commutes with Q since $Q \leq Z(P)$; and $\langle z \rangle \cap Q$ is cyclic so there is some element of Q not in this set.

Now let C be a connected component of $\Gamma(G)$ containing an element z of prime order p . Since $\Gamma(G)$ is invariant under $\text{Aut}(G)$, in particular it is normalized by all its elements, so $\langle C \rangle \leq N_G(C)$. In particular, C contains a Sylow p -subgroup of G (one containing the given element of order p in C).

If C contains an element of prime order r , and $\{r, s\}$ is an edge of the GK graph, then G contains an element g of order rs , then without loss of generality $g^s \in C$, and g^s is joined to g^r in $\Gamma(G)$, so also $g^r \in C$. Now connectedness of the GK graph shows that C contains a Sylow q -subgroup of G for every prime divisor of $|G|$. Hence $|N_G(C)|$ is divisible by every prime power divisor of $|G|$, whence $N_G(C) = G$.

Finally, let g be any non-identity element of G . Choose a maximal cyclic subgroup K containing g . If $C_G(K) = K$, then the generator of K commutes only with its powers, and is isolated in $\Gamma(G)$. If not, then there is an element of prime order in $C_G(K) \setminus K$. (If $h \in C_G(K) \setminus K$, then $\langle g, h \rangle$ is abelian but not cyclic, so contains a subgroup $\langle g \rangle \times C_m$ for some m ; choose an element of prime order in the second

factor.) This element is joined to g in the commuting graph but not in the power graph; so $g \in C$. We conclude that $C = G \setminus \{1\}$, and the proof is done. \square

Remarks. The theorem is probably not best possible. Let us consider the hypotheses.

We saw in Theorem 9.2 that, for groups with trivial centre (our main interest here), connectedness of the GK graph is equivalent to connectedness of the commuting graph, and so is clearly necessary for connectedness of $(\text{Com} - \text{Pow})(G)$.

The second condition (which is necessary for the above proof) is probably much too strong. Perhaps it can be weakened to say that the Sylow subgroups of G are not cyclic or generalized quaternion groups (or, subgroups of G are not cyclic or generalized quaternion groups (or, equivalently, that G has no subgroup $C_p \times C_p$ for prime p). Perhaps it is only necessary to assume this for one prime. More work needed.

9.3. Connectedness of the complement. As well as asking whether our graphs, after reduction (removing vertices joined to all others) are connected, we can ask the same question for the complementary graphs. For the commuting graph, there is a simple elegant argument, depending on the following result.

Proposition 9.7. *Let Γ be a graph whose vertex set is a group G , and suppose that for any vertex $g \in G$, the closed neighbourhood of g is a subgroup of G . Then the complementary graph has just one connected component of size larger than 1; this component has diameter at most 2.*

Proof. The isolated vertices in the complement of Γ are the vertices whose closed neighbourhood in Γ is the whole of G . Let g_1, g_2 be two elements of G which are not isolated in the complement of Γ . Then $H_1 = \{g_1\} \cup \Gamma(g_1)$ and $H_2 = \{g_2\} \cup \Gamma(g_2)$ are subgroups of G . Since a finite group cannot be written as the union of two proper subgroups (a simple consequence of Lagrange's Theorem), there is a vertex h outside these two subgroups, hence joined to g_1 and g_2 in the complement of Γ . \square

Corollary 9.8. *If G is a non-abelian finite group, then the complement of the reduced commuting graph of G is connected with diameter at most 2.*

Proof. The closed neighbourhood of a vertex g in the commuting graph is its centraliser. \square

What about our other graphs? The deep commuting graph of G is obtained from the commuting graph of a Schur cover of G by twin reduction; so its complement, restricted to the non-isolated vertices, is connected with diameter at most 2.

Theorem 9.9. *Let G be a finite group which is not a cyclic p -group. Then the complement of the power graph of G has just one connected component, apart from isolated vertices.*

Proof. Let Γ be the complement of the power graph of G . We begin with a few remarks.

- (a) By Theorem 9.1, we know the set of the isolated vertices in Γ (we have excluded the case where every vertex is isolated):

- if G is cyclic but not of prime power order, the identity and the generators of G ;
 - if G is generalised quaternion, $Z(G)$;
 - otherwise, just the identity.
- (b) Since every edge of the noncommuting graph is an edge of Γ , if G is nonabelian then $G \setminus Z(G)$ is contained in a single component, by Corollary 9.8. So any other component is contained in $Z(G) \setminus \{1\}$.
- (c) Suppose that G has more than one subgroup of prime order, and let S be the set of elements of prime order. Then the induced subgraph on S is connected (it is complete multipartite, with the cyclic subgroups as parts). Moreover, if g is an element whose order is not divisible by some prime divisor of $|G|$, then g has a neighbour in S .

Suppose first that G is not of prime power order. If $Z(G) = 1$, then the result follows from the second remark, so suppose not. Then $Z(G) \cap S$ is nonempty, so every element whose order fails to be divisible by some prime divisor of $|G|$ lies in $Z(G)$. In particular, the Sylow subgroups of G all lie in $Z(G)$, so G is abelian. Now we separate into two subcases:

- (a) Suppose that G is cyclic. If $g \in G$ and g is not a generator, then there is a prime p such that the power of p dividing the order of g is smaller than the power p^m which divides $|G|$. If the order of g is a p -power, it is joined to a vertex in S . Otherwise, there is an element of order p^m , joined both to g and to an element of S .
- (b) Suppose that G is not cyclic. There is a prime p such that G contains $C_{p^a} \times C_{p^b}$, where p^a is the p -part of the exponent of G . For any $h \in G$, h is joined to the generators of one or other of these two cyclic groups; and each of them is joined to an element of S .

So now suppose that G is a group of prime power order. If G has more than one subgroup of order p , then every element of G has a neighbour in S . Otherwise, by the theorem of Burnside [64, Theorem 12.5.2], G is either cyclic (which we have excluded) or generalised quaternion (in which case the vertices of $Z(G)$ are isolated, and the remainder is connected by the second opening remark). \square

I conclude with two questions which should not be difficult.

Question 19. What is the best possible upper bound for the diameter of non-trivial connected component of the complement of the power graph, and which groups attain the bound?

Question 20. Is it true that the complement of the enhanced power graph has just one connected component, apart from isolated vertices?

10. Automorphisms

Each type of graph in the hierarchy on a group G is preserved by the automorphism group of G . But in almost all cases, the automorphism group of the graph is much larger. This question has been considered in [13].

We saw in Proposition 7.4 that any of our hierarchy of graphs has non-trivial twin relation. So the first thing we need to do is to take a look at automorphisms of such graphs.

Let Γ be a graph. It is clear that its automorphism group $\text{Aut}(\Gamma)$ preserves twin relations on Γ , and that vertices in a twin equivalence class can be permuted arbitrarily. It follows by induction that the group induces an automorphism group on the cokernel Γ^* of Γ , say $\text{Aut}^-(\Gamma^*)$. We say that the twin reduction on Γ is *faithful* if $\text{Aut}^-(\Gamma^*) = \text{Aut}(\Gamma^*)$.

Trivially, if Γ is a cograph (so that its cokernel is the 1-vertex graph), the twin reduction is faithful; we ignore this case.

The reduction process is not always faithful. For a simple example, consider Figure 4.




FIGURE 4. Non-faithful twin reduction

In the left-hand graph, the two leaves on the right are twins, and twin reduction gives the right-hand graph as the cokernel. But the cokernel has an automorphism (reflection in the vertical axis of symmetry) not induced from an automorphism of the original.

Question 21. Given a finite group G and one of our types of graph (say X),

- When is twin reduction on $X(G)$ faithful?
- What is the automorphism group of the cokernel of $X(G)$?

Very little seems to be known about this question. I first discuss cographs, then give a couple of examples.

Proposition 10.1. *Let Γ be a cograph. Then the automorphism group of Γ can be built from the trivial group by the operations of direct product and wreath product with a symmetric group.*

Proof. Recall that a cograph can be built from the 1-vertex graph by the operations of complement and disjoint union. Now complementation does not change the automorphism group. If Γ is the disjoint union of m_1 copies of Δ_1, \dots, m_r copies of Δ_r , then

$$\text{Aut}(\Gamma) = (\text{Aut}(\Delta_1) \wr S_{m_1}) \times \cdots \times (\text{Aut}(\Delta_r) \wr S_{m_r}).$$

Assuming inductively that each of $\text{Aut}(\Delta_1), \dots, \text{Aut}(\Delta_r)$ can be built by direct products and wreath products with symmetric groups; then the same is true for $\text{Aut}(\Gamma)$. \square

Example. Let G be the alternating group A_5 , and consider the power graph of G . The identity is joined to all other vertices; after removing it we have six cliques of size 4 (corresponding to cyclic subgroups of order 5), ten of size 2 (corresponding to cyclic subgroups of order 3), and fifteen isolated vertices (corresponding to elements of order 2). This graph is easily seen to be a cograph, so its cokernel has

a single vertex. In fact, closed twin reduction contracts the cliques of sizes 2 and 4 to single vertices, giving a star on 32 vertices; then open twin reduction produces a single edge, and closed twin reduction reduces this to a single vertex.

Example. Let G be the Mathieu group M_{11} . The power graph of G has 7920 vertices. On removing the identity, we are left with a graph consisting of

- 144 complete graphs of size 10, corresponding to elements of order 11;
- 396 complete graphs of size 4, corresponding to elements of order 5;
- a single connected component Δ on the remaining 4895 vertices.

Two steps of twin reduction remove all the components which are complete. If we take Δ , and first factor out the relation “same closed neighbourhood”, and then factor out from the result the relation “same open neighbourhood”, we obtain a connected graph on 1210 vertices whose automorphism group is M_{11} . (This is shown by a GAP computation.) This group is induced by the automorphism group of the original power graph; so the reduction is faithful.

Exercise. Why is the number 1210 given above two less than the number of vertices of the cokernel of the power graph of M_{11} given in Table 1?

Question 22. For which non-abelian finite simple group G is it the case that the twin reduction on the power graph/enhanced power graph/deep commuting graph/commuting graph/generating graph of G is faithful?

Example. A curious example showing that this is not true for all such groups is described in [39]. Let G be the simple group $\text{PSL}(2, 16)$. The automorphism group of G is the group $\text{P}\Gamma\text{L}(2, 16)$, four times as large as G ; but the cokernel of the generating graph of G has an extra automorphism of order 2, interchanging the sets of vertices coming from elements of orders 3 and 5 in G . Twin reduction of this graph is thus not faithful. The cokernel is a graph on 784 vertices with automorphism group $C_2 \times \text{P}\Gamma\text{L}(2, 16)$.

Non-faithfulness means, as in this example, that extra automorphisms are introduced by twin reduction.

11. Above and beyond the hierarchy

This section contains some very brief comments on similar graphs.

11.1. Nilpotence, solvability and Engel graphs. Given a subgroup-closed class of graphs \mathcal{C} , we can define a graph on G in which x and y are joined if $\langle x, y \rangle$ belongs to \mathcal{C} .

For \mathcal{C} the class of cyclic groups, we obtain the enhanced power graph; and for the class of abelian groups, we obtain the commuting graph.

After these, the most natural classes to consider are those of nilpotent and solvable groups; let us denote the corresponding graphs by $\text{Nilp}(G)$ and $\text{Sol}(G)$ respectively.

A *Schmidt group* is a non-nilpotent group all of whose proper subgroups are nilpotent. These groups were characterised by Schmidt [98]; see [19] for an accessible account. All are 2-generated.

I do not know of a similar characterisation of the non-solvable groups all of whose proper subgroups are solvable. However, we can conclude that they are 2-generated, as follows. Let G be such a group, and S the solvable radical of G (the largest solvable normal subgroup). Then G/S is a non-abelian simple group. (For if H/S is a minimal normal subgroup of G/S , then H/S is a product of isomorphic simple groups, so H is not solvable, and by minimality $H = G$.) Now every finite simple group is 2-generated. If we take two cosets Sg, Sh which generate G/S , then $\langle g, h \rangle$ is a subgroup of G which projects onto G/S , and so is non-solvable; by minimality it is equal to G . (In fact we do not need the Classification of Finite Simple Groups here. For clearly G/S is a minimal simple group, and so is covered by Thompson's classification of N-groups [108].)

It follows that a group G is nilpotent (resp. solvable) if and only if every 2-generated subgroup of G is nilpotent (resp. solvable). For if every 2-generated subgroup of G is nilpotent, then G cannot contain a minimal non-nilpotent subgroup, and so G is nilpotent; similarly for solvability.

- Proposition 11.1.** (a) *For any finite group G , we have $E(\text{Com}(G)) \subseteq E(\text{Nilp}(G)) \subseteq E(\text{Sol}(G))$.*
 (b) *$E(\text{Com}(G)) = E(\text{Nilp}(G))$ if and only if all the Sylow subgroups of G are abelian.*
 (c) *$E(\text{Nilp}(G)) = E(\text{Sol}(G))$ if and only if G is nilpotent.*
 (d) *$E(\text{Com}(G)) = E(\text{Sol}(G))$ if and only if G is abelian.*
 (e) *If G is non-nilpotent, then $E(\text{Nilp}(G)) \subseteq E(\text{NGen}(G))$; equality holds if and only if G is a Schmidt group.*
 (f) *If G is non-solvable, then $E(\text{Sol}(G)) \subseteq E(\text{NGen}(G))$; equality holds if and only if G is a minimal non-solvable group.*

Proof. (a) The first point is clear from the definition.

(b) Suppose that $E(\text{Com}(G)) = E(\text{Nilp}(G))$. Then two elements from the same Sylow subgroup of G generate a nilpotent group; hence they commute. Conversely, if the Sylow subgroups are abelian, then a nilpotent subgroup is the product of its Sylow subgroups and hence is abelian.

(c) Suppose that $E(\text{Nilp}(G)) = E(\text{Sol}(G))$. If G is not nilpotent, it contains a minimal non-nilpotent subgroup, a Schmidt group, which is 2-generated and solvable, hence nilpotent, a contradiction. Conversely, if G is nilpotent, then $\text{Nilp}(G)$ is complete.

(d) If $\text{Com}(G)$ and $\text{Sol}(G)$ coincide, then G is nilpotent with abelian Sylow subgroups, hence is abelian. The converse is clear.

(e), (f) The forward direction in the last two points uses the fact that these groups are 2-generated, as remarked above. For if $\text{Nilp}(G)$ and $\text{NGen}(G)$ have the same edges, then two elements which do not generate G must generate a nilpotent group, and similarly for solvability. \square

Regarding (b), groups with all Sylow subgroups abelian are known, since Walter [110] classified the groups with abelian Sylow 2-subgroups. The simple groups arising here are $\text{PSL}(2, q)$ with q even or congruent to $\pm 3 \pmod{8}$ and the first Janko group J_1 .

The sets of vertices joined to all others in the nilpotency and solvability graphs have been characterized. The first part of this proposition is due to Abdollahi and Zarrin, the second part to Guralnick *et al.*

Theorem 11.2. *For any finite group G ,*

- (a) $Z_{\text{Nilp}}(G)$ *is the hypercentre of G ;*
- (b) $Z_{\text{Sol}}(G)$ *is the solvable radical of G .*

I will give a proof of the first statement after discussing the Engel graph below. I refer to the cited paper for the second. Note that $Z_{\text{Nilp}}(G)$ and $Z_{\text{Sol}}(G)$ are both subgroups of G .

This question has been recently studied in greater generality by Lucchini and Nemmi [85], who investigated the question of when the set of vertices in the \mathfrak{F} -graph of G which are joined to all other vertices is necessarily a subgroup, in the case where \mathfrak{F} is a saturated formation. Briefly, a *formation* is a class of groups closed under quotients and subdirect products; the formation \mathfrak{F} is *saturated* if $G/\Phi(G) \in \mathfrak{F}$ implies $G \in \mathfrak{F}$. It is easily shown that finite solvable groups, and finite cyclic groups, form saturated formations: the corresponding graphs are the enhanced power graph and the solvability graph. Also finite nilpotent groups form a saturated formation, see [97, Theorem 2.5.12]. However, finite abelian groups do not, as the dihedral and quaternion groups of order 8 show.

A class of graphs is *hereditary* if it is subgroup-closed. A group G is *critical* for a class \mathfrak{F} if $G \notin \mathfrak{F}$ but all proper subgroups of G are in \mathfrak{F} ; if in addition all proper quotients of G are in \mathfrak{F} , then G is *strongly critical*. Now Lucchini and Nemmi [85, Theorem 1] show:

Theorem 11.3. *Let \mathfrak{F} be a hereditary saturated formation, lying between the classes of abelian and solvable groups. The following are equivalent:*

- (a) *for any group G , the set*

$$Z_{\mathfrak{F}}(G) = \{x \in G : (\forall y \in G)(\langle x, y \rangle \in \mathfrak{F})\}$$

is equal to the intersection of all maximal \mathfrak{F} -subgroups of G (and hence is a subgroup of G);

- (b) *for every group G which is solvable and strongly critical for \mathfrak{F} , $G/\text{Soc}(G)$ is cyclic, where $\text{Soc}(G)$ is the socle of G .*

I conclude with the obvious question:

Question 23. Investigate analogues of the earlier results in this paper in the extended hierarchy of graphs containing $\text{Nilp}(G)$ and $\text{Sol}(G)$ (or to other graphs of this kind).

Universality is relatively straightforward, and both cases can be handled together. Recall from the proof of Theorem 5.7 that any graph Γ can be represented as the intersection graph of a linear

hypergraph, a family of sets with the property that two sets intersect in 1 point if the corresponding vertices are adjacent, and are disjoint otherwise. We can add some dummy points to ensure that all the sets in the collection have the same (prime) cardinality p at least 3. Now if we take cycles whose supports are these sets, then we see (as there) that the cycles corresponding to adjacent vertices generate the alternating group of degree $2p - 1$, while those corresponding to non-adjacent vertices generate $C_p \times C_p$. So both nilpotence and solvability graphs of finite groups embed all finite graphs.

Connectedness of the complement of the nilpotency graph has been investigated by Abdollahi and Zarrin [8]. There is a generalisation to \mathfrak{F} -groups, for saturated formations \mathfrak{F} , by Lucchini and Nemmi [85].

Each of these cases can be stratified: we can define the level- k nilpotence or solvability graph to have edges $\{x, y\}$ if $\langle x, y \rangle$ is nilpotent of class at most k (resp. solvable of derived length at most k).

Other classes of groups for which the corresponding graphs could be studied, for which the minimal groups not in the class have been considered, include the supersolvable groups (those for which every chief factor is cyclic) and the p -nilpotent groups (groups with normal p -complements). The groups minimal with respect to not lying in these classes are considered in [18] and [19] respectively.

A closely related graph is the *Engel graph* of a group, defined by Abdollahi [3]. Here is a brief account. We define, for each positive integer k , and all $x, y \in G$, the element $[x, {}_k y]$ of G to be the left-normed commutator of x and k copies of y ; more formally,

- $[x, {}_1 y] = [x, y] = x^{-1}y^{-1}xy$,
- for $k > 1$, $[x, {}_k y] = [[x, {}_{k-1} y], y]$.

Abdollahi defined x and y to be adjacent if $[x, {}_k y] \neq 1$ and $[y, {}_k x] \neq 1$ for all k . To fit with the philosophy of this paper, and at Abdollahi's suggestion, I will redefine it to be the complement of this graph. If we do this then we have a similar situation to that arising with the power graph. If we define the *directed Engel graph* to have an arc from x to y if $[y, {}_k x] = 1$ for some k , then the Engel graph (that is, the complement of the graph as defined in [3]) is the graph in which x and y are joined if there is an arc from one to the other. The directed graph may also have a role to play here.

A similar stratification to that for nilpotence and solvability graphs can also be defined in this case. In particular, the level 1 Engel graph is just the commuting graph.

Zorn [118] showed that, if a finite group G satisfies an Engel identity $[x, {}_k y] = 1$ for all x, y (for some k), then G is nilpotent; so the finite groups for which the directed Engel graph is complete are the same as those for which the nilpotency graph is complete. (For infinite groups, this is not true, though the result has been shown in a number of special cases.)

So there is a close connection between the Engel graph and the nilpotency graph. But they are not equal in general. For example, in the group S_3 , there is an arc of the directed Engel graph from each element of order 3 to each element of order 2, but not in the reverse direction.

Question 24. What can be said about the relation between the Engel and nilpotency graphs? In particular, in which groups are they equal?

Armed with this knowledge, we return briefly to the nilpotence graph. First, another definition. The *upper central series* of a group G is the sequence of subgroups defined by

$$Z^0(G) = \{1\}, \quad Z^{k+1}(G)/Z^k(G) = Z(G/Z^k(G)) \text{ for } k \geq 1.$$

The *hypercentre* of G is the union of these subgroups. Thus, if G is finite, then the hypercentre is $Z^k(G)$, where k is the smallest value such that $Z^k(G) = Z^{k+1}(G)$. It is clear that we have $Z^1(G) = Z(G)$, and subsequent terms are the subgroups of G that project onto the upper central series of $G/Z(G)$. Hence we can give an alternative definition:

$$Z^{k+1}(G)/Z(G) = Z^k(G/Z(G)).$$

Theorem 11.4. *Let G be a finite group. Then the set $Z_{\text{Nilp}}(G)$ of elements of G which are joined to all other elements in the nilpotence graph is equal to the hypercentre of G .*

Proof. Suppose that x is joined to all other elements of G , so that $\langle x, y \rangle$ is nilpotent for all $y \in G$. In particular, for all y , there exists k such that $[x, {}_k y] = 1$. Baer [15] showed that, in a group with the maximum condition on subgroups (and in particular a finite group), this implies that x is in the hypercentre of G .

For the reverse implication, we have to show that, if $x \in Z^k(G)$, where k is the least value such that $Z^k(G) = Z^{k+1}(G)$, then for any $y \in G$ we have $\langle x, y \rangle$ nilpotent. We prove this by induction on k . So assume that it is true with $k - 1$ replacing k in any group. Then for $x \in Z^k(G)$ we have $Z(G)x \in Z^{k-1}(G/Z(G))$, so $Z(G)\langle x, y \rangle/Z(G)$ is nilpotent. Thus $Z(G)\langle x, y \rangle$ is an extension of a central subgroup by a nilpotent group, and so is nilpotent; and so $\langle x, y \rangle$ is nilpotent, as required. \square

Question 25. What, if anything, can be said for infinite groups?

11.2. Other graphs. All the graphs studied so far have the property that two group elements which generate the same cyclic subgroup are closed twins. So it would be very natural to collapse them by factoring out this equivalence relation. Alternatively, one could simply remove edges between such pairs, so that they become open twins. Note that the original, the quotient, and the graph with edges removed all have the same cokernel; so, if one of them is a cograph, then they all are.

We could put a graph at the bottom of the hierarchy, in which $x \sim y$ if $\langle x \rangle = \langle y \rangle$; then the second possibility suggested above fits into our scheme as the difference between this graph and one of the others.

I end this section with a general question.

Question 26. For which types of graph, and which groups, is the relation \equiv given by $x \equiv y$ if $\langle x \rangle = \langle y \rangle$ definable directly from the graph without reference to the group?

In particular, if the cokernel of the graph is equal to the quotient by the equivalence relation \equiv , this will be true. For which groups is this the case?

12. Intersection graphs

There turns out to be a close connection between certain intersection graphs defined on G , and some of the graphs in our hierarchy. First I look briefly at the connection in the abstract, then discuss some particular cases.

12.1. Dual pairs. Let B be a bipartite graph. If it is connected, it has a unique bipartition: take a vertex v ; then the bipartite blocks are the sets of vertices at even (resp. odd) distance from v . If B is not connected, the bipartition is not unique; in fact, there are $2^{\kappa-1}$ bipartitions, where κ is the number of connected components, since we can make a bipartite block by choosing a bipartite block in each component and taking their union. However, I will always assume that the bipartition of B is given, and is part of its structure.

The *halved graphs* arising from B are the graphs Γ_1 and Γ_2 whose vertex set is a bipartite block, two vertices adjacent in the relevant graph if and only if they lie at distance 2 in B .

We call a pair of graphs Γ_1 and Γ_2 a *dual pair* if there is a bipartite graph B without isolated vertices such that Γ_1 and Γ_2 are the halved graphs of B .

I warn that this concept is not the same as the vague notion of duality which informed the name “dual enhanced power graph”. It is however closely connected with duality in design theory and geometry, or between a graph and the linear hypergraph (or partial linear space) that we used in Section 5.5.

Proposition 12.1. *Let Γ_1 and Γ_2 be a dual pair of graphs. Then Γ_1 is connected if and only if Γ_2 is connected. More generally, there is a natural bijection between connected components of Γ_1 and connected components of Γ_2 with the property that corresponding components have diameters which are either equal or differ by 1.*

Proof. Any vertex of Γ_1 is joined (by an edge of B) to a vertex of Γ_2 , and *vice versa*, since B has no isolated vertices. Now suppose that two vertices of Γ_1 are joined by a path of length d . Then there is a path of length $2d$ in B joining them. So a connected component of B is the union of corresponding connected components in Γ_1 and Γ_2 . Suppose that a component of Γ_1 has diameter d . Take two vertices v_1, v_2 in the corresponding component of Γ_2 . Choose vertices u_1 and u_2 of Γ_1 joined in B to v_1 and v_2 respectively. These two vertices lie at distance $r \leq d$, say; so there is a path of length at most $2r$ in B joining them. Thus v_1 and v_2 have distance at most $2r + 2$ in B , whence their distance in Γ_2 is at most $r + 1$, hence at most $d + 1$. So the diameter of a component of Γ_2 has diameter at most one more than the corresponding component of Γ_1 . Interchanging the roles of the dual pair completes the proof. \square

Question 27. What other relations hold between properties of a dual pair of graphs?

If the bipartite graph is *semiregular*, then a number of properties transfer between the corresponding dual pair, especially spectral properties, and (related to this) optimality properties of statistical designs [17].

12.2. Graphs on groups and intersection graphs. In order to apply this result, I give a general construction showing that certain graphs defined on the non-identity elements of a group form dual pairs with certain intersection graphs of families of subgroups.

Proposition 12.2. *Let G be a finite non-cyclic group, and let \mathcal{F} be a family of non-trivial proper subgroups of G with the property that its union is G . Let Γ be the graph defined on the non-identity elements of G by the rule that x is joined to y if and only if there is a subgroup $H \in \mathcal{F}$ with $x, y \in H$. Then Γ and the intersection graph of \mathcal{F} form a dual pair.*

Proof. We form the bipartite graph B whose vertex set is $(G \setminus \{1\}) \cup \mathcal{F}$, where a group element $x \neq 1$ is joined to a subgroup $H \in \mathcal{F}$ if and only if $x \in H$. We verify the conditions for a dual pair.

First, B has no isolated vertices: for each subgroup in \mathcal{F} is non-trivial, so contains an element of $G \setminus \{1\}$, and every such element is contained in a subgroup in \mathcal{F} , since the union of this family is G .

Next, two subgroups are joined in the intersection graph if and only if their intersection is non-trivial (that is, contains an element of $G \setminus \{1\}$; and, by assumption, two non-trivial elements are adjacent if and only if some element of \mathcal{F} contains both.

Note that we have assumed that G is non-cyclic; this in fact follows from the fact that it is a union of proper subgroups, since a generator would lie in no proper subgroup. \square

12.3. Applications. I will consider several cases. I begin with the “classical” case, where the vertices are all the non-trivial proper subgroups of G , joined if two vertices are adjacent. These were first investigated by Csákány and Pollák, who considered non-simple groups; they determined the groups for which the intersection graph is connected and showed that, in these cases, its diameter is at most 4. For simple groups, Shen [102] showed that the graph is connected and asked for an upper bound; Herzog *et al.* [68] gave a bound of 64, which was improved to 28 by Ma [86], and to the best possible 5 by Freedman [53], who showed that the upper bound is attained only by the Baby Monster and some unitary groups (it is not currently known exactly which).

Proposition 12.3. *Let G be a non-cyclic finite group. Then the induced subgraph of the non-generating graph of G on non-identity elements and the intersection graph of G form a dual pair.*

Proof. Take \mathcal{F} to be the family of all non-trivial proper subgroups of G . \square

So the reduced non-generating graph of a non-abelian finite simple group has diameter at most 6; this bound can be reduced to 5, and possibly to 4, perhaps with specified exceptions (Saul Freedman, personal communication).

Now we turn to the commuting graph.

Proposition 12.4. *Let G be a finite group with $Z(G) = 1$. Then the reduced commuting graph of G (on the vertex set $G \setminus \{1\}$) and the intersection graph of non-trivial abelian subgroups of G form a dual pair.*

Proof. The condition $Z(G) = 1$ ensures that the reduced commuting graph does have vertex set $G \setminus \{1\}$, and also implies that G is not cyclic. Take \mathcal{F} to be the family of all non-trivial abelian subgroups of G (all are proper subgroups since G is not abelian). Two elements are joined in the reduced commuting graph if and only if the group they generate is abelian. \square

Corollary 12.5. *For a finite group G with $Z(G) = \{1\}$, the following four conditions are equivalent:*

- (a) *the Gruenberg–Kegel graph of G is connected;*
- (b) *the reduced commuting graph of G is connected;*
- (c) *the intersection graph of non-trivial abelian subgroups of G is connected;*
- (d) *the intersection graph of maximal abelian subgroups of G is connected.*

Proof. The equivalence of (a) and (b) comes from Theorem 9.2, and that of (b) and (c) from Proposition 12.1. For the equivalence of (c) and (d), note that any non-trivial abelian subgroup is contained in a maximal abelian subgroup, to which it is joined, so (d) implies (c). The converse holds because any path in the intersection graph of non-trivial abelian subgroups can be lifted to a path in the intersection graph of maximal abelian subgroups. \square

Proposition 12.6. *Let G be a group which is not cyclic or generalised quaternion. Then the induced subgraph of the enhanced power graph of G on the set of non-identity elements and the intersection graph of non-trivial cyclic subgroups of G form a dual pair.*

In fact the theorem applies also to generalised quaternion groups; but for these, both the reduced enhanced power graph and the intersection graph are connected for the trivial reason that they contain a vertex joined to all others.

Proof. We take \mathcal{F} to be the family of non-trivial cyclic subgroups of G . \square

Note that, if we take the graph $\text{DEP}(G)$ and collapse the equivalence classes of the relation \equiv , where $x \equiv y$ if $\langle x \rangle = \langle y \rangle$, we obtain the intersection graph of non-trivial cyclic subgroups of G . (This is probably why it was called the “intersection graph” in [42].)

A study of intersection graphs of cyclic subgroups has been published by Rajkumar and Devi [94].

13. More general graphs

When we think about graphs on groups, we want there to be some connection between the graph and the group. This connection is mostly expressed in terms of invariance of the graph under something, either right translations or automorphisms of the group. The first gives rise to Cayley graphs, as discussed briefly in Section 1.2.

So the focus here is on graphs on a group G invariant under the automorphism group $\text{Aut}(G)$ of G . We have seen that all graphs in the hierarchy do satisfy this condition.

There are several ways we could approach the general case.

- Any graph invariant under $\text{Aut}(G)$ is a union of orbital graphs for $\text{Aut}(G)$.

- We could define the adjacency in the graph by a first-order formula with two free variables.
- We could define adjacency by some more recondite group-theoretic property.

We will see examples below.

However, it matters whether we are defining the graph on a single group, or defining it on the class of all groups.

13.1. On a specific group. If we are given a group G , and can compute $\text{Aut}(G)$, then the first procedure (taking unions of orbital graphs) obviously gives all orbital (di)graphs for G .

Theorem 13.1. *Given a group G , for every $\text{Aut}(G)$ -invariant graph, there is a formula ϕ in the first-order language of groups such that $x \sim y$ if and only if $G \models \phi(x, y)$.*

Proof. By the so-called Ryll-Nardzewski Theorem, proved also by Engeler and by Svenonius (see [70]), G is oligomorphic, so the G -orbits on n -tuples are n -types over G , that is, maximal sets of n -variable formulae consistent with the theory of G ; but all types are principal, so each is given by a single formula. \square

Question 28. Given G , is there a bound for the complexity of the formulae defining orbital graphs for $\text{Aut}(G)$ acting on G (for example, for the alternation of quantifiers)?

Clearly the commuting graph can be defined by the quantifier-free formula $xy = yx$. If G is an elementary abelian 2-group, there are only three (non-diagonal) orbital graphs, defined by the formulae $(x = 1) \wedge (y \neq 1)$, $(x \neq 1) \wedge (y = 1)$, and $(x \neq 1) \wedge (y \neq 1) \wedge (x \neq y)$ respectively.

13.2. For classes of groups. As we have seen, the commuting graph is defined uniformly for all groups by the quantifier-free formula $xy = yx$.

It seems unlikely that the other graphs listed earlier have uniform first-order definitions. The statement $\langle x, y \rangle = G$ seems to require quantification either over words in x, y or over subsets of G , and so to need some version of higher-order logic for its definition.

For example, suppose that there is a formula $\phi(x, y)$ which, in any finite group, specifies that x and y are joined in the power graph. Taking C_n with n even, with x a generator and y of order 2, the formula is always satisfied. So it should hold in an ultraproduct of such groups (see [21]). But in the ultraproduct, x has infinite order and y has order 2, so y cannot be a power of x .

13.3. Applications. If we are given a specific group G and know its automorphism group, then constructing all the orbital graphs is a simple polynomial-time procedure.

If we are given G and don't know (and maybe are trying to find out about) its automorphism group, then clearly some indication of which first-order formulae need to be considered would be helpful. Maybe, given $g, h \in G$, the type of (g, h) (the set of $\phi(x, y)$ such that $G \models \phi(g, h)$) could be described, and a formula generating the type found.

Another situation that might arise would be that we are given one or more graphs defined on general groups and are interested to know for which groups they have some property, e.g. two graphs equal. If

we had first-order descriptions of the graphs, we would just be looking for models of some first-order sentence.

14. Infinite groups

The definitions of the graphs in the hierarchy (with the exception of the deep commuting graph), and the inclusions among them, work without change for infinite groups. I will simply mention a few highlights here, as there is little in the way of general theory.

14.1. Power graph and directed power graph. Let p be a prime number. The *Prüfer group* $G = C_{p^\infty}$ is the group of rational numbers with p -power denominators mod 1, or the multiplicative group of p -power roots of unity. Every element of the group has p -power order, and the group has a unique subgroup of order p^n for any n . It follows that the power graph of G is a countable complete graph, independent of the choice of prime.

The directed power graph does determine the prime, since the class of elements immediately above the identity has size $p - 1$. This shows that the power graph does not determine the directed power graph for infinite groups in general.

However, the implication does hold for torsion-free groups. This was shown by Zahirović [116]; a preliminary result appears in [36]. In fact, the hypotheses in Zahirović's result are weaker; I refer to the paper for details, which show the important role played by the Prüfer groups in this problem.

14.2. Independence number. Perhaps the most striking result on the commuting graph of an infinite group is the following, due to Bernhard Neumann (answering a question of Paul Erdős):

Theorem 14.1. *Let G be an infinite group. Then the following are equivalent:*

- (a) $\text{Com}(G)$ has no infinite coclique;
- (b) there is a finite upper bound on the size of cocliques in $\text{Com}(G)$;
- (c) $Z(G)$ has finite index in G .

I have stated Neumann's result like this for comparison with what follows. He proved that (a) implies (b) and (c). Now (b) implies (a) is trivial, and (c) implies (b) because if (c) holds, then G is a finite union of abelian subgroups (since $\langle Z(G), g \rangle$ is abelian for all $g \in G$), and a coclique in $\text{Com}(G)$ can contain at most one vertex from each subgroup.

What about the power graph or enhanced power graph?

Certainly, if either of these graphs has no infinite coclique, then neither does $\text{Com}(G)$; so $Z(G)$ has finite index in G . But consider the group $G = C_{p^\infty} \times C_{q^\infty}$, where p and q are distinct primes. It is easy to show that $\text{Pow}(G)$ has no infinite coclique; but, if a_n has order p^n and b_n has order q^n , then

$$\{a_n b_0, a_{n-1} b_1, \dots, a_1 b_{n-1}, a_0 b_n\}$$

is a coclique of size $n + 1$, for any n .

However, if $\text{Pow}(G)$ has no infinite coclique, then G is the union of finitely many abelian subgroups. So we first ask, which abelian groups have no infinite coclique? Such a group must be a torsion group; for, if a were an element of infinite order, then $\{a^p : p \text{ prime}\}$ is an infinite coclique in the power graph. There can only be finitely many primes such that G contains element of order p . So G is the direct sum of its finitely many Sylow subgroups. Moreover, the Sylow subgroups must have finite rank. So we can conclude:

Theorem 14.2. *Let G be an infinite group. Then the following are equivalent:*

- (a) $\text{Pow}(G)$ has no infinite coclique;
- (b) $Z(G)$ has finite index in G and is a direct sum of finitely many p -torsion subgroups of finite rank, for primes p .

So G is locally finite, a result of Shitov [103].

If we make the stronger hypothesis that the size of cocliques is bounded, then we can strengthen the conclusion to assert that all but one of the Sylow subgroups of $Z(G)$ is finite.

For the enhanced power graph, Abdollahi and Hassanabadi [5] proved that the analogue of Neumann's Theorem does hold:

Theorem 14.3. *Let G be an infinite group. Then the following are equivalent:*

- (a) $\text{EPow}(G)$ has no infinite coclique;
- (b) there is a finite upper bound for the size of cocliques in $\text{EPow}(G)$;
- (c) $Z_{\text{EPow}}(G)$ has finite index in G .

(Recall that $Z_{\text{EPow}}(G)$ is the cyclicizer of G .)

14.3. Cliques and colourings of the power graph. Another significant body of work on power graphs concerns the clique parameters. Here are some striking results, which appear in [1, 37, 103].

Theorem 14.4. *The power graph of an infinite group has clique number and chromatic number at most countable.*

Theorem 14.5. *For an infinite group G , the following conditions are equivalent:*

- (a) $\text{Pow}(G)$ has finite clique number;
- (b) $\text{Pow}(G)$ has finite chromatic number;
- (c) $\text{EPow}(G)$ has finite clique number;
- (d) $\text{EPow}(G)$ has finite chromatic number;
- (e) G is a torsion group with finite exponent.

Proof. The power graph of an infinite cyclic group $\langle g \rangle$ contains an infinite clique $\{g^{2^n} : n \geq 0\}$. So a group satisfying any of the first four conditions is a torsion group. Now the results are proved just as for finite groups in Section 4. \square

The cited papers contain other miscellaneous results.

14.4. **Cographs.** Cographs work quite differently in the infinite case. The two definitions earlier (a graph containing no induced P_4 , and a graph built from the 1-vertex graph by disjoint union and complementation) are no longer equivalent, even if infinite disjoint unions are allowed. Covington [45] constructed a countable P_4 -free graph which is isomorphic to its complement (so it and its complement are both connected). This remarkable object also has a high degree of symmetry.

15. Beyond groups

The ideas behind some of these graphs can be extended to other algebraic structures.

A *magma* is a set with a binary operation. (The term *groupoid* is sometimes used, but I will avoid this since it is also used for a category in which every morphism is invertible.) Beyond groups, the two classes of magmas most studied are *semigroups* (satisfying the associative law) and *quasigroups* (in which left and right division are well-defined), and in particular *monoids* and *loops* (semigroups, resp. quasigroups, with identity elements).

Clearly the definition of commuting graph makes sense in any magma. For the power graph and its relatives, it is necessary to make sense of powers of an element. One can define *left powers* inductively by $a^1 = a$ and $a^{n+1} = a \circ a^n$ for $n \geq 1$. (This is the approach adopted in [109].) An alternative is to restrict to *power-associative magmas*, those in which the product of n terms each equal to an element a is independent of the bracketing used to evaluate it. Now, just as for groups, we have:

Proposition 15.1. *In a power-associative magma, the directed power graph is a partial preorder, and so the power graph is the comparability graph of a partial order.*

Question 29. For which magmas, or quasigroups, is the power graph (defined using left powers) a comparability graph of a partial order, or a perfect graph?

The power graph of a semigroup was defined early in their study of power graphs, see [77].

The commuting graphs of semigroups are considered by Araújo *et al.* [12], who pose a number of questions about them.

As for groups, the power graph of a semigroup is a spanning subgraph of its commuting graph. The enhanced power graph could be defined for any semigroup; to my knowledge this has not been studied. It is not clear whether the definition of the deep commuting graph could be adapted for semigroups.

Commuting graphs of semigroups are universal (and power graphs are universal for comparability graphs of partial orders), since these statements hold for groups.

The intersection graph of the subsemigroups of a semigroup had been studied much earlier: Bosák [27] raised the question of its connectedness in 1963, and the question was soon resolved by Lin [82] and Pondělíček [93]: for any finite semigroup, this graph is connected with diameter at most 3. These results preceded the investigation of the intersection graph for groups, mentioned earlier. (The intersection graphs of semigroups are not comparable with the intersection graphs of groups, since any subgroup of a group contains the identity, so adjacency requires their intersection to be non-trivial.)

The commuting graph (and its complement), the intersection graph of cyclic subgroups, and the power graph have also been studied for quasigroups and loops, especially for the classes of Moufang and Bol loops: see, for example, [10, 66, 67, 109]. Moufang loops form a class of loops which is perhaps closest to groups: a Moufang loop is a loop satisfying the identity $z(x(z y)) = ((z x) z) y$ (a weakening of the associative law). In particular, a 2-generated subloop of a Moufang loop is associative; so a Moufang loop is power-associative, and its power graph is the comparability graph of a partial order.

A question raised some time ago but to my knowledge not yet answered is:

Question 30. If the power graphs of two finite Moufang loops are isomorphic, are their directed power graphs isomorphic?

This question has been investigated by Nick Britten, who has found no counterexamples among the Moufang loops in the LOOPS package [89] for GAP (Michael Kinyon, personal communication).

Going beyond a single binary operation, we reach the class of rings. For these, the *zero-divisor graph* of a ring was introduced by Beck [20] in 1988: the vertices are the ring elements, with a and b joined whenever $ab = 0$. If the ring is commutative, the graph is undirected. Another graph associated with a ring is the *unit graph*, in which a and b are joined whenever $a + b$ is a unit [14].

REFERENCES

- [1] G. Aalipour, S. Akbari, P. J. Cameron, R. Nikandish and F. Shaveisi, On the structure of the power graph and the enhanced power graph of a group, *Electron. J. Combin.*, **24** (2017) PP. 18.
- [2] J. Abawajy, A. Kelarev and M. Chowdhury, Power graphs: a survey, *Elec. J. Graph Theory Appl.*, **1** (2013) 125–147.
- [3] A. Abdollahi, Engel graph associated with a group, *J. Algebra*, **318** (2007) 680–691.
- [4] A. Abdollahi, S. Akbari and H. R. Maimani, Noncommuting graph of a group, *J. Algebra*, **298** (2006) 468–492.
- [5] A. Abdollahi and A. Mohammadi Hassanabadi, Noncyclic graph of a group, *Commun. Algebra*, **35** (2007) 2057–2081.
- [6] A. Abdollahi and A. Mohammadi Hassanabadi, Non-cyclic graph associated with a group, *J. Algebra Appl.*, **8** (2009) 243–257.
- [7] A. Abdollahi and H. Shahverdi, Characterization of the alternating group by its non-commuting graph, *J. Algebra*, **357** (2012) 203–207.
- [8] A. Abdollahi and Mohammed Zarrin, Non-nilpotent graph of a group, *Commun. Algebra*, **38** (2010) 4390–4403.
- [9] M. Afkhami, M. Farrokhi and K. Khashyarmansh, Planar, toroidal and projective commuting and noncommuting graphs, *Comm. Algebra*, **43** (2015) 2964–2970.
- [10] Karim Ahmadidilir, On the non-commuting graph in finite Moufang loops, *J. Algebra Appl.*, **17** (2018) 1850070.
- [11] O. A. Alekseeva and A. S. Kondrat’ev, On recognizability of some finite simple orthogonal groups by spectrum, *Proc. Steklov Inst. Math.*, **266** (2009) 10–23.
- [12] J. Araújo, M. Kinyon and J. Konieczny, Minimal paths in the commuting graphs of semigroups, *Europ. J. Combinatorics*, **32** (2011) 178–197.
- [13] A. R. Ashrafi, A. Gholami and Z. Mehranian, Automorphism group of certain power graphs of finite groups, *Electronic J. Graph Theory Appl.*, **5** (2017) 70–82.
- [14] N. Ashrafi, H. R. Maimani, M. R. Pournaki and S. Yassemi, Unit graphs associated with rings, *Commun. Algebra*, **38** (2010) 2851–2871.
- [15] R. Baer, Engelsche Elemente Noetherscher Gruppen, *Math. Ann.*, **133** (1957) 256–270.

- [16] R. A. Bailey, *Association Schemes: Designed Experiments, Algebra and Combinatorics*, Cambridge Studies in Advanced Mathematics, **84**, Cambridge University Press, Cambridge, 2004.
- [17] R. A. Bailey and Peter J. Cameron, Using graphs to find the best block designs, in *Topics in Structural Graph Theory* (ed. L. W. Beineke and R. J. Wilson), Encyclopedia of Mathematics and its Applications, **147**, Cambridge University Press, Cambridge, (2013) 282–317.
- [18] A. Ballester-Bolinches and R. Esteban-Romero, On minimal non-supersoluble groups, *Rev. Mat. Iberoamericana*, **23** (2007) 127–142.
- [19] A. Ballester-Bolinches, R. Esteban-Romero and Derek J. S. Robinson, On finite minimal non-nilpotent groups, *Proc. Amer. Math. Soc.*, **133** (2005) 3455–3462
- [20] István Beck, Coloring of commutative rings, *J. Algebra*, **116** (1988) 208–226.
- [21] J. L. Bell and A. B. Slomson, *Models and Ultraproducts: An Introduction*, Dover Publ., New York, 2006 (reprint of 1974 ed.)
- [22] S. Bera, Hiranya Kishore Dey and Sajal Kumar Mukherjee, On the connectivity of enhanced power graphs of finite groups, *Graphs Combin.*, in press; doi: 10.1007/s00373-020-02267-5
- [23] S. R. Blackburn, P. M. Neumann and G. Venkataraman, *Enumeration of Finite Groups*, Cambridge Univ. Press, Cambridge, 2007.
- [24] A. Blass, G. Exoo and F. Harary, Paley graphs satisfy all firstorder adjacency axioms, *J. Graph Theory*, **5** (1981) 435–439.
- [25] F. A. Bogomolov, The Brauer group of quotient spaces by linear group actions, *Izv. Akad. Nauk SSSR Ser. Mat.*, **51** (1987) 485–516.
- [26] B. Bollobás and A. Thomason, Graphs which contain all small graphs, *Europ. J. Combinatorics*, **2** (1981) 13–15.
- [27] J. Bosák, The graphs of semigroups, *Theory of Graphs and its Applications* (Proc. Symp. Smolenice 1963), Academic Press, New York, 1965 119–125
- [28] R. Brauer and K. A. Fowler, On groups of even order, *Ann. Math.*, **62** (1955) 565–583.
- [29] T. Breuer, R. M. Guralnick and W. M. Kantor, Probabilistic generation of finite simple groups II, *J. Algebra*, **320** (2008) 443–494.
- [30] J. R. Britnell and N. Gill, Perfect commuting graphs, *J. Group Theory*, **20** (2017) 71–102.
- [31] T. C. Burness, R. M. Guralnick and S. Harper, The spread of a finite group, *Ann. of Math. (2)*, **193** (2021) 619–687. <https://arxiv.org/abs/2006.01421>
- [32] Peter J. Cameron, *Coherent configurations, association schemes, and permutation groups*, Groups, combinatorics geometry (Durham, 2001), World Sci. Publ., River Edge, NJ, 2003 55–71.
Coherent configurations, association schemes and permutation groups. Groups, combinatorics geometry (Durham, 2001), 55–71, World Sci. Publ., River Edge, NJ, 2003
- [33] Peter J. Cameron, The power graph of a finite group, II, *J. Group Theory*, **13** (2010) 779–783.
- [34] Peter J. Cameron, S. D. Freedman and C. M. Roney-Dougal, The non-commuting, non-generating graph of a nilpotent group, *Electronic J. Combinatorics*, **28** (2021) pp. 15.
- [35] Peter J. Cameron and S. Ghosh, The power graph of a finite group, *Discrete Math.*, **311** (2011) 1220–1222.
- [36] Peter J. Cameron, H. Guerra and S. Jurina, The power graph of a torsion-free group, *J. Algebraic Combin.*, **49** (2019) 83–98.
- [37] Peter J. Cameron and Sayyed Heidar Jafari, On the connectivity and independence number of power graphs of groups, *Graphs Combin.*, **36** (2020) 895–9904.
- [38] Peter J. Cameron and Bojan Kuzma, Between the enhanced power graph and the commuting graph, <https://arxiv.org/2012.03789>.
- [39] Peter J. Cameron, A. Lucchini and Colva M. Roney-Dougal, Generating sets of finite groups, *Trans. Amer. Math. Soc.*, **370** (2018) 6751–6770.

- [40] P. J. Cameron, P. Manna and R. Mehatari, Forbidden subgraphs of power graphs, <https://arxiv.org/abs/2010.05198>.
- [41] P. J. Cameron and N. Maslova, Criterion of unrecognizability of a finite group by its Gruenberg–Kegel graph, <https://arxiv.org/abs/2012.01482>.
- [42] T. T. Chelvam and M. Sattantathan, Subgroup intersection graph of finite abelian group, *Trans. Com.*, **1** (2012) 5–10.
- [43] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, *Ann. Math.*, **164** (2006) 51–229.
- [44] H. Cohn, *Number Theory*, Graduate Texts in Mathematics, **240**, Springer, New York, 2007.
- [45] J. Covington, A universal structure for N -free graphs, *Proc. London Math. Soc.* (3), **58** (1989) 1–16.
- [46] Ch. Garnet Cox, On the spread of infinite groups, <https://arxiv.org/abs/2012.12197>.
- [47] B. Csákny and G. Pollák, The graph of subgroups of a finite group. *Czechoslovak Math. J.*, **19** (1969) 241–247.
- [48] P. Diaconis and M. Simper, Statistical enumeration of groups by double cosets, <https://arxiv.org/abs/2102.04576>.
- [49] Robert P. Dilworth, A decomposition theorem for partially ordered sets, *Ann. Math.*, **51** (1950) 161–166.
- [50] S. Eberhard, Commuting probabilities of finite groups, *Bull. London Math. Soc.*, **47** (2015) 796–808.
- [51] G. Ellis, HAP, Homological Algebra Programming, Version 1.29 (2021), <https://gap-packages.github.io/hap>.
- [52] M. Feng, X. Ma, and K. Wang, The full automorphism group of the power (di)graph of a finite group, *Europ. J. Combinatorics* **52** (2016) 197–206. ,
- [53] S. D. Freedman, The intersection graph of a finite simple group has diameter at most 5, *Arch. Math.*, in press; doi [10.1007/s00013-021-01583-3](https://doi.org/10.1007/s00013-021-01583-3)
- [54] T. Gallai, Transitiv orientierbare Graphen, *Acta Math. Acad. Sci. Hung.*, **18** (1967) 25–66.
- [55] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.11.0; 2020, <https://www.gap-system.org>.
- [56] M. Giudici and C. W. Parker, There is no upper bound for the diameter of the commuting graph of a finite group, *J. Combinatorial Theory (A)*, **120** (2013) 1600–1603.
- [57] G. Glauberman, Central elements in core-free groups, *J. Algebra*, **4** (1996) 403–420.
- [58] C. D. Godsil, *Association Schemes*, <https://www.math.uwaterloo.ca/~cgodsil/pdfs/assoc2.pdf>.
- [59] D. Gorenstein and J. H. Walter, The characterization of groups with dihedral 2-subgroups. III., *J. Algebra*, **2** (1965) 354–393.
- [60] Robert L. Griess Jr., Schur multipliers of the known finite simple groups, *Bull. Amer. Math. Soc.*, **78** (1972) 68–71.
- [61] K. W. Gruenberg and K. W. Roggenkamp, Decomposition of the augmentation ideal and of the relation modules of a finite group, *Proc. London Math. Soc.* (3), **31** (1975) 149–166.
- [62] R. Guralnick, B. Kunyavskii, E. Plotin and A. Shalev, Thompson-line characterization of the solvable radical, *J. Algebra* **300** (2006) 363–375.
- [63] R. M. Guralnick and G. R. Robinson, On the commuting probability in finite groups, *J. Algebra*, **300** (2006) 509–528.
- [64] M. Hall, Jr., *The Theory of Groups*, Macmillan, New York, 1959.
- [65] Z. Han, G. Chen and X. Guo, A characterization theorem for sporadic simple groups, *Siberian Math. J.*, **49** (2008) 1138–1146.
- [66] H. Hasanzadeh Bashir and K. Ahmadidilir, Some structural graph properties of the commuting graph of a class of finite Moufang loops, *Electronic J. Graph Theory Appl.*, **8** (2020) 319–337.
- [67] Hamideh. Hasanzadeh Bashir and A. Iranmanesh, The intersection graph of a finite Moufang loop, *J. Math. Extension*, **14** (2020) 81–90.
- [68] M. Herzog, P. Longobardi and M. Maj, On a graph related to the maximal subgroups of a group, *Bull. Austral. Math. Soc.*, **81** (2010) 317–328.

- [69] G. Higman, *Odd characterizations of finite simple groups*, Lecture notes, University of Michigan, 1968 pp. 77.
- [70] W. Hodges, *A Shorter Model Theory*, Cambridge Univ. Press, Cambridge, 1997.
- [71] X. Huang, Q. Huang and S. M. Cioabă, The second eigenvalue of some normal Cayley graphs of high transitive groups, *Electronic J. Combinatorics*, **26** (2019) pp. 26.
- [72] A. Iranmanesh and A. A. Jafarzadeh, On the commuting graph associated with the symmetric and alternating groups, *J. Algebra Appl.*, **7** (2008) 129–146.
- [73] , M. Jerrum, *Computational Pólya theory*, in *Surveys in Combinatorics 1995* (ed. Peter Rowlinson), London Math. Soc. Lecture Note Series, **218**, Cambridge Univ. Press, Cambridge, (1995) 103–118.
- [74] U. Jezernik and P. Moravec, Commutativity preserving extensions of groups, *Proc. Roy. Soc. Edinburgh Sect. A*, **148** (2018) 575–592.
- [75] H. A. Jung, On a class of posets and the corresponding comparability graphs, *J. Combinatorial Theory Ser. B*, **24** (1978) 125–133.
- [76] A. V. Kelarev and S. J. Quinn, A combinatorial property and power graphs of groups, (Proc. of the Vienna Conference, Vienna, 1999), *Contrib. General Algebra*, **12** (2000) 229–235.
- [77] A. V. Kelarev and S. J. Quinn, Directed graph and combinatorial properties of semigroups, *J. Algebra*, **251** (2002) 16–26.
- [78] A. S. Kondrat'ev and V. D. Mazurov, Recognition of alternating groups of prime degree from the orders of their elements, *Siberian Math. J.*, **41** (2000) 294–302.
- [79] B. E. Kunyavskii, *The Bogomolov multiplier of finite simple groups*, In *Cohomological and geometric approaches to rationality problems*, Progress in Mathematics, **282** (2010) 209–217.
- [80] B. Larose, F. Laviolette and C. Tardif, On normal Cayley graphs and hom-independent graphs, *Europ. J. Combinatorics*, **19** (1998) 867–881.
- [81] M. W. Liebeck and A. Shalev, Simple groups, probabilistic methods, and a conjecture of Kantor and Lubotzky, *J. Algebra*, **184** (1996) 31–57.
- [82] Y.-F. Lin, A problem of Bosák concerning the graphs of semigroups, *Proc. Amer. Math. Soc.*, **21** (1969) 343–346.
- [83] L. Lovász, Balanced hypergraphs and the Perfect Graph Conjecture, *Discrete Math.*, **2** (1972) 253–267.
- [84] A. Lucchini and A. Maróti, On the clique number of the generating graph of a finite group, *Proc. Amer. Math. Soc.*, **137** (2009) 3207–3217.
- [85] A. Lucchini and D. Nemi, The diameter of the non-nilpotent graph of a finite group, *Trans. Comb.*, **9** (2020) 111–114.
- [86] X. Ma, On the diameter of the intersection graph of a finite simple group, *Czechoslovak Math. J.*, **66** (2016) 365–370.
- [87] G. A. Miller and H. C. Moreno, Non-abelian groups in which every subgroup is abelian, *Trans. Amer. Math. Soc.*, **27** (1903) 398–404.
- [88] G. L. Morgan and C. W. Parker, The diameter of the commuting graph of a finite group with trivial centre, *J. Algebra*, **393** (2013) 41–59.
- [89] G. Nagy and P. Vojtěchovský, The LOOPS package for GAP, <https://gap-packages.github.io/loops/>.
- [90] J. A. Nelder, The analysis of randomized experiments with orthogonal block structure, I; Block structure and the null analysis of variance, *Proc. Royal Soc. London (A)*, **283** (1965) 147–162.
- [91] B. H. Neumann, A problem of Paul Erdős on groups, *J. Austral. Math. Soc. (A)*, **21** (1976) 467–472.
- [92] D. Patrick and E. Wepsic, *Cyclicizers, centralizers and normalizers*, Tech. Report MS-TR 91-05, Rose-Hulman Institute of Technology, IN, USA, 1991.
- [93] B. Pondělíček, Diameter of a graph of a semigroup, *Časopis Pěst. Mat.*, **92** (1967) 206–211.
- [94] R. Rajkumar and P. Devi, Intersection graphs of cyclic subgroups of groups, *Electronic Notes Discrete Math.*, **53** (2016) 15–24.
- [95] G. Ravindra and K. R. Parthasarathy, Perfect product graphs, *Discrete Math.*, **20** (1977) 177–186.

- [96] B. Reed, A semi-strong perfect graph theorem, *J. Combinatorial Theory (B)*, **43** (1987) 223–240.
- [97] Derek J. S. Robinson, *A Course in the Theory of Groups*, (2nd edition), Springer, New York, 1996.
- [98] O. Yu. Schmidt, Groups all of whose subgroups are nilpotent, *Mat. Sbornik*, (Russian), **31** (1924) 366–372.
- [99] I. Schur, Über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen, *J. reine angew. Math.*, **127** (1904) 20–50.
- [100] Y. Segev, The commuting graph of minimal nonsolvable groups, *Geometriae Dedicata*, **88** (2001) 55–66.
- [101] D. Seinsche, On a property of the class of n -colorable graphs, *J. Combinatorial Theory (B)*, **16** (1974) 191–193.
- [102] R. Shen, Intersection graphs of subgroups of finite groups, *Czechoslovak Math. J.*, **60** (2010) 945–950.
- [103] Y. Shitov, Coloring the power graph of a semigroup, *Graphs Combin.*, **33** (2017) 485–487.
- [104] L. H. Soicher, The GRAPE package for GAP, Version 4.8.3, 2019, <https://gap-packages.github.io/grape>.
- [105] R. M. Solomon and A. J. Woldar, Simple groups are characterized by their non-commuting graphs, *J. Group Theory*, **16** (2013) 793–824.
- [106] T. P. Speed and R. A. Bailey, *On a class of association schemes derived from lattices of equivalence relations*, in: *Algebraic Structures and Applications*, (ed. P. Schulz, C. E. Praeger and R. P. Sullivan), M. Dekker, New York, (1982) 55–74.
- [107] D. P. Sumner, Dacey graphs, *J. Aust. Math. Soc.*, **18** (1974) 492–502.
- [108] J. G. Thompson, Nonsolvable finite groups all of whose local subgroups are solvable (Part I), *Bull. Amer. Math. Soc. (NS)*, **74** (1968) 383–437.
- [109] D. L. Walker, *Power Graphs of Quasigroups*, Graduate thesis, University of South Florida, 2019; <https://scholarcommons.usf.edu/etd/7984>.
- [110] J. H. Walter, The characterization of finite groups with abelian Sylow 2-subgroups, *Ann. Math.*, **89** (1969) 405–514.
- [111] H. Wielandt, *Finite Permutation Groups*, Academic Press, New York, 1964.
- [112] J. S. Williams, Prime graph components of finite groups, *J. Algebra*, **69** (1981) 487–513.
- [113] R. A. Wilson, et al., *Atlas of Finite Group Representations*, version 3, <http://brauer.maths.qmul.ac.uk/Atlas/v3/>.
- [114] A. Woldar, A combinatorial approach to the character theory of split metabelian groups, *J. Combinatorial Theory (A)*, **50** (1989) 100–109.
- [115] M.-Y. Xu, Automorphism groups and isomorphisms of Cayley digraphs *Discrete Math.*, **182** (1998) 309–319.
- [116] S. Zahirović, *The power graph of a torsion-free group determines the directed power graph*, <https://arxiv.org/abs/2006.01984>.
- [117] S. Zahirović, I. Bošnjak and R. Madarász, A study of enhanced power graphs of finite groups, *J. Algebra Appl.*, **19** (2020) pp. 20.
- [118] M. Zorn, Nilpotency of finite groups, *Bull. Amer. Math. Soc.*, **42** (1936) 485–486.

Peter J. Cameron

School of Mathematics and Statistics, University of St Andrews, UK

Email: pjc20@st-andrews.ac.uk