

Title: Cost burden of *Clostridioides difficile* infection to the health service: A retrospective cohort study in Scotland

Authors: Chris Robertson^{1,2}, Jiafeng Pan¹, Kim Kavanagh¹, Ian Ford³, Colin McCowan⁴, Marion Bennie^{5,6}, Charis Marwick⁷, Alistair Leanord^{8,9*}

Affiliations: ¹Department of Mathematics and Statistics, University of Strathclyde, Glasgow, UK; ²Health Protection Scotland, NHS National Services Scotland, Glasgow, UK; ³Robertson Centre for Biostatistics, University of Glasgow, Glasgow, UK; ⁴School of Medicine, University of St Andrews, St Andrews, UK; ⁵Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK; ⁶Public Health and Intelligence, NHS National Services Scotland, Edinburgh, UK; ⁷Division of Population Health and Genomics, School of Medicine, University of Dundee, Dundee, UK; ⁸Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK; ⁹NHS Greater Glasgow and Clyde, Glasgow, UK

Target journal: Journal of Hospital Infection

Word count (max. 4000 words; each figure and/or table contributes 200 words towards the total, therefore current number of figures/tables = 1200 words): 2899 words + 1200 allowance for figures/tables = 4099

Abstract (max. 250 words): 246

Number of proposed tables: 2

Number of proposed figures: 4

References (no max.): 24

Supplementary data: 4 tables

Running title (max. 40 characters and spaces): 34 characters and spaces; NHS cost burden of CDI in Scotland

Keywords (max. 6): *Clostridioides difficile* infection, cost burden, hospital infection, community infection, retrospective cohort

*Corresponding author:

Professor Alistair Leanord

Address:

Scottish Microbiology Reference Laboratories, Glasgow

Level 5, New Lister Building

Glasgow Royal Infirmary

10-16 Alexandra Parade

Glasgow, G31 2ER

Tel.: 0141 201 8633

Email: alistair.leanord@ggc.scot.nhs.uk

Abbreviations

CDI: *Clostridioides difficile* infection

CHI: Community Health Index

CI: confidence interval

ECOSS: Electronic Communication of Surveillance in Scotland

HOCA-CDI: hospital onset community-associated CDI

HOHA-CDI: hospital onset hospital-associated CDI

HR: hazard ratio

LOS: length of stay

LQ: lower quartile

NHS: National Health Service

PIS: Prescribing Information System

UQ: upper quartile

Summary

Word count: 246 / 250 words

Background: *Clostridioides difficile* infection (CDI) is associated with high healthcare demands and related costs.

Aim: To evaluate the healthcare and economic burden of CDI in hospitalised patients with community- (HOCA-CDI) or hospital-associated CDI (HOHA-CDI) in the National Health Service in Scotland.

Methods: A retrospective cohort study was conducted, examining data between August 2010 and July 2013 from four patient-level Scottish datasets, linked to death data. Data examined included prior antimicrobial prescriptions in the community, hospitalisations, length of stay and mortality. Each CDI case was matched to three hospital-based controls on the basis of age, gender, hospital and date of admission. Descriptive economic evaluations were based on bed-day costs for different types of wards.

Findings: Overall, 3304 CDI cases were included in the study. CDI was associated with additional median lengths of stay of 7.2 days for HOCA-CDI and 12.0 days for HOHA-CDI compared with their respective, matched controls. The 30-day mortality rate was 6.8% for HOCA-CDI and 12.4% for HOHA-CDI. Overall, recurrence within 90 days of the first CDI episode occurred in 373/2740 (13.6%) survivors. The median additional expenditure for each initial CDI case compared with matched controls was £1713. In the 6 months after the index hospitalisation, the cost associated with a CDI case was £5126 higher than for controls.

Conclusion: Using routinely collected national data, we demonstrate the substantial burden of CDI on healthcare services, including lengthy hospital stays and readmissions, which increase the costs of managing patients with CDI compared with matched controls.

Funded by Astellas Europe, Ltd.

Keywords (max. 6): *Clostridioides (Clostridium) difficile* infection, cost burden, hospital infection, community infection, retrospective cohort

Introduction

Clostridioides difficile infection (CDI) is the most common cause of healthcare-associated diarrhoea in developed countries and a major public health problem [1]. It is associated with high healthcare demands and related costs, with recurrence of infection and the need for readmission to hospital contributing to the burden of CDI [2]. Moreover, a recent study estimated an excess 30-day all-cause mortality of 10.9% in patients with CDI compared with matched patients without CDI [3]. While CDI is mainly a hospital-associated infection, a substantial proportion of cases have onset in or association with the community setting [4]. It is, therefore, important to explore both hospital- and community-associated CDI to obtain a comprehensive account of its impact on healthcare burden.

In a study from Germany, the direct treatment cost of CDI was €18,463, compared with €14,531 for controls, with CDI prolonging the mean overall length of stay (LOS) by 8 days [5]. A study from Scotland found that LOS was extended by ≥15.8% in patients aged ≥65 years with CDI compared with younger patients aged 15–64 years with CDI [6]. Furthermore, recurrence of CDI appears to prolong the LOS compared with an initial CDI episode, with some estimating a doubling of LOS [7]. Consequently, recurrent CDI substantially increases healthcare costs [7,8]. Dubberke et al. estimated the attributable inpatient cost of recurrent CDI to be US\$11,631 over the 6-month period after the initial CDI episode [9].

Owing to the considerable economic and healthcare burden of CDI, including recurrence, an accurate assessment of associated healthcare demands could help in the planning and prioritisation of resources.

Methods

Study design and data collection

This was a cohort study, which examined linked data from four patient-level Scottish datasets. Scotland has mandatory reporting of CDI cases to Health Protection Scotland, with data held on the Electronic Communication of Surveillance in Scotland (ECOSS) system. Healthcare encounters also have mandatory recording of a unique patient identifier [Community Health Index (CHI) number], enabling patient-level linkage of data. Each CDI case from the diagnostic laboratories in Scotland were linked to community-based antimicrobial prescriptions and hospital admissions data. Prescriptions for antimicrobials dispensed in primary care were obtained from the Prescribing Information System (PIS). The PIS captures 96% of CHIs for all prescriptions dispensed in the community in Scotland [10]. Hospital admissions data were obtained from the general/acute and inpatient day case Scottish Morbidity Record (SMR01) dataset [11]. Mortality data were obtained from National Records of Scotland and were linked to SMR01.

Using ECOSS data, all CDI cases aged ≥ 15 years diagnosed during a hospital stay were identified covering the period between 1 August 2010 and 31 July 2013. Each hospital-onset CDI case was matched to up to three hospital-based controls, who were hospitalised at the same time as the case, on the basis of age (± 5 years), gender, hospital and date of admission (± 7 days). The index date for comparisons of outcomes for the controls was taken to be the date of diagnosis for the matched case.

All data analysed were de-identified before being made available to researchers for analysis in the National Safe Haven, a secure repository for storing and sharing data for health research. Ethics permission for this study was not required as it used only non-identifiable routine data. Approval for the study was granted by the National Health Service (NHS) National Services Scotland Privacy Advisory Committee (study number XRB14037).

Definitions

A CDI case was defined as a hospitalised patient who tested positive for *C. difficile* toxin in stool concurrent with diarrhoea not attributable to any other cause, or a patient with *C. difficile* culture-positive stool concurrent with pseudomembranous colitis diagnosis. Cases were defined as hospital onset community-associated (HOCA-CDI) if the diagnostic sample was taken 0–2 days after admission. CDI diagnosed from samples taken from day 3 of admission was defined as hospital onset hospital-associated (HOHA-CDI). If a patient was diagnosed twice with CDI within a 28-day period, the second positive test was considered part of the same infection episode. CDI recurrence was defined as a second CDI diagnosis ≥ 28 days after the initial positive CDI test. Presumed cure of CDI occurred for patients who had a period of ≥ 28 days after the first positive *C. difficile* test with no positive result and no death. Any result reported in the ECOSS system as equivocal (i.e. positive for glutamate dehydrogenase but negative for *C. difficile* toxin) was removed from the final dataset.

The exact date of CDI symptom onset is not systematically reported. A proxy for the onset date of CDI from the ECOSS system was the date that specimens were collected from patients or, when this was unavailable, the date that specimens were received by the testing laboratory or the date that results were reported.

Assessments and statistical analyses

LOS, additional LOS owing to CDI, mortality rates for ≤ 2 months post-CDI diagnosis and the recorded cause of death were assessed in cases and matched controls. If cases had multiple CDI episodes, one episode was randomly selected for analysis of LOS, readmission and survival. Additional LOS for both cases and controls was analysed using a multivariable parametric survival regression model with a log-normal distribution assumed for time to event. A robust sandwich covariance matrix estimator was used to account for the matched design. Additional LOS due to CDI was calculated by comparing the median outcomes predicted by the survival regression model, adjusting for deprivation, comorbidities and time

in hospital before CDI, for the cases and controls. The 95% confidence intervals (CIs) for the additional LOS was obtained by parametric bootstrapping 500 times. Patients who were transferred to another hospital environment, ending the continuous inpatient stay, were censored at the date of transfer.

Analyses of CDI recurrence and its determinants included cases who survived the first CDI episode (i.e., did not die or were not censored within 28 days after the initial CDI diagnosis). Analysis of secondary CDI recurrence included all cases who survived at least two CDI recurrences. Multivariable competing risks models [12,13] were used to account for mortality when investigating recurrence. A sensitivity analysis was conducted in relation to CDI recurrence, replacing 28 days with 6 and 8 weeks as the presumed date of cure.

A multivariable Cox proportional hazards survival model was applied to investigate the determinants of hospital readmission. The analysis was stratified by whether the first discharge was pre- or post-30 days after CDI diagnosis. Readmission rates within 6 months after CDI diagnosis and factors that may predict readmission were assessed in cases who were discharged from hospital before censor/date of death.

A multivariable Cox proportional hazards survival model was used to compare survival between cases and matched controls. Matched cases and controls were included as strata in the model providing a different baseline hazard for each stratum to account for the matched design.

In the analysis of potential risk factors, the following were recorded: comorbidities for the index admission; comorbidities reported in hospital admissions in the previous 5 years; number of hospital admissions in the previous year; number of antimicrobial prescriptions in primary care, total number of prescriptions (all drugs) and number of different prescriptions (based on approved name) dispensed in primary care in the previous year; Charlson comorbidity index [14–16] based on ICD-10 discharge codes from all hospital admissions in the 5 years prior to the CDI episode; care home residency (yes/no); and Scottish Index of Multiple Deprivation (SIMD) quintile [17].

Cost analysis included LOS for both cases and matched controls. The first CDI episode was selected for the cost analysis if cases had multiple episodes. Costs were based on daily costs for different types of beds, such as beds on a general or critical care ward; see **Supplementary Table A1**. Descriptive economic evaluations were conducted for the NHS cost of managing the index CDI episode, the additional cost of managing the index CDI episode compared with matched controls, and NHS costs for hospital stay in the 6 months following the index CDI/matched control episodes.

All statistical analyses were stratified by HOHA-CDI and HOCA-CDI. Analyses were performed using statistical software R (version 3.2.1) [18].

Results

Cases and matched controls

There were 1297 HOCA-CDI cases and 3980 matched controls, and 2007 HOHA-CDI cases and 5536 matched controls. The identification and selection of cases and controls are shown in **Figure 1**.

A greater number of previous hospital admissions and prescriptions for any medications were recorded in the cases, but other covariates were reasonably balanced between cases and controls (**Table I** and **Supplementary Table A1**).

Outcomes

Length of stay

Adjusting for comorbidities on admission, previous admissions and social deprivation, the median LOS was more than double for CDI patients compared with controls and HOCA-CDI cases had an additional median LOS of 7.2 days (95% CI 4.7–12.0), while HOHA-CDI cases had an additional 12.0 days (95% CI 7.7–18.9) compared with their respective controls (**Table II**).

CDI recurrence

There were 1055 HOCA-CDI and 1685 HOHA-CDI cases eligible for analysis of recurrence (**Figure 1**). Overall, recurrence within 90 days of the first CDI episode occurred in 373/2740 (13.6%) survivors (**Figure 2**). Recurrence rates over the 1-year period after the initial CDI diagnosis were similar for patients with HOCA-CDI and HOHA-CDI (**Supplementary Table AIII**). Recurrence occurred in 161/1055 (15.3%) HOCA-CDI cases and 212/1685 (12.6%) HOHA-CDI cases. The hazard ratio (HR, 95% CI) for recurrence relative to patients aged ≤ 64 years was 1.51 (0.94–2.43) in patients aged 65–74 years, 1.98 (1.26–3.12) in patients aged 75–84 years and 2.16 (1.30–3.60) in patients aged ≥ 85 years.

After the first recurrent CDI episode, 371 cases were presumed cured. Rates of a second CDI recurrence in the next 12 months were 48/196 (24.6%) for cases with initial HOCA-CDI and 61/175 (34.8%) for cases with initial HOHA-CDI.

The sensitivity analyses revealed that the recurrence rates had little variation if replacing 28 days post-infection with 6 and 8 weeks as the presumed date of cure (**Supplementary Table AIII**).

Readmission to hospital

There were 868 HOCA-CDI and 844 HOHA-CDI cases that had been discharged from hospital within 30 days of the initial CDI episode and were eligible for analysis of readmission. Readmission within 6 months of the CDI episode occurred in 526/868 (60.6%) HOCA-CDI cases and 489/844 (57.9%) HOHA-CDI cases. The reason for readmission did not differ substantially from the reason for index admission (**Supplementary Table AIV**). For patients discharged from hospital ≥ 30 days after the initial CDI episode, 71/129 (55.0%) HOCA-CDI cases and 263/497 (52.9%) HOHA-CDI cases were readmitted to hospital within 6 months.

Mortality

Within 2 months of CDI diagnosis, 15.7% of cases with HOCA-CDI and 26.3% of cases with HOHA-CDI died in hospital, while the corresponding rates in control patients were 5.3% and 11.6%, respectively (**Table II**).

The in-hospital mortality rates and mortality rates within 2 months after CDI onset are shown in **Figure 3**. The 30-day mortality rate was 6.8% for HOCA-CDI cases and 12.4% for HOHA-CDI cases. Adjusting for comorbidities on admission, previous hospital admissions and social deprivation score, compared with the corresponding controls, the HR for mortality was 2.47 (95% CI 2.04–3.00) for HOCA-CDI cases and 2.01 (95% CI 1.79–2.25) for HOHA-CDI cases.

Regardless of the origin of CDI, only comorbidities associated with the current admission were positively associated with mortality. There was no impact of social deprivation, antimicrobial or any medication prescriptions, or the number of hospital admissions in the 12 months prior to admission. Admission from a care home was positively associated with mortality (HR 1.52, 95% CI 0.94–2.45 for HOCA-CDI and HR 1.57, 95% CI 1.19–2.08 for HOHA-CDI).

Economic evaluation

The median (lower quartile [LQ], upper quartile [UQ]) index inpatient cost was more than double for cases (£7456 [3728, 16,060]) compared to controls (£2796 [932, 8388]), (**Figure 4a**). The median (LQ, UQ) additional cost associated with the index inpatient stay, calculated as the difference in median costs between cases in the index inpatient stay and matched controls, was £1713 (-1864, 8396; **Figure 4b**). Median (95% CI) longer-term costs based on all hospital stays in the 6 months after the index inpatient stay were £11,650 (5126, 23750) for cases and £6524 (2324, 16170) for controls (**Figure 4c**).

Discussion

Using routinely collected national data, the present study identified and characterised the additional burden of CDI on the NHS in Scotland, not only in relation to patient outcome and hospital resource use, but also its impact on hospitalisation costs. Compared with matched controls, CDI is associated with additional median lengths of stay of 7.2 days for HOCA-CDI and 12.0 days for HOHA-CDI. The median additional expenditure for each CDI case is £1713. In the 6 months after the index hospitalisation, the cost associated with a CDI case is £5126 higher than for controls. As far as we are aware, this study offers the first European data on the economic burden of hospitalised cases of community-associated CDI [2]. It highlights the substantial healthcare and economic burden of CDI, including recurrent infection, and therefore provides a key rationale for campaigns that promote the early detection and treatment of cases and/or instigate infection control strategies to lessen the impact of CDI. Our findings could also help service providers to plan and prioritise resources.

From our study, patients with CDI had longer lengths of stay and poorer outcomes than matched controls, and 1 in 7 patients had a second recurrence within 90 days of the initial infection. The LOS for patients with CDI is comparable to that in other studies based on European data [19], including a recently published study of CDI cases over a 6-year period in Scotland [6]. While our study did not compare the costs associated with index CDI and recurrent CDI episodes, we calculated the costs of admissions with recurrent CDI, contributing to the evidence that recurrent CDI adds substantially to the total cost of treating CDI.

Our study also enabled a comparison of outcomes between community- and hospital-associated CDI, when identified in the hospital. It has long been assumed that community-associated CDI is less acute than hospital-associated CDI, but a substantial proportion of patients with community-associated CDI have serious sequelae, including severe infection, treatment failure, recurrent CDI and need for hospitalisation [20,21]. Although there is a

greater burden of hospital-associated disease, our study highlights that the mortality rate is similar in hospital- and community-associated CDI groups, when the latter is diagnosed in hospital. During the study period, ~801 community-associated, community-onset CDI cases were reported. The 30-day mortality rate of 12.4% for HOHA-CDI cases in our study was lower than the 17.5% reported for patients with hospital-associated CDI in a recent study from Scotland, although a greater proportion of patients in this other study had a higher Charlson comorbidity index compared with our study [6].

While a recently published study also reported on the impact of CDI on LOS and mortality in patients in Scotland [6], no data on costs, recurrent infection or readmissions were included. The present study considers the current economic burden in the NHS in the United Kingdom and highlights the substantial cost associated with the acute management of CDI. We demonstrate that the burden of financial cost continues over an extended period of time, such that 6 months after the acute episode there is still a clear and significant difference in cost between cases and controls. Recent data from both 2016/17 and 2017/18 suggest that 31% of all patients admitted to hospital have more than one admission within a year [22,23]; in our study, 57.7% of all hospital-onset CDI patients, from an earlier period (2010–2013), were readmitted within 6 months, suggesting that hospital-onset CDI is associated with particularly high rates of hospital readmission.

A key strength of this study was the matched design for estimating costs. The study design also minimised the impact of bias and confounding factors on our analyses. As a limitation, we recognise that the quality of the retrospective data relies on the accuracy and completeness of the medical records, although it is presumed that cases and controls were subject to the same issues. Additionally, and unlike the definition provided by the European Centre for Disease Prevention and Control (ECDC), our definition of HOCA-CDI did not take into account previous hospital discharges ≤ 4 weeks of the current admission [24], and therefore some HOCA-CDI cases may have been healthcare-associated CDI. Likewise, our definition of recurrent CDI incorporated a different timeline (second CDI diagnosis ≥ 28 days

after the initial positive CDI test) than that from the ECDC (second CDI >2 weeks and <8 weeks after onset of previous episode) [24]. Consequently, the current study may have under- or over-estimated HOCA-CDI and recurrent CDI. Also, the prescribing data used in our study were derived from primary care, rather than prescriptions received during the hospital stay. Furthermore, our analysis did not include cost data associated with prescribing or management within the community. Finally, by selecting the first CDI episode within the time period for individuals with multiple CDI episodes, we estimated the total additional costs of CDI within 6 months, which may have resulted in an overestimate compared with the selection of an episode at random, as had been done for the estimation of recurrence and readmission rates.

In conclusion, CDI poses a substantial burden on healthcare services, including lengthy hospital stays, and after the infection, patients are at greater risk of readmission to hospital, even for non-CDI reasons. Consequently, there are increased costs associated with the management of patients with CDI compared with matched controls.

Acknowledgements

We thank Andrew Walker, who supported the economic analysis for this study and the eDRIS team at National Services Scotland for the supply of data. This study uses data provided by patients and collected by the NHS as part of their care and support. This study was initiated and supported by Astellas Pharma Europe, Ltd. Medical writing support was provided by Rhian Harper Owen, PhD, for Cello Health MedErgy (Europe), funded by Astellas Pharma, Inc.

These data were presented in part as oral and poster presentations at the European Congress of Clinical Microbiology and Infectious Diseases (ECCMID), 22–25 April 2017, Vienna, Austria.

Transparency declarations

Conflicts of interest

All authors report financial support from Astellas Pharma for conduct of the study reported here and for medical writing support. In addition, Professor Leanord has received funding from Astellas Pharma relating to attendance at an educational meeting.

Author contributions

Conception: AL, CR, IF, CMcC, CM

Study design and conduct: AL, CR, IF, CMcC, KK, MB

Data acquisition: CR, KK, JP, CM, CMcC

Analysis and interpretation: JP, KK, CR, CMcC, IF, CM

Writing: AL, CR, JP, KK, IF, CMcC, MB, CM

Funding

This study was initiated and supported by Astellas Pharma Europe, Ltd. Medical writing support was funded by Astellas Pharma, Inc.

References

- [1] Bouza E. Consequences of *Clostridium difficile* infection: understanding the healthcare burden. Clin Microbiol Infect 2012;18 Suppl 6:5–12. doi:10.1111/1469-0691.12064.
- [2] Reigadas Ramirez E, Bouza ES. Economic burden of *Clostridium difficile* infection in European countries. Adv Exp Med Biol 2018;1050:1–12. doi:10.1007/978-3-319-72799-8_1.
- [3] Olsen MA, Stwalley D, Demont C, Dubberke ER. *Clostridium difficile* infection increases acute and chronic morbidity and mortality. Infect Control Hosp Epidemiol 2019;40:65–71. doi:10.1017/ice.2018.280.
- [4] van Dorp SM, Kinross P, Gastmeier P, Behnke M, Kola A, Delmée M, et al. Standardised surveillance of *Clostridium difficile* infection in European acute care hospitals: a pilot study, 2013. Euro Surveill 2016;21:pii=30293. doi:10.2807/1560-7917.ES.2016.21.29.30293.
- [5] Heimann SM, Vehreschild JJ, Cornely OA, Wisplinghoff H, Hallek M, Goldbrunner R, et al. Economic burden of *Clostridium difficile*-associated diarrhoea: a cost-of-illness study from a German tertiary care hospital. Infection 2015;43:707–14. doi:10.1007/s15010-015-0810-x.
- [6] Banks A, Moore EK, Bishop J, Coia JE, Brown D, Mather H, et al. Trends in mortality following *Clostridium difficile* infection in Scotland, 2010-2016: a retrospective cohort and case-control study. J Hosp Infect 2018;100:133–41. doi:10.1016/j.jhin.2018.07.023.
- [7] Shah DN, Aitken SL, Barragan LF, Bozorgui S, Goddu S, Navarro ME, et al. Economic burden of primary compared with recurrent *Clostridium difficile* infection in hospitalized patients: a prospective cohort study. J Hosp Infect 2016;93:286–9.

doi:10.1016/j.jhin.2016.04.004.

- [8] Tresman R, Goldenberg SD. Healthcare resource use and attributable cost of *Clostridium difficile* infection: a micro-costing analysis comparing first and recurrent episodes. *J Antimicrob Chemother* 2018;73:2851–5. doi:10.1093/jac/dky250.
- [9] Dubberke ER, Schaefer E, Reske KA, Zilberberg M, Hollenbeak CS, Olsen MA. Attributable inpatient costs of recurrent *Clostridium difficile* infections. *Infect Control Hosp Epidemiol* 2014;35:1400–7. doi:10.1086/678428.
- [10] Alvarez-Madrado S, McTaggart S, Nangle C, Nicholson E, Bennie M. Data Resource Profile: The Scottish National Prescribing Information System (PIS). *Int J Epidemiol* 2016;45:714-715f. doi:10.1093/ije/dyw060.
- [11] NHS National Services Scotland. Scottish Morbidity Record (SMR) Datasets. Information Services Division (ISD) Scotland Data Dictionary. 2019. <https://www.ndc.scot.nhs.uk/Copyright.asp> (accessed August 16, 2019).
- [12] Gray RJ. A Class of K-Sample Tests for Comparing the Cumulative Incidence of a Competing Risk. *Ann Stat* 1988;16:1141–54.
- [13] Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc* 1999;94:496–509.
- [14] Quan H, Sundararajan V, Halfon P, Fong A, Burnand B, Luthi J-C, et al. Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005;43:1130–9.
- [15] Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373–83.
- [16] Sundararajan V, Henderson T, Perry C, Muggivan A, Quan H, Ghali WA. New ICD-10 version of the Charlson comorbidity index predicted in-hospital mortality. *J Clin*

- Epidemiol 2004;57:1288–94. doi:10.1016/j.jclinepi.2004.03.012.
- [17] Scottish Government. Scottish Index of Multiple Deprivation 2012 2012. <https://www2.gov.scot/Topics/Statistics/SIMD> (accessed January 23, 2018).
- [18] R Development Core Team. R: A language and environment for statistical computing. 2015.
- [19] Pirson M, Poirrier J-E, Joubert S, Van den Bulcke J, Leclercq P, Avena L, et al. Evaluation of the cost and length of hospital stays related to the management of an intestinal *Clostridium difficile* infection. Acta Gastroenterol Belg 2018;81:263–8.
- [20] Gupta A, Khanna S. Community-acquired *Clostridium difficile* infection: an increasing public health threat. Infect Drug Resist 2014;7:63–72. doi:10.2147/IDR.S46780.
- [21] Khanna S, Pardi DS, Aronson SL, Kammer PP, Baddour LM. Outcomes in community-acquired *Clostridium difficile* infection. Aliment Pharmacol Ther 2012;35:613–8. doi:10.1111/j.1365-2036.2011.04984.x.
- [22] NHS National Services Scotland. Acute Hospital Activity and NHS Beds Information in Scotland 2017. <https://www.isdscotland.org/Health-Topics/Hospital-Care/Publications/2017-11-07/2017-11-07-Annual-QuarterlyAcuteActivity-Report.pdf> (accessed October 24, 2019).
- [23] NHS National Services Scotland. Acute Hospital Activity and NHS Beds Information in Scotland 2018. <https://www.isdscotland.org/Health-Topics/Hospital-Care/Publications/2017-11-07/2017-11-07-Annual-QuarterlyAcuteActivity-Report.pdf> (accessed October 24, 2019).
- [24] European Centre for Disease Prevention and Control (ECDC). European Surveillance of *Clostridioides (Clostridium) difficile* infections. Surveillance protocol version 2.4 2019. <https://www.ecdc.europa.eu/sites/default/files/documents/clostridium-difficile-infections-EU-surveillance-protocol-vers2.4.pdf>.

Tables

Table I. Demographics and characteristics of *Clostridioides difficile* infection cases and controls

Variable, n (%)	Cases N=3304	Controls N=9516
Gender		
Male	1385 (41.9)	3907 (41.1)
Female	1919 (58.0)	5609 (58.9)
Age group, years		
15–64	807 (24.4)	2263 (23.8)
65–74	730 (22.1)	2080 (21.9)
75–84	1049 (31.8)	3159 (33.2)
≥85	718 (21.7)	2014 (21.2)
Origin of CDI		
HOCA	1297 (39.3)	-
HOHA	2007 (60.7)	-
Admitted from a care home		
No	3132 (94.8)	9125 (95.9)
Yes	172 (5.2)	391 (4.1)
Charlson comorbidity index		
0	1650 (49.9)	4676 (49.1)
1	524 (15.9)	1399 (14.7)

2	600 (18.2)	1658 (17.4)
3	218 (6.6)	535 (5.6)
≥4	180 (5.5)	385 (4.1)
Unknown	132 (4.0)	863 (9.1)
Characteristic, median (IQR)		
Antimicrobial prescriptions in previous year	1 (0–3)	1 (0–3)
Number of admissions in previous year	2 (1–3)	1 (0–3)
Total prescriptions in previous year	58 (30–95)	48 (22–82)
Different prescriptions in previous year	13 (8–18)	11 (6–16)
Length of stay (days) before the date of CDI	5 (1–19)	4 (1–15)

CDI, *Clostridioides difficile* infection; HOCA, hospital onset, community-associated; HOHA, hospital onset, hospital-associated; IQR, interquartile range

Table II. Additional length of stay, in-hospital mortality and discharge from hospital according to *Clostridioides difficile* infection type

	HOCA-CDI		HOHA-CDI	
	Cases N=1297	Controls N=3980	Cases N=2007	Controls N=5536
Median (95% CI) additional length of stay, days ^a	11.6 (7.7–18.8)	4.4 (3.0–6.9)	22.8 (14.8–35.5)	10.8 (7.0–16.2)
Died in hospital, n (%)	204 (15.7)	211 (5.3)	527 (26.3)	644 (11.6)
Discharged within 2 months, n (%)	966 (74.5)	3535 (88.8)	1056 (52.6)	4007 (72.4)

HOCA-CDI, hospital onset, community-associated *Clostridioides difficile* infection; HOHA-CDI, hospital onset, hospital-associated *Clostridioides difficile* infection; CI, confidence interval

^aAdditional length of stay was adjusted according to comorbidities on admission, previous admissions, length of stay in hospital before CDI, and social deprivation

Figures


Figure 1. Identification and selection of *Clostridioides difficile* infection cases and controls

Figure 2. Recurrence of *Clostridioides difficile* infection and readmission to hospital for any reason

Figure 3. Mortality by origin of *Clostridioides difficile* infection


Figure 4. Economic evaluation of *Clostridioides difficile* infection (CDI) cases and controls in relation to a) costs associated with first continuous inpatient stay after CDI diagnosis, b) additional costs of cases versus controls, and c) longer term costs in the 6 months after CDI diagnosis. For a), all costs valued above £40,000 were revalued as £40,000, while for b) and c), all costs valued above £60,000 were revalued as £60,000

Figure 1. Identification and selection of *Clostridioides difficile* infection cases and controls




CDI, *Clostridioides difficile* infection; HO, hospital onset; HOCA-CDI, hospital onset, community-associated CDI; HOHA-CDI, hospital onset hospital-associated CDI; SMR, Scottish Morbidity Record

Figure 2. Recurrence of *Clostridioides difficile* infection and readmission to hospital for any reason



CDI, *Clostridioides difficile* infection

Figure 3. Mortality by origin of *Clostridioides difficile* infection




No. events/ Total no. patients	204/ 1297	211/ 3980	527/ 2007	644/ 5536		290/ 1297	381/ 3980	663/ 2007	981/ 5536
-----------------------------------	--------------	--------------	--------------	--------------	--	--------------	--------------	--------------	--------------


CDI, *Clostridioides difficile* infection; HOCA-CDI, hospital onset, community-associated CDI; HOHA-CDI, hospital onset, hospital-associated CDI

Figure 4. Economic evaluation of *Clostridioides difficile* infection (CDI) cases and controls



a)



b)



c)



Supplementary information

Supplementary Table A1. Cost per day for each ward used in the calculation of costs in the analysis

Ward type	Proxy for:	Cost per case (£)	LOS	Cost per day (£)
Cardiac surgery	Transplant unit	7,857	3.6	2,183
Intensive care unit		9,105	4.2	2,168
Burns unit		4,613	2.8	1,648
A&E ward		961	0.6	1,602
Medical paediatrics	Children's unit	1,418	1.1	1,289
Cardiac care unit		1,893	1.9	996
High dependency unit		2,265	2.5	906
Spinal paralysis	Spinal unit	27,072	49.8	544
General medicine	Palliative care Stroke unit Assessment unit Other	1,584	3.4	466
Rehabilitation ward	GORU Convalescent unit	16,710	37.5	446
Young chronic sick	Younger physically disabled	3,039 per week		434
Geriatric long stay	Geriatric assessment	1,732 per week		247

A&E, accident and emergency; GORU, geriatric orthopaedic rehabilitation unit; LOS, length of stay

Supplementary Table AII. Detailed demographics and characteristics of cases and controls

Variable	Level	HOCA-CDI		HOHA-CDI	
		Cases (1297)	Controls (3980)	Cases (2007)	Controls (5536)
		N (%)	N (%)	N (%)	N (%)
Gender	Female	805 (62.1)	2468 (62.0)	1114 (55.5)	3141 (56.7)
Scottish Index of Multiple Deprivation	1: most deprived	391 (30.1)	1104 (27.7)	571 (28.5)	1430 (25.8)
	2	318 (24.5)	917 (23.0)	463 (23.1)	1252 (22.6)
	3	250 (19.3)	739 (18.6)	374 (18.6)	1051 (19.0)
	4	182 (14.0)	623 (15.7)	306 (15.2)	951 (17.2)
	5: least deprived	152 (11.7)	583 (14.6)	284 (14.2)	830 (15.0)
	unknown	4 (0.3)	14 (0.4)	9 (0.4)	22 (0.4)
Care home	Yes	64 (4.9)	129 (3.2)	108 (5.4)	262 (4.7)
Comorbidity - admission	1 Myocardial infarction	62 (4.8)	210 (5.3)	124 (6.2)	415 (7.5)
	2 Congestive heart failure, cardiomyopathy etc.	70 (5.4)	245 (6.2)	179 (8.9)	502 (9.1)
	3 Atherosclerosis, aortic aneurysm, vascular disease	49 (3.8)	116 (2.9)	136 (6.8)	290 (5.2)
	4 Stroke etc.	34 (2.6)	185 (4.6)	178 (8.9)	498 (9)
	5 Dementia	54 (4.2)	192 (4.8)	139 (6.9)	458 (8.3)
	6 Bronchitis, pneumoconiosis etc.	173 (13.4)	510 (12.8)	262 (13.1)	812 (14.7)
	7 Gout, lupus, rheumatoid arthritis etc.	19 (1.5)	67 (1.7)	43 (2.1)	107 (1.9)
	8 Gastro ulcers	13 (1)	33 (0.8)	59 (2.9)	76 (1.4)

	9 Liver problems	60 (4.6)	67 (1.7)	92 (4.6)	132 (2.4)
	10 Diabetes	140 (10.8)	373 (9.4)	244 (12.2)	613 (11.1)
	11 Diabetes with complications	9 (0.7)	17 (0.4)	35 (1.7)	64 (1.2)
	12 Renal problems	148 (11.4)	233 (5.9)	281 (14)	536 (9.7)
	13 Cancer	220 (17)	600 (15.1)	359 (17.9)	853 (15.4)
	14 Alcohol-related liver failure, varices etc.	26 (2)	26 (0.7)	58 (2.9)	59 (1.1)
	15 Metastatic cancer	56 (4.3)	240 (6)	127 (6.3)	332 (6)
Comorbidity - previous 5 years	1 Myocardial infarction	150 (11.6)	373 (9.4)	195 (9.7)	577 (10.4)
	2 Congestive heart failure, cardiomyopathy etc.	167 (12.9)	341 (8.6)	252 (12.6)	617 (11.1)
	3 Atherosclerosis, aortic aneurysm, vascular disease	110 (8.5)	249 (6.3)	194 (9.7)	447 (8.1)
	4 Stroke etc.	131 (10.1)	324 (8.1)	255 (12.7)	617 (11.1)
	5 Dementia	52 (4)	148 (3.7)	105 (5.2)	322 (5.8)
	6 Bronchitis, pneumoconiosis etc.	286 (22.1)	739 (18.6)	357 (17.8)	1006 (18.2)
	7 Gout, lupus, rheumatoid arthritis etc.	63 (4.9)	146 (3.7)	75 (3.7)	184 (3.3)
	8 Gastro ulcers	74 (5.7)	106 (2.7)	73 (3.6)	121 (2.2)
	9 Liver problems	94 (7.3)	117 (2.9)	109 (5.4)	168 (3)
	10 Diabetes	206 (15.9)	519 (13)	346 (17.2)	775 (14)
	11 Diabetes with complications	33 (2.6)	68 (1.7)	86 (4.3)	137 (2.5)
	12 Renal problems	214 (16.6)	307 (7.7)	340 (16.9)	626 (11.3)
	13 Cancer	283 (21.9)	685 (17.2)	369 (18.4)	826 (14.9)
	14 Alcohol-related liver failure, varices etc.	53 (4.1)	55 (1.4)	58 (2.9)	99 (1.8)
	15 Metastatic cancer	80 (6.2)	242 (6.1)	79 (3.9)	264 (4.8)

	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)
Age	73 (61–81)	73 (61–82)	77 (67–85)	78 (68–85)
Antimicrobial prescriptions in previous 12 months	1 (0–4)	1 (0–3)	1 (0–3)	1 (0–3)
Length of stay before the date of CDI	1 (0–1)	1 (0–1)	14 (7–32)	12 (6–25)
Number of admissions in previous 12 months	2 (1–4)	1 (0–3)	1 (0–3)	1 (0–2)
Total prescriptions in previous 12 months	59 (29–97)	44 (19–78)	58 (31–92)	50 (25–84)
Different prescriptions in previous 12 months	14 (8–19)	10 (6–16)	13 (8–18)	11 (7–17)
Charlson comorbidity index - previous 5 years	2 (0–4)	1 (0–2)	2 (0–3)	1 (0–3)

CDI, *Clostridioides difficile* infection; HOCA-CDI, hospital onset, community-associated CDI; HOHA-CDI, hospital onset, hospital-associated CDI

Supplementary Table AIII. Recurrence rates (per 100-patient-years) of *Clostridioides difficile* infection (CDI) during the year after the date of cure, assuming the date of cure was 4, 6 or 8 weeks after initial CDI diagnosis

Time point	HOCA-CDI (%)	HOHA-CDI (%)
Week 4	19.4	16.4
Week 6	18.8	16.1
Week 8	18.2	15.2

CDI, *Clostridioides difficile* infection; HOCA-CDI, hospital onset, community-associated CDI; HOHA-CDI, hospital onset, hospital-associated CDI

Supplementary Table AIV. Principal reason for initial hospitalisation and readmission to hospital within 6 months

Reason ^a	Admission (N=3304), %	Readmission ≤180 days (N=349), %
1 Myocardial infarction	5.6	5.2
2 Congestive heart failure, cardiomyopathy etc.	7.5	6.4
3 Atherosclerosis, aortic aneurysm, vascular disease	5.6	4.2
4 Stroke etc.	6.4	3.1
5 Dementia	5.8	3.7
6 Bronchitis, pneumoconiosis etc.	13.2	11.0
7 Gout, lupus, rheumatoid arthritis etc.	1.9	1.9
8 Gastrointestinal or ulcers	2.2	1.1
9 Liver problems	4.6	4.8
10 Diabetes	11.6	12.2
11 Diabetes with complications	1.3	1.9
12 Hemiplegia, paraplegia etc.	0.2	0.6
13 Renal problems	13.0	11.8
14 Cancer	17.5	18.8
15 Alcohol-related liver failure, varices etc.	2.5	2.0
16 Metastatic cancer	5.5	4.0
17 HIV related	0.2	0.6

^aPercentages of patients admitted for 17 principal reasons. Percentages do not add up to 100 as some reasons are not included. Furthermore, the percentages associated with small numbers are subject to disclosure control so that exact numbers were not released when fewer than five patients were involved