

Access to Diarylmethanols by Wittig Rearrangement of *ortho*-, *meta*-, and *para*-Benzyloxy-*N*-Butylbenzamides

R. Alan Aitken,* Andrew D. Harper, Ryan A. Inwood, and Alexandra M. Z. Slawin



Cite This: *J. Org. Chem.* 2022, 87, 4692–4701



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information


ABSTRACT: The *N*-butyl amide group, CONHBu, has been found to be an effective promoter of the [1,2]-Wittig rearrangement of aryl benzyloxy ethers and thus allow the two-step synthesis of isomerically pure substituted diarylmethanols starting from simple hydroxybenzoic acid derivatives. The method is compatible with a wide range of functional groups including methyl, methoxy, and fluoro, although not with nitro and, unexpectedly, is applicable to *meta* as well as *ortho* and *para* isomeric series.



INTRODUCTION

Directed *ortho*-metalation is now well established as a powerful method for regioselective functionalization of aromatic compounds,¹ and subsequent reaction with an electrophile such as an aromatic aldehyde allows facile construction of *ortho*-substituted diarylmethanols (Scheme 1). In contrast,

Scheme 1. Alternative Approaches to Diarylmethanols



methods for the analogous *meta*- or *para*-functionalization are nowhere near so well developed despite some recent progress.² In view of the low cost and ready availability of the three isomeric hydroxybenzoic acids, an attractive alternative strategy to access specifically substituted diarylmethanols would be to transform the carboxylic acid into a suitable activating group, *O*-benzylate the phenolic OH, and then conduct a [1,2]-Wittig rearrangement (Scheme 1). Although this is a well-known aromatic rearrangement,³ it has not been widely exploited in synthesis, most likely due to the strongly basic conditions required, which limit functional group compatibility, and there have been few recent examples of its

use.⁴ Some highlights in recent use of the Wittig rearrangement include rearrangement of benzyl butadienyl ethers,⁵ benzyl pyridyl ethers,⁶ tandem anion translocation—Wittig rearrangement and tandem Wittig rearrangement—aldol reaction,⁷ as well as the study of systems where there is competition between 1,2- and 2,3- or 1,2- and 1,4-Wittig rearrangements.⁸ The Wittig rearrangement has also been used to access a range of chiral binaphthyl ligands that have been applied in asymmetric catalysis.⁹ The rearrangement has been carried out in an enantioselective way by adding a chiral bis(oxazoline) catalyst,¹⁰ and diastereoselective Wittig rearrangements have been reported directed by adjacent carbohydrate and α -alkoxyamide functions.¹¹ Finally, it is also possible to suppress the Wittig rearrangement if desired in order to make use of the unrearranged α -lithiobenzyloxy group in the synthesis of benzofurans and other heterocycles.¹² In this paper, we describe the discovery and development of the facile Wittig rearrangement of isomeric benzyloxy-*N*-butylbenzamides to furnish the corresponding diarylmethanols in the *ortho*-, *meta*-, and *para*-series.

RESULTS AND DISCUSSION

Our entry into this area came from a serendipitous discovery during attempted ring bromination of the 2,4-bis(benzyloxy)-phenyloxazoline **1**. Treatment with *n*-butyllithium followed by bromine gave not the expected product **2** but instead a product identified by spectroscopic methods and X-ray diffraction (see Supporting Information) as **4**, presumably formed by Wittig


Received: December 30, 2021

Published: March 14, 2022




rearrangement to afford the intermediate **3**, which was then oxidized by bromine¹³ to the ketone (**Scheme 2**).

Scheme 2. Unexpected Reaction Leading to **4**



We then examined the reactivity of the simpler 2-benzyloxyphenyl compound **5** and found that, depending upon the reaction conditions, varying mixtures of the Wittig rearrangement product **6**, the 3-aminobenzofuran **7** resulting from intramolecular nucleophilic ring opening of the oxazoline by the benzyl anion, and the *O*-dealkylation product **8** were formed (**Scheme 3**). As we have already reported elsewhere,¹⁴


Scheme 3. Behavior of Mono(benzyloxy)phenyloxazolines



the process could be optimized toward the formation of **7** using *n*-butyllithium/potassium *tert*-butoxide. However, optimized conditions for the formation of **6**, namely 2.2 equiv *n*-BuLi in THF at rt for 1 h, resulted in an isolated yield after chromatographic purification of just 29%. Under the same conditions, the *para*-isomer **9** gave the rearrangement product **10** but in only 11% isolated yield, and the *meta*-isomer **11** was recovered unchanged. It was clear that the 4,4-dimethyloxazoline group was not an optimal promoter of the Wittig rearrangement.


We next investigated carbamates and tertiary amides, both classic *ortho*-directing groups,¹ as promoters of Wittig rearrangement, but this was uniformly unsuccessful. The diethylcarbamate **12** suffered a nucleophilic attack at carbonyl to give **13** and **14** in low yield while the phenylcarbamate **15** was recovered unchanged (**Scheme 4**). The *N,N*-diethyl amide

Scheme 4. Reaction of Carbamates and Tertiary Benzamides



16 reacted at the carbonyl group to give the ketone **17**. In a previous case where an undesired attack of butyllithium at a diethyl amide was encountered,¹⁵ this could be suppressed by changing to the *N,N*-diisopropyl amide but the reaction of **18** took a different course, giving 2-hydroxybenzil **19** in THF, but the 3-aminobenzofuran **20** in toluene. There are only a few other synthetic routes to substituted 3-aminobenzofurans. The formation of both these products involves an initial intramolecular nucleophilic attack of benzyloxy anion on the amide carbonyl with hydrolysis and dehydration giving **20**, while loss of diisopropylamine, hydrolysis, and oxidation affords **19**. The closely analogous formation of **19** by base treatment and oxidation of methyl 2-benzyloxybenzoate has already been reported.¹⁶

Success was finally achieved by moving to the secondary *N*-butylbenzamides and reaction of **21a** with 3.3 equiv *n*-butyllithium in THF at rt for 2 h followed by workup gave an essentially quantitative yield of the diarylmethanol **22a** (**Scheme 5**). This product was, however, found to be unstable and slowly cyclized over a period of weeks to give mainly 3-phenylphthalide **23a** (80%) accompanied by a low yield of anthraquinone **24a** resulting from an alternative mode of cyclization and subsequent oxidation. Alternatively, treating the crude product **22a** with *p*-toluenesulfonic acid in boiling toluene for 1 h,¹⁷ followed by aqueous workup and chromatographic purification led directly to **23a** in 90% isolated yield. The need for 3 equiv of *n*-BuLi is, we believe, due to the first two equivalents reacting to deprotonate the NH and then bring about an amide-directed *ortho*-metalation. Only with the third equivalent of base is the benzyl group deprotonated. Use of less *n*-BuLi resulted in progressively lower yields and recovery of unreacted starting material.¹⁸

Scheme 5. Wittig Rearrangement of 2-Benzyloxy-*N*-butylbenzamide

With these optimized conditions in hand, the scope was now explored, and analogues **21b–w** were prepared in good yield by *O*-alkylation of *N*-butylsalicylamide **25** with benzylic and other halides and potassium carbonate in DMF (Scheme 6).


These were now subjected to the Wittig rearrangement conditions used for **21a** and a varied pattern of reactivity emerged (Scheme 7). The systems containing methyl, methoxy, and fluoro substituents all reacted to give the corresponding secondary alcohols **22b–j**, which were fully characterized (see Supporting Information) but cyclized upon storage or *p*-toluenesulfonic acid treatment to give the corresponding phthalides **23**. In the cases of the fluorobenzyl compounds **21h** and **21j**, the corresponding anthraquinone products **24h** and **24j** were also isolated in low yield. The thienyl compound **21q**, the α -methylbenzyl compound **21s**, and the prenyl compound **21u** also underwent rearrangement. However, the remaining compounds bearing more electron-withdrawing substituents did not, and compounds **21k–p**, **r**, **v**, and **w** either underwent decomposition or were recovered unchanged. While all the products shown in Scheme 7 were obtained by spontaneous cyclization, those marked *, in addition to **23a**, were also prepared by the *p*-toluenesulfonic acid method.

The case of the allyl compound **21t** was particularly interesting. It afforded an inseparable mixture of the expected rearrangement product **22t** and an isomer, which proved to be the 3-ethyl-3-hydroxyisoindolinone **27**. Over a period of months, the mixture converted entirely into the latter, whose structure was confirmed by X-ray diffraction (Scheme 8).

We believe this to involve double-bond migration in **22t** to give the enol, which tautomerizes to ketone **26**, which then undergoes ring closure. Such isomerization of allyl carbinols to ethyl ketones occurs under a range of basic conditions.¹⁹



Attention was now turned to the isomeric *para*-system, and a range of substrates **29a–p** were prepared in good yield by *O*-alkylation of *N*-butyl-*p*-hydroxybenzamide **28** (Scheme 9).

These were subjected to the normal rearrangement conditions, and a similar pattern emerged as for the *ortho* series. The unsubstituted system as well as those with methyl, methoxy, and fluoro substituents rearranged to give the diarylmethanols **30a–h** mostly in good yield (Scheme 10). The α -methylbenzyl compound **29o** and the prenyl compound **29p** also reacted to give **30o** and **30p** although with a low yield in the latter case. In the case of **30c**, the molecular structure was confirmed by X-ray diffraction and the crystal structure featured each molecule involved in two donor and two acceptor interactions, with head-to-tail hydrogen-bonded dimers linked by C=O...H–O interactions, which were then further linked by N–H...O(H)–C interactions with adjacent

Scheme 6. Preparation of *ortho*-substituted *N*-butylbenzamides



molecules (see Supporting Information). Again the three isomeric nitro compounds **29j–l** as well as the pentafluorophenyl compound **29m**, the diphenylmethyl compound **29n** and the methoxynaphthyl system **29i** either decomposed or were recovered unchanged.

Having established the viability of the rearrangement for both *ortho* and *para* isomeric systems, it was of interest to compare the relative ease of the two processes, and for this, the 2,4-bis(benzyloxy)benzamide **31** was prepared. When this was treated with 3.3 equiv *n*-BuLi, the product was mainly **32** resulting from rearrangement of the *ortho* group with just a trace of the isomer **33** from the reaction of the *para* group (Scheme 11). This appears to be the first example of an amide directing lithiation onto an *ortho*-alkoxy group. Storage of compound **32** over a period of months resulted in spontaneous cyclization to give phthalide **34** in good yield. On the other hand, treatment of **31** with 4.4 equiv of *n*-BuLi resulted in


Scheme 7. Products from Wittig Rearrangement of *ortho*-Substituted BenzamidesScheme 8. Isomerization of **22t**

rearrangement of both groups to give the diol **35** as a 1:1 mixture of diastereomers.

Based on the earlier finding that the *meta*-benzyloxy oxazoline **11** did not react, in contrast to *ortho* and *para*-isomers **5** and **9**, and assuming that the activating effect of the amide group in the Wittig rearrangement would be via a spiro anionic intermediate (see below, Scheme 17), we expected that *meta*-benzyloxy *N*-butylbenzamides **37** would not react. However, to our surprise, the unsubstituted compound **37a** did rearrange under the normal conditions to afford **38a**, albeit in a somewhat lower yield than for *ortho* or *para*-isomers. The structure of **38a** was confirmed by X-ray diffraction and again the crystal structure featured each molecule involved in two

Scheme 9. Preparation of *para*-substituted *N*-butylbenzamidesScheme 10. Products from Wittig Rearrangement of *para*-Substituted Benzamides


Scheme 11. Reactivity of bis(benzyloxy) Compound 31



donor and two acceptor interactions, but in contrast to 30c, this involved head-to-tail hydrogen-bonded dimers linked by N–H...O(H)–C interactions, which were then further linked by C=O...H–O interactions with adjacent molecules (see Supporting Information). Based on this result, a range of substituted examples 37b–n were prepared in good-to-moderate yield by *O*-alkylation of 36 (Scheme 12). When these were subjected to the standard rearrangement conditions, a somewhat more restricted pattern of reactivity emerged with successful rearrangement only being observed for alkyl, methoxy, and fluoro substituents as well as the thienyl compound 37m and, in all cases, the isolated yields were lower than for the isomeric systems (Scheme 13).


So far substituent effects have only been examined in the benzyl as opposed to the aryl ring. To further examine the scope, we returned to the *ortho* system and investigated the effect of substituents on the other aryl ring. Starting from the 5-nitrosalicylamide 39, a range of three *O*-benzyl derivatives 40a–c were prepared, while the corresponding 5-dimethylamino compound 41 led to derivatives 42a–c. However, when these six compounds were subjected to the usual rearrangement conditions, only in the cases of 42a and 42c were the corresponding rearrangement products 43a and 43c obtained (Scheme 14). It is clear that the presence of a nitro group on either ring is sufficient to prevent the reaction. Since this is most likely due to incompatibility of the nitro function with the reaction conditions, additional evidence was sought from isomeric compounds with methoxy and fluoro substituents, both of which were compatible with the reaction conditions. Thus the 6-fluoro compound 44 in which *ortho*-metalation is impossible was treated with 2.2 equiv *n*-BuLi to give the expected rearrangement product 45, isolated after *p*-toluenesulfonic acid-mediated cyclization as the 7-fluoro-3-phenylphthalide (27%), together with the *n*-butyl compound 47 resulting from nucleophilic aromatic substitution in low yield (Scheme 15). Methoxy substituents in either the 4- or 5-position were also compatible with the rearrangement and 48 reacted to form 49, isolated as the phthalide 50 while 51 reacted via 52 to give the phthalide product 53 in moderate yield. In contrast to this, the 5-fluoro compound 54 (Scheme 16) underwent decomposition under the normal rearrangement conditions, perhaps due to intervention of an aryne process, while the 3,5-dimethyl-4-benzyloxy compound 55 in which the supposed spiro anion intermediate is sterically disfavored, was recovered largely unchanged.

To further probe the mechanism, the 4-trimethylsilyl compound 56 was prepared and was found to undergo rearrangement readily with *n*-BuLi to afford the alcohol 57. Finally, compounds 58 and 59 in which the benzyloxy group of 21a and 29a is replaced by the isomeric phenoxyethyl group


Scheme 12. Preparation of *meta*-substituted *N*-butylbenzamides

were prepared, and these were also found to readily undergo the rearrangement under the normal conditions, giving, respectively, 22a and 30a.


Although there have been a good number of mechanistic studies on the [1,2]-Wittig rearrangement, some early suggestions involving the intermediacy of arynes and carbonyl compounds were later disproven.²⁰ The most recent and detailed mechanistic study on aryl benzyl ethers employing both experimental and theoretical methods,²¹ quite clearly points to two major mechanistic possibilities: an anionic mechanism via a spiro-epoxide intermediate that is the normal route for neutral and electron-poor aryl rings, and a radical dissociation/recombination route that is more likely to be important for electron-rich aryl systems. In our system, we envisage initial amide NH deprotonation and *ortho*-metalation in each case before the third equivalent of *n*-BuLi deprotonates the benzyl group. The resulting intermediates 60, 62, and 64 can each cyclize to the spiro epoxides implicated in the Wittig rearrangement (Scheme 17) but, while 61 and 65 are be

Scheme 13. Products from Wittig Rearrangement of *meta*-Substituted Benzamides

Scheme 14. Nitro and Dimethylamino Substituted Systems




Scheme 15. Rearrangement with 4-, 5-, or 6-Substituents




stabilized by the negative charge being on nitrogen, this is not possible for **63** derived from the *meta* compound thus explaining the lower yields obtained in that case.

The recent mechanistic study²¹ was focused on substituent effects on the aryl ring rather than the benzyl ring, and substituent effects on the benzyl ring do not seem to have been examined in detail until now. Overall our results on the three

Scheme 16. Further Mechanistic Probes Examined



Scheme 17. Opportunities for Charge Stabilization in the Three Isomeric Systems




isomeric benzyloxybenzamide systems show that a single *N*-butylbenzamide group on the aryl ring is sufficient to facilitate the rearrangement of *ortho* and *para* benzyloxy systems, presumably via the anionic spiro-epoxide intermediates. The *meta* system is slightly less prone to rearrangement presumably reflecting the reduced capacity for delocalization of the negative charge to a *meta* disposed electron-withdrawing amide. In all three series, the reaction proceeds with a wide range of both electron-donating and electron-withdrawing benzyl substituents. Only nitro compounds were uniformly unsuccessful due to the incompatibility of that group with butyllithium. The fact that the silyl compound **56** rearranges much more rapidly than benzyl phenyl ether is consistent with the anionic mechanism in that case where the stabilizing effect of silicon upon the α -anion is key. For the isomeric phoxymethyl compounds **58** and **59** where the spiro-epoxide intermediate cannot be stabilized, the amide group nevertheless promotes the reaction perhaps by favoring the benzylic deprotonation, but the rearrangement must necessarily proceed by the radical route in these cases.

The fact that **48** gives a substantially lower yield than the isomer **51** implies that the rearrangement is discouraged by the presence of a second inductively electron-withdrawing (but mesomerically electron-donating) group in the *para* position to the rearranging group. The occurrence of the rearrangement for **42a** and **42c**, which also have an inductively electron-

withdrawing group *para* to the reaction site, can perhaps be taken to indicate that the electron-donating mesomeric effect outweighs the inductive effect in these cases.

Finally, we examined briefly whether the process could be extended from the ethers to the corresponding sulfides or amines since both thia- and aza-[1,2]-Wittig rearrangements are known.^{22,23} The 2-benzylthio-*N*-butylbenzamide **66** was readily prepared, but upon treatment with butyllithium, it underwent dehydrative cyclization to afford the 3-amino-benzothiophene **67** in moderate yield, in a reaction analogous to the formation of **20** from **18** (Scheme 18). In an attempt to

Scheme 18. Attempted Extension to thia- and aza-Analogues



suppress this process, the more bulky *tert*-butylbenzamide **68** was prepared, but it was recovered unreacted from BuLi treatment as was the 2-(benzylmethylamino) analogue **69**.

EXPERIMENTAL SECTION

¹H and ¹³C NMR spectra were recorded in CDCl₃ unless otherwise stated with internal TMS as reference. IR spectra were recorded using the ATR technique. HRMS measurements were made either using ESI with TOF analyzer or NSI with an ion trap analyzer.

The following procedures are illustrative; full experimental procedures and characterization data are given in the Supporting Information.

2-(Benzyloxy)-*N*-butylbenzamide 21a. A solution of 2-(benzyloxy)benzoyl chloride¹³ (10.85 g, 44.0 mmol) in toluene (60 mL) was added dropwise to a stirred 0 °C solution of *n*-butylamine (12.1 mL, 8.95 g, 0.122 mol) in toluene (60 mL). Once the addition was complete, the reaction mixture was allowed to warm to rt for 1 h before being poured into water and washed with 2 M NaOH and brine. The organic layer was dried and evaporated to afford, after recrystallization (EtOAc/hexane), **21a** (9.34 g, 75%) as colorless crystals, mp 52–54 °C; IR 3380, 1648, 1599, 1558, 1292, 1238, 1164, 1101, 1005, 865, 752, 700 cm⁻¹; ¹H NMR (500 MHz) δ 8.25 (dd, *J* = 7.8, 1.8 Hz, 1H, ArH), 7.88 (br s, 1H, NH), 7.47–7.38 (m, 6H, ArH and Ph), 7.12–7.08 (m, 1H, ArH), 7.06 (d, *J* = 8.5 Hz, 1H, ArH), 5.15 (s, 2H, OCH₂), 3.34 (td, *J* = 7.0, 5.5 Hz, 2H, NCH₂), 1.35–1.29 (m, 2H, NCH₂CH₂), 1.19–1.11 (m, 2H, CH₂CH₃), 0.80 (t, *J* = 7.3 Hz, 3 H, CH₃); ¹³C NMR (75 MHz) δ 164.9(C), 156.8(C), 135.5(C), 132.5 (CH), 132.4 (CH), 128.9 (2CH), 128.8 (CH), 128.2 (2CH), 122.0(C), 121.6 (CH), 112.4 (CH), 71.4 (OCH₂), 39.4 (NCH₂), 31.2 (CH₂), 20.0 (CH₂), and 13.7 (CH₃); HRMS (ESI⁺) *m/z* [M + Na⁺] calcd for C₁₈H₂₁NaNO₂ 306.1465, found 306.1455.

***N*-Butyl-2-(hydroxy(phenyl)methyl)benzamide 22a, Anthraquinone 24a and 3-Phenylphthalide 23a.** Under a nitrogen atmosphere, *n*-butyllithium (2.6 mL, 6.50 mmol) was added dropwise to a stirred solution of 2-(benzyloxy)-*N*-butylbenzamide **21a** (0.5678 g, 2.00 mmol) in dry THF (20 mL). After stirring at rt for 2 h, the reaction mixture was quenched by addition of sat. aq. NH₄Cl and extracted with Et₂O (×3). The combined organic layers were washed with 2 M NaOH and water before being dried and evaporated to give **22a** as a pale yellow oil: IR 3296, 3064, 2931, 1635, 1540, 1450, 1303, 1228, 1104, 1024, 757, 699 cm⁻¹; ¹H NMR (500 MHz) δ 7.38 (t, *J* =

7.3 Hz, 2H, ArH), 7.29–7.17 (m, 7H, ArH and Ph), 6.31 (t, *J* = 5.3 Hz, 1H, NH), 5.79 (s, 1H, CHOH), 3.21–3.14 (m, 1H, NCH₂), 3.12–3.05 (m, 1H, NCH₂), 1.29–1.23 (m, 2H, NCH₂CH₂), 1.22–1.15 (m, 2H, CH₂CH₃), 0.85 (t, *J* = 7.0 Hz, 3H, CH₃); ¹³C NMR (125 MHz) δ 170.8(C), 143.1(C), 142.7(C), 135.8(C), 130.6 (CH), 129.9 (CH), 127.74 (CH), 127.71 (2CH), 127.69 (CH), 126.7 (CH), 126.2 (2CH), 74.9 (CHOH), 39.7 (NCH₂), 31.1 (CH₂), 19.9 (CH₂), 13.6 (CH₃); HRMS (ESI⁺) *m/z* [M + Na⁺] calcd for C₁₈H₂₁NaNO₂ 306.1465, found 306.1456.

On standing at rt in EtOAc solution for 2–3 months, an intramolecular cyclization occurred to give, after purification by column chromatography (SiO₂, Et₂O/hexane 2:3), at R_f 0.80, **24a** (17.5 mg, 4%) as yellow needles, mp 275–279 °C (lit.²⁴ 275 °C); ¹H NMR (500 MHz) δ 8.34–8.30 (m, 4 H, ArH), 7.83–7.79 (m, 4 H, ArH). The ¹H NMR spectral data were in accordance with those previously reported.²⁵

This was followed by a second fraction to give, at R_f 0.55, **23a** (0.3350 g, 80%) as tan-colored crystals, mp 113–116 °C (lit.²⁶ 115.5 °C); ¹H NMR (500 MHz) δ 7.96 (d, *J* = 7.5 Hz, 1H, ArH), 7.65 (td, *J* = 7.5, 1.0 Hz, 1H, ArH), 7.55 (t, *J* = 7.5 Hz, 1H, ArH), 7.39–7.36 (m, 3H, ArH), 7.33 (dd, *J* = 7.8, 0.8 Hz, 1H, ArH), 7.29–7.26 (m, 2H, ArH), 6.41 (s, 1H, CHPh). The ¹H NMR spectral data were in accordance with those previously reported.²⁷

Alternatively, the following literature procedure²⁸ may be employed: A mixture of *N*-butyl-2-(hydroxy(phenyl)methyl)benzamide **22a** (prepared as above from 1.14 g **21a**, assuming 4.02 mmol) and *p*-toluenesulfonic acid monohydrate (1.55 g, 8.15 mmol) in toluene (80 mL) was heated at reflux for 1 h. After cooling to rt, the reaction mixture was washed with water (50 mL), 2 M NaOH (50 mL), and brine (50 mL) before being dried and evaporated. The crude residue was purified by column chromatography (SiO₂, gradient elution, Et₂O/hexane 1:4 to Et₂O) to give **23a** (0.76 g, 90%) as tan-colored crystals.

***N*-Butyl-4-hydroxybenzamide 28.** A mixture of methyl 4-hydroxybenzoate (30.43 g, 0.20 mol) and *n*-butylamine (100 mL, 74.00 g, 1.01 mol) was heated at reflux for 4 d before being concentrated *in vacuo*. The residue was acidified to pH 1 by the addition of 2 M HCl and extracted with Et₂O (3 × 100 mL). The combined organic layers were washed with water (100 mL) before being dried and evaporated. The crude residue was recrystallized (EtOAc/PhMe) to give **28** (28.01 g, 72%) as colorless crystals, mp 118–120 °C (lit.²⁹ 118.5–119.5 °C); ¹H NMR (500 MHz) δ 7.95 (br s, 1H, OH), 7.62 (d, *J* = 8.8 Hz, 2H, 2,6-H), 6.86 (d, *J* = 8.8 Hz, 2H, 3,5-H), 6.17 (t, *J* = 5.5 Hz, 1H, NH), 3.44 (td, *J* = 7.0, 5.5 Hz, 2H, NCH₂), 1.62–1.56 (m, 2H, NCH₂CH₂), 1.44–1.36 (m, 2H, CH₂CH₃), 0.94 (t, *J* = 7.3 Hz, 3H, CH₃). The ¹H NMR spectral data were in accordance with those previously reported.³⁰

4-(Benzyloxy)-*N*-butylbenzamide 29a. *N*-Butyl-4-hydroxybenzamide **28** (3.87 g, 20.0 mmol) was added to a stirred suspension of sodium hydride (60% in mineral oil, prewashed with hexane, 0.82 g, 20.5 mmol) in DMF (20 mL), and the mixture was stirred at rt for 15 min before benzyl bromide (2.4 mL, 3.45 g, 20.2 mmol) was added. After stirring for 18 h at rt, the reaction mixture was poured into water and extracted with CH₂Cl₂ followed by Et₂O (×3). The combined organic layers were washed with brine (×5) and 2 M NaOH before being dried and evaporated. Recrystallization of the residue (EtOAc/hexane) gave **29a** (4.68 g, 82%) as colorless crystals, mp 126–128 °C; (lit.³¹ 119.1–119.7 °C); ¹H NMR (500 MHz) δ 7.72 (d, *J* = 8.8 Hz, 2H, 2,6-H), 7.44–7.38 (m, 4H, Ph), 7.35–7.32 (m, 1H, Ph), 6.99 (d, *J* = 8.8 Hz, 2H, 3,5-H), 6.03 (t, *J* = 5.5 Hz, 1H, NH), 5.11 (s, 2H, OCH₂), 3.44 (td, *J* = 7.3, 5.5 Hz, 2H, NCH₂), 1.62–1.56 (m, 2H, NCH₂CH₂), 1.45–1.37 (m, 2H, CH₂CH₃), and 0.95 (t, *J* = 7.3 Hz, 3H, CH₃). The ¹H NMR spectral data were in accordance with those previously reported.³¹

***N*-Butyl-4-(hydroxy(phenyl)methyl)benzamide 30a.** Under a nitrogen atmosphere, *n*-butyllithium (2.5 M in hexane, 6.6 mL, 16.5 mmol) was added dropwise to a stirred solution of 4-(benzyloxy)-*N*-butylbenzamide **29a** (1.41 g, 4.98 mmol) in dry THF (50 mL). After stirring at rt for 2 h, the reaction mixture was quenched by addition of sat. aq. NH₄Cl and extracted with Et₂O (×3). The combined organic

layers were washed with 2 M NaOH and water before being dried and evaporated. Purification of the residue by column chromatography (SiO₂, gradient elution, Et₂O/hexane 7:3 to Et₂O) and subsequent recrystallization (EtOAc/hexane) gave **30a** (1.11 g, 79%) as colorless crystals, mp 113–114 °C; IR 3432, 3337, 2953, 2926, 1616, 1542, 1448, 1303, 1228, 1045, 736, 694 cm⁻¹; ¹H NMR (400 MHz) δ 7.67 (d, *J* = 8.2 Hz, 2H, ArH), 7.41 (d, *J* = 8.2 Hz, 2H, ArH), 7.36–7.30 (m, 4H, Ph), 7.28–7.24 (m, 1H, Ph), 6.14 (t, *J* = 5.6 Hz, 1H, NH), 5.85 (s, 1H, CHOH), 3.41 (td, *J* = 7.2, 5.6 Hz, 2H, NCH₂), 2.76 (d, *J* = 3.2 Hz, 1H, OH), 1.61–1.53 (m, 2H, NCH₂CH₂), 1.43–1.34 (m, 2H, CH₂CH₃), 0.94 (t, *J* = 7.4 Hz, 3H, CH₃); ¹³C NMR (125 MHz) δ 167.3(C), 147.1(C), 143.4(C), 133.8(C), 128.6 (2CH), 127.8 (CH), 126.9 (2CH), 126.6 (2CH), 126.5 (2CH), 75.7 (CHOH), 39.8 (NCH₂), 31.7 (CH₂), 20.1 (CH₂), 13.8 (CH₃); HRMS (ESI⁺) *m/z* [M + H⁺] calcd for C₁₈H₂₂NO₂ 284.1645, found 284.1644.

3-(Benzyloxy)-N-butylbenzamide 37a. To a stirred solution of 3-(benzyloxy)benzoyl chloride³² (6.82 g, 27.6 mmol) in CH₂Cl₂ (100 mL) at 0 °C, Et₃N (3.85 mL, 27.6 mmol) was added dropwise. The solution was stirred for 5 min, and then *n*-butylamine (2.73 mL, 27.6 mmol) was added dropwise and the mixture stirred at rt for 18 h. The reaction mixture was poured into H₂O, extracted (×3) with CH₂Cl₂, and the combined organic fractions were dried over MgSO₄ and concentrated to afford, after recrystallization (EtOH) **37a** (5.80 g, 74%) as off-white crystals, mp 83–86 °C; IR 3298, 3229, 2961, 1626, 1603, 1580, 1553, 1016, 698 cm⁻¹; ¹H NMR (400 MHz) δ 7.47–7.38 (m, 5H, ArH), 7.38–7.32 (m, 2H, ArH), 7.32–7.28 (m, 1H, ArH), 7.09 (ddd, *J* = 7.8, 2.6, 1.5 Hz, 1H, ArH), 6.07 (br s, 1H, NH), 5.11 (s, 2H, OCH₂), 3.45 (td, *J* = 7.1, 5.7 Hz, 2H, NHCH₂), 1.65–1.53 (m, 2H, NHCH₂CH₂), 1.51–1.30 (m, 2H, CH₂CH₃), 0.96 (t, *J* = 7.3 Hz, 3H, CH₂CH₃); ¹³C NMR (100 MHz) δ 167.2(C), 158.9(C), 136.5(C), 136.4(C), 129.6 (CH), 128.6 (CH), 128.1 (CH), 127.5 (CH), 118.8 (CH), 118.1 (CH), 113.3 (CH), 70.1 (OCH₂), 39.8 (NHCH₂), 31.7 (CH₂), 20.1 (CH₂), 13.8 (CH₃); HRMS (ESI⁺) *m/z* [M + H⁺] calcd for C₁₈H₂₂NO₂ 284.1651, found 284.1636.

N-Butyl-3-(hydroxy(phenyl)methyl)benzamide 38a. Under a nitrogen atmosphere, *n*-butyllithium (2.5 M in hexane, 2.64 mL, 6.60 mmol) was added dropwise to a stirred solution of 3-(benzyloxy)-*N*-butylbenzamide **37a** (567 mg, 2.0 mmol) in dry THF (20 mL). After stirring at rt for 2 h, the reaction mixture was quenched by addition of sat. aq. NH₄Cl and extracted with Et₂O (×3). The combined organic layers were dried and evaporated. Purification of the residue by column chromatography (gradient elution, Et₂O/hexane 1:1 to Et₂O/hexane 7:3) gave **38a** (298 mg, 53%) as colorless crystals, mp 97–100 °C; IR 3298, 3229, 2961, 2932, 2857, 1626, 1553, 1418, 1327, 1016, 746, 698 cm⁻¹; ¹H NMR (400 MHz) δ 7.75 (t, *J* = 1.8 Hz, 1H, ArH), 7.57 (dt, *J* = 7.7, 1.5 Hz, 1H, ArH), 7.42–7.35 (m, 1H, ArH), 7.31–7.24 (m, 5H, ArH), 7.24–7.21 (m, 1H, ArH), 6.43 (t, *J* = 5.7 Hz, 1H, NH), 5.74 (s, 1H, CHOH), 3.71 (s, 1H, CHOH), 3.31 (td, *J* = 7.2, 5.7 Hz, 2H, NHCH₂), 1.54–1.45 (m, 2H, NHCH₂CH₂), 1.39–1.27 (m, 2H, CH₂CH₃), 0.90 (t, *J* = 7.3 Hz, 3H, CH₂CH₃); ¹³C NMR (100 MHz) δ 167.6(C), 144.5(C), 143.5(C), 134.7(C), 129.5 (CH), 128.5 (CH), 128.4 (2CH), 127.5 (CH), 126.5 (2CH), 125.9 (CH), 124.7 (CH), 75.6 (CHOH), 39.8 (NHCH₂), 31.5 (CH₂), 20.1 (CH₂), 13.7 (CH₃); HRMS (ESI⁺) *m/z* [M + H⁺] calcd for C₁₈H₂₂NO₂ 284.1651, found 284.1637.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.1c03160>.

Experimental procedures, characterization data, NMR spectra of all newly synthesized compounds, and X-ray crystallographic details (PDF)

Accession Codes

CCDC 2105490–2105493 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by

emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

R. Alan Aitken – *EaStCHEM School of Chemistry, University of St Andrews, Fife KY16 9ST, U.K.*; orcid.org/0000-0001-6959-5311; Email: raa@st-and.ac.uk

Authors

Andrew D. Harper – *EaStCHEM School of Chemistry, University of St Andrews, Fife KY16 9ST, U.K.*

Ryan A. Inwood – *EaStCHEM School of Chemistry, University of St Andrews, Fife KY16 9ST, U.K.*

Alexandra M. Z. Slawin – *EaStCHEM School of Chemistry, University of St Andrews, Fife KY16 9ST, U.K.*; orcid.org/0000-0002-9527-6418

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.joc.1c03160>

Author Contributions

The manuscript was written through the contributions of all authors. A.D.H. and R.A.I. contributed equally to the synthetic work, and A.M.Z.S. performed the crystallography. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We dedicate this paper to the memory of Victor Snieckus (1937–2020) and Peter Beak (1936–2021). We thank EPSRC (UK) for a DTA studentship to A.D.H. (Grant EP/L505079/1), EPSRC (UK), and CRITICAT Centre for Doctoral Training for a studentship to R.A.I. (Grant code: EP/L016419/1) and the EPSRC UK National Mass Spectrometry Facility at Swansea University.

■ REFERENCES

- (1) (a) Gschwend, H. W.; Rodriguez, H. R. Heteroatom-Facilitated Lithiations. *Org. React.* **1979**, *26*, 1–360. (b) Beak, P.; Snieckus, V. Directed Lithiation of Aromatic Tertiary Amides: An Evolving Synthetic Methodology for Polysubstituted Aromatics. *Acc. Chem. Res.* **1982**, *15*, 306–312. (c) Reuman, M.; Meyers, A. I. The Synthetic Utility of Oxazolines in Aromatic Substitution. *Tetrahedron* **1985**, *41*, 837–860. (d) Snieckus, V. Directed Ortho Metalation. Tertiary Amide and *O*-Carbamate Directors in Synthetic Strategies for Polysubstituted Aromatics. *Chem. Rev.* **1990**, *90*, 879–933. (e) Matsui, S.; Uejima, A.; Suzukia, Y.; Tanaka, K. Asymmetric Synthesis of Optically Active Phthalides via *ortho*-Lithiation and Cyclization of Chiral *N*-Monosubstituted Benzamides. *J. Chem. Soc. Perkin. Trans. 1* **1993**, 701–704. (f) Nguyen, T.-H.; Castanet, A.-S.; Mortier, J. Directed *Ortho*-Metalation of Unprotected Benzoic Acids. Methodology and Regioselective Synthesis of Useful Contiguously 3- and 6-Substituted 2-Methoxybenzoic Acid Building Blocks. *Org. Lett.* **2006**, *8*, 765–768. (g) Jia, B.; Yang, Y.; Jin, X.; Mao, G.; Wang, C. Rhenium-Catalyzed Phthalide Synthesis from Benzamides and Aldehydes via C–H Bond Activation. *Org. Lett.* **2019**, *21*, 6259–6263.
- (2) (a) Korvorapun, K.; Samanta, R. C.; Rogge, T.; Ackermann, L. Remote C–H Functionalizations by Ruthenium Catalysis. *Synthesis* **2021**, *53*, 2911–2934. (b) Dey, A.; Agasti, S.; Maiti, D. Palladium catalysed *meta*-C–H-functionalization reactions. *Org. Biomol. Chem.* **2016**, *14*, 5440–5453. (c) Mihai, M. T.; Genov, G. R.; Phipps, R. J. Access to the *meta* position of arenes through transition metal catalysed C–H bond functionalisation: a focus on metals other than

- palladium. *Chem. Soc. Rev.* **2018**, *47*, 149–171. (d) Mihai, M. T.; Williams, B. D.; Phipps, R. J. Para-Selective C–H Borylation of Common Arene Building Blocks Enabled by Ion-Pairing with a Bulky Counteranion. *J. Am. Chem. Soc.* **2019**, *141*, 15477–15482. (e) Liu, L.-Y.; Qiao, J. X.; Yeung, K.-S.; Ewing, W. R.; Yu, J.-Q. meta-Selective C–H Arylation of Fluoroarenes and Simple Arenes. *Angew. Chem., Int. Ed.* **2020**, *59*, 13831–13835.
- (3) (a) Schorigin, P. Über die Carbinol-Umlagerung von Benzyläthern. *Ber. Dtsch. Chem. Ges.* **1924**, *57*, 1634–1637. (b) Wittig, G.; Löhmann, L. Über die kationotrope Isomerisation gewisser Benzyläther bei Einwirkung von Phenyl-lithium. *Liebigs Ann. Chem.* **1942**, *550*, 260–268.
- (4) Wang, F.; Wang, J.; Zhang, Y.; Yang, J. The [1,2]- and [1,4]-Wittig rearrangement. *Tetrahedron* **2020**, *76*, No. 130857.
- (5) Nakano, T.; Soeta, T.; Endo, K.; Inomata, K.; Ukaji, Y. Stereoselective synthesis of (2Z,4E)-2,4-pentadien-1-ols via sequential 1,4-elimination reaction and [1,2]-Wittig rearrangement starting from (E)-4-alkoxy-2-butenyl benzoates. *J. Org. Chem.* **2013**, *78*, 12654–12661.
- (6) (a) Yang, J.; Dudley, G. B. [1,2]-Anionic rearrangement of 2-benzoyloxy-pyridine and related pyridyl ethers. *J. Org. Chem.* **2009**, *74*, 7998–8000. (b) Yang, J.; Dudley, G. B. Pyridine-directed organolithium addition to an enol ether. *Adv. Synth. Catal.* **2010**, *352*, 3438–3442.
- (7) (a) Barluenga, J.; Fañanás, F. J.; Sanz, R.; Marcos, C.; Trabada, M. On the reactivity of *o*-lithiated ethers: tandem anion translocation and Wittig rearrangement. *Org. Lett.* **2002**, *4*, 1587–1590. (b) Bertrand, M. B.; Wolfe, J. P. Tandem Wittig rearrangement/aldol reactions for the synthesis of glycolate aldols. *Org. Lett.* **2006**, *8*, 4661–4663.
- (8) (a) Capriati, V.; Florio, S.; Ingrosso, G.; Granito, C.; Troisi, L. Stereoselective and competitive [1,2]- and [2,3]-Wittig rearrangements of allyl heteroarylalkyl ethers. *Eur. J. Org. Chem.* **2002**, *2002*, 478–484. (b) Mori-Quiroz, L. M.; Maleczka, R. E., Jr. Stereoconvergent [1,2]- and [1,4]-Wittig rearrangements of 2-silyl-6-aryl-5,6-dihydropyran: a tale of steric vs. electronic regiocontrol of divergent pathways. *J. Org. Chem.* **2015**, *80*, 1163–1191.
- (9) (a) Gao, G.; Gu, F.-L.; Jiang, J.-X.; Jiang, K.; Sheng, C.-Q.; Lai, G.-Q.; Xu, L.-W. Neighboring lithium-assisted [1,2]-Wittig rearrangement: practical access to diarylmethanol-based 1,4-diols and optically active BINOL derivatives with axial and sp³-central chirality. *Chem. – Eur. J.* **2011**, *17*, 2698–2703. (b) Zheng, L.-S.; Jiang, K.-Z.; Deng, Y.; Bai, X.-F.; Gao, G.; Gu, F.-L.; Xu, L.-W. Synthesis of Ar-BINMOL ligands by [1,2]-Wittig rearrangement to probe their catalytic activity in 1,2-addition reactions of aldehydes with Grignard reagents. *Eur. J. Org. Chem.* **2013**, *2013*, 748–755. (c) Song, T.; Zheng, L.-S.; Ye, F.; Deng, W.-H.; Wei, Y.-L.; Jiang, K.-Z.; Xu, L.-W. Modular synthesis of Ar-BINMOL-Phos for catalytic asymmetric alkylation of aromatic aldehydes with unexpected reversal of enantioselectivity. *Adv. Synth. Catal.* **2014**, *356*, 1708–1718. (d) Song, T.; Li, L.; Zhou, W.; Zheng, Z.-J.; Deng, Y.; Xu, Z.; Xu, L.-W. Enantioselective copper-catalyzed azide-alkyne click cycloaddition to desymmetrization of maleimide-based bis(alkynes). *Chem. – Eur. J.* **2015**, *21*, 554–558. (e) Wang, C.-Y.; Dong, C.; Zheng, Z.-J.; Xu, Z.; Yang, K.-F.; Xu, L.-W. Construction of an all-substituted pyrrolidine derivative with multiple stereogenic centers and Betti-base-derived γ -amino alcohols by [1,2]-Wittig rearrangement. *RSC Adv.* **2015**, *5*, 55819–55824.
- (10) Tomooka, K.; Yamamoto, K.; Nakai, T. Enantioselective [1,2] Wittig rearrangement using an external chiral ligand. *Angew. Chem., Int. Ed.* **1999**, *38*, 3741–3743.
- (11) (a) Tomooka, K.; Yamamoto, K.; Nakai, T. [1,2]-Wittig rearrangement of acetal systems: a highly stereocontrolled conversion of *O*-glycosides to *C*-glycosides. *J. Am. Chem. Soc.* **1996**, *118*, 3317–3318. (b) Tomooka, K.; Yamamoto, H.; Nakai, T. Stereoselective synthesis of highly functionalized *C*-glycosides based on acetal [1,2] and [1,4] Wittig rearrangements. *Angew. Chem., Int. Ed.* **2000**, *39*, 4500–4502. (c) Kitagawa, O.; Momose, S.; Yamada, Y.; Shiro, M.; Taguchi, T. Synthesis of optically active aldol derivatives through chirality transfer type 1,2-Wittig rearrangement of α -alkoxycarboxamides. *Tetrahedron Lett.* **2001**, *42*, 4865–4868.
- (12) (a) Velasco, R.; Feberero, C.; Sanz, R. α -Lithiated aryl benzyl ethers: inhibition of [1,2]-Wittig rearrangement and application to the synthesis of benzo[*b*]furan derivatives. *Org. Lett.* **2015**, *17*, 4416–4419. (b) Sedano, C.; Velasco, R.; Feberero, C.; Suárez-Pantiga, S.; Sanz, R. α -Lithiobenzoyloxy as a directed metalation group in *ortho*-lithiation reactions. *Org. Lett.* **2020**, *22*, 6365–6369.
- (13) Joshi, G.; Patil, R. D.; Adimurthy, S. Green bromine: *in situ* generated catalyst for the selective oxidation of alcohols using H₂O₂ as a benign oxidant. *RSC Adv.* **2012**, *2*, 2235–2239.
- (14) Aitken, R. A.; Harper, A. D.; Slawin, A. M. Z. Base-Induced Cyclisation of *ortho*-Substituted 2-Phenylloxazolines to Give 3-Aminobenzofurans and Related Heterocycles. *Synlett* **2017**, *28*, 1738–1742.
- (15) Beak, P.; Brown, R. A. The Tertiary Amide as an Effective Director of *Ortho* Lithiation. *J. Org. Chem.* **1982**, *47*, 34–46.
- (16) Worayuthakarn, R.; Boonya-udtayan, S.; Arom-oon, E.; Ploypradith, P.; Ruchirawat, S.; Thasana, N. Synthesis of Unsymmetrical Benzil Licoagrodione. *J. Org. Chem.* **2008**, *73*, 7432–7435.
- (17) Kawasaki, T.; Kimachi, T. Sparteine-Mediated Enantioselective [2,3]-Wittig Rearrangement of Allyl *ortho*-Substituted Benzyl Prenyl Ethers. *Tetrahedron* **1999**, *55*, 6847–6862.
- (18) We are grateful to a reviewer for suggesting that the extra equivalent of butyllithium may be required not for *ortho* metalation, but rather for either coordination to the amide oxygen or to form a bimetallic complex required for the rearrangement. We are unable to exclude these possibilities.
- (19) (a) Tiffeneau, M. Mécanisme des transpositions phényles chez les iodhydrines aromatiques. *Bull. Soc. Chim. Fr.* **1907**, *4*, 1205–1220. (b) Meisenheimer, J.; Link, J. Über die Verschiebung in der Allyl-Gruppe. 3. Mitteilung über Substitution und Addition. *Liebigs Ann. Chem.* **1930**, *479*, 211–277. (c) Zheng, H.-X.; Xiao, Z.-F.; Yao, C.-Z.; Li, Q.-Q.; Ning, X.-S.; Kang, Y.-B.; Tang, Y. Transition-Metal-Free Self-Hydrogen-Transferring Allylic Isomerization. *Org. Lett.* **2015**, *17*, 6102–6105. (d) Martínez-Erro, S.; Sanz-Marco, A.; Bermejo Gómez, A.; Vázquez-Romero, A.; Ahlquist, M. S. G.; Martín-Matute, B. Base-Catalyzed Stereospecific Isomerization of Electron-Deficient Allylic Alcohols and Ethers through Ion-Pairing. *J. Am. Chem. Soc.* **2016**, *138*, 13408–13414.
- (20) (a) Lansbury, P. T.; Pattison, V. A. The Wittig Rearrangement of the Benzyl Ethers of Cyclobutanol and Cyclopropylcarbinol. *J. Am. Chem. Soc.* **1962**, *84*, 4295–4298. (b) Lansbury, P. T.; Pattison, V. A.; Sidler, J. D.; Bieber, J. B. Mechanistic Aspects of the Rearrangement and Elimination Reactions of α -Metalated Benzyl Alkyl Ethers. *J. Am. Chem. Soc.* **1966**, *88*, 78–84. (c) Schäfer, H.; Schöllkopf, U.; Walter, D. Wanderungstendenz von Alkylgruppen bei der Wittig-Umlagerung. *Tetrahedron Lett.* **1968**, *9*, 2809–2814. (d) Eisch, J. J.; Kovacs, C. A.; Rhee, S.-G. Rearrangements of organometallic compounds X. The mechanism of 1,2-aryl migration in the Wittig rearrangement of α -metallated benzyl aryl ethers. *J. Organomet. Chem.* **1974**, *65*, 289–301. (e) Rautenstrauch, V.; Büchi, G.; Wuest, H. Vinyl Migration in Wittig Rearrangements. *J. Am. Chem. Soc.* **1974**, *96*, 2576–2580. (f) Strunk, S.; Schlosser, M. Wittig Rearrangement of Lithiated Allyl Aryl Ethers: A Mechanistic Study. *Eur. J. Org. Chem.* **2006**, *2006*, 4393–4397.
- (21) Velasco, R.; Silva López, C.; Nieto Faza, O.; Sanz, R. Exploring the Reactivity of α -Lithiated Aryl Benzyl Ethers: Inhibition of the [1,2]-Wittig Rearrangement and the Mechanistic Proposal Revisited. *Chem. – Eur. J.* **2016**, *22*, 15058–15068.
- (22) Biellmann, J. F.; Schmitt, J. L. Solvation and rearrangement of a carbanion. *Tetrahedron Lett.* **1973**, *14*, 4615–4618.
- (23) Eisch, J. J.; Kovacs, C. A. The π -orbital overlap requirement in 1,2-anionic rearrangements. *J. Organomet. Chem.* **1971**, *30*, C97–C100.
- (24) Kekulé, A.; Franchimont, A. Ueber das Benzophenonchlorid und die Bildung von Anthrachinon bei der Darstellung von Benzophenon. *Ber. Dtsch. Chem. Ges.* **1872**, *5*, 908–910.

(25) Barton, D. H. R.; Hui, R. A. H. F.; Ley, S. V. Oxidation of benzylic hydrocarbons with benzeneseleninic anhydride and related reactions. *J. Chem. Soc., Perkin. Trans. 1* **1982**, 2179–2185.

(26) Mermod, E.; Simonis, H. Ueber einige Phthalide und Mekonine. *Ber. Dtsch. Chem. Ges.* **1908**, *41*, 982–985.

(27) Weeks, D. P.; Cella, J. Preparation of 3-arylphthalides by the Clemmensen reduction of 2-arylbenzoic acids. *J. Org. Chem.* **1969**, *34*, 3713.

(28) Kawasaki, T.; Kimachi, T. Sparteine-mediated Enantioselective [2,3]-Wittig Rearrangement of Allyl *ortho*-Substituted Benzyl Ethers and *ortho*-Substituted Benzyl Prenyl Ethers. *Tetrahedron* **1999**, *55*, 6847–6862.

(29) Abramovitz, A. S.; Massey, V. Purification of intact old yellow enzyme using an affinity matrix for the sole chromatographic step. *J. Biol. Chem.* **1976**, *251*, 5321–5326.

(30) Mamidi, N.; Manna, D. Zn(OTf)₂-Promoted Chemoselective Esterification of Hydroxyl Group Bearing Carboxylic Acids. *J. Org. Chem.* **2013**, *78*, 2386–2396.

(31) Kajitani, T.; Kohmoto, S.; Yamamoto, M.; Kishikawa, K. Liquid crystalline amides: linear arrangement of rod-like molecules by lateral intermolecular hydrogen bonding and molecular shape effect. *J. Mater. Chem.* **2004**, *14*, 3449–3456.

(32) Myllymäki, M.; Castillo-Melendez, J.; Koskinen, A.; Minkkilä, A.; Saario, S.; Nevalainen, T.; Järvinen, T.; Poso, A.; Salo-Ahen, O. Heterocyclic phenyl carbamates as novel FAAH-inhibitors. *PCT Int. Appl. WO*, 129129 A1, 2008.