

Linking chromospheric activity and magnetic field properties for late-type dwarf stars

E. L. Brown^{1*}, S. V. Jeffers², S. C. Marsden¹, J. Morin³, S. Boro Saikia⁴, P. Petit⁵, M. M. Jardine⁶, V. See^{7,8}, A. A. Vidotto^{9,10}, M. W. Mengel¹, M. N. Dahlkemper^{11,12} and the BCool Collaboration¹³

¹University of Southern Queensland, Centre for Astrophysics, Toowoomba, QLD, 4350, Australia

²Max Planck Institut for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany

³LUPM, Université de Montpellier, CNRS, Place Eugène Bataillon, F-34095 Montpellier, France

⁴University of Vienna, Department of Astrophysics, Turkenchanzstrasse 17, 1180 Vienna, Austria

⁵Institut de Recherche en Astrophysique et Planétologie, Université de Toulouse, CNRS, CNES, 31400 Toulouse, France

⁶SUPA, School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK

⁷European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands

⁸University of Exeter, Department of Physics & Astronomy, Stoker Road, Devon, Exeter, EX4 4QL, UK

⁹School of Physics, Trinity College Dublin, The University of Dublin, College Green, Dublin-2, Ireland

¹⁰Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

¹¹University of Göttingen, Physics Education Research, Friedrich Hund Platz 1, 37077, Göttingen, Germany

¹²CERN, Teacher and Student Programmes, Esplanade de Particules 1, CH-1211, Genève 23, Switzerland

¹³<https://bcool.irap.omp.eu/>

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Spectropolarimetric data allow for simultaneous monitoring of stellar chromospheric $\log R'_{\text{HK}}$ activity and the surface-averaged longitudinal magnetic field, B_l , giving the opportunity to probe the relationship between large-scale stellar magnetic fields and chromospheric manifestations of magnetism. We present $\log R'_{\text{HK}}$ and/or B_l measurements for 954 mid-F to mid-M stars derived from spectropolarimetric observations contained within the PolarBase database. Our magnetically active sample complements previous stellar activity surveys that focus on inactive planet-search targets. We find a positive correlation between mean $\log R'_{\text{HK}}$ and mean $\log |B_l|$, but for G stars the relationship may undergo a change between $\log R'_{\text{HK}} \sim -4.4$ and -4.8 . The mean $\log R'_{\text{HK}}$ shows a similar change with respect to the $\log R'_{\text{HK}}$ variability amplitude for intermediately-active G stars. We also combine our results with archival chromospheric activity data and published observations of large-scale magnetic field geometries derived using Zeeman Doppler Imaging. The chromospheric activity data indicate a slight under-density of late-F to early-K stars with $-4.75 \leq \log R'_{\text{HK}} \leq -4.5$. This is not as prominent as the original Vaughan–Preston gap, and we do not detect similar under-populated regions in the distributions of the mean $|B_l|$, or the B_l and $\log R'_{\text{HK}}$ variability amplitudes. Chromospheric activity, activity variability and toroidal field strength decrease on the main sequence as rotation slows. For G stars, the disappearance of dominant toroidal fields occurs at a similar chromospheric activity level as the change in the relationships between chromospheric activity, activity variability and mean field strength.

Key words: stars:late-type; stars:activity; stars:magnetic field

1 INTRODUCTION

The dynamo-driven magnetic fields of cool stars generate a range of activity phenomena with widespread impacts. Magnetized stellar winds influence circumstellar environments, including the forma-

tion, evolution and potential habitability of surrounding planets (e.g. Ó Fionnagáin & Vidotto 2018). The detectability and parameterization of these planets is also complicated by the temporal evolution of magnetic features such as faculae and starspots on the stellar surface (Queiroz et al. 2001; Huélamo et al. 2008). Magnetic activity is intricately linked to the rotational evolution of a star, since the coupling of the magnetic field with magnetic wind causes the loss of stellar angular momentum (Skumanich 1972; Gallet & Bouvier

* E-mail: emma.brown@usq.edu.au

2015; Finley & Matt 2018; See et al. 2019b). For this reason, activity is often used as a proxy to estimate the rotation periods (i.e. Noyes et al. 1984) and ages (Lorenzo-Oliveira et al. 2018) of stars. Thus, improving our understanding of the stellar dynamo, its evolution throughout the life of a star, interplay with stellar rotation and other stellar properties, and the induced surface activity are important goals.

Stellar dynamo theory is largely informed by observations of the Sun. It is well established that the cyclic, solar magnetic dynamo is powered by convection and rotation within the outer convective layers of the Sun (Charbonneau 2005). Differential rotation is a key mechanism that converts an initially poloidal field to a toroidal field, and then the poloidal field is regenerated through cyclonic convection or the Babcock-Leighton mechanism (Charbonneau 2005). Some features of solar magnetic cycles are still not well explained nor reliably predicted using current dynamo theory, such as the amplitudes and lengths of cycles (Petrovay 2020), so monitoring of the magnetism of other cool stars has become an important adjunct to studies of the Sun. Through long-term monitoring of the magnetic fields and surface activity of late-type stars with a range of properties, we can provide context for the Sun’s magnetic behaviour and can work toward untangling the intricate relationships between stellar properties and the magnetic dynamo.

Chromospheric Ca II H&K spectral line emissions provide an important, indirect diagnostic of stellar magnetic activity. Magnetic heating in the chromosphere causes increased emissions in the cores of the Ca II H&K spectral lines, both in the Sun and other cool stars (Eberhard & Schwarzschild 1913). Ca II H&K activity is usually represented by the S-index (Vaughan et al. 1978), where the core emissions in the H and K lines are normalised to the continuum. Alternatively, when comparing the activity of stars across spectral types, photospheric and basal flux contributions are removed to provide a purely chromospheric activity index, $\log R'_{\text{HK}}$. Decades-long monitoring programs, such as the Mount Wilson Program (Wilson 1978; Duncan et al. 1991; Baliunas et al. 1995), have revealed S-index variability on a range of timescales and amplitudes across the HR diagram (Wilson 1978; Duncan et al. 1991; Baliunas et al. 1995; Hall et al. 2007; Lehtinen et al. 2016; Boro Saikia et al. 2018a; Gomes da Silva et al. 2020), with some stars showing cyclic S-index variability similar to the Sun’s 11-year activity cycle. The relationships between mean activity levels, chromospheric activity cycle periods and stellar properties have been previously investigated by Noyes et al. (1984); Böhm-Vitense (2007); Jenkins et al. (2011); Brandenburg et al. (2017); Boro Saikia et al. (2018a) and Gomes da Silva et al. (2020), among others. These studies suggest that cool stars may fall into distinct activity groups based on their mean chromospheric activity and activity cycle periods. Noyes et al. (1984) first reported that main sequence stars fall into ‘active’ and ‘inactive’ groups when their mean $\log R'_{\text{HK}}$ is plotted against stellar B-V. They reported an apparent lack of F and G-type stars with intermediate mean chromospheric activity, known as the ‘Vaughan-Preston gap’. More recently, Boro Saikia et al. (2018a) determined that the Vaughan-Preston gap is not prominent in their chromospheric activity study of 4454 cool stars drawn from multiple surveys. Gomes da Silva et al. (2020), in their sample of 1674 FGK stars from the AMBRE-HARPS sample (de Laverny et al. 2013), identified up to three different activity regimes for F-K stars, each separated by reduced populations of stars and occurring at different activity levels for different stellar types. Explanations for the reduced populations between activity groups range from sample biases to phases of rapid stellar evolution. As yet, there is no consensus on the existence of distinct activity groups, nor their physical basis.

Chromospheric activity variability has been less studied compared to mean activity levels or the periods of activity cycles. Saar & Brandenburg (2002) studied chromospheric activity cycle amplitudes for 31 stars from the Mount Wilson Project with well-defined chromospheric activity cycles, finding that peak-to-peak cycle amplitudes increase with mean activity. Gomes da Silva et al. (2020) also studied the relationship between mean chromospheric activity and activity variability amplitudes for their 1674 AMBRE-HARPS stars, but their data consisted of less-dense time series’ of observations, covering shorter time spans, compared to that used by Saar & Brandenburg (2002). Gomes da Silva et al. (2020) found that, although the upper envelope of chromospheric activity variability scales with mean activity level, low activity stars do not necessarily have low chromospheric variability amplitudes. This implies that planet-search stars, which are targeted for low chromospheric activity variability, cannot be identified based on their mean chromospheric activity level alone. Several studies have also compared the amplitudes of stellar photometric variability to chromospheric variability amplitudes and mean chromospheric activity levels (i.e. Radick et al. 1998; Lockwood et al. 2007; Radick et al. 2018). Stellar photometric variability amplitudes generally increase with mean chromospheric activity levels, but the Sun appears to have a lower photometric variability compared to other stars with similar mean chromospheric activity (Radick et al. 2018; Reinhold et al. 2020). Young, active stars tend to become darker as their chromospheric emissions increase throughout activity cycles, while mature and less active stars, including the Sun, become brighter with increasing chromospheric emissions. This suggests a transition from spot dominated to faculae dominated photometric variability at around middle age (Radick et al. 2018).

In this paper we analyse time-series spectropolarimetric observations for a sample of main sequence and youthful F-M stars from the PolarBase¹ database (Petit et al. 2014). Spectropolarimetry uniquely allows for the simultaneous measurement of the strength of the surface-averaged, large-scale stellar magnetic field (B_l) and the chromospheric emissions in the cores of the Ca II H&K lines. Thus, the data provide an opportunity to compare large-scale magnetic fields with the properties of smaller-scale magnetic surface features. These are known to be related (Lehmann et al. 2018) but the nature of the relation is not yet clear.

The PolarBase data span the years 2005 through to 2019, allowing for long-term monitoring of magnetic and chromospheric activity variability. We present both the average chromospheric activity and magnetic field strengths, and their variability amplitudes. We also combine our data with archival $\log R'_{\text{HK}}$ measurements from Gomes da Silva et al. (2020) and Boro Saikia et al. (2018a), as well as details of the magnetic field geometries for ~ 60 of our PolarBase targets which have been the subjects of previous Zeeman Doppler Imaging (ZDI, Semel 1989) studies.

The key aims of this work are to (i) compile a database of chromospheric activity measurements and surface-averaged large-scale magnetic field strengths for cool stars in the PolarBase sample, (ii) compare the distributions of chromospheric activity, B_l and their variability across the sample, (iii) search for distinct activity groups in the activity distribution, as well as any B_l counterparts, and (iv) use published ZDI results to gain new insights into the relationships between chromospheric activity, activity variability, magnetic field geometry and stellar properties.

¹ <http://polarbase.irap.omp.eu/>

2 THE STELLAR SAMPLE

2.1 Sample selection

PolarBase contains spectra for >2000 stellar objects, sourced from the high-resolution spectropolarimeters ESPaDOnS (Donati 2004; Donati et al. 2006), coupled with the 3.6 m Canada-France-Hawaii Telescope at Mauna Kea Observatory, and NARVAL (Aurière 2003), coupled with the 2 m Telescope Bernard Lyot at Pic du Midi Observatory. Targets cover the spectral types O4 to M9, with 60 percent of targets being cooler than F5 (Petit et al. 2014). We filtered the PolarBase targets to include only stars that are within the GAIA catalogue (DR2, Gaia Collaboration et al. 2016, 2018). We included only stars with effective temperatures between 3200 and 6700 K, which correspond roughly to spectral types M5 through to F5, and aimed to include pre-main sequence and main sequence stars only. We initially filtered giants and more evolved stars from the sample based on their luminosity classification within SIMBAD², where available. We then further filtered out subgiant stars using the method described in section 2.2.4. From our filtered target list, we include in this paper only the 954 stars for which we were able to measure the chromospheric $\log R'_{\text{HK}}$ (section 4.3.2) and/or the longitudinal magnetic field, B_l (sections 4.1 and 4.4). A HR diagram of the sample is shown in Figure 1. PolarBase stars are generally targeted because they are magnetically active, so our sample is comparatively more active and complements previous chromospheric activity surveys that derive from samples of planet-search targets (e.g. Gomes da Silva et al. 2020).

2.2 Stellar properties

Stellar properties are shown in Table 1. The full Table 1 is available online.

2.2.1 Effective temperature, luminosity, surface gravity, radius and B-V

We estimated the effective temperature, luminosity, surface gravity ($\log g$), stellar radius and Tycho B_T and V_T magnitudes by interpolating a grid of MIST (MESA Isochrones & Stellar Tracks) evolutionary tracks (Paxton et al. 2011, 2013, 2015, 2018; Choi et al. 2016; Dotter 2016) overlaid on the GAIA colour-magnitude diagram (G versus BP-RP magnitudes) for our sample. BP and RP magnitudes were taken directly from GAIA DR2 (Gaia Collaboration et al. 2016, 2018), and the absolute G-band magnitudes were derived using the apparent G magnitudes and parallaxes from GAIA DR2. We assumed solar metallicity (as in e.g. Arun et al. 2019). We converted the Tycho B_T and V_T magnitudes to the Johnson B-V magnitude shown in Table 1 according to Høg et al. (2000).

2.2.2 Rotational velocity

The line-of-sight projected rotational velocity ($v \sin i$) was used to estimate the rotation period of each target (section 4.5). We adopted $v \sin i$ from Valenti & Fischer (2005), Głęboccki & Gnaniński (2005) or the median of published values listed in SIMBAD, as indicated in Table 1.




Figure 1. Bottom: HR diagram of the sample of PolarBase stars selected for our study. Circles denote single objects, stars denote binary or multiple star systems where the measured $\log R'_{\text{HK}}$ may be impacted by the presence of companions, and triangles indicate binary or multiple systems where the B_l and $\log R'_{\text{HK}}$ are known to be impacted by blending. Filled markers are taken to be main sequence stars. Open markers have significant Li I absorption lines present in their spectra, suggesting that they may be young stars. Markers with reduced opacity are stars with noisy spectra, for which it is not clear if the stars have strong Li I lines. The Sun is indicated with the usual symbol in red. Top: Histogram showing the breakdown of the sample by spectral classification.

2.2.3 Binary/multiple systems

We identified binary and multiple systems using the SIMBAD database. We also inspected the LSD Stokes I spectral line profiles to identify any additional binary/multiple systems. The presence of multiple stellar signals in spectra and LSD profiles can result in the overestimation of the $\log R'_{\text{HK}}$ and/or B_l because of blending between components. Systems for which the measured B_l is impacted by the presence of companion stars are flagged in Table 1. We assume that the $\log R'_{\text{HK}}$ for all binaries/multiples may be contaminated by contributions from companions. This is because the cores of the Ca II H and K lines are intrinsically very broad compared to most metallic lines, so the contributions of companions are usually blended.

² <http://simbad.u-strasbg.fr/simbad/>

Table 1: Stellar properties for the PolarBase stars selected for this study. Columns (left to right) show the main identifier for each star, its GAIA ID, spectral type and luminosity classification according to SIMBAD, whether the presence of binary/multiple companion stars impact on our observations, whether significant Li I absorption lines were detected in spectra, stellar effective temperature (T_{eff}), luminosity ($\log \frac{L}{L_{\odot}}$), color index (B-V), surface gravity ($\log g$), radius (Rad) and projected rotational velocity ($v \sin i$). Asterisks (*) denote binaries/multiples for which our B_I measurements are not likely to be impacted by the presence of companions, and crosses † denote stars where the B_I measurement is impacted by companions. We assume that the measured $\log R'_{\text{HK}}$ for all binary/multiple stars may be impacted by the presence of companions. The full table is available online.

Main ID	GAIA ID	Spectral Type	Binary impact	Li I present	T_{eff} (K)	$\log(L/L_{\odot})$	B-V	$\log g$ (cm s $^{-2}$)	Rad (R_{\odot})	$v \sin i$ (km s $^{-1}$)
54 Psc	2802397960855105920	K0.5V			5107	-0.266	0.823	4.37	0.94	1.1 ¹
61 Cyg A	1872046574983507456	K5V	*		4275	-0.795	1.164	4.43	0.73	2.0 ²
61 UMa	4025850731201819392	G8V	*		5458	-0.198	0.704	4.49	0.89	2.4 ¹
χ^1 Ori	3399063235753901952	G0V	*		5854	0.084	0.557	4.38	1.07	9.8 ¹
ϵ Eri	5164707970261630080	K2V	*		4871	-0.420	0.910	4.39	0.87	2.4 ¹
κ^1 Cet	3269362645115584640	G5V	*		5630	-0.050	0.632	4.41	0.99	5.2 ¹
ξ Boo A	1237090738916392704	G7Ve	*		5420	-0.238	0.713	4.51	0.86	4.6 ¹
ρ^1 Cnc	704967037090946688	K0IV-V	*		5133	-0.193	0.813	4.31	1.01	2.5 ¹
τ Boo	1244571953471006720	F7IV-V	*		6301	0.529	0.434	4.17	1.54	15.0 ¹
ν And	348020448377061376	F9V	*		6061	0.555	0.492	4.08	1.72	9.6 ¹
EMSR 9	6049153921054413440	K5e		yes	3861	-0.023	1.278	3.48	2.18	14.2 ²
HD 166435	4590489981163831296	G1IV			5728	-0.021	0.601	4.43	0.99	7.9 ¹
HD 189733	1827242816201846144	K2V	*		4907	-0.462	0.898	4.46	0.81	2.9 ²
HD 190771	2061876742143320064	G2V	*		5680	0.026	0.617	4.36	1.06	4.3 ¹
HD 206860	1772187382746856320	G0V+	*	yes	5862	0.054	0.556	4.41	1.03	10.6 ¹
HD 224085	2855095251072482432	K2+IVeFe-I	*	yes	4331	0.345	1.184	3.52	2.64	23.0 ²
HD 75332	716109930305855360	F7Vs			6148	0.310	0.467	4.28	1.26	9.0 ¹
HD 76151	5760701787150565888	G2V			5663	-0.013	0.620	4.38	1.02	1.2 ¹
HD 78366	714116137767540096	G0IV-V			5904	0.103	0.542	4.38	1.08	3.9 ¹
HD 9986	2585856120791459584	G2V			5689	0.037	0.610	4.35	1.07	2.6 ¹
BP Tau	164832740220756608	K5/7Ve		yes	4014	-0.279	1.260	3.95	1.50	10.9 ²
EK Dra	1668690628102524672	G1.5V	*	yes	5528	-0.054	0.671	4.39	1.02	16.8 ¹
TW Hya	5401795662560500352	K6Ve		yes	4022	-0.549	1.248	4.26	1.09	14.0 ²
V410 Tau	164518589131083136	K3Ve	†	yes	4108	0.151	1.267	3.52	2.35	72.7 ²
V471 Tau	43789772861265792	K2V+DA	*	yes	4946	-0.415	0.882	4.43	0.84	
V830 Tau	147831571737487488	M0-1		yes	3841	-0.308	1.256	3.80	1.58	33.3 ³
V889 Her	4524317420039164544	G2V		yes	5581	0.027	0.647	4.32	1.10	38.0 ²
π^1 UMa	1092545710514654464	G0.5V			5763	-0.009	0.593	4.44	0.99	9.3 ²
.....										

¹ Valenti & Fischer (2005); ² Głęboccki & Gnaniński (2005); ³ SIMBAD.

2.2.4 Li I absorption lines to distinguish young stars from evolved subgiants

The presence of a strong Li I line (centered at 6707.8Å) in stellar spectra is an indicator of youth for single stars (Soderblom 2010). We visually inspected the spectra to check for Li I lines. For stars that were clearly located away from the main sequence, and with no/weak Li I absorption lines, we assumed that the stars are evolving off the main sequence and excluded them from our sample. For the remaining sample, we assumed stars are youthful if they have a Li I line as strong as the nearby 6717.7Å Ca I line. In some cases, the spectra were noisy and we could not clearly detect Li I or Ca I lines, in which case we included the stars in our sample, and have indicated throughout our results that we are unsure if a strong Li I line is present. Although this is a rudimentary approach to classifying stars as ‘youthful’, we consider it to be sufficient for this study.

3 DATA

3.1 Spectropolarimetry with ESPaDOnS and NARVAL

The ESPaDOnS and NARVAL spectropolarimeters have been gathering observations since 2005 and 2006 respectively. PolarBase includes observations that continue until mid-2019, prior to the commissioning of the NeoNARVAL spectropolarimeter at the TBL. Both instruments have a spectral coverage of 3700 to 10480 Å with a resolution of ~65000 in polarimetric mode. When operating in non-polarimetric mode, the instruments provide high-resolution Stokes I (intensity) spectra. When in polarimetric mode, they can also provide Stokes V (circularly polarized) spectra or Stokes Q and U (linearly polarized) spectra. Stokes V generally have stronger Zeeman polarization signatures compared to Stokes Q and U spectra for late-type stars, and are therefore more commonly collected for PolarBase targets. Stokes V spectra are obtained from a series of 4 individual ‘sub-exposures’, each of which measures flux in two orthogonal polarization states, $I + V$ and $I - V$. The positions of the orthogonally polarized beams are switched between sub-exposures by rotating the retarding rhombs of the polarimeter. A Stokes I spectrum can be extracted from each raw sub-exposure, and the polarized spectrum is derived from the full series of 4 sub-exposures, as described in section 3.2.

3.2 Data reduction and calibration

PolarBase provides ESPaDOnS and NARVAL observations in their reduced and calibrated format. The automatic reduction software package LIBRE-ESPRIT, based on ESPRIT (‘Echelle Spectra Reduction: an Interactive Tool’, Donati et al. 1997), is applied to reduce and calibrate each sub-exposure using bias and flat-field exposures taken during the observing runs. A first wavelength calibration is carried out using a Thorium-Argon arc-lamp exposure taken on the observing night, and is then refined using telluric lines as radial velocity (RV) references. This results in a final RV accuracy of ~ 20 to 30 m s⁻¹ (Moutou et al. 2007). Stokes I spectra are extracted from each sub-exposure, continuum normalized, and are made available on PolarBase at this point. Stokes V spectra are extracted from the series of calibrated sub-exposures by dividing sub-exposures with orthogonal polarization states, which removes instrumental polarization signals (Donati et al. 1997). Stokes V observations available on PolarBase also include a ‘mean’ Stokes I spectrum, which is derived by adding the series of 4 Stokes I sub-exposures together, and a ‘null’ spectrum which is calculated by dividing spectra with

Table 2. Table of normalization parameters used for LSD (extension of Table 4 from Marsden et al. 2014).

T_{eff} (K)	g_0	d_0	λ_0
3000	1.22	0.55	690
3500	1.22	0.55	670
4000	1.22	0.55	650
4500	1.22	0.55	630
5000	1.22	0.54	610
5500	1.22	0.53	590
6000	1.22	0.51	570
6500	1.21	0.49	560

identical polarization states, giving a measure of instrumental polarization (noise) and indicating the reliability of the polarimetric measurement.

4 ANALYSIS

4.1 Least-squares deconvolution

Least-squares deconvolution (LSD, Donati et al. 1997; Kochukhov et al. 2010) is a technique that uses the thousands of lines within a spectrum to derive a single, ‘mean’ spectral line profile with an improved SNR. LSD assumes that most spectral lines have almost identical shapes, including identical starspot-induced intensity contrasts in Stokes I spectra, and polarization signatures in Stokes V spectra. The entire spectrum is taken to be a convolution of the mean Stokes I or V line profile, and a line pattern function that describes the locations and relative strengths of the individual spectral lines. The line pattern function is modelled using a synthetic line mask for a quiet star having similar properties to the target, and each line in the mask is weighted by a factor w :

$$w_j = \frac{d_j}{d_0} \text{ for Stokes } I \text{ data, or} \quad (1)$$

$$w_j = \frac{g_j \lambda_j d_j}{g_0 \lambda_0 d_0} \text{ for Stokes } V \text{ data} \quad (2)$$

where g_j , λ_j and d_j are the j^{th} line’s effective Landé factor, central wavelength and central depth. The factors g_0 , λ_0 and d_0 are the normalization parameters from Table 2 (an extension of Table 4 from Marsden et al. 2014). The line pattern function can thus be deconvolved from the observed spectrum to determine the mean Stokes I or V profile, as described in Donati et al. (1997) and Kochukhov et al. (2010).

We performed LSD using the automated technique of Donati et al. (1997). We generated synthetic spectral line masks using stellar atmospheric models from the Vienna Atomic Line Database (VALD, Kupka et al. 2000), for stellar temperatures ranging between 3000 and 6500 K in steps of 500 K, and $\log g$ between 2.0 and 5.0 cm s⁻² in steps of 0.5 cm s⁻². Line masks excluded lines with depth ≤ 10 per cent of the continuum. For each target we selected the line mask with the nearest T_{eff} and $\log g$, and used the line normalization parameters from Table 2. An example LSD line profile is shown in Figure 2, and further details of the LSD technique can be found in Donati et al. (1997) and Kochukhov et al. (2010).




Figure 2. Example LSD profile for 61 Cygni A. Axes (top to bottom) show the mean Stokes V profile, Null profile and mean Stokes I profile. Note that the y-axis scale has been enlarged for the Stokes V and Null profiles for clarity.

4.2 Radial velocity

We used the LSD line profiles to measure the precise radial velocity of each target during each observation. We fitted a pseudo-Voigt profile (convolution of a Gaussian and a Lorentzian) to each Stokes I LSD profile and took the central velocity of the fitted model as the RV. We then RV-corrected each LSD profile and the corresponding spectra. In Table 3 we show the error-weighted average RV and standard deviation for each target. Note that the particularly large standard deviations shown in Table 3, such as for V471 Tau, are generally related to the changes in the RV throughout binary orbits.

4.3 Chromospheric activity and variability

4.3.1 Chromospheric S-index

The S-index (Vaughan et al. 1978) is the ratio of the fluxes in the Ca II H and K lines to the continuum flux either side of the H and K lines. It is calculated as

$$S_{MW} = \frac{aF_H + bF_K}{cF_{RHK} + dF_{VHK}} + e, \quad (3)$$

where F_H and F_K are the fluxes in two triangular bandpasses centred on the cores of the Ca II H and K lines (3968.469 Å and 3933.663 Å) with widths of 2.18 Å (at the base), and F_{RHK} and F_{VHK} are the fluxes in two rectangular 20 Å bandpasses centered on the continuum either side of the H and K lines at 3901.07 Å and 4001.07 Å. The coefficients a , b , c , d and e are derived in Marsden et al. (2014) to convert S-indices from NARVAL and ESPaDOnS to the Mount Wilson S-index scale.

We calculated S-indices using individual Stokes I sub-exposures as these were available on PolarBase for a large majority of our targets, as opposed to ‘mean’ Stokes I profiles which were only available for stars with full series’ of 4 sub-exposures. S-index errors were calculated by propagating the uncertainties computed during the reduction process for each spectral bin of the normalized Stokes I spectra through Equation 3.

For targets with ≥ 10 observations we removed outliers in the S-indices by excluding values outside 3 standard deviations from

the mean (as in Gomes da Silva et al. 2020). For smaller data sets we found this to inappropriately exclude values that were not outliers, so we did no σ clipping for these stars. Some observations measured intensities below zero in the region of the Ca II H&K lines. We filtered out observations where $>1\%$ of pixels within the H and K bandpasses had intensities < 0 .

Table 3 shows the mean S-index for each target ($\langle S \rangle$), weighted by the errors in individual S-index observations. We propagated S-index errors through the weighted mean equation, and they are so small that we have excluded them from Table 3. We have also measured the peak-to-peak S-index range (ΔS) as a representation of the amplitude of activity variability, as in Saar & Brandenburg (2002). We found the peak-to-peak amplitude to be a more realistic representation of activity variability compared to the standard deviation, since chromospheric activity cycles are not strictly periodic nor symmetrical about the mean.

4.3.2 Conversion from S-index to R'_{HK}

Chromospheric Ca II H&K emissions include a basal flux contribution (Schrijver 1987), unrelated to magnetic activity, and a photospheric contribution due to magnetic heating (Noyes et al. 1984). Both can be estimated based on the B-V colour of a star (Middelkoop 1982; Rutten 1984), and must be removed when comparing the chromospheric activity of stars across the HR diagram, to obtain a purely chromospheric activity index, R'_{HK} . This is often done using the method of Noyes et al. (1984, equation 4),

$$R'_{HK} = 1.34 \times 10^{-4} \cdot C_{cf} \cdot S_{MW} - R_{phot}, \quad (4)$$

where C_{cf} is the bolometric correction factor and R_{phot} is the photospheric contribution to the H and K bandpasses. However, the original calibration of the photospheric correction is valid only for $0.44 \leq B - V \leq 0.82$, while our sample cover $0.37 \leq B - V \leq 1.4$. Others have calibrated the Noyes et al. (1984) photospheric and bolometric corrections to a larger spectral range, such as Suárez Mascareño et al. (2015) ($0.4 \leq B - V \leq 1.9$), Astudillo-Defru et al. (2017) ($0.54 \leq B - V \leq 1.9$) and Boro Saikia et al. (2018a). We used the Suárez Mascareño et al. (2015) conversion, where C_{cf} and R_{phot} are calculated as

$$\log_{10} C_{cf} = 0.668(B - V)^3 - 1.270(B - V)^2 - 0.645(B - V) - 0.443 \quad (5)$$

$$\log_{10} R_{phot} = 1.48 \times 10^{-4} \cdot \exp[-4.3658(B - V)]. \quad (6)$$

Three stars within our sample are slightly outside the calibrated B-V range of the Suárez Mascareño et al. (2015) conversion, with B-V between 0.37 and 0.40. We still used the Suárez Mascareño et al. (2015) conversion for these stars, and we do not expect that this has any significant impact on the key findings of this study. We calculated the mean $\log R'_{HK}$ and $\log \Delta R'_{HK}$ range ($\log \Delta R'_{HK}$) based on the mean S-index and S-index range from Table 3, and the B-V from Table 1.

4.4 Longitudinal magnetic field

The longitudinal magnetic field, B_l , is the line-of-sight magnetic field averaged over the entire stellar disk. For the stars within our

sample that have Stokes V profiles available in PolarBase, we measured B_I using

$$B_I = -2.14 \times 10^{11} \frac{\int vV(v)dv}{\lambda g c \int [1 - I(v)]dv} \quad (7)$$

from Donati et al. (1997), where B_I is in gauss, $V(v)$ and $I(v)$ are the RV-corrected Stokes V and I LSD profiles respectively, which are normalised by the unpolarized continuum (I_c), λ and g are the normalization parameters from Table 2, c is the speed of light in km s^{-1} and v is velocity in km s^{-1} .

The velocity domain over which to integrate is unique for each target, and was chosen to include the entire Stokes V polarization signal while minimizing noise. We used the mean full-width at half-maximum (FWHM) of the Stokes I LSD profile (Table 3) to determine an appropriate domain according to the equation

$$v \in (\pm \alpha FWHM), \quad (8)$$

where $\alpha = 1.3$ for $FWHM < 40 \text{ km s}^{-1}$. For a small selection of stars with very broad profiles ($FWHM \geq 40 \text{ km s}^{-1}$) $\alpha = 0.8$ was more appropriate.

The error for each B_I observation was determined by propagating the uncertainties computed during the reduction process for each spectral bin of the normalized spectrum through equation 7. To determine the mean and range of the B_I , we first filtered out observations with very large uncertainties ($>200G$), and those for which the uncertainty in the B_I was greater than the measured value. For stars with >10 observations, we also removed B_I values that were more than 3 standard deviations from the mean. We measured the B_I from the Null profile, $B_{I,N}$, using the same method as above. Non-zero values of $B_{I,N}$ that are significant with respect to their uncertainties, and significant with respect to the B_I measured from the Stokes V profile, may indicate that our magnetic field detection is spurious. Therefore, we also filtered out observations with large detections in the Null spectrum ($|B_{I,N}| \geq |B_I|$). Table 3 shows the error-weighted mean, unsigned longitudinal field strength from the Stokes V and Null profiles ($\langle |B_I| \rangle$ and $\langle |B_{I,N}| \rangle$ respectively), as well as the peak-to-peak B_I range (ΔB_I). Uncertainties in the $\langle |B_I| \rangle$ and $\langle |B_{I,N}| \rangle$ were determined by propagating the errors in individual measurements through the weighted mean equation. We also show the mean uncertainty in the $B_{I,N}$ values, $\langle \sigma_{B_{I,N}} \rangle$, which provides an indication of the significance of the average detection in the Null spectrum. There are a few stars with large $\langle |B_{I,N}| \rangle$ values, but these are not significant relative to their average uncertainties, $\langle \sigma_{B_{I,N}} \rangle$.

4.5 Stellar rotation

We estimated stellar rotation periods ($P_{\text{est}, i=60^\circ}$) by assuming solid-body rotation, taking a constant inclination angle for all stars of 60° , and using the stellar radius and $v \sin i$ from Table 1.

5 RESULTS AND DISCUSSION

We measured chromospheric $\log R'_{\text{HK}}$ indices for 913 stars. For 621 of these we had multiple $\log R'_{\text{HK}}$ observations and could estimate the amplitude of the chromospheric activity variability, $\log \Delta R'_{\text{HK}}$. We measured the mean, unsigned, surface-averaged magnetic field strength, $\langle |B_I| \rangle$, for 425 stars, and estimated the B_I variability amplitude for 233 of this sample. Stars for which we were able to measure both the $\log R'_{\text{HK}}$ and $\langle |B_I| \rangle$ totalled 383. Table 3 provides

the measured parameters (the full table is available online only), which are also shown on the HR diagram in Figure 3.

The sample of stars with measured $\langle |B_I| \rangle$ includes a greater number of very cool stars compared to the $\log R'_{\text{HK}}$ sample, as shown in Figure 3. This is because cooler stars emit weakly in the blue spectral wavelengths, where the Ca II H&K lines lie, so SNRs in this domain are particularly poor. The $\log R'_{\text{HK}}$ sample consists of a greater number of mature, magnetically and chromospherically inactive stars compared to B_I sample, due to the lower-limit for detecting the large-scale magnetic fields for inactive stars.

Many of the stars we identified as young stars are located away from the main sequence in Figure 3, but several also appear to populate the main sequence. Given that those on the main sequence are highly active and have strong Li I absorption lines, it is possible that they are zero-age main sequence stars. Gomes da Silva et al. (2020) similarly found that their main sequence population included a small number of highly active stars, which had uncertain isochronal ages. They also concluded that the stars were probably younger compared to others on the main sequence. It is also possible that the positions of these stars on the main sequence is a reflection of our assumption of solar-metallicity when deriving the stellar temperatures and luminosities (section 2.2.1).

For stars with a large number of observations (e.g. ≥ 100), the means and amplitudes we report in Table 3 characterize the stellar variability on a rotational time-scale, as well as modulations related to magnetic and chromospheric activity cycles. For stars with fewer observations, the reported mean levels of chromospheric activity and magnetic field strength will be relatively poorly constrained. The $\log \Delta R'_{\text{HK}}$ and ΔB_I will be underestimated for targets with few observations, and are likely to only characterise the rotational modulation of chromospheric activity and magnetic field strength.

5.1 Mean chromospheric activities and large-scale magnetic field strengths

Figure 4 compares the mean $\log R'_{\text{HK}}$ with mean $\log |B_I|$ across sub-samples of F, G, K and M stars. The number of stars in each sub-sample is shown at the top of each plot, along with the Pearson correlation coefficient, ρ . Marker colour scales with the number of observations for each star, and marker shapes/styles are the same as in Figures 1 and 3. The black-line histograms indicate the distributions of the mean $\log |B_I|$ and mean $\log R'_{\text{HK}}$ across each sample of young stars, while the grey shaded histograms indicate the data distributions for mature stars and those with noisy spectra for which we could not determine if significant Li I lines were present. In Figure A1 we also show histograms of our chromospheric activity data combined with the data presented by Boro Saikia et al. (2018a) and Gomes da Silva et al. (2020).

5.1.1 Correlation between mean $\log R'_{\text{HK}}$ and $\log |B_I|$

Chromospheric activity across our sample ranges from $\log R'_{\text{HK}} \sim -3.5$ to -5.0 , and magnetic field strengths range from $\log |B_I| \sim -0.7$ to 2.7 (~ 0.2 to $500G$). Across all spectral types we find a positive correlation between mean $\log R'_{\text{HK}}$ and mean $\log |B_I|$, although the correlation is only marginal for F and M stars. For all spectral types, the correlation is improved when considering only stars with a high number of observations (≥ 100 observations, indicated by dark blue markers). Binary/multiple stellar systems generally lie along the same trend as single stars. Marsden et al. (2014) also found a positive correlation between chromospheric activity and the upper envelope

Table 3: Activity and magnetic field results for the stars shown in Table 1. The full table is available online. Columns show the weighted mean RV measured from the LSD profiles for each star, where the uncertainty in the RV is its standard deviation across observations, the mean LSD full-width at half-maximum (FWHM), the number of S-index observations (N_S), error-weighted mean and peak-to-peak range of the S-indices ($\langle S \rangle$ and ΔS), which are also converted to an R'_{HK} mean and range ($\log \langle R'_{\text{HK}} \rangle$ and $\log \Delta R'_{\text{HK}}$). Also shown are the number of B_I observations (N_{B_I}), the velocity range used to measure B_I , the mean unsigned B_I ($\langle |B_I| \rangle$) and peak-to-peak B_I range (ΔB_I), mean strength of the magnetic field in the Null profile ($\langle |B_{I,N}| \rangle$), mean error on the $B_{I,N}$ measurements ($\langle \sigma_{B_{I,N}} \rangle$) and estimated stellar rotation period, $P_{\text{est},t=60^\circ}$.

ID	RV (km s ⁻¹)	LSD FWHM (km s ⁻¹)	N_S	$\langle S \rangle$	ΔS	$\log \langle R'_{\text{HK}} \rangle$	$\log \Delta R'_{\text{HK}}$	N_{B_I}	Vel. range (km s ⁻¹)	$\langle B_I \rangle$ (G)	ΔB_I (G)	$\langle B_{I,N} \rangle$ (G)	$\langle \sigma_{B_{I,N}} \rangle$ (G)	$P_{\text{est},t=60^\circ}$ (d)
54 Psc	-32.7 ± 0.1	7.8	273	0.187	0.079	-4.914	-5.162	61	-43:-23	2.1 ± 0.1	8.8	0.3 ± 0.1	0.4	37.4
61 Cyg A	-65.5 ± 0.4	7.7	720	0.637	0.326	-4.725	-4.995	157	-76:-56	5.4 ± 0.0	25.6	0.3 ± 0.0	0.4	16.0
61 UMa	-5.4 ± 0.1	8.1	323	0.313	0.107	-4.529	-4.903	69	-15:5	2.7 ± 0.1	22.9	0.3 ± 0.1	0.4	16.2
χ^1 Ori	-13.8 ± 3.2	14.7	559	0.326	0.088	-4.397	-4.846	101	-31:7	3.1 ± 0.1	16.8	0.7 ± 0.1	0.9	4.8
ϵ Eri	16.5 ± 0.5	8.2	664	0.498	0.167	-4.500	-4.938	155	6:26	4.0 ± 0.0	25.1	0.2 ± 0.0	0.3	15.8
κ^1 Cet	19.2 ± 0.1	10.3	298	0.351	0.112	-4.410	-4.812	67	6:32	4.6 ± 0.1	22.3	0.6 ± 0.1	0.7	8.4
ξ Boo A	2.0 ± 0.3	9.7	675	0.452	0.109	-4.347	-4.905	167	-10:14	7.2 ± 0.0	30.8	0.4 ± 0.0	0.6	8.2
ρ^1 Cnc	27.5 ± 0.3	7.9	399	0.201	0.128	-4.865	-4.943	45	17:37	2.2 ± 0.0	11.7	0.2 ± 0.0	0.3	17.8
τ Boo	-16.3 ± 0.7	22.1	1634	0.190	0.099	-4.737	-4.675	142	-44:12	2.0 ± 0.1	14.6	0.9 ± 0.1	1.0	4.5
ν And	-28.4 ± 0.1	15.2	486	0.152	0.020	-4.951	-5.421	1	-48:-10	0.8 ± 0.5	-	0.4 ± 0.5	0.5	7.8
EMSR 9	-6.8 ± 1.6	22.5	6	5.819	4.122	-3.936	-4.084	138	-36:22	131.3 ± 1.2	558.3	11.3 ± 1.2	14.3	6.7
HD 166435	-14.2 ± 0.1	13.1	290	0.393	0.068	-4.326	-4.997	65	-29:3	7.1 ± 0.1	29.7	0.7 ± 0.1	1.0	5.5
HD 189733	-2.1 ± 0.3	8.9	583	0.541	0.300	-4.447	-4.669	133	-14:8	4.5 ± 0.0	26.2	0.4 ± 0.0	0.6	12.3
HD 190771	-25.8 ± 1.1	9.7	603	0.328	0.059	-4.435	-5.078	139	-38:-14	4.1 ± 0.1	25.0	0.5 ± 0.1	0.6	10.8
HD 206860	-16.6 ± 0.1	15.8	638	0.321	0.067	-4.407	-4.961	131	-37:3	6.4 ± 0.1	22.7	1.1 ± 0.1	1.6	4.3
HD 224085	-17.0 ± 56.8	30.7	483	2.960	2.010	-4.073	-4.237	63	-22:56	61.9 ± 0.5	325.4	3.4 ± 0.5	4.2	5.0
HD 75332	4.4 ± 0.4	14.6	416	0.270	0.038	-4.465	-5.118	73	-13:23	3.7 ± 0.2	18.6	1.2 ± 0.2	1.7	6.1
HD 76151	32.2 ± 0.1	7.7	271	0.235	0.056	-4.633	-5.102	60	23:43	2.9 ± 0.1	8.9	0.3 ± 0.1	0.4	37.4
HD 78366	26.3 ± 0.1	9.3	384	0.248	0.046	-4.555	-5.110	83	14:38	3.8 ± 0.1	16.5	0.5 ± 0.1	0.6	12.1
HD 9986	-20.8 ± 0.2	8.6	335	0.177	0.035	-4.815	-5.296	48	-32:-10	1.3 ± 0.1	6.0	0.4 ± 0.1	0.6	18.1
BP Tau	15.6 ± 1.5	13.4	158	8.170	10.245	-3.757	-3.657	85	-1:33	308.6 ± 1.9	576.3	12.4 ± 1.7	15.4	6.0
EK Dra	-20.3 ± 0.4	24.5	319	0.608	0.236	-4.163	-4.528	72	-51:11	20.8 ± 0.7	96.8	3.8 ± 0.7	6.0	2.7
TW Hya	12.7 ± 0.4	11.0	161	12.185	14.324	-3.562	-3.491	65	-1:27	262.2 ± 1.0	552.5	4.6 ± 0.8	6.6	3.4
V410 Tau	16.1 ± 5.5	98.1	118	2.208	1.327	-4.342	-4.557	84	-62:94	136.9 ± 5.0	655.3	34.6 ± 5.0	45.4	1.4
V471 Tau	1.8 ± 190.0	120.5	622	1.097	1.235	-4.104	-4.035	71	72:264	89.9 ± 4.6	457.7	25.1 ± 4.6	38.6	-
V830 Tau	17.2 ± 8.5	43.5	8	5.552	0.487	-3.918	-4.972	102	-17:51	76.9 ± 1.8	145.4	13.6 ± 1.8	17.7	2.1
V889 Her	-23.1 ± 0.7	53.3	283	0.613	0.172	-4.139	-4.640	72	-65:19	33.0 ± 0.9	119.3	5.4 ± 0.9	7.7	1.3
π^1 UMa	-12.6 ± 0.1	15.3	337	0.372	0.065	-4.349	-5.014	66	-31:7	6.1 ± 0.1	27.6	0.9 ± 0.1	1.2	4.7
.....														




Figure 3. Top: HR diagrams showing (left to right) mean chromospheric activity ($\log R'_{\text{HK}}$), amplitude of chromospheric activity variability ($\log \Delta R'_{\text{HK}}$) and number of observations used to measure the mean $\log R'_{\text{HK}}$. Bottom: HR diagrams showing (left to right) the mean, unsigned surface-averaged longitudinal magnetic field strength ($\log |B_l|$), amplitude of magnetic field strength variability ($\log \Delta B_l$) and number of observations used to measure the mean $|B_l|$. For all plots, the marker shapes and fill-styles are the same as in Figure 1. Li I abundant stars, which we take to be young stars, are indicated by open markers, while mature stars are shown by filled markers. Stars for which we were unable to determine if significant Li I lines were present are shown with a reduced opacity. Circles denote single objects, while star symbols indicate binaries for which B_l measurements do not appear to be impacted by blending between companions, and triangles represent binaries for which B_l is known to be impacted.

of the longitudinal field strength in their snapshot study of 170 F, G and K stars from the BCool sample, and Reiners et al. (2022) found an approximately linear power-law relation between nonthermal Ca II H&K emissions and magnetic flux (measured using Zeeman broadening) for their sample of 292 M dwarfs.

Interestingly, our results suggest that the relationship between the mean $\log R'_{\text{HK}}$ and mean $\log |B_l|$ may be marked by multiple regimes. These are the most clear for the sample of G stars in Figure 4, possibly because this sample is the largest compared to the other stellar types. For illustrative purposes in Figure 4 we show a continuous, piece-wise linear function (black line) which we fitted to the G-star data using a weighted least-squares fit (wherein data is weighted by the number of observations). For G stars with $\log |B_l|$ between ~ 2 and 0.4 , and $\log R'_{\text{HK}}$ between ~ -4.0 and -4.4 , chromospheric activity and magnetic field strengths decrease simultaneously. At $\log |B_l| \sim 0.4$, the field strength remains almost constant while $\log R'_{\text{HK}}$ decreases from ~ -4.4 to -4.8 . The relation then reverts back to decreasing $\log R'_{\text{HK}}$ with decreasing $|B_l|$ for $\log |B_l| < 0.4$ and $\log R'_{\text{HK}} < -4.8$.

We considered in detail if the multiple regimes could be related to some systematic issue with the estimation of the mean magnetic field strength. For example, it is possible that the mean $|B_l|$ could be overestimated for stars with $\log R'_{\text{HK}} \leq -4.4$ because we filter out observations where the measured B_l is below the noise level

(measured in the Null profile). Thus, for inactive stars, the mean $|B_l|$ could relate to magnetic maxima only, and this might explain why inactive stars appear to be ‘shifted’ to the right in $|B_l|$ space. However, when we plotted the maximum $|B_l|$ against the mean $\log R'_{\text{HK}}$ for the sample shown in Figure 4, distinct regimes were still evident. Therefore, overestimation of the mean $|B_l|$ for inactive stars is unlikely to fully explain the relationship shown in Figure 4.

Similarly, a systematic underestimation of the mean $|B_l|$ for low activity stars would not fully explain the observed shape of the $\log R'_{\text{HK}} - \log |B_l|$ relation. The $|B_l|$ could be underestimated for inactive stars due to their low rotational velocity, which may limit the spatial resolution of the Stokes V and I spectral line profiles. Or, if the magnetic field were concentrated into smaller spatial scales with decreasing activity (Petit et al. 2008), the B_l may be unable to reliably recover the field due to cancellations between small-scale, oppositely polarized regions. These scenarios could explain why we are unable to recover any increase in B_l as $\log R'_{\text{HK}}$ increases between -4.8 and -4.4 . However, they do not explain why we are able to recover increasing magnetic field strength with increasing chromospheric activity for stars with $\log R'_{\text{HK}} \leq -4.8$. For all inactive stars, we would instead expect to see a vertical stacking of stars having continually decreasing chromospheric activity and an almost constant, minimum detectable B_l .

Perhaps the strongest argument against a systematic cause for




Figure 4. Mean chromospheric activity versus mean unsigned longitudinal field strength for our sample, separated by spectral type. Marker shapes and fill-styles are the same as in Figure 1. Marker color scales with the number of observations. We take the number of observations as the average of the number of $\log R'_{\text{HK}}$ observations and $4 \times B_l$ observations (since $4 \times$ Stokes I observations are required to measure B_l). The grey shaded histograms above and to the right of each plot show the distributions of the mean $\log |B_l|$ and $\log R'_{\text{HK}}$ across each sample of main sequence stars. The black line histograms indicate the distributions of data for young, lithium rich stars only. The dashed lines indicate the limits of possible reduced populations in the activity distributions; the lines are red for the F and K stars because we did not find the reduced populations to be significant with respect to the histogram uncertainties in Figure A1. The solid black line shown for G stars indicates a continuous piece-wise linear fit to the data.

the three regimes in the $\log R'_{\text{HK}} - \log |B_l|$ relation, is the fact that we also see a similar and possibly related ‘step’ in the relationship between chromospheric activity variability amplitudes and mean chromospheric activity, which we discuss further in section 5.2.1. If the distinct regimes in the $\log R'_{\text{HK}} - \log |B_l|$ relation are real, they may relate to a change in the surface properties of the magnetic dynamo at around $\log |B_l| \sim 0.4$. One possibility is a change in the area ratio of chromospheric plages to photospheric spots, as suggested by Foukal (2018). Foukal (2018) proposed that stellar plage/spot ratios may be lower for stars that are slightly more active compared to the Sun ($\log R'_{\text{HK}} \sim -4.8$ in cycle 23, Lehmann et al. (2021)), based on the fact that the Sun exhibits a reduced plage/spot area ratio when at its most active (Foukal 1998). This phenomenon could also relate to the well-observed change from spot-dominated to faculae-dominated photospheric variability with decreasing mean chromospheric activity and increasing stellar age (Radick et al. 1998; Lockwood et al. 2007; Radick et al. 2018), which occurs at $\log R'_{\text{HK}} \sim -4.65$ to -4.75 . If stars with $\log |B_l| \sim 0.4$ are impacted by a reduced plage/spot ratio then they would appear weaker

in terms of their chromospheric activity (which relates to chromospheric plages) relative to their $|B_l|$ (which relates to the magnetic fields in both spots and faculae). The affected stars would drop vertically down in our Figure 4, which would be consistent with our observations. Interestingly, Nichols-Fleming & Blackman (2020) observed a similar, sudden decrease in X-ray luminosity variability (which is known to correlate strongly with chromospheric Ca II H&K emissions) for G stars, which occurred at a rotation period of ~ 15 d, and this rotation period also coincides with our observed $\log R'_{\text{HK}}$ ‘step down’ for G stars (see Figure B1).

5.1.2 Trends with stellar age and spectral type

For both G and K stars, chromospheric activity and magnetic field strengths are higher for young stars (open circles) compared to the more mature main sequence stars (filled circles). For F and M-type stars, there appears to be a similar decrease in activity and magnetic field strengths from young to mature stars, but our sample size, particularly the samples of young stars, are small compared to G

and K stars. As a result, the distributions of chromospheric activity and magnetic field strengths for young F and M stars are dominated by statistical noise.

The lower-limits of chromospheric activity and magnetic field strengths for young G and K stars are similar ($\log R'_{\text{HK}} \sim -4.3$ and $\log |B_I| \sim 1.0$), but the young K stars extend to higher mean $\log R'_{\text{HK}}$ and $\log |B_I|$ compared to the G stars. This is likely to be related (at least partly) to a sample bias; young and highly active K (and M) stars are targeted in priority for spectropolarimetric observations to increase the likelihood of a magnetic detection. The population of young stars appears to correspond to the ‘very-active’ group of stars with $\log R'_{\text{HK}} \geq -4.2$ observed by [Gomes da Silva et al. \(2020\)](#).

For both G and K stars, the populations of young stars are separated in both the $\log R'_{\text{HK}}$ and $\log |B_I|$ histograms from the populations of mature, main sequence stars, but again it is possible that the apparent grouping of stars is an effect of targeted observing programs with NARVAL and ESPaDOnS. If it is a real phenomenon, the under-density of stars between the young and main sequence populations resembles the ‘gap’ in stellar rotation periods described by [Barnes \(2003\)](#) that occurs around the transition from the pre-main sequence to the main sequence. [Barnes \(2003\)](#) propose that the core and envelope of a star re-couple in this region, and the magnetic field may change from a ‘convective’ field to an ‘interface’ field produced at the tachocline. The presence of the gap in both the chromospheric activity and magnetic field strengths is consistent with a rapid evolution between these pre-main sequence and main sequence phases.

For main sequence F, G and K stars, the lower and upper-limits of chromospheric activity and magnetic field strengths are similar, while our sample of M stars appears to generally have higher $\log R'_{\text{HK}}$ and $\log |B_I|$. Again, this may be partly related to a sample bias, whereby active M stars are preferentially targeted. Or it could be related to the longer main sequence lifetimes of mid-M stars, their less-efficient activity/rotation spin down or the difference in convective turnover timescales between stellar types.

The $\log R'_{\text{HK}}$ histograms for main sequence F, G and K stars (grey shaded histograms) show that chromospheric activity may be bimodally distributed, or could be skewed toward inactive stars with an extended tail toward higher activity stars. [Figure 4](#) shows slightly under-populated regions of F stars between $\log R'_{\text{HK}} = -4.7$ and -4.8 , G stars between -4.4 and -4.8 , and K stars between -4.5 and -4.7 . In each case, the under-density of stars is not significant for our sample when we consider the uncertainties in the histogram bins ([Figure A1](#)), but a bimodal distribution is significant for G stars when we combine our data with that from [Boro Saikia et al. \(2018a\)](#) and [Gomes da Silva et al. \(2020\)](#). Similar to [Boro Saikia et al. \(2018a\)](#), the possible reduced population of intermediately-active G stars in [Figure A1](#) is less prominent compared to the ‘Vaughan-Preston gap’, originally shown by [Noyes et al. \(1984\)](#) to be completely devoid of F and G stars between $\log R'_{\text{HK}} = -4.6$ and -4.9 . This is likely due to our larger sample size compared to [Noyes et al. \(1984\)](#). We do not observe any distinct, ‘very-inactive’ group of stars in any of the chromospheric activity distributions, with $\log R'_{\text{HK}} \leq -5.0$, as was detected by [Gomes da Silva et al. \(2020\)](#). This is probably because their data includes stars evolving off the main sequence, while our sample excludes these stars.

In contrast to the chromospheric activity distributions, the $\log |B_I|$ data for main sequence stars seem to be normally distributed, particularly across each of the G and K samples. It is interesting that the low-population region that separates young stars from old for both the G and K samples, is evident in both $\log |B_I|$ and $\log R'_{\text{HK}}$ distributions, while the reduced population of main

sequence G stars with intermediate activity is shown in only the $\log R'_{\text{HK}}$ distribution. The fact that we see no discontinuity in $\log |B_I|$ for G stars at intermediate activity supports a change in dynamo properties on the main sequence, such as those we discussed in [section 5.1.1](#), rather than a rapid evolutionary phase.

5.2 Chromospheric activity and magnetic field strength variability amplitudes

[Figures 5](#) and [6](#) compare the peak-to-peak amplitudes of $\log R'_{\text{HK}}$ and $\log |B_I|$ variability with the mean $\log R'_{\text{HK}}$ and mean $\log |B_I|$ respectively. In [Figure 5](#), stars are grouped by spectral type, and the marker color scales with the number of observations. In [Figure 6](#) we show all spectral types together, and in [6\(a\)](#) marker color scales with the number of observations, while in [\(b\)](#) it indicates stellar temperature. As in [Figure 4](#), the number of stars in each sub-sample and the Pearson correlation coefficient are shown at the top of each plot. Filled markers represent mature stars, open markers show young stars and the black line histograms indicate the distributions of activity and magnetic field strength for the young sample. Due to the significant scatter in these plots, we show the distributions of $\log \Delta R'_{\text{HK}}$ and $\log \Delta |B_I|$ for each entire main sequence sample in orange. The grey histograms relate to filtered samples of main sequence stars that have ≥ 10 observations.

5.2.1 $\log \Delta R'_{\text{HK}}$ versus mean $\log R'_{\text{HK}}$

For G, K and M stars, the amplitude of $\log R'_{\text{HK}}$ variability shows a moderate, positive correlation with the mean $\log R'_{\text{HK}}$. [Saar & Brandenburg \(2002\)](#) similarly observed an increase in chromospheric activity amplitudes with mean activity above $\log R'_{\text{HK}} = -5.0$. Meanwhile, for F stars there is only a weak correlation between chromospheric activity variability and the mean activity level. The positive correlations for G and K stars are strongest for stars with a high number of observations. Unlike in [Figure 4](#), where stars with few observations are scattered both above and below the main trend, it is clear in [Figure 5](#) (and [6](#)) that stars with few observations are scattered mostly below the trend. This is because variability amplitudes are inevitably underestimated for poorly observed stars. Some of the scatter will also be due to the fact that activity is restricted in latitude, such that stars with different inclinations to the observer will have different apparent activity amplitudes ([Saar & Brandenburg 2002](#)).

Considering targets with at least ~ 50 obs. (green to dark blue markers), there appear to be at least two regimes of activity variability for G stars, and possibly for K stars. G stars show decreasing chromospheric activity variability with decreasing mean activity between $\log R'_{\text{HK}} \sim -3.8$ and -4.4 , but below $\log R'_{\text{HK}} \sim -4.4$ there is only a minimal change in chromospheric activity variability with decreasing mean activity. It is possible that chromospheric variability again begins to decrease with mean activity below $\log R'_{\text{HK}} \sim -4.8$, but this is based on a very small sample of well-observed stars. Our chromospheric activity variability data combined with the results from [Gomes da Silva et al. \(2020, Figure C1\)](#) are also consistent with a non-linear relationship between $\log \Delta R'_{\text{HK}}$ and $\log R'_{\text{HK}}$. Although the functional form of the relationship is poorly constrained, the combined data set supports multiple regimes of chromospheric activity variability. [Gomes da Silva et al. \(2020\)](#) interpreted their results as three possible chromospheric variability regimes but they found that only the upper envelope of chromospheric activity variability scales with mean activity across all regimes. Given that the




Figure 5. Amplitude of chromospheric activity variability versus mean chromospheric activity for samples of F, G, K and M stars. Marker shapes and fill-styles are the same as in Figure 1. Marker color scales with the number of observations. The histograms to the right of the plots indicate the distributions of $\log \Delta R'_{HK}$ for the entire main sequence sample (orange), main sequence stars with ≥ 10 observations (grey), and young stars (black line). The dashed lines correspond to the reduced population of G stars from Figure 4.

significant scatter in Figure 5 seems to be mostly related to stars with low numbers of observations, we find that both the upper and lower levels of activity variability scale well with mean activity.

It is interesting that the possible changes in the $\log \Delta R'_{HK}$ - mean $\log R'_{HK}$ relation seem to occur at a similar activity level as changes in the mean $\log R'_{HK}$ - $\log |B_l|$ relation, and may coincide with the reduced-population region of intermediately-active G stars from Figure 4 (indicated by dashed lines in Figure 5). This suggests that the change in the mean $\log R'_{HK}$ - $\log |B_l|$ relation is not related to some systematic over or underestimation of $|B_l|$ for low-activity stars. Rather, it supports a change in the properties of the magnetic dynamo at the stellar surface.

For our samples of G and K-type stars, the mature main sequence stars generally have lower mean chromospheric activity and variability compared to the more youthful stars. Variability amplitudes appear to have similar ranges across spectral types F to K, for both the young and main sequence samples, and the variability amplitudes of young stars are not distinctly separated from the variability of mature stars. The chromospheric activity variability of main sequence F, G and K stars appears to be normally distributed, similar to the $\log |B_l|$ distributions shown in Figure 4. There is no reduced population in the $\log \Delta R'_{HK}$ distribution that corresponds to the reduced population of main sequence G stars with intermediate mean chromospheric activity in Figure 4. Although the orange histograms in Figure 5 show possibly bimodal distributions of $\log \Delta R'_{HK}$, this is clearly related to the scattering of data below the main trend.

5.2.2 Magnetic field variability versus mean field strength

Figure 6 shows a strong, positive correlation between $\log \Delta B_l$ and mean $\log |B_l|$ for well observed stars (green to dark blue markers). Young stars typically have stronger magnetic fields with greater variability amplitudes compared to mature stars, and there is clear separation between our samples of young and mature stars in the $\log \Delta B_l$ distribution. K and M stars generally show higher magnetic field strengths and magnetic field variability compared to the warmer F and G stars, but this is somewhat impacted by the fact that active K and M stars are preferentially targeted for spectropolarimetric observations.

5.3 Chromospheric activity versus magnetic field strength from ZDI

Figure 7 compares the surface-averaged large scale magnetic field strength from published ZDI maps³ to the mean $\log R'_{HK}$ activity we calculated from PolarBase observations. For stars that have been observed using ZDI over multiple epochs, we show the mean field strength from all ZDI maps. We have removed outliers by excluding stars for which the mean $\log R'_{HK}$ is derived from ≤ 20 Stokes I observations.

The data indicate that mean chromospheric activity is directly related to the strength of the large-scale magnetic field recovered from ZDI. This is consistent with previous work by Petit et al.

³ Boro Saikia et al. (2015, 2016, 2018b); Brown et al. (2021); do Nascimento et al. (2016); Donati et al. (2008a,b,c, 2010, 2011a,b, 2013, 2014, 2015); Fares et al. (2009, 2010, 2012, 2013); Folsom et al. (2016, 2018a,b, 2020); Hébrard et al. (2016); Jeffers et al. (2014, 2017, 2018); Marsden et al. (2006); Mengel et al. (2016); Morgenthaler et al. (2011); Morin et al. (2008, 2010); Petit et al. (2008); See et al. (2019a); Waite et al. (2015, 2017)




Figure 6. Magnetic field variability amplitude versus mean magnetic field strength. Marker shapes and fill-styles are the same as in Figure 1. In (a) the marker color scales with the number of observations, while in (b) marker color scales with stellar effective temperature. The histograms to the right of the plots indicate the distributions of ΔB_l for the entire main sequence sample (orange), main sequence stars for which the variability amplitude is based on ≥ 10 observations (grey), and young stars (black line).




Figure 7. Mean chromospheric activity from our study versus surface-averaged large-scale magnetic field strength from published ZDI work. Marker fill-styles are the same as in Figure 1; open circles represent young stars with strong spectral Li I lines and filled circles are mature stars assumed to be on the main sequence. The black regression line has a slope of 0.44 ± 0.04 , and the grey shaded region shows its 95% confidence interval. The red line shows the previous results of Petit et al. (2008).

(2008), although they found a lower slope, $R'_{\text{HK}} \propto \langle |B| \rangle_{\text{ZDI}}^{0.33}$, compared to $R'_{\text{HK}} \propto \langle |B| \rangle_{\text{ZDI}}^{0.44 \pm 0.04}$ for our data. If we consider only main sequence F-K stars we obtain a slope of ~ 0.29 , closer to the results of Petit et al. (2008). The relationship between $\log R'_{\text{HK}}$ and $\log \langle |B| \rangle_{\text{ZDI}}$ is consistent with the dependence of $\log R'_{\text{HK}}$ on the surface-averaged longitudinal magnetic field shown in Figure 4, although we do not find evidence of a change in slope between $\log R'_{\text{HK}}$ and $\log \langle |B| \rangle_{\text{ZDI}}$. This is probably due to the small sample size in Figure 7 compared to Figure 4.

5.4 Chromospheric activity and magnetic field geometry versus effective temperature

Figure 8 shows mean chromospheric activity versus stellar effective temperature for our sample (dark grey markers). We have also included the publicly available chromospheric activity data from Gomes da Silva et al. (2020), and activity data compiled by Boro Saikia et al. (2018a) (light grey markers). Activity amplitudes (sym-

bol size) and the poloidal fraction of the large-scale magnetic field (symbol colour) are also shown in Figure 8 for a selection of stars with published ZDI maps. We show the Sun with the usual symbol, which is scaled for its activity variability during cycle 23 (Lehmann et al. 2021). The fractional poloidal field of the Sun we show here is the minimum value measured by Lehmann et al. (2021); note that since the Sun's magnetic field has been observed in much higher cadence compared to the other stars, its large-scale toroidal component has been better recovered. For the combined sample shown in Figure 8, we also computed a two-dimensional histogram which is shown in Figure 9.

5.4.1 Young stars

Youthful stars, shown by open circles, are clearly more active compared to main sequence stars across all temperatures, and have greater activity variability, as has been shown in the previous sections. In the upper right of Figure 8 there are a number of very-cool, highly active stars that are marked as main sequence age (filled circles), for which we did not detect strong Li I lines within the spectra. According to SIMBAD these are all known to be young, pre-main sequence stars (OT Ser, DS Leo, BD+132618, BD+61195 and HD209290).

Young stars with $\log R'_{\text{HK}} \geq -4.0$ and $3800 \leq T_{\text{eff}} \leq 4000$ K appear to have strongly poloidal magnetic field geometries. This contrasts against young, active stars with $T_{\text{eff}} \leq 3800$ K, which show mixed field geometries and comparatively lower variability in chromospheric activity. Morin et al. (2011) suggested that the presence of both dominantly toroidal and dominantly poloidal fields in active M stars could indicate dynamo bi-stability. Another interpretation is that such stars have similar magnetic cycles to the solar cycle, and that those observed to have dominant toroidal fields have simply been observed at times of a magnetic field inversion, which is when the solar magnetic field becomes its least poloidal (Kitchatinov et al. 2014).

Young stars with $T_{\text{eff}} \geq 4000$ K also show mixed field geometries, and have $\log R'_{\text{HK}}$ between ~ -4.0 and -4.3 . The change in field geometry at ~ 4000 K from poloidal to mixed geometries has been well observed (Donati et al. 2011c; Gregory et al. 2012; Folsom et al. 2018a; Hill et al. 2019), and is thought to occur around the




Figure 8. Mean chromospheric activity versus stellar effective temperature for our sample of stars (dark grey) and published data (light grey) from [Boro Saikia et al. \(2018a\)](#) and [Gomes da Silva et al. \(2020\)](#). The properties of the large-scale magnetic field are shown for a selection of stars with published ZDI maps. Marker color represents the fraction of the large-scale magnetic field that is stored in the poloidal component and marker size represents our measured activity variability amplitude, $\log \Delta R'_{HK}$. Marker fill-styles are the same as in Figure 1, and the Sun is indicated with the usual symbol. The black curve corresponds to the black curve indicated in Figure 9.




Figure 9. Two-dimensional histogram indicating the number of stars in Figure 8 populating equal-sized temperature and $\log R'_{HK}$ bins. For targets that are common to both PolarBase and the published data we take the mean of all available $\log R'_{HK}$ values for Figure 9. The black curve we have drawn is intended to draw the eye to the densely populated region we discuss in section 5.4, and corresponds to the black curve shown in Figure 8.

transition from fully-convective to partially-convective stars (Folsom et al. 2018a; Hill et al. 2019; Villebrun et al. 2019). For the young stars with $T_{\text{eff}} \geq 4000\text{K}$ there is no clear dependence of the strength of the poloidal field on stellar temperature or mean activity level. This is also shown in Figure 10, where we directly compare $\log R'_{\text{HK}}$ and $\log \Delta R'_{\text{HK}}$ to the fraction of the magnetic field stored in the poloidal component, and the fraction of the poloidal field stored in high order modes. For young stars, the poloidal field may reach a possible minimum fraction of 20-40 percent at around $\log R'_{\text{HK}} \sim -4.25$, before stars reach the main sequence.

5.4.2 Main sequence stars

For main sequence stars with $4200 \leq T_{\text{eff}} \leq 6300\text{K}$, the upper level of chromospheric activity has little dependence on T_{eff} . This suggests that late-F to late-K stars begin their main sequence lives with a similar level of chromospheric activity, $\log R'_{\text{HK}} \sim -4.3$. The basal level of chromospheric activity also appears to be independent of T_{eff} for $4500 \leq T_{\text{eff}} \leq 6500\text{K}$, apart from a small very-low activity group of stars at around 5000 K, which are from the Gomes da Silva et al. (2020) sample and are likely to be subgiant stars evolving off the main sequence. The basal activity increases for stars with $T_{\text{eff}} \approx 4500$ toward 4200 K. Mittag et al. (2013) found a similar lack of low activity stars in the B-V range of 1.1 to 1.5. A simple explanation for this may be that K and M stars have different convective properties compared to F and G stars, and/or longer main sequence lifetimes with less-efficient loss of activity as they spin-down, such that they have not reached a similarly low activity level. For $T_{\text{eff}} \leq 4200\text{K}$, the upper and lower activity levels for main sequence stars decrease toward lower temperatures. It is not clear why these very cool stars begin the main sequence with lower chromospheric activity compared to F, G and K stars, but decreasing $\log R'_{\text{HK}}$ with decreasing T_{eff} is consistent with previous work (Mohanty & Basri 2003; Reiners & Basri 2008; Astudillo-Defru et al. 2017). For main sequence stars with $T_{\text{eff}} \geq 6300\text{K}$, the population of very low activity stars ($\log R'_{\text{HK}} \sim -5.0$) begins to disappear. It is possible that this is because the chromospheric activity for such low-activity stars is difficult to detect, or could be related to differences in convective properties for F stars. It should also be noted that the stars with $T_{\text{eff}} \geq 6300\text{K}$ and $\leq 4200\text{K}$ are mostly taken from Boro Saikia et al. (2018a), and extend beyond the S-index to $\log R'_{\text{HK}}$ calibration range we use here, which should be taken into account when comparing trends.

The ZDI results show clear evolution of the large-scale surface magnetic field on the main sequence. Figures 8 and 10 indicate that for main sequence F, G and K stars, the poloidal component of the magnetic field increases as the mean $\log R'_{\text{HK}}$, and possibly the $\log \Delta R'_{\text{HK}}$, decrease. This is consistent with previous studies, which showed that both the toroidal energy fraction (Donati & Landstreet 2009; Folsom et al. 2016; See et al. 2016) and chromospheric activity cycle amplitudes (Saar & Brandenburg 2002) decrease with the Rossby Number (Ro , ratio of rotation period to convective turnover time, see Appendix D1). Figure 10 also shows a wide range in the complexity of the poloidal field component for main sequence F-K stars, from fairly simple fields with almost all of the poloidal field energy stored in octopolar or lower modes, to as much as 80 percent of the poloidal field stored in higher order modes.

The late-F to late-G stars shown in Figure 8 appear to transition at $\log R'_{\text{HK}} \sim -4.5$ from being capable of generating significant toroidal fields to having dominantly poloidal fields. Our data are also consistent with K stars undergoing a shift from significantly toroidal to dominantly poloidal fields at a similar activity level to G

and late-F stars, although published ZDI maps are sparse for K stars around $\log R'_{\text{HK}} = -4.5$. For mid-F stars, strong toroidal fields may prevail toward lower activity levels, suggesting that the transition to dominantly poloidal fields may occur at a lower $\log R'_{\text{HK}}$. It is not clear from our sample if M stars transition to dominantly poloidal field structures as their chromospheric activity and activity variability decrease.

The 2-dimensional histogram in Figure 9 indicates a densely populated, low activity region with $5000 \leq T_{\text{eff}} \leq 6300\text{K}$ and $-5.05 \leq \log R'_{\text{HK}} \leq -4.75$ (marked by a black curve in Figures 8 and 9). Above this region, there is a slightly reduced population of late-F to early-K stars with activity between -4.50 and -4.75 . The reduced population is less-prominent compared to the original Vaughan-Preston gap, which is consistent with the findings of Boro Saikia et al. (2018a). Above the reduced population, there is also a slightly higher-density band of active main sequence stars, with $\log R'_{\text{HK}}$ between ~ -4.30 and -4.50 . This band covers a smaller temperature range compared to the densely populated region of inactive stars (marked by the black curve), and may extend from ~ 6100 to 5300K . These features are consistent with the bimodal distribution of main sequence G stars shown in Figure A1, and suggest that the bimodal distribution could extend to slightly higher and lower temperatures to include late-F to early-K stars. For stars with T_{eff} below $\sim 5000\text{K}$, the distribution of activity appears to be fairly flat and dominated by statistical noise. Gomes da Silva et al. (2020) observed a triple-peaked distribution in their sample of stars with $4350 \leq T_{\text{eff}} \leq 5280\text{K}$ (which they classify as K stars), but we do not find this to be clearly evident in Figure 9, nor significant with respect to the histogram uncertainties in Figure A1.

Similar to the findings of Metcalfe et al. (2016), our results indicate that the transition from strong toroidal fields to dominantly poloidal fields occurs at a similar activity level as the upper-boundary of the under-populated region of intermediately-active main sequence stars we identified in Figure 9. Metcalfe et al. (2016) suggested that the underpopulated region could be explained by a shift in the character of differential rotation, possibly resulting in a period of rapid evolution. Considering that the under-density also occurs at a similar activity level as the possible changes in the relationships between mean activity, magnetic field strength and activity variability that we described in the previous sections, we find that the under-populated region is more likely related to a change in magnetic surface properties, such as a change in the plage-spot ratio, rather than a period of rapid evolution.

5.5 Chromospheric activity and magnetic field geometry versus rotation period

Figure 11 compares chromospheric activity, activity variability (symbol size) and magnetic field geometry (symbol colour) to the stellar rotation period for subsets of main sequence F, G, K and M stars, as well as a separate plot for young, Li I abundant stars. Note that the rotation periods used here are estimates only, based on our derived stellar radius, $v \sin i$ and assuming an inclination angle of 60° .

The data show that mean activity decreases as stellar rotation slows for all stars on the main sequence. This is consistent with stellar spin-down theory (Skumanich 1972), where young stars have high magnetic field strengths and chromospheric activity, which diminish throughout their main sequence lifetimes due to angular momentum loss from the coupling of the magnetic field and stellar wind. The chromospheric activity of F, G and K stars appears to plateau at $\log R'_{\text{HK}} \approx -5.0$ for slower rotating stars. This levelling-




Figure 10. Left: $\log R'_{\text{HK}}$ and $\log \Delta R'_{\text{HK}}$ versus the fraction of the large scale magnetic field stored in the poloidal component. Right: $\log R'_{\text{HK}}$ and $\log \Delta R'_{\text{HK}}$ versus the fraction of the poloidal field stored in octopolar or higher energy modes. Marker color scales with stellar effective temperature, and marker styles are the same as in Figure 1. We have removed outliers by excluding stars for which $\log R'_{\text{HK}}$ and $\log \Delta R'_{\text{HK}}$ are derived from ≤ 20 high SNR observations.

off of chromospheric activity as rotation slows is not obvious for our sample of M stars, possibly because our sample is biased toward active M stars, or because low mass M stars have longer spin-down timescales (West et al. 2008; Johnstone et al. 2021). Petit et al. (2008) found for their sample of G stars that a rotation period lower than ~ 12 d is necessary for the toroidal magnetic energy to dominate over the poloidal component. This is consistent with our results for main sequence G-type stars. Our data also suggest that dominant toroidal fields are only present for main sequence F stars with estimated rotation periods below ~ 7 d, and for main sequence K stars with estimated rotation periods below ~ 16 d. This is likely to be related to convection zone depth, which decreases from K to F stars. The change from strongly toroidal to dominantly poloidal fields is hypothesized to occur when the rotation period becomes close to the convective turnover time ($Ro \sim 1$, see Appendix D1), which would occur at shorter rotation periods for F stars with shallow convective zones, and at longer rotation periods for later-type stars with deeper convection zones.

Young stars with greater chromospheric activity and variability, and dominantly poloidal field structures, have longer rotation periods compared to stars those with weaker activity variability and which are capable of generating both dominantly poloidal and toroidal fields. This is consistent with previous work on young stars (Folsom et al. 2018a; Hill et al. 2019; Villedun et al. 2019), which indicate that magnetic fields become more complex as young stars contract toward the main sequence and their rotation rate increases. The transition from dominantly poloidal to significantly toroidal field structures appears to occur at an estimated rotation period of ~ 4.5 d.

6 CONCLUSIONS

We provide a catalog of new chromospheric activity and surface-averaged large-scale magnetic field measurements for 954 main

sequence and youthful stars with spectral types from mid-F to mid-M. This includes mean values and peak-to-peak variability amplitudes for both the $\log R'_{\text{HK}}$ and B_l , which were derived using time-series spectropolarimetric observations from the PolarBase database. In this work we compare chromospheric activity, magnetic field strengths and their variability amplitudes. We also use previously published chromospheric activity data and ZDI results to diagnose relationships between chromospheric activity, activity variability, large-scale magnetic field geometries and stellar properties.

The data confirm that chromospheric activity, magnetic field strengths, and their variability amplitudes are generally higher for youthful stars compared to more mature, main sequence stars. Stars from late-F through to mid-K type appear to have similar levels of activity and similar magnetic field strengths as they begin their main sequence lives. For F, K and M stars, chromospheric activity decreases fairly smoothly with decreasing mean, unsigned magnetic field strength. Conversely for G stars, our data suggest that the $\log R'_{\text{HK}} - \log |B_l|$ relationship may have three distinct phases; mean chromospheric activity and field strength decrease together until $\log R'_{\text{HK}} \sim -4.4$ and $\log |B_l| \sim 0.4$, at which point there is a significant step down in chromospheric activity but minimal change in magnetic field strength. This is followed by another decreasing phase for both chromospheric activity and magnetic field strength for $\log R'_{\text{HK}} \leq -4.8$ and $\log |B_l| \leq 0.4$. The distinct phases in the $\log R'_{\text{HK}}$ and $\log |B_l|$ relationship could be related to a change in the surface properties of the magnetic dynamo on the main sequence, such as a change in the area ratio of plages and spots. Further long-term observations of the magnetic fields and chromospheric activity of stars across a range of activity levels will be required to determine if the three phases we observed are real, and if they are unique to G stars.

The amplitudes of chromospheric activity and magnetic field variability show clear dependence on the mean chromospheric activity and magnetic field strengths. However, the proportionality




Figure 11. Mean $\log R'_{\text{HK}}$ versus the estimated stellar rotation period for our sample of stars (grey). Rotation periods were derived from the published stellar radius and $v \sin i$ from Table 1 and assuming an inclination angle of 60° . The properties of the large-scale magnetic field are also shown for a selection of stars with published ZDI maps. Markers are the same as in Figure 8. Note that for young stars the plot shows a reduced range of rotation periods. The vertical dashed lines indicate the estimated transition from stars capable of generating significant toroidal field geometries, to those with dominantly poloidal fields.

between $\log \Delta R'_{\text{HK}}$ and mean $\log R'_{\text{HK}}$ appears to undergo a change at around the middle of the main sequence, similar to the relation between the mean $\log R'_{\text{HK}}$ and mean $\log |B_l|$. Stars with both high and low mean $\log R'_{\text{HK}}$ can show similar levels of chromospheric activity variability.

The distribution of $\log R'_{\text{HK}}$ data from our study, combined with data from Gomes da Silva et al. (2020) and Boro Saikia et al. (2018a), indicates a slightly under-populated region of intermediately-active, main sequence stars with spectral types from late-F to early-K and mean $\log R'_{\text{HK}}$ between -4.5 and -4.75. The under-density of stars is not as distinct as the original Vaughan-Preston gap, which is consistent with the findings of Boro Saikia et al. (2018a). We do not find evidence for similar under-populated regions in the distributions of chromospheric activity variability, magnetic field strength nor magnetic field variability across our sample of stars. If the bimodal distribution of chromospheric activity we observed for late-F to early-K, main sequence stars is real, the fact that we do not see similar distributions in the magnetic field strength or chromospheric activity variability across our sample supports a change in dynamo properties on the main sequence, rather than a period of rapid stellar evolution.

The ZDI data reveal that young stars are able to produce both dominantly poloidal fields and generate strong toroidal fields, like the most active main sequence stars. As stars spin down on the main sequence, mean chromospheric activity and activity variability amplitudes decrease as the large-scale magnetic field becomes more dominantly poloidal. Our results are consistent with mid-to-late F stars, G and K stars losing their ability to generate dominantly toroidal fields at estimated rotation periods of ~ 7 , 12 and 16 d respectively. For G-type stars, this occurs at $\log R'_{\text{HK}} \sim -4.5$, and may roughly coincide with changes in the relationships between mean chromospheric activity, activity variability amplitudes and mean large-scale magnetic field strengths.

ACKNOWLEDGEMENTS

We thank the referee for their constructive and thorough review, which has helped us to improve the quality of this manuscript.

This work is based on observations obtained at the TBL and CFHT. The TBL is operated by the Institut National des Sciences de l'Univers of the Centre National de la Recherche Scientifique of France (INSU/CNRS). The CFHT is operated by the the National Research Council of Canada, the INSU/CNRS and the University of Hawaii. We thank the staff at the TBL and CFHT for their time and data. The observations at the CFHT were performed with care and respect from the summit of Maunakea which is a significant cultural and historic site. We also acknowledge the use of the PolarBase database, which makes TBL and CFHT observations publicly available, and is operated by the CNRS, Observatoire Midi-Pyrénées and Université Toulouse III - Paul Sabatier.

We acknowledge use of the SIMBAD and VIZIER data bases operated at CDS, Strasbourg, France. This work has also made use of the VALD database, operated at Uppsala University, the Institute of Astronomy RAS in Moscow, and the University of Vienna.

ELB is supported by an Australian Postgraduate Award Scholarship. SVJ acknowledges the support of the German Science Foundation (DFG) priority program SPP 1992 'Exploring the Diversity of Extrasolar Planets' (JE 701/5-1). AAV acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 817540, ASTROFLOW). MMJ acknowledges fund-

ing from STFC consolidated grant ST/M001296/1. SBS acknowledges the support of the Austrian Science Fund (FWF) Lise Meitner project M2829-N. VS acknowledges funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement No. 682393 AWESoMeStars) and support from the European Space Agency (ESA) as an ESA Research Fellow.

DATA AVAILABILITY

All NARVAL and ESPaDOnS data presented here are publicly available through the PolarBase data base (<http://polarbase.irap.omp.eu/>). Archival chromospheric activity data used for this study are available via VIZIER.

REFERENCES

- Arriagada P., 2011, *ApJ*, **734**, 70
- Arun R., Mathew B., Manoj P., Ujjwal K., Kartha S. S., Viswanath G., Narang M., Paul K. T., 2019, *AJ*, **157**, 159
- Astudillo-Defru N., Delfosse X., Bonfils X., Forveille T., Lovis C., Rameau J., 2017, *A&A*, **600**, A13
- Aurière M., 2003, in Arnaud J., Meunier N., eds, *EAS Publications Series Vol. 9*, EAS Publications Series. p. 105
- Baliunas S. L., et al., 1995, *ApJ*, **438**, 269
- Ballesteros F. J., 2012, *EPL (Europhysics Letters)*, **97**, 34008
- Barnes S. A., 2003, *ApJ*, **586**, 464
- Böhm-Vitense E., 2007, *ApJ*, **657**, 486
- Bonfils X., et al., 2013, *A&A*, **549**, A109
- Boro Saikia S., Jeffers S. V., Petit P., Marsden S., Morin J., Folsom C. P., 2015, *A&A*, **573**, A17
- Boro Saikia S., et al., 2016, *A&A*, **594**, A29
- Boro Saikia S., et al., 2018a, *A&A*, **616**, A108
- Boro Saikia S., et al., 2018b, *A&A*, **620**, L11
- Brandenburg A., Mathur S., Metcalfe T. S., 2017, *ApJ*, **845**, 79
- Brown E. L., et al., 2021, *MNRAS*, **501**, 3981
- Charbonneau P., 2005, *Living Reviews in Solar Physics*, **2**, 2
- Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *ApJ*, **823**, 102
- de Laverny P., Recio-Blanco A., Worley C. C., De Pascale M., Hill V., Bijaoui A., 2013, *The Messenger*, **153**, 18
- do Nascimento J. D. J., et al., 2016, *ApJ*, **820**, L15
- Donati J. F., 2004, in Combes F., Barret D., Contini T., Meynadier F., Pagani L., eds, *SF2A-2004: Semaine de l'Astrophysique Française*. p. 217
- Donati J. F., Landstreet J. D., 2009, *ARA&A*, **47**, 333
- Donati J. F., Semel M., Carter B. D., Rees D. E., Collier Cameron A., 1997, *MNRAS*, **291**, 658
- Donati J. F., Catala C., Landstreet J. D., Petit P., 2006, in Casini R., Lites B. W., eds, *Astronomical Society of the Pacific Conference Series Vol. 358*, *Solar Polarization 4*. p. 362
- Donati J. F., et al., 2008a, *MNRAS*, **385**, 1179
- Donati J. F., et al., 2008b, *MNRAS*, **386**, 1234
- Donati J. F., et al., 2008c, *MNRAS*, **390**, 545
- Donati J. F., et al., 2010, *MNRAS*, **409**, 1347
- Donati J. F., et al., 2011a, *MNRAS*, **412**, 2454
- Donati J. F., et al., 2011b, *MNRAS*, **417**, 472
- Donati J. F., et al., 2011c, *MNRAS*, **417**, 1747
- Donati J. F., et al., 2013, *MNRAS*, **436**, 881
- Donati J. F., et al., 2014, *MNRAS*, **444**, 3220
- Donati J. F., et al., 2015, *MNRAS*, **453**, 3706
- Dotter A., 2016, *ApJS*, **222**, 8
- Duncan D. K., et al., 1991, *ApJS*, **76**, 383
- Eberhard G., Schwarzschild K., 1913, *ApJ*, **38**, 292
- Fares R., et al., 2009, *MNRAS*, **398**, 1383
- Fares R., et al., 2010, *MNRAS*, **406**, 409
- Fares R., et al., 2012, *MNRAS*, **423**, 1006
- Fares R., Moutou C., Donati J. F., Catala C., Shkolnik E. L., Jardine M. M., Cameron A. C., Deleuil M., 2013, *MNRAS*, **435**, 1451
- Finley A. J., Matt S. P., 2018, *ApJ*, **854**, 78
- Folsom C. P., et al., 2016, *MNRAS*, **457**, 580
- Folsom C. P., et al., 2018a, *MNRAS*, **474**, 4956
- Folsom C. P., et al., 2018b, *MNRAS*, **481**, 5286
- Folsom C. P., Ó Fionnagáin D., Fossati L., Vidotto A. A., Moutou C., Petit P., Dragomir D., Donati J. F., 2020, *A&A*, **633**, A48
- Foukal P., 1998, *ApJ*, **500**, 958
- Foukal P., 2018, arXiv e-prints, p. [arXiv:1810.06558](https://arxiv.org/abs/1810.06558)
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Gaia Collaboration et al., 2018, *A&A*, **616**, A1
- Gallet F., Bouvier J., 2015, *A&A*, **577**, A98
- Głębocki R., Gnaniński P., 2005, in Favata F., Hussain G. A. J., Battrick B., eds, *ESA Special Publication Vol. 560*, *13th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun*. p. 571
- Gomes da Silva J., et al., 2020, arXiv e-prints, p. [arXiv:2012.10199](https://arxiv.org/abs/2012.10199)
- Gray R. O., Corbally C. J., Garrison R. F., McFadden M. T., Bubar E. J., O'Donoghue A. A., Knox E. R., McGahee C. E., 2005, in *American Astronomical Society Meeting Abstracts*. p. 25.08
- Gregory S. G., Donati J. F., Morin J., Hussain G. A. J., Mayne N. J., Hillenbrand L. A., Jardine M., 2012, *ApJ*, **755**, 97
- Hall J. C., Lockwood G. W., Skiff B. A., 2007, *AJ*, **133**, 862
- Hall J. C., Henry G. W., Lockwood G. W., Skiff B. A., Saar S. H., 2009, *AJ*, **138**, 312
- Hébrard É. M., Donati J. F., Delfosse X., Morin J., Moutou C., Boisse I., 2016, *MNRAS*, **461**, 1465
- Henry T. J., Soderblom D. R., Donahue R. A., Baliunas S. L., 1996, *AJ*, **111**, 439
- Hill C. A., Folsom C. P., Donati J. F., Herczeg G. J., Hussain G. A. J., Alencar S. H. P., Gregory S. G., Matysse Collaboration 2019, *MNRAS*, **484**, 5810
- Høg E., et al., 2000, *A&A*, **355**, L27
- Huélamo N., et al., 2008, *A&A*, **489**, L9
- Isaacson H., Fischer D., 2010, *ApJ*, **725**, 875
- Jeffers S. V., Petit P., Marsden S. C., Morin J., Donati J. F., Folsom C. P., 2014, *A&A*, **569**, A79
- Jeffers S. V., Boro Saikia S., Barnes J. R., Petit P., Marsden S. C., Jardine M. M., Vidotto A. A., BCool Collaboration 2017, *MNRAS*, **471**, L96
- Jeffers S. V., et al., 2018, *MNRAS*, **479**, 5266
- Jenkins J. S., et al., 2011, *A&A*, **531**, A8
- Johnstone C. P., Bartel M., Güdel M., 2021, *A&A*, **649**, A96
- Kitchatinov L. L., Moss D., Sokoloff D., 2014, *MNRAS*, **442**, L1
- Kochukhov O., Makaganiuk V., Piskunov N., 2010, *A&A*, **524**, A5
- Kupka F. G., Ryabchikova T. A., Piskunov N. E., Stempels H. C., Weiss W. W., 2000, *Baltic Astronomy*, **9**, 590
- Lehmann L. T., Jardine M. M., Mackay D. H., Vidotto A. A., 2018, *MNRAS*, **478**, 4390
- Lehmann L. T., Hussain G. A. J., Vidotto A. A., Jardine M. M., Mackay D. H., 2021, *MNRAS*, **500**, 1243
- Lehtinen J., Jetsu L., Hackman T., Kajatkari P., Henry G. W., 2016, *A&A*, **588**, A38
- Lockwood G. W., Skiff B. A., Henry G. W., Henry S., Radick R. R., Baliunas S. L., Donahue R. A., Soon W., 2007, *ApJS*, **171**, 260
- Lorenzo-Oliveira D., et al., 2018, *A&A*, **619**, A73
- Lovis C., et al., 2011, arXiv e-prints, p. [arXiv:1107.5325](https://arxiv.org/abs/1107.5325)
- Marsden S. C., Donati J. F., Semel M., Petit P., Carter B. D., 2006, *MNRAS*, **370**, 468
- Marsden S. C., et al., 2014, *MNRAS*, **444**, 3517
- Mengel M. W., Fares R., Marsden S. C., Carter B. D., Jeffers S. V., Petit P., Donati J.-F., Folsom C. P., 2016, *MNRAS*, **459**, 4325–4342
- Metcalfe T. S., Egeland R., van Saders J., 2016, *ApJ*, **826**, L2
- Middelkoop F., 1982, *A&A*, **107**, 31
- Mittag M., Schmitt J. H. M. M., Schröder K. P., 2013, *A&A*, **549**, A117
- Mohanty S., Basri G., 2003, in Brown A., Harper G. M., Ayres T. R., eds, *Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun Vol.*

- 12, The Future of Cool-Star Astrophysics: 12th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun. pp 683–688
- Morgenthaler A., Petit P., Morin J., Aurière M., Dintrans B., Konstantinova-Antova R., Marsden S., 2011, *Astronomische Nachrichten*, **332**, 866
- Morin J., et al., 2008, *MNRAS*, **390**, 567
- Morin J., Donati J. F., Petit P., Delfosse X., Forveille T., Jardine M. M., 2010, *MNRAS*, **407**, 2269
- Morin J., Delfosse X., Donati J. F., Dormy E., Forveille T., Jardine M. M., Petit P., Schirmer M., 2011, in Alecian G., Belkacem K., Samadi R., Valls-Gabaud D., eds, SF2A-2011: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics. pp 503–508 ([arXiv:1208.3341](https://arxiv.org/abs/1208.3341))
- Moutou C., et al., 2007, *A&A*, **473**, 651
- Nichols-Fleming F., Blackman E. G., 2020, *MNRAS*, **491**, 2706
- Noyes R. W., Hartmann L. W., Baliunas S. L., Duncan D. K., Vaughan A. H., 1984, *ApJ*, **279**, 763
- Ó Fionnagáin D., Vidotto A. A., 2018, *MNRAS*, **476**, 2465
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, *ApJS*, **192**, 3
- Paxton B., et al., 2013, *ApJS*, **208**, 4
- Paxton B., et al., 2015, *ApJS*, **220**, 15
- Paxton B., et al., 2018, *ApJS*, **234**, 34
- Petit P., et al., 2008, *MNRAS*, **388**, 80
- Petit P., Louge T., Théado S., Paletou F., Manset N., Morin J., Marsden S. C., Jeffers S. V., 2014, *PASP*, **126**, 469
- Petrovay K., 2020, *Living Reviews in Solar Physics*, **17**, 2
- Queloz D., et al., 2001, *A&A*, **379**, 279
- Radick R. R., Lockwood G. W., Skiff B. A., Baliunas S. L., 1998, *ApJS*, **118**, 239
- Radick R. R., Lockwood G. W., Henry G. W., Hall J. C., Pevtsov A. A., 2018, *ApJ*, **855**, 75
- Reiners A., Basri G., 2008, *ApJ*, **684**, 1390
- Reiners A., et al., 2022, arXiv e-prints, [p. arXiv:2204.00342](https://arxiv.org/abs/2204.00342)
- Reinhold T., Shapiro A. I., Solanki S. K., Montet B. T., Krivova N. A., Cameron R. H., Amazo-Gómez E. M., 2020, *Science*, **368**, 518
- Rutten R. G. M., 1984, *A&A*, **130**, 353
- Saar S. H., Brandenburg A., 2002, *Astronomische Nachrichten*, **323**, 357
- Schrijver C. J., 1987, *A&A*, **172**, 111
- See V., et al., 2016, *MNRAS*, **462**, 4442
- See V., et al., 2019a, *ApJ*, **876**, 118
- See V., et al., 2019b, *ApJ*, **886**, 120
- Semel M., 1989, *A&A*, **225**, 456
- Skumanich A., 1972, *ApJ*, **171**, 565
- Soderblom D. R., 2010, *ARA&A*, **48**, 581
- Suárez Mascareño A., Rebolo R., González Hernández J. I., Esposito M., 2015, *MNRAS*, **452**, 2745
- Valenti J. A., Fischer D. A., 2005, *ApJS*, **159**, 141
- Vaughan A. H., Preston G. W., Wilson O. C., 1978, *PASP*, **90**, 267
- Villebrun F., et al., 2019, *A&A*, **622**, A72
- Waite I. A., Marsden S. C., Carter B. D., Petit P., Donati J. F., Jeffers S. V., Boro Saikia S., 2015, *MNRAS*, **449**, 8
- Waite I. A., et al., 2017, *MNRAS*, **465**, 2076
- West A. A., Hawley S. L., Bochanski J. J., Covey K. R., Reid I. N., Dhital S., Hilton E. J., Masuda M., 2008, *AJ*, **135**, 785
- Wilson O. C., 1978, *ApJ*, **226**, 379
- Wright J. T., Marcy G. W., Butler R. P., Vogt S. S., 2004, *ApJS*, **152**, 261

APPENDIX A: SIGNIFICANCE OF $\log R'_{HK}$ ACTIVITY DISTRIBUTIONS

Figure A1 shows the distributions of $\log R'_{HK}$ activity across samples of F, G, K and M stars, including both young and mature stars. On the left we show the data from our study only, and on the right we combine our data with the publicly accessible $\log R'_{HK}$ measurements of Gomes da Silva et al. (2020), and activity data compiled by Boro Saikia et al. (2018a) from the Mount Wilson




Figure A1. Histograms showing the distributions of $\log R'_{HK}$ activity across F, G, K and M stars, with error bars showing the 1σ uncertainty for each bin.

Project (Duncan et al. 1991; Baliunas et al. 1995), HARPS (Lovis et al. 2011; Bonfils et al. 2013), Lowell (Hall et al. 2009), CRIS (Wright et al. 2004; Isaacson & Fischer 2010), Magellan (Arriagada 2011) and Southern Stars (Henry et al. 1996; Gray et al. 2005) surveys. The combined sample is made up of 5179 stars. Both the Boro Saikia et al. (2018a) and Gomes da Silva et al. (2020) data include stellar B-V, which we used to convert their mean S-indices to R'_{HK} values using the method described in section 4.3.1. Gomes da Silva et al. (2020) provides stellar effective temperatures, and for the Boro Saikia et al. (2018a) data we converted B-V to T_{eff} according to Ballesteros (2012). We note that since the stellar B-V and temperature were taken from the published data tables, and were not calculated consistently with section 2.2.1, they will have some impact on the data distributions for the combined sample. The error bars in Figure A1 indicate the 1σ uncertainties in the height of each histogram bin, which we determined from the Poisson uncertainty (\sqrt{N}). They indicate that a bimodal distribution is significant with respect to the 1σ uncertainties for the combined sample of G stars. For F and K stars in the combined sample, bimodal/triple peaked activity distributions are not statistically significant.

APPENDIX B: MEAN CHROMOSPHERIC ACTIVITY, MAGNETIC FIELD STRENGTH AND ESTIMATED ROTATION PERIOD FOR G STARS

Figure B1 shows the same activity and magnetic field data as Figure 4 for G stars, but marker colour scales with the estimated rotation period, $P_{\text{est}, i=60^\circ}$. The ‘step down’ in chromospheric activity at




Figure B1. Mean chromospheric activity versus mean longitudinal magnetic field strength for G stars, where color scales with the estimated rotation period, $P_{\text{est}, i=60^\circ}$. Marker shapes/styles are the same as in Figure 1.

around $\log |B_l| \sim 0.4$ occurs as G stars reach estimated rotation periods of around 15d.

APPENDIX C: CHROMOSPHERIC ACTIVITY VARIABILITY DATA FROM GOMEZ DA SILVA ET AL. (2020)

Gomes da Silva et al. (2020) studied chromospheric activity and activity variability amplitudes for HARPS planet search stars, and we combine their data with our results in Figure C1. The method used by Gomes da Silva et al. (2020) to convert S-indices to R'_{HK} values was different compared to our study. While we use the photospheric and bolometric correction factors from Suárez Mascareño et al. (2015) due to the large B-V range of our sample, Gomes da Silva et al. (2020) derived photospheric contributions according to Noyes et al. (1984) and bolometric corrections from Rutten (1984). For Figure C1 we have adapted our R'_{HK} conversion to follow Gomes da Silva et al. (2020). Rather than showing the peak-to-peak activity variability amplitudes we present in section 5.2, Figure C1 shows the standard deviation of $R'_{HK} \times 10^5$ for each target, consistent with Gomes da Silva et al. (2020).

APPENDIX D: CHROMOSPHERIC ACTIVITY AND MAGNETIC FIELD GEOMETRY VERSUS ROSSBY NUMBER

Figure D1 compares chromospheric activity, activity variability (symbol size) and magnetic field geometry (symbol colour) to the theoretical Rossby number, taken from See et al. (2019a). F, G, K and M stars are shown together, with main sequence stars indicated by filled circles and young stars indicated by open circles. Chromospheric activity variability is reduced and the fraction of the magnetic field stored in the poloidal component becomes stronger as $\log Ro$ increases. All stars with $\log Ro \geq 1$ show dominantly poloidal magnetic field geometries.




Figure C1. Activity variability versus mean activity level for our data and the HARPS sample studied by Gomes da Silva et al. (2020). Squares indicate data from Gomes da Silva et al. (2020), while circles, stars and triangles show our data, with marker styles the same as in Figure 1. For stars common to both studies we have shown our results only. We have filtered the data to show only stars with ≥ 50 observations, and marker color scales with the number of observations.




Figure D1. Mean $\log R'_{HK}$ versus the theoretical Rossby number taken from [See et al. \(2019a\)](#), where marker colour shows the fractional poloidal field and marker size scales with chromospheric activity variability.

ORIGINAL UNEDITED MANUSCRIPT