

SYSTEMATIC REVIEW / META - ANALYSIS

What is the association between childhood adversity and subsequent chronic pain in adulthood? A systematic review

Karen P. Nicolson^{1,*}, Sarah E. E. Mills², Dhaneesha N. S. Senaratne¹, Lesley A. Colvin¹ and Blair H. Smith¹

¹Division of Population Health & Genomics, University of Dundee, Dundee, UK and ²School of Medicine, University of St Andrews, St Andrews, UK

*Corresponding author. E-mail: knicolson@dundee.ac.uk



Abstract

Background: Adverse childhood experiences and chronic pain are complex problems affecting millions of people worldwide, and result in significant healthcare utilisation. Our review aimed to determine known associations between adversity in childhood and chronic pain in adulthood.

Methods: We performed a prospectively registered systematic review (PROSPERO ID: 135625). Six electronic databases (Pubmed, Medline, Cochrane, Scopus, APA PsycNet, Web of Science) were searched from January 1, 2009 until May 30, 2022. Titles and abstracts were screened, and all original research studies examining associations between adverse childhood experiences and chronic pain in adulthood were considered for inclusion. Full texts were reviewed, and a narrative synthesis was used to identify themes from extracted data. Ten percent of studies were dual reviewed to assess inter-rater reliability. Quality assessment of study methodology was undertaken using recognised tools.

Results: Sixty-eight eligible studies describing 196 130 participants were included. Studies covered 15 different types of childhood adversity and 10 different chronic pain diagnoses. Dual reviewed papers had a Cohen's kappa reliability rating of 0.71. Most studies were of retrospective nature and of good quality. There were consistent associations between adverse childhood experiences and chronic pain in adulthood, with a 'dose'-dependent relationship. Poor mental health was found to mediate the detrimental connection between adverse childhood experiences and chronic pain.

Conclusion: A strong association was found between adverse childhood experiences and chronic pain in adulthood. Adverse childhood experiences should be considered in patient assessment, and early intervention to prevent adverse childhood experiences may help reduce the genesis of chronic pain. Further research into assessment and interventions to address adverse childhood experiences is needed.

Keywords: adverse childhood experiences; childhood adversity; childhood trauma; chronic pain; systematic review


Chronic pain and adverse childhood experiences are common and complex issues globally. Adverse events in childhood have been shown to be detrimental to adult mental and physical well-being,^{1,2} with lifelong consequences.³ Adverse childhood experiences may be defined as traumatic experiences occurring before the age of 18, and include (but are not limited to) physical/emotional/sexual abuse, physical/emotional neglect, and familial/household dysfunction (e.g. parental incarceration, parental mental health disorder,

parental separation).⁴ Adverse childhood experiences have been called a public health crisis, and it is widely accepted that they have a detrimental impact on social and health outcomes.^{5,6} In the UK, 47% of people were found to have lived through, or with, at least one type of adverse childhood experience; however, 12.3% had lived with four or more.⁷ WHO data from a Mental Health Survey of nine countries showed that the mean number of adverse childhood experiences is 2.5–2.9.^{4,8,9}

Received: 6 October 2022; Accepted: 28 March 2023

Crown Copyright © 2023 Published by Elsevier Ltd on behalf of British Journal of Anaesthesia. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

For Permissions, please email: permissions@elsevier.com



Chronic pain is a major long-term condition, affecting 19% of adults in Europe.¹⁰ In the UK, chronic pain affects between a third and a half of all adults, approximately 28 million people.¹¹ Chronic pain is pain that persists or recurs for longer than 3 months.^{12,13} Rather than being considered a symptom of other diseases, chronic pain is a distinct clinical entity and is a diagnosis in its own right, recognised in the most recent International Classification of Diseases, 11th Revision (ICD-11) classification.¹⁴ It is associated with poor general health and quality of life, and has detrimental impacts on financial stability, relationship dynamics, and healthcare utilisation; similarly, people who have experienced adverse childhood experiences have also been shown to have high levels of healthcare utilisation.^{15,16}

Previous systematic reviews have analysed chronic pain experience and previously self-reported childhood abuse, concluding that there was an increased risk of chronic pain in those who had experienced abuse and neglect.¹⁷ Other reviews have focused on a specific type of chronic pain^{18,19} or a specific type of adverse childhood experience.²⁰ This current systematic review and narrative synthesis aimed to identify and describe associations between any adverse childhood experience and any/all types of chronic pain reported in adulthood (i.e. after the age of 18).

Methods

Childhood adversity is described and classified in many ways throughout the literature. For the purposes of this review, we defined 'adverse childhood experiences' as traumatic experiences endured by children before the age of 18 yr.^{4,21,22} The term 'chronic pain' was used to capture any type of chronic pain that had been studied alongside any adverse childhood experience. Although recognising that chronic pain is a heterogeneous condition, for the purposes of this paper all forms

of chronic pain were considered together. The review was prospectively registered with PROSPERO (ID: 135625).

Search strategy

To optimise study identification, search strategies were developed by the research team with librarian support. Electronic databases (Pubmed, Medline, Cochrane, Scopus, APA PsycNet, Web of Science) were searched for papers published from 1 January 2009 until 30 May 2022. Further studies were identified by hand searching bibliographies of any relevant systematic reviews and searching tables of contents of relevant journals for articles published in the same time period. The search terms used were:

1. Child OR childhood and child/childhood abuse and/or childhood maltreatment and/or child/childhood adversity and/or childhood trauma and/or childhood neglect;
2. Adult survivors of abuse;
3. 'Chronic Pain'; 'Long-term pain'; 'Physical suffering'; 'Persistent Pain'
4. 1 AND 2 AND 3.

Inclusion/exclusion criteria

Studies were eligible for inclusion if they included at least one type of adverse childhood experience and examined possible biological, psychological, or social associations between these adversities and the subsequent presence of chronic pain in adulthood (aged 18 yr or older). No limits were applied for country of origin, type of adverse childhood experience, type of chronic pain, or study design. Articles published in languages other than English were excluded owing to lack of access to translation services. Articles for which no full-text version was available were also excluded. Review articles and any papers reporting duplicate data/studies were excluded,

Table 1 Type of childhood adversity documented in study.

Type of childhood adversity	Number of studies	Authors and years of studies
Emotional/physical/sexual abuse	33	Gerber and colleagues (2018), ²⁵ Gunduz and colleagues (2018), ²⁶ Karas and colleagues (2017), ²⁷ Riedl and colleagues (2019), ²⁸ Schrepf and colleagues (2018), ²⁹ Poli-Neto and colleagues (2018), ³⁰ Coppens and colleagues (2017), ³¹ Eriksen and colleagues (2016), ³² Brown and colleagues (2018), ³³ Generaal and colleagues (2016), ³⁴ Tietjen and colleagues (2010), ³⁵ Bohn and colleagues (2013), ³⁶ Fishbain and colleagues (2014), ³⁷ Hauser and colleagues (2019), ³⁸ De Roa and colleagues (2018), ³⁹ Prangnell and colleagues (2020), ⁴⁰ Kascakova and colleagues (2020), ⁴¹ Bayram and Erol (2014), ⁴² Tietjen and colleagues (2010), ⁴³ Macedo and colleagues (2019), ⁴⁴ Alhalal and colleagues (2018), ⁴⁵ Coppens and colleagues (2018), ⁴⁶ Achenbach and colleagues (2022), ⁴⁷ Yeung and colleagues (2016), ⁴⁸ Häuser and colleagues (2012), ⁴⁹ Liebermann and colleagues (2018), ⁵⁰ Maatta and colleagues (2019), ⁵¹ Nicolson and colleagues (2010), ⁵² Powers and colleagues (2014), ⁵³ Tesarz and colleagues (2016), ⁵⁴ Tietjen and colleagues (2010) ⁵⁵
Parental attachment/affection/bonding	1	Anno and colleagues (2015) ⁵⁶
Physical abuse	1	As-Sanie and colleagues (2014) ⁵⁷
Traumatic experience, physical abuse, and emotional abuse	1	Bottioli and colleagues (2018) ⁵⁸
Emotional, physical, and sexual abuse + Violence (Witnessing and Domestic)	2	Chiu and colleagues (2017), ⁵⁹ Santo and colleagues (2022) ⁶⁰
Adverse childhood experiences	3	Sachs-Ericsson and colleagues (2017), ⁶¹ Anda and colleagues (2010), ⁶² Fowler and colleagues (2020) ⁶³
Childhood adversities	4	Scott and colleagues (2011), ⁶⁴ Salonsalmi and colleagues (2021), ⁶⁵ Varinen and colleagues (2017), ⁶⁶ Piontek and colleagues (2021) ⁶⁷
Physical and sexual abuse, parental mental health, poverty, and family function	1	Gonzalez and colleagues (2012) ⁵⁸
Hospitalisation, surgical operations, maternal separation, parental death, family difficulties	1	Jones and colleagues (2009) ⁶⁹
Sexual abuse	1	Kamiya and colleagues (2016) ⁷⁰
Sexual abuse, punishment, and neglect	1	Johnson and colleagues (2020) ⁷¹
Physical + sexual abuse and fear of danger	1	Khandker and colleagues (2014) ⁷²
Physical and sexual abuse	4	Harris and colleagues (2018), ⁷³ Taghian and colleagues (2021), ⁷⁴ McBeth and colleagues (2015), ⁷⁵ Hart-Johnson and colleagues (2012) ⁷⁶
Serious illness before age 24 yr	1	Muthuri and colleagues (2016) ⁷⁷
Emotional/physical/sexual abuse and primary care giver attachment	2	Nacak and colleagues (2017), ⁷⁸ Waller and colleagues (2016) ⁷⁹
Emotional/physical/sexual abuse and household dysfunction	1	McKernan and colleagues (2019) ⁸⁰
Traumatic events	1	Naliboff and colleagues (2015) ⁸¹
Emotional/physical/sexual abuse and bullying	1	Nault and colleagues (2016) ⁸²
Emotional/physical abuse	1	Ortiz and colleagues (2016) ⁸³
Sexual/emotional/physical abuse and family/ household dysfunction, and parental ill health	2	Stickley and colleagues (2015), ⁸⁴ Reuchlein and colleagues (2016) ⁸⁵
Emotional abuse	1	Saariaho and colleagues (2011) ⁸⁶
Emotional/physical/sexual abuse and traumatic events	2	Schrepf and colleagues (2018), ²⁹ You and colleagues (2019) ⁸⁷
Traumatic events, road traffic accidents, natural disasters, witnessing death, child abuse	1	Smith and colleagues (2010) ⁸⁸
Bullying in childhood	1	Varinen and colleagues (2019) ⁸⁹
Perceived childhood neglect	1	Ziadni and colleagues (2020) ⁹⁰

although the referenced original source papers were reviewed for inclusion.

Screening and data extraction

From the initial longlist, study titles and abstracts were independently screened by two reviewers (KPN and SEEM). Full text review was performed by KPN, and 10% of included

studies were double-reviewed by DS to assess inter-rater reliability. Disagreements were resolved through discussion with BHS.

The following data were extracted using a standardised form: country of origin, cohort size, demography, type of chronic pain, type of childhood adversity, questionnaire type used to determine the adversity experienced, primary and




Fig 2. Number of studies describing each type of chronic pain, for studies included in the systematic review and narrative synthesis.

secondary adverse outcomes reported, association with adversity, conclusions, and key concepts.

Narrative synthesis

A narrative synthesis was used to identify common themes reflecting the associations. Two reviewers (KPN and SEEM/DS) independently identified thematic associations between childhood adversity and chronic pain. Emerging themes were combined iteratively into a narrative synthesis to summarise and describe the findings of included studies. Decisions on identification and grouping of themes were agreed by all authors.

Quality assessment

Each study's risk of bias was independently assessed by two reviewers (KPN and SEEM/DS) using standardised tools selected for each study design. Tools used for quality assessment of included papers were selected from the tools recommended by the Scottish Intercollegiate Guideline Network (SIGN).²³ SIGN is an internationally recognised organisation that produces evidence-based clinical guidelines and is accredited as part of the Guidelines International Network. AXIS was chosen as the best available tool for cross-sectional studies, as SIGN does not provide a tool for this study design. The quality assessment tools used in this analysis were: cross-sectional studies – AXIS appraisal tool²⁴; cohort studies – SIGN methodology checklist tool 3²³; case-control studies – SIGN methodology checklist tool 4.²³

Results

Overview of results

We identified 2904 records from the search strategy, and 33 records were identified through other sources (Fig 1). Titles, abstracts, and full-text papers were screened, identifying 68 studies for inclusion with data from 196 130 participants. There were 50 cross-sectional studies, 14 case-control studies, and four cohort studies: 10% of studies were double-reviewed for inter-rater reliability with Cohen's kappa of 0.71. Demographic information from all included studies is shown in Supplementary Table S1.

The combination of emotional/physical/sexual abuse was the most commonly studied adverse childhood experience, followed by general 'adverse childhood experiences', and 'physical and sexual abuse' as shown in Table 1. Of the 10 types of chronic pain identified in included studies as being associated with adverse childhood experiences, generalised chronic pain, fibromyalgia, and chronic pelvic pain were the most common (Fig 2). Supplementary Table S2 demonstrates the impact of childhood adversity findings from each study and the recommended outcomes. The main themes identified in this narrative synthesis, with the corresponding evidence sources, are summarised in Table 2.

Quality assessment

Sixty-one of the 68 included studies were assessed as high quality. Seven papers were assessed as having moderate risk

Table 2 Summary of main themes.

Main findings from study	Study reference	Number of studies
Greater number of adversities are associated with worse pain experience.	25–35,57,59,61,62,64,65,73,87,36,39,56,58,66,69,72 41–43,46,80,88,90,84,81,82,91,63,68,76,70,50,83,85,55,92	46
Current poor mental health mediates association between childhood adversity and pain.	30,31,72,42,45,93,50,53	8
In those with chronic pain, previous childhood adversity should be assessed.	26,28,29,57,61,71,39,74,60,41,42,67,45 84,91,82,49,76,93,52,85,92	23
Screening for mental health should be considered for those with chronic pain.	57,31,39,41,42,67,45,80,82,93,70	9
Education on the awareness of childhood adversity on pain needs to be more widespread.	42,53	2
Early intervention is key to prevent childhood adversities from occurring.	28,65,66,89,94,68	6

of bias. Elements of weakness that were identified in quality analysis included small sample size (11 studies had fewer than 100 participants); limited demographic information; exact age of participants was missing (13 studies); and further details on breakdown of gender was not provided (seven studies).

Most papers were consistent in noting recall bias as a potential weakness. The main confounder documented was mental health in the form of anxiety, depression, current stress, and post-traumatic stress disorder (PTSD). In cross-sectional studies, a common weakness was lack of information about non-responders. One paper lacked analysis to determine the statistical significance of associations reported; however, it did not add any factors that had not been identified by other studies included in the review.²⁵ Further details of quality assessment are available in [Appendix 1](#).

Narrative synthesis

Adverse childhood experiences were shown to increase the risk of subsequent chronic pain as an adult and in one study, 75% of participants with chronic pain had experienced early life adversity.²⁶ Childhood violence was associated with multiple physical diseases, of which chronic pain was the most prevalent.^{27,28,73} Adverse childhood experiences were shown to lead to greater dysfunction and disability for those with chronic pain when compared with healthy controls, or those with pain but no adversity.^{29,57} The more adversity experienced in childhood, the more likely an adult was to experience chronic pain, and the greater the chance that the chronic pain would not improve^{29,30,31,64,87} adding to the chronicity of pain.⁵⁹ Specific adverse childhood experiences in the form of parental loss early in life, along with verbal and sexual abuse were independently associated with the subsequent onset of painful conditions. Adverse childhood experiences led to a higher risk of developing mental health and painful conditions.^{54,61} Those who experienced adverse childhood experiences were shown to have more sites of chronic pain and pain of greater severity than those who did not experience such adversity.^{32–35,62,65} However, although stress that is experienced in childhood is associated with increased risk of developing chronic pain in adulthood, stress that is experienced in adulthood does not show the same correlation with developing chronic pain.⁷¹ Those with chronic pain who experienced adverse childhood experiences had greater sensitivity and a lower tolerance to pain than those without a history of adverse childhood experiences.⁸⁷ ‘Catastrophising

and hypervigilance to somatic symptoms in chronic pain were also found to be related to previous adverse childhood experiences.^{36,95} Two studies did not find a connection between adverse childhood experiences and chronic pain; however, each of these studies concentrated on only one form of adversity and type of pain.^{37,38}

An insecure environment where children were emotionally deprived was also related to chronic pain development. Those who experienced emotional neglect, affectionless parenting, parental loss, or overprotective parenting in childhood were more likely to experience chronic pain.^{30,39,56,86} A combination of adverse childhood experiences, ‘alexithymia’ (an inability to determine the cause of feelings in your body) and ongoing life stress may lead to chronic pain.^{58,78} Somatoform pain disorder was also linked to previous emotional neglect, poor awareness of body signals in the context of depression, and poor attachment to primary caregiver.⁸⁷ Emotional abuse was related to higher pain sensitisation and was found to be associated with chronic pain in people who injected drugs.⁴⁰ In the population of people who inject drugs, physical and sexual abuse led to doubling of risk of developing chronic pain in adulthood. In those with substance misuse, pain severity was greater. ‘Catastrophisation’ and detrimental pain interference in life was also more prevalent for those who had experienced physical and sexual abuse when with those without these particular adversities.⁷⁴ In those with chronic pain, a larger number of adverse childhood experiences were found to lead to a greater risk of substance misuse and mental health disorders compared with those with fewer adverse childhood experiences.⁶⁰ Living in a house with financial strain; experiencing parental divorce, parental mental ill health, or alcohol problems; and/or living in fear of a family member were associated with a greater chance of developing chronic pain in adulthood compared with those with no adversity.⁶⁶ Poor social situations, physically traumatic events such as accidents, and living in fear during childhood were also associated with chronic pain.^{69,72} Experiencing bullying as a child may be related to chronic pain, either directly or mediated by depression.^{33,87}

Current poor mental health as an adult is thought to mediate the development and severity of chronic pain in the presence of childhood adversity,⁴¹ particularly with depression and anxiety.^{39,42,43,67} Borderline personality disorder and childhood adversity were also associated with chronic pain; childhood adversity was associated with experiencing pain and particular pain syndromes such as fibromyalgia in those with borderline personality disorder.⁷¹ However, even after

controlling for mental health, the dose-response relationship between adverse childhood experiences and subsequent chronic pain persisted.⁶¹ Depression can worsen an experience of pain,⁶² and those with depression and pain were found to have experienced more emotional neglect than those who had depression and no chronic pain.^{27,44} Anxiety may be caused by adverse childhood experiences which in turn contributes to chronic pain prevalence.⁹⁴ PTSD was found to play a part in chronic pain experience^{45,80} causing a lower level of functioning and higher levels of pain when adverse childhood experiences had been experienced. Perceived stress and injustice as an adult have been found to mediate the relationship between chronic pain and adverse childhood experiences.^{88,90}

A few studies explored potential biological mechanisms driving the identified associations. The direct physiological changes caused by adverse childhood experiences that influence chronic pain are still debated. Hypotheses include disruption in cortisol regulation or that maternal neglect causes neurobiological changes.^{31,42} Irregularities of the hypothalamic pituitary axis have also been postulated, whereas leptin and endocannabinoid dysfunction after adversity may be related.^{46,47} An association was also found connecting serious illness, social adversity, and physical trauma before the age of 25 to chronic widespread pain, more than chronic regional pain, raising the possibility that different mechanisms may underlie different types of chronic pain. Central sensitisation leading to pain sensitivity and longevity has also been postulated.⁸⁷

Discussion

Summary

Studies in this review consistently documented high levels of adverse childhood experiences in adults with chronic pain. Many studies used a pre-existing formal chronic pain diagnosis as an inclusion criterion, rather than asking specifically about any pain lasting longer than 3 months. A single type of adversity was rarely found in isolation; where there was one form of adversity, there were usually others.^{57,56,84} This review shows that adverse childhood experiences, in its many forms, can have a detrimental impact in the form, presence, severity, and extent of chronic pain, the greater number of adversities experience the worse this effect is.

Strengths and limitations

This is the first comprehensive systematic review to describe known associations between any form of adverse childhood experience and any type of adult chronic pain. Strengths include detailed analysis of a wide range of studies from many different countries covering a wide range of adverse childhood experiences and types of chronic pain. Limitations include exclusion of papers not available in full or in English (for reasons of resource), which may produce a cultural and geographical bias; however, only four papers were excluded for this reason, so the impact of this limitation is likely to have been minimal. Further limitations include terminology for adverse childhood experiences; the heterogeneity of adversity a child can experience and consequent range of descriptive terms, means studies may have been missed. We used a broad search strategy across multiple databases, and extensive hand-searching to supplement the initial search strategy, so we hope to have minimised the risk of missing relevant studies. However, given the heterogeneity of terminology in this field, we

cannot be certain that a small number of relevant studies were not missed. The majority of studies were retrospective; therefore, recall bias was also noted. This is a limitation of the methods used to address the research questions. Most of the studies included were of high quality, with only six noted to be of medium quality and two of poor quality. Any studies of poorer methodological quality were included in the final synthesis for completeness, and although their conclusions were given less emphasis in the final synthesis, their findings were consistent with higher quality studies (see Table 2). The complexity and heterogeneity of data meant that we opted for a narrative analysis. If consistent definitions and measurements of adverse childhood experiences can be developed, then future work could involve substantial meta-analysis.

Comparison with existing literature

The more types of adverse childhood experiences, the greater the detrimental impact on development and experience of pain in adult life. However, much of the published research has a relatively narrow focus on emotional, physical, and sexual abuse. In previous systematic reviews, fibromyalgia has been shown to be associated with prior psychological or physical trauma.¹⁹ A meta-analysis examining only two types of adversity (physical or sexual abuse) found that those who recalled adverse childhood experiences were at greater risk of chronic pain as an adult when compared with those who had not reported experiencing such events in their childhood.¹⁷ In addition, a previous meta-analysis found that sexual abuse was linked to non-specific chronic pain and chronic pelvic pain.⁹⁶ Poor mental health may be a contributory factor to chronic pain, and the negative impact of childhood adversity on mental health is already documented.^{97,96} This current systematic review examines associations with a much broader range of adversities such as household dysfunction, experiencing bullying, generalised living in fear, traumatic events, and poor parental attachment, all of which were found to have adverse associations with chronic pain in adulthood.^{29,64,59,61,62,65,56,58,78,66,69,89,88,84,63,77,81,82,91} Studies focusing on adolescents with chronic pain have documented a dose response: the more adverse childhood experience a child experienced the more chronic pain and in a population of adolescents with pain the more adverse childhood experience the greater the mental health impact of anxiety and depression as well of fear of pain itself. These findings are consistent with those from this review of adult chronic pain.^{98,99}

Implications for research and practice

Despite the high prevalence of adverse childhood experiences documented in the included studies, not every person with chronic pain has experienced an adverse childhood experience, or vice versa. There was general agreement about the need for more prospective studies to determine the nature and extent of the links between adverse childhood experiences and chronic pain,^{20,100} and the reasons behind these. Data linkage, where adverse childhood experiences were previously recorded, or prospective cohort studies will provide more evidence.^{101,102} Most studies examined emotional, physical, and sexual abuse, but other traumatic events also need to be studied in detail. Emotional abuse was also noted to be both common and overlooked, requiring further research. This review demonstrates variability in acquisition of information on

adverse childhood experiences. This review highlights the need to enquire about the possibility of adverse childhood experiences in those with chronic pain, and enquiring about current mental health symptoms. Understanding the potential for a history of adversity in people with chronic pain is an opportunity to consider management options.^{31,36,48} Further research into how this information is obtained should be undertaken to allow for consistency of approach. A history of childhood adversity should be considered in the assessment of chronic pain so that relevant management options can be determined (although further research into effective interventions is also required). Enquiring about childhood adversity has already been shown to be acceptable, although the sensitivity of this subject must always be respected and psychological safety maintained.¹⁰³ Currently, National Institute for Health and Care Excellence in the UK do not recommend routinely assessing for adverse childhood experiences in chronic pain.¹⁰³ Scottish Intercollegiate National Guidelines on management of chronic pain suggest a biopsychological assessment but the consistency of enquiring about adverse childhood experiences within clinical practice is unknown.^{104,105} An adverse childhood experience assessment¹⁰⁶ has been recommended during a chronic pain assessment.

Conclusions

This review identified that a wide range of adverse childhood experiences are associated with the development, severity, and impact of chronic pain in adulthood, and that this association with adverse childhood experiences is consistent across a wide range of chronic pain conditions. The importance of adverse childhood experiences when assessing and managing chronic pain needs to be recognised and integrated into clinical practice. Further research on the underlying mechanisms of these associations is required, along with the assessment and management of childhood adversity in the context of chronic pain.

Authors' contributions

Planning and development of the review: KN, SEEM, LAC, BHS.
Initial plan for the review: KN.
Conception and design: KN, SEEM, DS.
Acquisition of data: KN, SEEM, DS.
Analysis and interpretation of data: KN, SEEM, DS.
Selection of tools for quality assessment: KN, SEEM.
Quality assessment: KN, SEEM, DS.
Identification of themes from included papers for narrative synthesis: KN, SEEM, DS.
Writing of the first draft of this paper: KN.
Overall research supervisors: LAC, BHS.
Development, thematic synthesis, interpretation, and overall conclusions: LAC, BHS.
Writing of the final version of the paper: KN, SEEM, DS, LAC, BHS.
Final approval of the version to be published: KN, SEEM, DS, LAC, BHS.

Acknowledgements

We thank Mr Scott Macgregor, librarian, at The University of Dundee for his assistance.

Declarations of interest

LAC was past editor of *British Journal of Anaesthesia* (2010–21), and is currently a member of the editorial board of the *BJA* and *BJA Open*. BHS was Scottish Government Lead Clinician for Chronic Pain (2014–2021) and is Speciality Adviser (Chronic Pain) to the Chief Medical Officer of Scotland. The authors have no other competing interests to declare.

Funding

NHS Education for Scotland. KN was academic fellow from 2018 to 2022. SEEM was funded through an academic clinical fellowship from the Scottish Government's Chief Scientist Office (CSO grant number CAF_17_06). DS is a fellow on the Multimorbidity Doctoral Training Programme for Health Professionals, which is supported by the Wellcome Trust [grant number 223499/Z/21/Z].

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bjao.2023.100139>.

References

- Ehlert U. Enduring psychobiological effects of childhood adversity. *Psychoneuroendocrinology* 2013; **38**: 1850–7
- Gilbert LK, Breiding MJ, Merrick MT, et al. Childhood adversity and adult chronic disease: an update from ten states and the District of Columbia, 2010. *Am J Prev Med* 2015; **48**: 345–9
- World Health Organisation. Global status report on violence prevention 2014. Available from, www.who.int/publications/item/9789241564793. [Accessed 13 January 2023]
- Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the Adverse Childhood Experiences (ACE) Study. *Am J Prev Med* 1998; **14**: 245–58
- Dube SR. Continuing conversations about adverse childhood experiences (ACEs) screening: a public health perspective. *Child Abuse Negl* 2018; **85**: 180–4
- Pearce J, Murray C, Larkin W. Childhood adversity and trauma: experiences of professionals trained to routinely enquire about childhood adversity. *Heliyon* 2019; **5**, e01900
- Bellis MA, Lowey H, Leckenby N, Hughes K, Harrison D. Adverse childhood experiences: retrospective study to determine their impact on adult health behaviours and health outcomes in a UK population. *J Public Health (Oxf)* 2014; **36**: 81–91
- Felitti VJ. Health appraisal and the adverse childhood experiences study: national implications for health care, cost, and utilization. *Perm J* 2019; **23**
- Kessler RC, McLaughlin KA, Green JG, et al. Childhood adversities and adult psychopathology in the WHO world mental health surveys. *Br J Psychiatry* 2010; **197**: 378–85
- Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain* 2006; **10**: 287–333

11. Fayaz A, Croft P, Langford RM, Donaldson LJ, Jones GT. Prevalence of chronic pain in the UK: a systematic review and meta-analysis of population studies. *BMJ Open* 2016; **6**, e010364
12. Treede RD, Rief W, Barke A, et al. Chronic pain as a symptom or a disease: the IASP classification of chronic pain for the international classification of diseases (ICD-11). *Pain* 2019; **160**: 19–27
13. Treede RD, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain* 2015; **156**: 1003–7
14. Nicholas M, Vlaeyen JW, Rief W, et al. The IASP classification of chronic pain for ICD-11: chronic primary pain. *Pain* 2019; **160**: 28–37
15. Bellis M, Hughes K, Hardcastle K, et al. The impact of adverse childhood experiences on health service use across the life course using a retrospective cohort study. *J Health Serv Res Policy* 2017; **22**: 168–77
16. Andersson HI, Ejlertsson G, Leden I, Scherstén B. Impact of chronic pain on health care seeking, self care, and medication. Results from a population-based Swedish study. *J Epidemiol Community Health* 1999; **53**: 503–9
17. Davis DA, Luecken LJ, Zautra AJ. Are reports of childhood abuse related to the experience of chronic pain in adulthood? A meta-analytic review of the literature. *Clin J Pain* 2005; **21**: 398–405
18. Yavne Y, Amital D, Watad A, Tiosano S, Amital H. A systematic review of precipitating physical and psychological traumatic events in the development of fibromyalgia. *Semin Arthritis Rheum* 2018; **48**: 121–33
19. Hauser W, Kosseva M, Uceyler N, Klose P, Sommer C. Emotional, physical, and sexual abuse in fibromyalgia syndrome: a systematic review with meta-analysis. *Arthritis Care Res* 2011; **63**: 808–20
20. Paras ML, Murad MH, Chen LP, et al. Sexual abuse and lifetime diagnosis of somatic disorders: a systematic review and meta-analysis. *JAMA* 2009; **302**: 550–61
21. Hughes K, Bellis MA, Hardcastle KA, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2017; **2**, e356–e66
22. Grummitt LR, Kreski NT, Kim SG, Platt J, Keyes KM, McLaughlin KA. Association of childhood adversity with morbidity and mortality in US adults: a systematic review. *JAMA Pediatrics* 2021; **175**: 1269–78
23. SIGN. <https://www.sign.ac.uk/what-we-do/methodology/checklists/>
24. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical appraisal tool to assess the quality of cross-sectional studies (AXIS). *BMJ Open* 2016; **6**, e011458
25. Gerber MR, Bogdan KM, Haskell SG, Scioli ER. Experience of childhood abuse and military sexual trauma among women veterans with fibromyalgia. *J Gen Intern Med* 2018; **33**: 2030–1
26. Gunduz N, Polat A, Erzincan E, Turan H, Sade I, Tural U. Psychiatric comorbidity and childhood trauma in fibromyalgia syndrome. *Turk J Phys Med Rehabil* 2018; **64**: 91–9
27. Karas H, Yildirim EA, Kucukgoncu S, Yakut U. The relationship of childhood trauma, dissociative experiences and depression with pain in female patients with fibromyalgia: a cross-sectional study. *Dusunen Adam* 2017; **30**: 86–94
28. Riedl D, Beck T, Exenberger S, et al. Violence from childhood to adulthood: the influence of child victimization and domestic violence on physical health in later life. *J Psychosom Res* 2019; **116**: 68–74
29. Schrepf A, Naliboff B, Williams DA, et al. Adverse childhood experiences and symptoms of urologic chronic pelvic pain syndrome: a multidisciplinary approach to the study of chronic pelvic pain research network study. *Ann Behav Med* 2018; **52**: 865–77
30. Poli-Neto OB, Tawasha KAS, Romao A, et al. History of childhood maltreatment and symptoms of anxiety and depression in women with chronic pelvic pain. *J Psychosom Obstet Gynaecol* 2018; **39**: 83–9
31. Coppens E, Van Wambeke P, Morlion B, et al. Prevalence and impact of childhood adversities and post-traumatic stress disorder in women with fibromyalgia and chronic widespread pain. *Eur J Pain* 2017; **21**: 1582–90
32. Eriksen AM, Schei B, Hansen KL, Sorlie T, Fleten N, Javo C. Childhood violence and adult chronic pain among indigenous Sami and non-Sami populations in Norway: a SAMINOR 2 questionnaire study. *Int J Circumpolar Health* 2016; **75**, 32798
33. Brown RC, Plener PL, Braehler E, Fegert JM, Huber-Lang M. Associations of adverse childhood experiences and bullying on physical pain in the general population of Germany. *J Pain Res* 2018; **11**: 3099–108
34. Generaal E, Milaneschi Y, Jansen R, Elzinga BM, Dekker J, Penninx B. The brain-derived neurotrophic factor pathway, life stress, and chronic multi-site musculoskeletal pain. *Mol Pain* 2016; **12**
35. Tietjen GE, Brandes JL, Peterlin BL, et al. Childhood maltreatment and migraine: Part II. Emotional abuse as a risk factor for headache chronification. *Headache* 2010; **50**: 32–41
36. Bohn D, Bernardy K, Wolfe F, Häuser W. The association among childhood maltreatment, somatic symptom intensity, depression, and somatoform dissociative symptoms in patients with fibromyalgia syndrome: a single-center cohort study. *J Trauma Dissociation* 2013; **14**: 342–58
37. Fishbain DA, Bruns D, Meyer LJ, Lewis JE, Gao J, Disorbio JM. Exploration of affirmation of childhood molestation (sexual abuse) in chronic pain patients, acute pain patients, community patients with pain and community nonpatients without pain. *Pain Pract* 2014; **14**: 515–25
38. Hauser W, Brahler E, Schmutzler G, Glaesmer H. The association of adverse childhood experiences and of resilience with chronic noncancer pain in the German adult population — a cross-sectional survey. *Eur J Pain* 2019; **23**: 555–64
39. De Roa P, Paris P, Poindessous JL, Maillet O, Heron A. Subjective experiences and sensitivities in women with fibromyalgia: a quantitative and comparative study. *Pain Res Manag* 2018; **2018**, 8269564
40. Prangnell A, Shoveller J, Voon P, et al. The impact of childhood emotional abuse on pain interference among people with chronic pain who inject drugs in Vancouver, Canada. *Pain Med* 2020; **21**: 704–13
41. Kascakova N, Furstova J, Hasto J, Madarasova-Geckova A, Tavel P. When a head is about to burst: Attachment mediates the relationship between childhood trauma and migraine. *Int J Environ Res Public Health* 2020; **17**: 4579
42. Bayram K, Erol A. Childhood traumatic experiences, anxiety, and depression levels in fibromyalgia and rheumatoid arthritis. *Noro Psikiyatri Arsivi* 2014; **51**: 344–9

43. Tietjen GE, Brandes JL, Peterlin BL, et al. Childhood maltreatment and migraine (part I). Prevalence and adult revictimization: a multicenter headache clinic survey. *Headache* 2010; **50**: 20–31
44. Macedo BBD, von Werne Baes C, Menezes IC, Juruena MF. Child abuse and neglect as risk factors for comorbidity between depression and chronic pain in adulthood. *J Nerv Ment Dis* 2019; **207**: 538–45
45. Alhalal E, Ford-Gilboe M, Wong C, AlBuhairan F. Factors mediating the impacts of child abuse and intimate partner violence on chronic pain: a cross-sectional study. *BMC Womens Health* 2018; **18**: 160
46. Coppens E, Kempke S, Van Wambeke P, et al. Cortisol and subjective stress responses to acute psychosocial stress in fibromyalgia patients and control participants. *Psychosom Med* 2018; **80**: 317–26
47. Achenbach J, Volkmann L, Tran A-T, et al. Neurohumoral profiles and childhood adversity of patients with multisomatoform disorder and pain as the leading bodily symptom. *Dis Markers* 2022; **2022**, 7958375
48. Yeung EW, Davis MC, Ciaramitaro MC. Cortisol profile mediates the relation between childhood neglect and pain and emotional symptoms among patients with fibromyalgia. *Ann Behav Med* 2016; **50**: 87–97
49. Häuser W, Bohn D, Kühn-Becker H, Erdkönig R, Brähler E, Glaesmer H. Is the association of self-reported childhood maltreatments and adult fibromyalgia syndrome attributable to depression? A case control study. *Clin Exp Rheumatol* 2012; **30**: 59–64
50. Liebermann C, Kohl Schwartz AS, Charpidou T, et al. Maltreatment during childhood: a risk factor for the development of endometriosis? *Hum Reprod* 2018; **33**: 1449–58
51. Maatta J, Martikainen A, Pakarinen M, et al. High level of childhood trauma predicts a poor response to spinal cord stimulation in chronic neuropathic pain. *Pain Physician* 2019; **22**: E37–44
52. Nicolson NA, Davis MC, Kruszewski D, Zautra AJ. Childhood maltreatment and diurnal cortisol patterns in women with chronic pain. *Psychosom Med* 2010; **72**: 471–80
53. Powers A, Fani N, Pallos A, Stevens J, Ressler KJ, Bradley B. Childhood abuse and the experience of pain in adulthood: the mediating effects of PTSD and emotion dysregulation on pain levels and pain-related functional impairment. *Psychosomatics* 2014; **55**: 491–9
54. Tesarz J, Eich W, Treede RD, Gerhardt A. Altered pressure pain thresholds and increased wind-up in adult patients with chronic back pain with a history of childhood maltreatment: a quantitative sensory testing study. *Pain* 2016; **157**: 1799–809
55. Tietjen GE, Brandes JL, Peterlin BL, et al. Childhood maltreatment and migraine: Part III. Association with comorbid pain conditions. *Headache* 2010; **50**: 42–51
56. Anno K, Shibata M, Ninomiya T, et al. Paternal and maternal bonding styles in childhood are associated with the prevalence of chronic pain in a general adult population: the Hisayama Study. *BMC Psychiatry* 2015; **15**: 181
57. As-Sanie S, Clevenger LA, Geisser ME, Williams DA, Roth RS. History of abuse and its relationship to pain experience and depression in women with chronic pelvic pain. *Am J Obstet Gynecol* 2014; **210**, 317.e1–8
58. Bottiroli S, Galli F, Viana M, Sances G, Tassorelli C. Traumatic experiences, stressful events, and alexithymia in chronic migraine with medication overuse. *Front Psychol* 2018; **9**: 704
59. Chiu CD, Lee MH, Chen WC, Ho HL, Wu HC. Childhood trauma perpetrated by close others, psychiatric dysfunction, and urological symptoms in patients with interstitial cystitis/bladder pain syndrome. *J Psychosom Res* 2017; **93**: 90–5
60. Santo Jr T, Campbell G, Gisev N, Degenhardt L. Exposure to childhood trauma increases risk of opioid use disorder among people prescribed opioids for chronic non-cancer pain. *Drug Alcohol Depend* 2022; **230**, 109199
61. Sachs-Ericsson NJ, Sheffler JL, Stanley IH, Piazza JR, Preacher KJ. When emotional pain becomes physical: adverse childhood experiences, pain, and the role of mood and anxiety disorders. *J Clin Psychol* 2017; **73**: 1403–28
62. Anda R, Tietjen G, Schulman E, Felitti V, Croft J. Adverse childhood experiences and frequent headaches in adults. *Headache* 2010; **50**: 1473–81
63. Fowler C, Homandberg L, Steele C, et al. Adult correlates of adverse childhood experiences in Ukraine. *Child Abuse Negl* 2020; **107**, 104617
64. Scott KM, Von Korff M, Angermeyer MC, et al. Association of childhood adversities and early-onset mental disorders with adult-onset chronic physical conditions. *Arch Gen Psychiatry* 2011; **68**: 838–44
65. Salonsalmi A, Pietiläinen O, Lahelma E, Rahkonen O, Lallukka T. Contributions of childhood adversities to chronic pain among mid-life employees. *Scand J Public Health* 2021; **50**: 333–9
66. Varinen A, Kosunen E, Mattila K, Koskela T, Sumanen M. The relationship between childhood adversities and fibromyalgia in the general population. *J Psychosom Res* 2017; **99**: 137–42
67. Piontek K, Apfelbacher C, Ketels G, Brünahl C, Löwe B. Depression partially mediates the association of adverse childhood experiences with pain intensity in patients with chronic pelvic pain syndrome: results from a cross-sectional patient survey. *Pain Med* 2021; **22**: 1174–84
68. Gonzalez A, Boyle MH, Kyu HH, Georgiades K, Duncan L, MacMillan HL. Childhood and family influences on depression, chronic physical conditions, and their comorbidity: findings from the Ontario Child Health Study. *J Psychiatr Res* 2012; **46**: 1475–82
69. Jones GT, Power C, Macfarlane GJ. Adverse events in childhood and chronic widespread pain in adult life: results from the 1958 British Birth Cohort Study. *Pain* 2009; **143**: 92–6
70. Kamiya Y, Timonen V, Kenny RA. The impact of childhood sexual abuse on the mental and physical health, and healthcare utilization of older adults. *Int Psychogeriatr* 2016; **28**: 415–22
71. Johnson BN, Lumley MA, Cheavens JS, McKernan LC. Exploring the links among borderline personality disorder symptoms, trauma, and pain in patients with chronic pain disorders. *J Psychosom Res* 2020; **135**, 110164

72. Khandker M, Brady SS, Stewart EG, Harlow BL. Is chronic stress during childhood associated with adult-onset vulvodynia? *J Womens Health (Larchmt)* 2014; **23**: 649–56
73. Harris HR, Wieser F, Vitonis AF, et al. Early life abuse and risk of endometriosis. *Hum Reprod* 2018; **33**: 1657–68
74. Taghian NR, McHugh RK, Griffin ML, Chase AR, Greenfield SF, Weiss RD. Associations between childhood abuse and chronic pain in adults with substance use disorders. *Subst Use Misuse* 2021; **56**: 87–92
75. McBeth J, Tomenson B, Chew-Graham CA, et al. Common and unique associated factors for medically unexplained chronic widespread pain and chronic fatigue. *J Psychosom Res* 2015; **79**: 484–91
76. Hart-Johnson T, Green CR. The impact of sexual or physical abuse history on pain-related outcomes among blacks and whites with chronic pain: gender influence. *Pain Med* 2012; **13**: 229–42
77. Muthuri SG, Kuh D, Bendayan R, Macfarlane GJ, Cooper R. Chronic physical illness in early life and risk of chronic widespread and regional pain at age 68: evidence from the 1946 British birth cohort. *Pain* 2016; **157**: 2382–9
78. Nacak Y, Morawa E, Tuffner D, Erim Y. Insecure attachment style and cumulative traumatic life events in patients with somatoform pain disorder: a cross-sectional study. *J Psychosom Res* 2017; **103**: 77–82
79. Waller E, Scheidt CE, Endorf K, Hartmann A, Zimmermann P. Unresolved trauma in fibromyalgia: a cross-sectional study. *J Health Psychol* 2016; **21**: 2457–65
80. McKernan LC, Johnson BN, Reynolds WS, et al. Post-traumatic stress disorder in interstitial cystitis/bladder pain syndrome: relationship to patient phenotype and clinical practice implications. *Neurorol Urodyn* 2019; **38**: 353–62
81. Naliboff BD, Stephens AJ, Afari N, et al. Widespread psychosocial difficulties in men and women with urologic chronic pelvic pain syndromes: case-control findings from the multidisciplinary approach to the Study of Chronic Pelvic Pain Research Network. *Urology* 2015; **85**: 1319–27
82. Nault T, Gupta P, Ehler M, et al. Does a history of bullying and abuse predict lower urinary tract symptoms, chronic pain, and sexual dysfunction? *Int Urol Nephrol* 2016; **48**: 1783–8
83. Ortiz R, Ballard ED, Machado-Vieira R, Saligan LN, Walitt B. Quantifying the influence of child abuse history on the cardinal symptoms of fibromyalgia. *Clin Exp Rheumatol* 2016; **34**: S59–66
84. Stickley A, Koyanagi A, Kawakami N. Childhood adversities and adult-onset chronic pain: results from the World Mental Health Survey, Japan. *Eur J Pain* 2015; **19**: 1418–27
85. Reuchlein B, Henn L, Brian T, Schier K, Hardt J. Childhood adversities and adult headache in Poland and Germany. *PLoS One* 2016; **11**, e0148162
86. Saariaho TH, Saariaho AS, Karila IA, Joukamaa MI. Early maladaptive schemas in Finnish adult chronic pain patients and a control sample. *Scand J Psychol* 2011; **52**: 146–53
87. You DS, Albu S, Lisenbardt H, Meagher MW. Cumulative childhood adversity as a risk factor for common chronic pain conditions in young adults. *Pain Med* 2019; **20**: 486–94
88. Smith BW, Papp ZZ, Tooley EM, Montague EQ, Robinson AE, Cosper CJ. Traumatic events, perceived stress and health in women with fibromyalgia and healthy controls. *Stress Health* 2010; **26**: 83–93
89. Varinen A, Kosunen E, Mattila K, Suominen S, Sillanmaki L, Sumanen M. The association between bullying victimization in childhood and fibromyalgia. Data from the nationwide Finnish health and social support (HeSSup) study based on a sample of 64,797 individuals. *J Psychosom Res* 2019; **117**: 48–53
90. Ziadni MS, You DS, Sturgeon JA, Mackey SC, Darnall BD. Perceived injustice mediates the relationship between perceived childhood neglect and current function in patients with chronic pain: a preliminary pilot study. *J Clin Psychol Med Settings* 2021; **28**: 349–60
91. Etner S, Stingl K, Schlegel AK, Wichmann M, Nickenig A. Biopsychosocial correlations in patients with chronic oro-facial pain: Part II. Experiences of pain and dramatic events before the 16th year of life. *J Oral Rehabil* 2009; **36**: 408–14
92. Prangnell A, Voon P, Shulha H, et al. The relationship between childhood emotional abuse and chronic pain among people who inject drugs in Vancouver, Canada. *Child Abuse Negl* 2019; **93**: 119–27
93. Häuser W, Hoffmann E-M, Wolfe F, et al. Self-reported childhood maltreatment, lifelong traumatic events and mental disorders in fibromyalgia syndrome: a comparison of US and German outpatients. *Clin Exp Rheumatol* 2015; **33**: S86
94. Kascakova N, Furstova J, Hasto J, Madarasova Geckova A, Tavel P. The unholy trinity: childhood trauma, adulthood anxiety, and long-term pain. *Int J Environ Res Public Health* 2020; **17**: 414
95. Crombez G, De Paepe AL, Veirman E, Eccleston C, Verleyen G, Van Ryckeghem DML. Let's talk about pain catastrophizing measures: an item content analysis. *PeerJ* 2020; **8**, e8643
96. Raposo SM, Mackenzie CS, Henriksen CA, Afifi TO. Time does not heal all wounds: older adults who experienced childhood adversities have higher odds of mood, anxiety, and personality disorders. *Am J Geriatr Psychiatry* 2014; **22**: 1241–50
97. Kalmakis KA, Chandler GE. Health consequences of adverse childhood experiences: a systematic review. *J Am Assoc Nurse Pract* 2015; **27**: 457–65
98. Groenewald CB, Murray CB, Palermo TM. Adverse childhood experiences and chronic pain among children and adolescents in the United States. *Pain Rep* 2020; **5**: e839
99. Nelson S, Simons LE, Logan D. The incidence of adverse childhood experiences (ACEs) and their association with pain-related and psychosocial impairment in youth with chronic pain. *Clin J Pain* 2018; **34**: 402–8
100. Marin TJ, Lewinson RE, Hayden JA, et al. A systematic review of the prospective relationship between child maltreatment and chronic pain. *Children (Basel)* 2021; **8**: 806
101. Walker SM, Melbourne A, O'Reilly H, et al. Somatosensory function and pain in extremely preterm young adults from the UK EPICure cohort: sex-dependent differences and impact of neonatal surgery. *Br J Anaesth* 2018; **121**: 623–35

102. Boyd A, Golding J, Macleod J, et al. Cohort profile: the 'children of the 90s'—the index offspring of the avon longitudinal study of parents and children. *Int J Epidemiol* 2013; **42**: 111–27
103. Hardcastle K, Bellis M. *Routine enquiry for history of adverse childhood experiences (ACEs) in the adult patient population in a general practice setting: a pathfinder study*. Wrexham: Public Health Wales NHS Trust; 2018
104. National Institute for Health and Care Excellence. *Chronic pain (primary and secondary) in over 16s: assessment of all chronic pain and management of chronic primary pain*.
- Available from, <https://www.nice.org.uk/guidance/ng193>. [Accessed 7 December 2022]
105. Scottish Intercollegiate Guidelines Network. *Management of chronic pain 2019*. Available from, <https://www.sign.ac.uk/our-guidelines/management-of-chronic-pain/>. [Accessed 7 December 2022]
106. Craner JR, Lake ES, Barr AC, Kirby KE, O'Neill M. Childhood adversity among adults with chronic pain: prevalence and association with pain-related outcomes. *Clin J Pain* 2022; **38**: 551–61

Handling Editor: Phil Hopkins