

HERSCHEL EVIDENCE FOR DISK FLATTENING OR GAS DEPLETION IN TRANSITIONAL DISKS*

J. T. KEANE¹, I. PASCUCCI¹, C. ESPAILLAT², P. WOITKE³, S. ANDREWS⁴, I. KAMP⁵, W.-F. THI⁶, G. MEEUS⁷, AND W. R. F. DENT⁸

¹ Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721, USA

² Department of Astronomy, Boston University, Boston, MA 02215, USA

³ SUPA, School of Physics & Astronomy, University of St. Andrews, North Haugh, St. Andrews KY16 9SS, UK

⁴ Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

⁵ Kapteyn Astronomical Institute, Postbus 800, 9700 AV Groningen, The Netherlands

⁶ Institut de Planétologie et d'Astrophysique (IPAG) UMR 5274, Université Joseph Fourier Grenoble-1, CNRS-INSU, F-38041 Grenoble, France

⁷ Departamento de Física Teórica, Universidad Autónoma de Madrid, Campus Cantoblanco, E-28049 Madrid, Spain

⁸ ALMA SCO, Alonso de Cordova 3107, Vitacura, Santiago, Chile

Received 2014 January 8; accepted 2014 March 28; published 2014 May 14

ABSTRACT

Transitional disks are protoplanetary disks characterized by reduced near- and mid-infrared emission, with respect to full disks. This characteristic spectral energy distribution indicates the presence of an optically thin inner cavity within the dust disk believed to mark the disappearance of the primordial massive disk. We present new *Herschel* Space Observatory PACS spectra of [O I] 63.18 μm for 21 transitional disks. Our survey complements the larger *Herschel* GASPS program (“Gas in Protoplanetary Systems”) by quadrupling the number of transitional disks observed with PACS in this wavelength. [O I] 63.18 μm traces material in the outer regions of the disk, beyond the inner cavity of most transitional disks. We find that transitional disks have [O I] 63.18 μm line luminosities ~ 2 times fainter than their full disk counterparts. We self-consistently determine various stellar properties (e.g., bolometric luminosity, FUV excess, etc.) and disk properties (e.g., disk dust mass, etc.) that could influence the [O I] 63.18 μm line luminosity, and we find no correlations that can explain the lower [O I] 63.18 μm line luminosities in transitional disks. Using a grid of thermo-chemical protoplanetary disk models, we conclude that either transitional disks are less flared than full disks or they possess lower gas-to-dust ratios due to a depletion of gas mass. This result suggests that transitional disks are more evolved than their full disk counterparts, possibly even at large radii.

Key words: accretion, accretion disks – circumstellar matter – infrared: stars – protoplanetary disks – stars: pre-main sequence

Online-only material: color figures, extended figures

1. INTRODUCTION

Protoplanetary disks (gas-rich dust disks around young stars) provide the raw building blocks for solar systems. While significant progress has been made in understanding the relevant evolutionary timescales of protoplanetary disks, little is known about the physical mechanisms driving the eventual dispersal of dust and gas about these young systems (for review, see Pascucci & Tachibana 2010). The goal of this paper is to gain insight into these dispersal processes by investigating a special type of protoplanetary disk that is thought to be in the process of losing its primordial dust disk: the transitional disks.

Transitional disks, like their full protoplanetary disk cousins, are often identified by their spectral energy distributions (SEDs). While there is significant variation in the SEDs of young star systems, transitional disks appear as a distinct subgroup of protoplanetary disks: their SEDs show reduced near- and mid-infrared emission, with respect to full disks (Strom et al. 1989). This characteristic SED points to the presence of an optically thin inner cavity, extending from the star out to $1 \sim 20$ AU. The excavation of this cavity is believed to mark the early stages of the dispersal of the primordial, massive dust disk, whose continuous dust disk extended as close as a few stellar radii to the central star (e.g., Calvet et al. 2002; Espaillat et al. 2007). The existence of inner cavities has been directly confirmed for a few transitional disks via sensitive, high-resolution millimeter

observations which detect reduced (or absent) dust emission from the inner disk as a result of a deficit of millimeter size grains (e.g., Andrews et al. 2009; Brown et al. 2009). While transitional disks may possess dust cavities, it is known that, in most cases, these dust cavities are not devoid of gas. Transitional disks are still actively accreting (e.g., Najita et al. 2007), and various optical emission lines (e.g., CO lines, [O I] 6300 Å and 5577 Å, etc.) indicate the presence of gas within the dust cavity region, though it may be depleted (e.g., TW Hya; Gorti et al. 2011).

There are three leading hypotheses for the driving mechanism behind the formation of cavities in transitional disks (for review, see Espaillat et al. 2014).

1. *Dust coagulation.* As disks evolve, submicron-sized dust grains coagulate into larger aggregates which have little emission at infrared wavelengths and thus reduce the disk opacity. These larger aggregates would eventually coalesce into planetesimals and planetary embryos. Since dynamical timescales increase with increasing radial distance from the central star, grain growth occurs inside-out and leads to the development of an expanding optically thin inner cavity, although the total mass of this inner disk region is not necessarily lower (e.g., Dullemond & Dominik 2005).
2. *Photoevaporation.* High-energy photons from the central star can drive photoevaporative winds, particularly from the outer regions of the protoplanetary disk (beyond \sim few AU). As the viscous accretion rate drops below the photoevaporation mass loss rate, a gap opens in the disk and the inner disk viscously accretes onto the star, resulting in an inner

* *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

cavity (e.g., Alexander et al. 2014). Direct irradiation of the cavity wall is expected to rapidly disperse the outer disk (Alexander et al. 2006). Photoevaporative winds have been detected for select protoplanetary disks via blueshifted (\sim few km s⁻¹) [Ne II] 12.81 μ m lines, which traces unbound winds within the inner \lesssim 10's of AU (Pascucci & Sterzik 2009).

3. *Dynamical clearing by giant planets.* Dynamical interactions between the disk and an embedded giant planet (with masses roughly equal to that of Jupiter) can open gaps within the disk (e.g., Lubow et al. 1999). Gas from the inner disk (within the planet's orbit) can continue to accrete onto the central star, while most of the gas from the outer disk (beyond the planet's orbit) accretes onto the planet, and only a small amount of gas flows past the planet into the inner disk. In addition to the physical gap created by the planet, pressure gradients setup at the outer edge of the gap can act as a dust filter, allowing only grains below a critical size to reach the inner disk and perhaps forming an optically thin inner cavity (Rice et al. 2006).

While these different mechanisms can produce qualitatively similar SEDs, they predict distinctive differences in the distribution of disk gas. Furthermore, these different processes can, and probably do, operate simultaneously.

In this paper, we use *Herschel Space Observatory* far-infrared data to examine whether full disks and transitional disks are different in their outer disk regions, beyond 10's of AU. We use the [O I] 63.18 μ m emission line and the nearby 63 μ m continuum emission to trace the gas and dust components, respectively, beyond 10 \sim 100 AU (e.g., Aresu et al. 2012). In addition, we use ancillary data to characterize our sample at different wavelengths. In Section 2, we provide a short description of our sample, the *Herschel*/PACS observations and data reduction, and the ancillary stellar and disk properties used to characterize our sample. In Section 3, we summarize our [O I] 63.18 μ m line 63 μ m continuum results. Most notably, we find that transitional disks possess [O I] 63.18 μ m line luminosities a factor of 2 \sim 3 lower than full disks, despite having similar 63 μ m continuum luminosities, a trend previously identified by Howard et al. (2013), though expanded in this work with quadruple the number of transitional disks. In Section 4, we rule out various observable stellar and disk properties (e.g., FUV and X-ray luminosity) as the potential cause for this [O I] 63.18 μ m line luminosity difference between full disks and transitional disks. In Section 5, we use the results of the DENT grid (a grid of 300,000 thermochemical protoplanetary disk models, by Woitke et al. 2010), to examine other possible causes for the [O I] 63.18 μ m line luminosity difference. We conclude that the lower [O I] 63.18 μ m line luminosity of transitional disks could be due to transitional disks either being less flared or by having lower gas-to-dust ratios. In Section 6, we discuss the implications of this result for disk evolution models and potential follow-up observations.

2. OBSERVATIONS AND DATA REDUCTION

2.1. Sample Description

We selected 21 transitional disks from predominantly young (a few Myr old) and nearby (\leq 200 pc) star-forming regions. Five additional transitional disks were selected from the GASPS sample (“Gas in Protoplanetary Systems”; Dent et al. 2013), resulting in a total of 26 transitional disks. Our sample is listed in Table 1. The transitional disks were identified by significant dips in their *Spitzer*/IRS spectra. (For the relevant spectra used

to identify each transitional disk as transitional, see Brown et al. 2007; Calvet et al. 2002; Espaillat et al. 2010; Furlan et al. 2009; Kim et al. 2009; Merín et al. 2008, 2010.) Because only 10% of protoplanetary disks are transitional (e.g., Williams & Cieza 2011; Muzerolle et al. 2010), we selected targets from a number of star-forming regions, including Taurus-Auriga, Ophiuchus, Chameleon, and Lupus. Targets that have been previously modeled either with continuum radiative transfer codes or simple prescriptions for the disk inner cavity were given preference, as were objects with archival measurements of accretion rates and infrared and millimeter observations.

For comparison with our sample of transitional disks, we selected an additional 33 protoplanetary disks from the GASPS (“Gas Survey of Protoplanetary Systems”; Dent et al. 2013) survey of the Taurus-Auriga star-forming region.⁹ These disks were selected as being typical protoplanetary disks, with IRS spectra close to the Taurus-Auriga mean (D’Alessio et al. 2006), and were also selected to sample similar spectral types to those of the transitional disks. Like the transitional disk subsample, we took preference for objects with known accretion rates and millimeter observations. Of these 33 protoplanetary disks, 15 have jets/outflows as identified by a combination of optical and near-IR spectroscopy and imaging (see Kenyon et al. 2008 and references therein). We will refer to this subsample as “outflow” sources. The remaining 18 disks without outflows will be referred to as “full” disks. This distinction between outflow disks and full disks varies slightly from Howard et al. (2013), who identified outflow disks as objects with either directly imaged jets in H α , [O I] λ 6300, [S II] λ 6371, being associated with Herbig–Haro objects, or having very broad [O I] λ 6300 emission line profiles. This slight difference in definition only changes the classification of two disks (AA Tau and DL Tau). None of our full disks were noted to have broadened or spatially extended [O I] 63.18 μ m emission in Howard et al. (2013).

Targets with binary companions represent a possible source of contamination within our samples given the large spaxel size of PACS (9'.4 on a side, which corresponds to a projected separation of \sim 1300 AU at the distance of Taurus). Close binary companions can interact with the primary star's disk and produce transitional SEDs (e.g., CoKu Tau/4; Ireland & Kraus 2008). Medium and large separation binaries (projected separations $>$ 10 AU) do not seem to strongly affect the first steps of planet formation (grain growth and dust settling; Pascucci et al. 2008). However, binaries with separations \leq 40 AU significantly hasten the process of disk dispersal (Kraus et al. 2012). While it might be best to remove all binaries from our sample and just focus on single stars, this approach could bias our samples and, as such, our results. The Taurus-Auriga star-forming region, from which we draw most of the full disks, has been well surveyed for multiplicity. However, Ophiuchus, Chameleon, and the Lupus star-forming regions, from which we draw our sample of transitional disks, have not been as well studied. Thus, there are very likely undetected multiple systems within our transitional disk sample. We opted to retain full disks within multiple systems, as long as the mid-infrared flux (\sim 10 μ m) ratio between members is large ($L_{\text{IR,primary}}/L_{\text{IR,secondary}} \gtrsim 3$), and the protoplanetary disks are not circumbinary. Targets that do not meet this criterion are excluded from all analysis (though they are included in tables and figures, for reference). Table 1 lists the multiplicity status for all targets, as well as the relevant

⁹ An additional full disk, SZ 50, from the Cha I star-forming region, was included in order to more properly match the spectral type distribution between full disks and transitional disks.

Table 1
Herschel/PACS Sample and Observations

ID	Name	R.A.	Decl.	Association	SpTy	Ref.	Multiplicity	Ref.	ObsID	Duration (s)
Transition Disks										
T1	16201-2410F-1*	16 23 09.23	-24 17 04.70	Ophiuchus	G0	F09	1	...	1342250127	8212
T2	CHX 22*	11 12 42 69	-77 22 23.00	Chameleon	G8	L07	2	D13	1342233474	8212
T3	CHX 7*	11 06 15 41	-77 21 56.90	Chameleon	G5	L07	1342233477	8212
T4	CR Cha	10 59 06 99	-77 01 40.40	Chameleon	K2	E11	2	G07	1342232614	8212
T5	CS Cha*	11 02 24 91	-77 33 35.70	Chameleon	K6	E11	1342233480	8212
T6	DM Tau	04 33 48.72	+18 10 09.99	Taurus	M1	KH95	1342225825*	6628
T7	DoAr 28	16 26 47.42	-23 14 52.20	Ophiuchus	K5	M92	1342241707	8212
T8	DoAr 44	16 31 33.46	-24 27 37.30	Ophiuchus	K3	M92	1342250578	8212
T9	GM Aur	04 55 10.99	+30 21 59.25	Taurus	K5.5	E11	1342191357*	1252
T10	Hn 24*	13 04 55 75	-77 39 49.50	Chameleon	M0.5	M10	2	B96	1342235656	8212
T11	LkCa 15	04 39 17.80	+22 21 03.48	Taurus	K5	KH95	1342225798*	6628
T12	LkHalpha 330*	3 45 48 28	+32 24 11.90	Perseus	G3	BR07	1342238377	8212
T13	RXJ1615.3-3255	16 15 20 23	-32 55 05.10	Lupus	K4	M10	1342229825	8212
T14	SSTLup	16 10 29.60	-39 22 15.00	Lupus	M5	M10	1342241709	8212
T15	Sz 111	16 08 54 69	-39 37 43.10	Lupus	M1.5	H94	1342220928	8212
T16	Sz 18	11 07 19 15	-76 03 04.80	Chameleon	M2.5	L07	1342232585	8212
T17	Sz 27	11 08 39 05	-77 16 04.20	Chameleon	K8	L07	1342233476	8212
T18	Sz 45	11 17 37 01	-77 04 38.10	Chameleon	M0.5	L07	1342233475	8212
T19	Sz 84	15 58 02 53	-37 36 02.70	Lupus	M5.5	M10	1342229826	8212
T20	Sz 91	16 07 11 61	-39 03 47.10	Lupus	M0.5	H94	1342229827	8212
T21	Sz Cha	10 58 16 77	-77 17 17.10	Chameleon	K0	E11	2	D13	1342233478	8212
T22	T Cha*	11 57 13 53	-79 21 31.50	Chameleon	G8	BR07	1342232294	2068
T23	TW Hya	11 01 52 03	-34 42 18.60	TW Hydra	K6	R06	1342187127*	1252
T24	UX Tau	04 30 03.76	+18 13 49.88	Taurus	K2	KH95	3	M06	1342214357*	1252
T25	WSB60	16 28 16.51	-24 36 58.00	Ophiuchus	M4.5	WMRG05	1342250128	8212
T26	YLW8*	16 27 10 28	-24 19 12.70	Ophiuchus	G2.5	BR07	2	M06	1342229824	2068
Full Disks										
F1	AA Tau	04 34 55.42	+24 28 53.16	Taurus	K7	KH95	1342225758*	6628
F2	BP Tau*	04 19 15.84	+29 06 26.94	Taurus	K7	KH95	1342225728*	3316
F3	CI Tau	04 33 52.00	+22 50 30.18	Taurus	K7	KH95	1342192125*	1252
F4	CY Tau	04 17 33.73	+28 20 46.85	Taurus	M1	KH95	1342192794*	1252
F5	DE Tau	04 21 55.64	+27 55 06.06	Taurus	M2	KH95	1342192797*	1252
F6	DK Tau	12 53 17.23	-77 07 10.70	Taurus	K7	KH95	2	WG01	1342225732*	3316
F7	DL Tau	04 33 39.06	+25 20 38.23	Taurus	K7	KH95	1342225800*	6628
F8	DN Tau	04 35 27.37	+24 14 58.94	Taurus	M0	KH95	1342225757*	3316
F9	DQ Tau*	04 46 53.04	+17 00 00.50	Taurus	M0	KH95	2	AW05	1342225806*	1252
F10	DS Tau	04 47 48.21	29 25 13.83	Taurus	K5	KH95	2	AW05	1342225851*	3316
F11	GG Tau*	04 32 30.35	+17 31 40.60	Taurus	K7	KH95	2	WG01	1342192121*	1252
F12	GO Tau	04 43 03.10	+25 20 18.75	Taurus	M0	KH95	1342225826*	3316
F13	HBC 347	03 29 38.24	+24 30 37.74	Taurus	1342192136*	1252
F14	HK Tau	04 31 50.67	+24 24 17.44	Taurus	M0.5	KH95	2	WG01	1342225736*	3316
F15	IQ Tau	04 29 51.56	+26 06 44.89	Taurus	M0.5	KH95	1342225733*	3316
F16	SU Aur	04 55 59.38	+30 34 01.56	Taurus	G2	KH95	1342217844*	3316
F17	SZ 50	13 00 55.36	-77 10 22.10	Chameleon	M3	HH92	1342226008*	3316
F18	V836 Tau	05 03 06.60	+25 23 19.71	Taurus	K7	KH95	1342227634*	3316
Outflow Disks										
O1	CW Tau	04 14 17.00	+28 10 57.83	Taurus	K3	KH95	1342216221*	1252
O2	DF Tau*	04 27 02.80	+25 42 22.30	Taurus	M3	KH95	2	P08	1342190359*	1252
O3	DG Tau	04 27 04.698	+26 06 16.31	Taurus	K6	KH95	1342190382*	1252
O4	DG Tau B	04 27 02.41	+26 05 31.76	Taurus	M0	KH95	1342192798*	1252
O5	DO Tau	04 38 28.58	+26 10 49.44	Taurus	M0	KH95	1342190385*	1252
O6	DP Tau	04 42 37.56	+25 15 39.62	Taurus	M0.5	KH95	1342191362*	1252
O7	GI/GK Tau	04 55 10.85	+30 22 01.69	Taurus	K6	KH95	2	AW05	1342225760*	1252
O8	Haro 6-13	04 32 15.41	+24 28 59.75	Taurus	M0	RM12	1342192128*	1252
O9	HN Tau	04 33 39.44	+17 51 52.24	Taurus	K5	KH95	2	WG01	1342225796*	3316
O10	HV Tau	04 38 35.38	+26 10 37.80	Taurus	M1	KH95	2	WG01	1342225801*	3316
O11	RW Aur	05 07 49.41	+30 24 07.65	Taurus	K3	KH95	2	WG01	1342191359*	1252
O12	RY Tau	04 21 57.40	+28 26 35.54	Taurus	K1	KH95	1342190361*	1252
O13	T Tau	04 21 59.30	+19 32 08.53	Taurus	K0	KH95	2	WG01	1342190353*	1252
O14	UY Aur *	04 51 47.15	+30 47 14.44	Taurus	K7	KH95	2	WG01	1342215699*	1252
O15	UZ Tau*	04 32 42.73	+25 52 35.00	Taurus	M1	KH95	4	WG01	1342192131*	1252

Table 1
(Continued)

Notes. Targets tagged with an asterisk were excluded from statistical tests due to either being a binary that does not meet the criteria in Section 2.1, or having a spectral type earlier than K-type. BP Tau was also excluded from statistical tests, due to its nature as an “evolved” full disk. ObsIDs tagged with a star (★) were observed by the GASPS team and were previously reported in Howard et al. (2013), Meeus et al. (2012), and Podio et al. (2012), although they were re-reduced here using an updated version of the Herschel pipeline. Distances for each star-forming association (from Reipurth 2008a, 2008b): Chameleon, 160 pc; Lupus, 155 pc; Ophiuchus, 120 pc; Perseus, 250 pc; Taurus, 140 pc; TW Hya, 56 pc.

References. Andrews & Williams 2005 (AW05); Brandner et al. 1996 (B96); Brown et al. 2007 (BR07); Daemgen et al. 2013 (D13); Espaillat et al. 2011 (E11); Furlan et al. 2009 (F09); Guenther et al. 2007 (G07); Hughes et al. 1994 (H94); Kenyon & Hartmann 1995 (KH95); Luhman 2007 (L07); Magazzu et al. 1992 (M92); McCabe et al. 2006 (M06); Merín et al. 2010 (M10); Pascucci et al. 2008 (P08); Riaz et al. 2006 (R06); White & Ghez 2001 (WG01); Wilking et al. 2005 (WMRG05).

references for the projected separations and mid-infrared flux ratio for binaries.

2.2. *Herschel* PACS Spectroscopy

We obtained *Herschel Space Observatory* PACS (Poglitsch et al. 2010) spectroscopy for our sample of 21 transitional disks. The relevant *Herschel* observation identification numbers (ObsIDs), exposure times, and dates of our observations are listed in Table 1. The five additional transitional disks (DM Tau, LkCa 15, GM Aur, TW Hya, and UX Tau) and the entire sample of full disks and outflow disks were previously observed as part of the *Herschel* Key Program: GASPS (PI, W. Dent). We used the line spectroscopy mode (“PacsLineSpec”) to take spectra centered on the [O I] 63 μ m line, between 62.93 and 63.43 μ m. All of the observations were executed in “ChopNod” mode, in order to remove telescope emission and background.

We reduced our original observations and re-reduced the GASPS archival data with the *Herschel* Interactive Processing Environment (HIPE; Ott 2010) version 9.0.0. We used the default “ChopNodLineScan” pipeline along with the most recent calibration tree (CalTree 32). The data reduction process included: removal of saturated and overly-noisy pixels; differencing the on-source and off-source observations; spectral response function division; rebinning to the native resolution of the instrument (oversample = 2, upsample = 1); spectral flat fielding; and averaging over the two nod positions. We extracted our spectrum from the central spaxel and accounted for diffraction losses to neighboring spaxels with an aperture correction provided in HIPE. Since outflow targets generally can have extended emission spanning multiple spaxels (Podio et al. 2012; Howard et al. 2013), our measured fluxes for these outflow targets will generally underestimate their true fluxes. We verified that for all of our transitional disks and full disks, there was no appreciable emission in neighboring spaxels. This lack of extended emission also suggests that there is no significant mispointing in the observations of our transitional disks and full disks.

2.3. *Stellar and Disk Properties*

To interpret our *Herschel* PACS observations of [O I] and its relationship to the protoplanetary disk environment, we aggregated stellar properties (effective temperature, bolometric luminosity, FUV, and X-ray luminosities, etc.) as well as disk properties (disk mass, disk structure, accretion rates) that, through past work, are known to affect the [O I] 63.18 μ m emission. In this section, we explain the methods by which we derived these stellar and disk properties. We will relate these to our *Herschel* observations in Section 4.

2.3.1. *Effective Temperature and Bolometric Luminosity*

We determined stellar effective temperatures by relating the host star’s spectral type (from the literature) to the corresponding effective temperature (Luhman 1999), as listed in Table 2. Generally, we do not assume that the effective temperatures are accurate to more than one spectral subtype (~ 100 K).

We self-consistently derived bolometric luminosities for all targets by performing a bolometric correction on de-reddened, literature-available *I*-band photometry listed in Table 2. *I*-band photometry is preferential, as it is less affected by intervening dust. We de-reddened all of our *I*-band fluxes by relating *V*-band extinctions (which are more commonly reported in the literature) to *I*-band extinctions using relationships from Mathis (1990), assuming R_V values typical of the interstellar medium ($R_V = 3.1$). The de-reddened continuum fluxes were converted to luminosities using the known distances to each different star-forming region (see Table 1 and references therein). Finally, bolometric corrections from Luhman (1999) were used to calculate bolometric luminosities for each target. These effective temperatures and bolometric luminosities are listed in Table 3.

2.3.2. *FUV Luminosities and Accretion Rates*

While ultraviolet observations of T Tauri stars would provide the most direct measurement of the FUV luminosity, these observations are notoriously difficult. Instead, we made use of the well-known correlation between accretion rate and FUV excess emission to derive FUV luminosities from stellar accretion rates (e.g., Dahm 2008; Herczeg & Hillenbrand 2008). Accretion rates can be determined from a large number of other more commonly measured emission lines (e.g., Rigliaco et al. 2011).

We used the H α emission line at 6563 Å to estimate stellar accretion rates. H α is advantageous because it is a very commonly reported observational diagnostic, and it correlates well with accretion luminosities (the excess luminosity arising from the infall and accretion of material onto the central star), as derived from other accretion tracers (e.g., Rigliaco et al. 2011). To determine accretion luminosities, we calculated the H α line fluxes from literature H α equivalent widths (listed in Table 2). For targets with multiple H α equivalent widths available in the literature, we used the mean equivalent width.¹⁰ To determine H α line fluxes, we combined the H α equivalent width with the nearest available photometric point in the literature: *R*-band. De-reddening was done similarly as for our bolometric luminosity analysis: using *V*-band extinctions converted to

¹⁰ While H α is known to be variable, it has been shown that the variability does not introduce significant scatter in H α -derived accretion rates (Biazzo et al. 2012). For our targets with multiple H α equivalent widths, using either the maximum or minimum H α equivalent width changes the resulting accretion rate on average only 0.12 dex. This variation is less than the uncertainty that results from the empirical relationships used to convert H α luminosity to accretion luminosity (Fang et al. 2009).

Table 2
Literature Data

ID	R_{mag}	Ref.	I_{mag}	Ref.	A_V	Ref.	H α EW (Å)	Ref.	$\log L_X$ (L_{\odot})	Ref.
Transition Disks										
T1	14.20	C03	12.88	DENIS	6.90	MFM10
T2	10.45	GS92	10.02	GS92	1.21	GS92	1.5	GS92	-3.45	FK89
T3	10.28	GS92	7.64	GS92	3.39	GS92	1.2	B08	-3.02	FK89
T4	10.46	GS92	9.73	GS92	1.50	E11	38.1	GE97	-2.94	FK89
T5	10.92	GS92	9.11	GS92	0.85	GS92	13.3	GS92	-3.36	FK89
T6	12.92	KH95	11.77	KH95	0.00	KH95	138.7	CK79	-4.33	G07
T7	12.10	C03	11.69	DENIS	2.10	CMLW95	36	M92
T8	11.70	BA92	10.80	BA92	2.20	A11	68.3	BA92	-3.69	A11
T9	11.20	B93	10.70	B93	0.14	KH95	96.5 109 71	CK79, E94, C90	\leq -3.89	A11
T10	13.00	C03	11.95	S07	2.00	M10	0.2	M10	\leq -3.02	A00
T11	11.58	KH95	10.79	KH95	0.62	KH95	18.05	SB09	\leq -3.99	A11
T12	11.20	C03	10.80	F95	1.55	OB95	16	SB09
T13	11.28	M10	10.54	M10	1.00	M10	26	M10	-3.19	A11
T14	15.79	M10	13.90	M10	1.00	M10	18	M10
T15	13.34	M08	12.17	M03	0.10	H94	145.2	H94
T16	14.15	GS92	12.69	GS92	1.60	E11	5	L04
T17	14.96	GS92	13.41	GS92	3.50	E11	100	L04	-4.79	W00
T18	12.57	GS92	11.59	GS92	0.60	E11	56	L04	-3.69	W00
T19	14.53	M10	12.94	M10	0.50	M10	44	M10
T20	14.28	H94	12.92	H94	2.00	H94	95.9	H94
T21	11.21	GS92	9.25	GS92	1.88	GS92	12	GS92	-3.99	F93
T22	11.07	S09	10.28	DENIS	1.70	S09	7.8	S09
T23	11.40	DENIS	9.38	DENIS	1.00	K99	213.8	R06	-3.85	H07
T24	10.48	KH95	9.75	KH95	0.21	KH95	3.9	T09	-3.33	D95
T25	16.55	WMRG05	14.33	DENIS	2.00	WMRG05	81	WMRG05
T26	12.20	DENIS	11.29	DENIS	9.00	PGS03	4	SB09	-3.59	A11
Full Disks										
F1	12.06	KH95	10.99	HHG94	0.49	KH95	37.1 80 21	CK79, E94, C90	-3.49	G07
F2	11.31	KH95	10.45	KH95	0.49	KH95	40.1 55 47 49.4	CK79, E94, C90, MCH01	-3.45	G07
F3	12.22	KH95	11.12	KH95	1.77	KH95	102.1 64	CK79, C90	-4.30	G07
F4	12.35	KH95	11.18	KH95	0.10	KH95	69.5	CK79	-4.46	G07
F5	11.66	KH95	10.75	HHG94	0.59	KH95	54 76	CK79, C90	\leq -3.78	D95
F6	11.43	KH95	10.46	KH95	0.76	KH95	19.4 13 28	CK79, E94, C90	-3.62	G07
F7	11.85	KH95	10.89	KH95	1.70	HEG95	105 111 138	WG01, CK79 C90	\leq -3.59	D95
F8	11.49	KH95	10.49	KH95	0.49	KH95	11.9 22 15 11.1	CK79, E94, C90, MCH01	-3.52	G07
F9	12.40	KH95	11.27	KH95	0.97	KH95	112.9	CK79
F10	11.56	KH95	10.80	KH95	0.31	KH95	38.5	K98
F11	11.31	WG01	10.44	WG01	1.03	WG01	56 43 52	WG01, CK79, C90	\leq -3.55	D95
F12	13.62	KH95	12.30	KH95	1.18	KH95	80.8	CK79	-4.19	G07
F13	-3.65	D95
F14	13.93	KH95	12.37	KH95	2.32	KH95	53.5	K98
F15	12.28	KH95	11.11	KH95	1.25	KH95	7.8	CK79	-3.97	G07
F16	8.62	KH95	8.10	KH95	0.90	KH95	3.5 5	CK79, C90	-2.61	G07
F17	14.30	HH92	12.50	HH92	2.14	HH92	66 46	SA11, CK79
F18	12.17	KH95	11.19	KH95	0.59	KH95	9 5	B90, C90	-3.54	N95
Outflow Disks										
O1	12.33	KH95	11.42	KH95	2.29	KH95	137.9	R10	-3.13	G07
O2	11.07	KH95	9.87	KH95	0.21	KH95	53.9	CK79	-3.78	D95
O3	11.51	KH95	10.54	KH95	3.20	HEG95	112.8 73 110	CK79, E94, C90	\leq -4.39	...
O4	-2.60	G07
O5	12.41	KH95	11.37	HHG94	2.64	KH95	108.9 101	CK79, C90	\leq -4.27	B99
O6	13.09	KH95	11.95	KH95	1.46	KH95	85.4	CK79	-4.58	G07
O7	12.15	KH95	11.06	KH95	0.87	KH95	22.5 17	K98, CK79	-3.66	G07
O8	14.85	KGW08	13.54	L00	11.90	K09	88.2	CK79	-3.68	G07
O9	12.96	KH95	12.17	KH95	0.52	KH95	158	E87	-4.40	G07
O10	12.68	KH95	9.87	KH95	1.91	KH95	8.5	E87	\leq -4.46	N95
O11	9.95	KH95	9.34	KH95	0.50	F09	84.2	CK79	-4.03	D95
O12	9.53	KH95	8.80	KH95	1.84	KH95	21	B90	-2.87	G07
O13	9.19	KH95	8.50	KH95	1.39	KH95	38	T09	-2.79	C98
O14	11.92	KH95	10.83	KH95	1.35	KH95	47 72.8	E87, CK79
O15	11.20	KH95	10.28	M03	1.49	KH95	73.5 98.1	E87, CK79	-3.64	G07

Table 2
(Continued)

References. Alcalá et al. 2000 (A00); Andrews et al. 2011 (A11); Antonucci et al. 2011 (SA11); Bary et al. 2008 (B08); Beckwith et al. 1990 (B90); Bouvier & Appenzeller 1992 (BA92); Bouvier et al. 1993 (B93); Briceño et al. 1999 (B99); Cabrit et al. 1990 (C90); Carkner et al. 1998 (C98); Chen et al. 1995 (CMLW95); Cohen & Kuhl 1979 (CK79); Cutri et al. 2003 (C03); Damiani et al. 1995 (D95); DENIS Consortium 2005 (DENIS); Edwards et al. 1987 (E87); Edwards et al. 1994 (E94); Espaillat et al. 2011 (E11); Feigelson & Kriss 1989 (FK89); Feigelson et al. 1993 (F93); Fernandez 1995 (F95); Furlan et al. 2009 (F09); Gauvin & Strom 1992 (GS92); Güdel et al. 2007 (G07); Guenther & Emerson 1997 (GE97); Hartigan et al. 1995 (HEG95); Herbst et al. 1994 (HHG94); Herczeg et al. 2007 (H07); Hughes et al. 1994 (H94); Kastner et al. 1999 (K99); Kenyon & Hartmann 1995 (KH95); Kenyon et al. 1998 (K98); Kenyon et al. 2008 (KGW08); Kraus & Hillenbrand 2009 (K09); Luhman 2000 (L00); Luhman 2004 (L04); Magazzu et al. 1992 (M92); McClure et al. 2010 (MFM10); Merín et al. 2010 (M10); Monet et al. 2003 (M03); Muzerolle et al. 2001 (MCH01); Neuhäuser et al. 1995 (N95); Osterloh & Beckwith 1995 (OB95); Prato et al. 2003 (PGS03); Rebull et al. 2010 (R10); Riaz et al. 2006 (R06); Salyk et al. 2009 (SB09); Schisano et al. 2009 (S09); Spezzi et al. 2007 (S07); Taguchi et al. 2009 (T09); White & Ghez 2001 (WG01); White et al. 2000 (W00); Wilking et al. 2005 (WMRG05).

R-band extinctions via the relationships of Mathis (1990). The de-reddened line fluxes were converted to line luminosities using the known distances to each different star-forming region (see Table 1 and references therein). The $H\alpha$ line luminosities, $L_{H\alpha}$, were then converted into accretion luminosities, L_{acc} , with the empirical relationships of Fang et al. (2009):

$$\log(L_{acc}/L_{\odot}) = (2.27 \pm 0.23) + (1.25 \pm 0.007) \times \log(L_{H\alpha}/L_{\odot}). \quad (1)$$

Our derived accretion luminosities are listed in Table 3. From our accretion luminosities, we then used the empirical relationships of Yang et al. (2012) to relate accretion luminosities to FUV luminosities:

$$\log(L_{FUV}/L_{\odot}) = -1.670 + 0.836 \times \log(L_{acc}/L_{\odot}). \quad (2)$$

Our derived FUV luminosities are also listed in Table 3. For the 12 disks shared between this study and Yang et al. (2012), we found that our FUV luminosities agreed to those derived by Yang et al. (2012) within 0.35 dex. We found no systematic shift between our $H\alpha$ -derived FUV luminosities and their directly measured FUV luminosities. Stellar chromospheric activity can also result in $H\alpha$ emission, so we used the spectral type dependent, equivalent width cutoffs of White & Basri (2003) to distinguish between chromospheric activity and accretion. For targets where the $H\alpha$ equivalent width fell below these cutoffs, we report accretion and FUV luminosity upper limits.

Converting accretion luminosities to accretion rates requires some physical knowledge of the system and the processes of accretion. Gullbring et al. (1998) developed a simple magnetospheric accretion model whereby the accretion luminosity is generated by the release of potential energy as gas falls from the inner edge of the disk onto the surface of the star along stellar magnetic field lines. In this model, the accretion rate, \dot{M} , is related to accretion luminosity by

$$\dot{M} = \frac{L_{acc}R_{\star}}{GM_{\star}\left(1 - \frac{R_{\star}}{R_{in}}\right)}, \quad (3)$$

where R_{\star} and M_{\star} are the radius and mass of the star, G is Newton's gravitational constant, and R_{in} is the inner truncation radius of the disk. R_{in} is generally unknown, but is usually assumed to be $\approx 5R_{\star}$, which corresponds to the typical co-rotation distance (Gullbring et al. 1998; Shu et al. 1994). The stellar radius is determined from the star's effective temperature and bolometric luminosity via the Stefan–Boltzmann Law. We used pre-main sequence evolutionary tracks from Siess et al. (2000) to relate the effective temperature and bolometric

luminosity to specific stellar masses. Our final accretion rates (as well as stellar masses) are listed in Table 3.

To test the validity of our self-consistently derived accretion rates, we compared our results with an array of other studies, including Najita et al. (2007), Gullbring et al. (1998), Hartmann et al. (1998), White & Ghez (2001), and Hartigan et al. (1995). While differences between accretion rates can develop from several factors (including the use of different accretion tracers, different estimates of extinction, different bolometric corrections, use of non-contemporaneous photometry, etc.), we find that our accretion rates generally agree with past studies to within ~ 0.5 dex. This level of variation between accretion rates computed from different tracers is typical, even if observations are contemporaneous (Rigliaco et al. 2012). Our accretion rates are also not significantly offset from past studies of accretion rates, with the exception of Hartigan et al. (1995), who find systematically higher accretion rates (although this systematic offset from other estimates has been noted in previous studies; e.g., Gullbring et al. 1998).

One of the major advantages of our study, as compared to many past studies, is that our accretion rates are self-consistently derived using all the same metric, instead of being aggregated from different literature sources which adopt different methods.

2.3.3. Disk Structure and Disk Mass

Many transitional disks in our sample have been previously modeled with radiative transfer codes in order to reproduce near- and mid-infrared disk spectra and resolved millimeter images. While the exact nature of these disk models can vary between papers, they all involve the creation of a simple, axisymmetric model disk with a prescribed dust and gas surface density. Models specific to transitional disks include gas and dust cavities within a specified radius: r_{cavity} . At the outer edge of this dust cavity, the frontally illuminated disk wall puffs up to a wall height of h_{wall} , which can significantly affect the near-infrared emission of transitional disks (both due to excess emission and shadowing of the outer disk; Espaillat et al. 2011). These disk models are then subject to simulated observations, and the relevant model spectra or resolved images are calculated (for some specified viewing angle) and fit to observations. We aggregated values for the cavity size (r_{cavity}) and the wall height (h_{wall}) from the literature. These disk properties are listed in Table 4. While the individual models can vary between papers, the majority of these cavity sizes and wall heights are taken from Andrews et al. (2011) and Espaillat et al. (2011), which both use similar disk models.

In addition to looking at the cavity size and wall height, we used self-consistently calculated estimates of the total disk

Table 3
Stellar and Accretion Properties

ID	T_{eff} (K)	M_{\star} (M_{\odot})	L_{bol} (L_{\odot})	R_{\star} (R_{\odot})	$\log L_{\text{acc}}$ (L_{\odot})	$\log \dot{M}$ ($M_{\odot} \text{ yr}^{-1}$)	$\log L_{\text{FUUV}}$ (L_{\odot})
Transition Disks							
T1	6030	1.15	1.01	0.92
T2	5520	1.34	1.98	1.54	-2.0	-9.3	-3.3
T3	5770	3.7	47.60	6.92	-1.2	-8.3	-2.7
T4	4900	1.98	2.83	2.34	-0.1	-7.4	-1.8
T5	4205	1.62	3.76	3.66	-1.2	-8.2	-2.6
T6	3705	0.45	0.18	1.02	-1.3	-8.4	-2.8
T7	4350	0.95	0.33	1.01	-1.0	-8.4	-2.5
T8	4730	1.26	0.82	1.35	-0.5	-7.8	-2.1
T9	4277.5	1.00	0.50	1.28	-0.7	-7.9	-2.2
T10	3777.5	0.52	0.46	1.59	≤ -4.0	≤ -11.0	≤ -5.0
T11	4350	1.05	0.53	1.29	-1.5	-8.9	-3.0
T12	5830	1.25	2.78	1.64	-0.4	-7.7	-2.0
T13	4590	1.28	1.01	1.59	-0.9	-8.2	-2.5
T14	3125	0.16	0.08	0.94	≤ -3.4	≤ -10.0	≤ -4.5
T15	3632.5	0.40	0.16	1.00	-1.4	-8.4	-2.8
T16	3487.5	0.35	0.21	1.26	≤ -3.0	≤ -9.9	≤ -4.2
T17	4060	0.80	0.23	0.97	-1.1	-8.4	-2.6
T18	3777.5	0.52	0.35	1.38	-1.3	-8.3	-2.7
T19	3060	0.17	0.17	1.45	-2.5	-8.9	-3.7
T20	3777.5	0.52	0.18	0.99	-1.4	-8.5	-2.8
T21	5250	2.18	5.50	2.84	-1.0	-8.2	-2.5
T22	5520	1.31	0.89	1.04	-1.6	-9.1	-3.0
T23	3850	0.58	0.39	1.40	-1.0	-8.0	-2.5
T24	4900	1.3	1.21	1.53	-2.0	-9.3	-3.3
T25	3197.5	0.17	0.04	0.64	-2.9	-9.7	-4.1
T26	5845	2.20	11.06	3.25	0.3	-6.9	-1.4
Full Disks							
F1	4060	0.8	0.43	1.32	-1.3	-8.5	-2.8
F2	4060	0.78	0.70	1.70	-0.9	-8.0	-2.4
F3	4060	0.78	0.67	1.65	-0.6	-7.7	-2.2
F4	3705	0.46	0.32	1.37	-1.4	-8.3	-2.8
F5	3560	0.38	0.59	2.03	-0.9	-7.6	-2.4
F6	4060	0.76	0.78	1.79	-1.4	-8.4	-2.8
F7	4060	0.76	0.80	1.81	-0.3	-7.3	-1.9
F8	3850	0.56	0.70	1.88	-1.7	-8.5	-3.0
F9	3850	0.57	0.42	1.46	-0.8	-7.8	-2.4
F10	4350	1.2	0.46	1.20	-1.2	-8.6	-2.7
F11	4060	0.76	0.90	1.92	-0.7	-7.7	-2.3
F12	3850	0.57	0.18	0.95	-1.5	-8.7	-3.0
F13
F14	3777.5	0.52	0.28	1.23	-1.5	-8.5	-2.9
F15	3777.5	0.52	0.55	1.74	-2.1	-9.0	-3.4
F16	5860	1.8	7.96	2.74	-0.7	-8.0	-2.3
F17	3415	0.33	0.34	1.67	-1.6	-8.3	-3.0
F18	4060	0.72	0.37	1.23	≤ -2.4	≤ -9.5	≤ -3.6
Outflow Disks							
O1	4730	1.07	0.66	1.21	-0.2	-7.5	-1.8
O2	3415	0.33	1.25	3.20	-0.9	-7.3	-2.4
O3	4205	0.9	2.06	2.71	0.4	-6.5	-1.4
O4	3850
O5	3850	0.56	0.80	2.01	-0.2	-7.1	-1.9
O6	3777.5	0.52	0.28	1.24	-1.1	-8.2	-2.6
O7	4205	0.95	0.46	1.27	-1.7	-9.0	-3.1
O8	3850	0.55	6.42	5.71	1.9	-4.5	-0.1
O9	4350	0.7	0.14	0.67	-1.1	-8.5	-2.6
O10	3705	0.45	2.34	3.72	≤ -2.0	≤ -8.5	≤ -3.4
O11	4730	1.49	2.03	2.13	0.1	-7.2	-1.6
O12	5080	2.2	6.15	3.21	0.0	-7.2	-1.7
O13	5250	2.18	6.77	3.15	0.3	-6.9	-1.4
O14	4060	0.77	0.72	1.72	-0.8	-7.8	-2.3
O15	3705	0.47	1.34	2.81	-0.2	-6.8	-1.8

Table 4
Disk Properties

ID	r_{gap} (AU)	Ref.	h_{wall} (AU)	Ref.	$f_{850\mu\text{m}}$ (mJy)	$f_{1.3\text{mm}}$ (mJy)	Ref.	$M_{850\mu\text{m}}$ (M_{Jup})	$M_{1.3\text{mm}}$ (M_{Jup})
Transition Disks									
T1
T2	37.1	KM09	≤ 118	H93	...	≤ 25.8
T3	146.7	KM09	≤ 143	H93	...	≤ 32.5
T4	10	E11	124.9	H93	...	28.4
T5	38	E11	7	E11	...	128.4	H93	...	29.1
T6	19	A11	5.7	A11	237	109	M13	13.6	19.0
T7	≤ 75	M13	...	≤ 9.6
T8	30	A11	9	A11	181	105	M13	7.6	13.4
T9	23	E11	2.9	E11	...	253	M13	...	44.1
T10
T11	39	E11	5	E11	428	167	M13	24.5	29.1
T12	68	A11	6.8	A11	...	70	OB95	...	38.9
T13	30	A11	2	A11
T14
T15
T16	13	E11	2	E11	...	105	H93	...	23.9
T17	5	E11	4	E11	...	100	H93	...	22.8
T18	20	E11	4	E11	...	47.8	H93	...	10.9
T19	55	M10	≤ 36	N97	...	≤ 7.7
T20	≤ 27	N97	...	≤ 5.8
T21	18	E11	4	E11	...	77.5	H93	...	17.7
T22	15	BR07	105.2	H93	...	11.1
T23	4	T10
T24	≤ 173	≤ 63	M13	≤ 3.6	≤ 11.0
T25	15	A11	0.8	A11	149	89	M13	6.3	11.4
T26	36	A11	8.2	A11	397	95	M13	16.7	12.2
Full Disks									
F1	144	88	M13	8.3	15.3
F2	130	47	M13	7.5	8.2
F3	324	190	M13	18.6	33.1
F4	111	G11	...	19.3
F5	90	36	M13	5.2	6.3
F6	80	35	M13	4.6	6.1
F7	440	230	M13	25.2	40.1
F8	201	84	M13	11.5	14.6
F9	208	91	M13	11.9	15.9
F10
F11	1255	593	M13	72.0	103.3
F12	173	83	M13	9.9	14.5
F13
F14
F15	178	87	M13	5.0	15.2
F16	74	≤ 30	M13	4.2	≤ 5.2
F17
F18	74	37	M13	4.2	6.4
Outflow Disks									
O1	66	96	M13	3.8	16.7
O2	8.8	≤ 25	M13	0.5	≤ 4.4
O3	389.9	G11	...	67.9
O4
O5	248	136	M13	14.2	23.7
O6	≤ 10	≤ 27	M13	≤ 0.6	≤ 4.7
O7	33	≤ 21	M13	1.9	≤ 3.7
O8	34.2	G11	...	5.9
O9	29	≤ 15	M13	1.7	≤ 2.6
O10	47	40	A05	2.7	7.0
O11	79	42	M13	4.5	7.3
O12	560	229	M13	32.1	39.9
O13	628	280	M13	36.0	48.8
O14	102	29	M13	5.9	5.1
O15	560	172	M13	32.1	30.0

References. Andrews & Williams 2005 and references therein (A05); Andrews et al. 2011 (A11); Brown et al. 2007 (BR07); Espaillat et al. 2011 (E11); Guilloteau et al. 2011 (G11); Henning et al. 1993 (H93); Kim et al. 2009 (KM09); Merín et al. 2010 (M10); Mohanty et al. 2013 and references therein (M13); Nuernberger et al. 1997 (N97); Osterloh & Beckwith 1995 (OB95); Thi et al. 2010 (T10).

mass, from 1.3 mm and 850 μm photometry available in the literature. Following Beckwith et al. (1990), it is possible to invert observed millimeter flux into an apparent disk dust mass if we assume that the emission is (1) optically thin, (2) arises from an isothermal region of the disk of known temperature, and (3) is due to material with a known opacity. (See Beckwith et al. 1990 and Mohanty et al. 2013 for a more detailed explanation of this process, and the assumed dust temperatures and dust opacities.) Using the canonical gas-to-dust ratio of 100-to-1, we then converted dust masses into total gas masses. The resulting total disk masses are listed in Table 4. It is important to note that even if we disregard uncertainties in the dust temperature or opacity and the questionable gas-to-dust ratio, these disk masses are likely lower limits. Millimeter observations are only sensitive to small dust grains, less than ~ 1 cm in size. It is possible that substantial mass may be in larger grains, planetesimals, or even protoplanets.

3. RESULTS: DETECTION OF [O I] 63.18 μm AND O-H₂O EMISSION

We detect [O I] 63.18 μm emission from 17 of our 21 transitional disks. Coupling these new results with our reanalysis of select disks from the GASPS sample (Dent et al. 2013; Howard et al. 2013), we report [O I] 63.18 μm emission from 21 of 26 transitional disks, 12 of 18 full disks, and emission from all of the outflow disks. We fit all observed emission lines to Gaussians using an original MATLAB fitting routine. To mitigate noise in the PACS spectrum, we fit the lines over a range of wavelength baselines (the minimum wavelength range: 63.13–63.23 μm ; the maximum wavelength range spanned the entire PACS spectrum: 62.93–63.43 μm). The best-fitting spectrum was deemed as the spectrum closest to the median of all line fits for a given target. The line flux of this best-fitting spectrum was calculated from the (continuum-subtracted) Gaussian line profile (flux = amplitude $\cdot \sigma_{\text{Gaussian}} \sqrt{2\pi}$). For [O I] 63.18 μm non-detections, we derive 3σ upper limits assuming a Gaussian profile with a $3\sigma_{\text{rms}}$ peak height (where σ_{rms} is the standard deviation of the continuum linear-fit) and a 98 km s⁻¹ line width corresponding to the FWHM of an unresolved line in PACS (*PACS Observer's Manual*). Continuum fluxes at 63 μm were also found from the best-fitting Gaussian line profile, as the constant baseline flux term. Continuum emission at 63 μm was detected for all targets, with the exception of DS Tau (an upper limit of 0.037 Jy). The [O I] 63 μm line fluxes and 63 μm continuum fluxes are reported in Table 5, and the spectra are provided in the Appendix. To validate our data reduction, we compared our resulting [O I] 63 μm line fluxes and 63 μm continuum fluxes to the fluxes reported by Howard et al. (2013). Despite using a more recent version of HIPE (version 9, rather than version 4), our fluxes generally agree with those of Howard et al. (2013) to within $\sim 30\%$, which is comparable to the absolute flux accuracy of PACS (which has a peak-to-peak accuracy $\sim 30\%$, and rms accuracy of $\sim 10\%$; *PACS Observer's Manual*). Compared to this pipeline uncertainty, the uncertainties in our line fits are negligible. Representative error bars for both of these types of flux calibration uncertainty are shown in Figures 1(a) and (b).

Typical line fluxes (normalized to the distance of the Taurus-Auriga star-forming region, at 140 pc) are on the order of 10^{-16} – 10^{-17} W m⁻², corresponding to line luminosity of 10^{-7} – $10^{-5} L_{\odot}$. Continuum fluxes (again, normalized to 140 pc) range from 0.1–100 Jy, corresponding to continuum

luminosities ($L_{\text{continuum}} = f_{\nu} v 4\pi d^2$) of 10^{-2} – $1 L_{\odot}$. Figure 1(a) shows the [O I] 63.18 μm line luminosity as a function of 63 μm continuum luminosity for all of our targets. Figure 1(b) shows the ratio of [O I] 63.18 μm line luminosity to 63 μm continuum luminosity as a function of 63 μm continuum luminosity for all of our targets.

We used the Astronomy SURVival package (ASURV; LaValley et al. 1992) to perform linear regressions and correlation tests between the line and continuum luminosities for each subsample. ASURV is particularly useful as it allows for the incorporation of censored data points (i.e., non-detection, line flux upper limits). Compared to the other subsamples, we oversample G-type stars in transitional disks (five G-type transitional disks; one G-type full disk; zero G-type outflow disks). Because of this oversampling and the seemingly chaotic nature of the G-type line and continuum fluxes, we have omitted them from many of our statistical tests.¹¹ Additionally, as discussed in Section 2.1, we also exclude multiple systems where the multiplicity likely strongly affects our *Herschel*/PACS observations. Tables 6–8 summarize the results from a variety of statistics and fitting routines that were used to characterize differences between the three subsamples. There are a number of important trends in our [O I] 63.18 μm line and 63 μm continuum luminosity data.

- [O I] 63.18 μm line luminosities are positively correlated with 63 μm continuum luminosities, both for the sample as a whole and for each individual subsample, as shown in Table 6. This correlation was previously recognized in the *Herschel*/PACS GASPS survey of Taurus-Auriga protoplanetary disks (Howard et al. 2013) and Herbig Ae/Be stars (Meeus et al. 2012), though our study extends this result to a significantly larger sample of transitional disks.
- Outflow disks tend to have [O I] 63.18 μm line luminosities and 63.18 μm continuum luminosities that differ markedly from full disks and transitional disks. This is simply demonstrated in Table 7, which shows that both the line and continuum luminosities of outflow disks are not likely from the same parent population as either the full disks or transitional disks. As shown in Table 8, outflow disks tend to have higher [O I] 63.18 μm line luminosities (by 0.5–1 dex), higher 63 μm continuum luminosities (by ~ 0.5 dex), and higher line-to-continuum luminosity ratios (by ~ 0.5 dex). This was previously recognized by Podio et al. (2012) and Howard et al. (2013).
- Full disks and transitional disks have similar 63 μm continuum luminosities. This is most easily shown in Table 7, which shows that the 63 μm continuum luminosities of transitional disks and full disks are effectively indistinguishable.
- Given the same 63.18 μm continuum luminosity, full disks tend to have larger [O I] 63.18 μm line luminosities than transitional disks,¹² by a factor of ~ 2 . While this is visually evident in Figure 1(a), there is sufficient scatter (and non-detections) to make this difficult to quantify, and the

¹¹ Our oversampling of G-type transitional disks is not intentional. Due to the rarity of transitional disks, we cannot discriminate transitional disks by spectral type in order to populate our transitional disk subsample. Simultaneously, it is difficult to populate subsamples of full or outflow disks with G-type stars from the GASPS Taurus-Auriga survey, since Taurus-Auriga is a low-mass star-forming region.

¹² The one notable exception is BP Tau (C2). BP Tau has a significantly lower [O I] 63.18 μm line luminosity, compared to other full disks

Table 5
Herschel PACS Results

ID	[O I] 63.18 μm Line Flux ($10^{-17} \text{ W m}^{-2}$)	o-H ₂ O 63.32 μm Line Flux ($10^{-17} \text{ W m}^{-2}$)	63 μm Continuum Flux (Jy)
Transition Disks			
T1	2.340 \pm 0.143	\leq 0.716	3.029 \pm 0.010
T2	4.492 \pm 0.200	\leq 1.192	0.596 \pm 0.020
T3	11.284 \pm 0.851	\leq 5.325	2.262 \pm 0.055
T4	1.530 \pm 0.150	\leq 0.658	1.618 \pm 0.015
T5	2.200 \pm 0.134	\leq 0.702	4.017 \pm 0.010
T6	1.276 \pm 0.305	\leq 0.960	0.972 \pm 0.016
T7	0.949 \pm 0.076	\leq 0.541	0.829 \pm 0.013
T8	2.536 \pm 0.189	1.014 \pm 0.145	5.315 \pm 0.012
T9	3.793 \pm 0.513	\leq 2.435	2.810 \pm 0.036
T10	0.764 \pm 0.101	\leq 0.523	0.195 \pm 0.008
T11	1.306 \pm 0.165	\leq 0.770	1.285 \pm 0.012
T12	\leq 1.276	\leq 1.276	13.239 \pm 0.019
T13	1.934 \pm 0.118	\leq 0.534	1.317 \pm 0.009
T14	\leq 1.078	\leq 1.078	0.303 \pm 0.016
T15	0.910 \pm 0.112	\leq 0.540	1.406 \pm 0.008
T16	0.823 \pm 0.140	\leq 0.817	0.652 \pm 0.012
T17	1.390 \pm 0.124	\leq 0.755	0.556 \pm 0.011
T18	0.354 \pm 0.097	\leq 0.637	0.810 \pm 0.008
T19	\leq 0.770	\leq 0.770	0.490 \pm 0.013
T20	1.035 \pm 0.112	\leq 0.730	0.779 \pm 0.016
T21	1.777 \pm 0.100	\leq 0.793	4.199 \pm 0.008
T22	5.455 \pm 0.291	\leq 1.493	7.318 \pm 0.021
T23	4.239 \pm 0.345	\leq 1.556	3.675 \pm 0.024
T24	3.894 \pm 0.266	\leq 1.671	4.021 \pm 0.024
T25	\leq 0.590	\leq 0.590	0.690 \pm 0.009
T26	1.543 \pm 0.296	\leq 2.534	39.540 \pm 0.023
Full Disks			
F1	2.606 \pm 0.109	0.956 \pm 0.131	1.106 \pm 0.016
F2	0.647 \pm 0.146	0.898 \pm 0.173	0.490 \pm 0.024
F3	2.016 \pm 0.183	\leq 0.921	0.979 \pm 0.014
F4	1.489 \pm 0.416	\leq 1.573	0.126 \pm 0.027
F5	0.712 \pm 0.384	\leq 2.150	1.946 \pm 0.039
F6	1.748 \pm 0.155	0.455 \pm 0.157	0.932 \pm 0.012
F7	2.792 \pm 0.208	0.640 \pm 0.176	1.268 \pm 0.012
F8	\leq 1.023	\leq 1.023	0.780 \pm 0.015
F9	2.505 \pm 0.351	\leq 1.413	1.292 \pm 0.021
F10	\leq 0.782	\leq 0.782	\leq 0.037
F11	6.185 \pm 0.381	\leq 1.850	3.756 \pm 0.028
F12	\leq 0.969	\leq 0.969	0.343 \pm 0.019
F13	\leq 1.585	\leq 1.585	0.086 \pm 0.024
F14	3.847 \pm 0.262	\leq 1.234	2.428 \pm 0.023
F15	1.512 \pm 0.256	0.968 \pm 0.294	0.744 \pm 0.019
F16	12.650 \pm 0.336	\leq 1.474	9.043 \pm 0.027
F17	\leq 0.866	\leq 0.866	0.719 \pm 0.014
F18	\leq 1.259	\leq 1.259	0.370 \pm 0.015
Outflow Disks			
O1	9.061 \pm 0.364	\leq 1.601	1.707 \pm 0.024
O2	4.541 \pm 0.333	\leq 1.410	0.369 \pm 0.035
O3	187.160 \pm 3.075	\leq 10.125	18.015 \pm 0.224
O4	80.446 \pm 1.891	\leq 6.355	13.600 \pm 0.113
O5	39.268 \pm 1.971	\leq 7.357	3.932 \pm 0.129
O6	4.069 \pm 1.085	\leq 3.880	0.411 \pm 0.060
O7	2.592 \pm 0.803	\leq 4.039	1.286 \pm 0.057
O8	8.075 \pm 0.397	\leq 2.109	6.240 \pm 0.030
O9	5.272 \pm 0.221	\leq 1.130	1.020 \pm 0.017
O10	10.000 \pm 0.548	\leq 2.385	1.376 \pm 0.034
O11	17.452 \pm 0.701	0.947 \pm 0.320	2.109 \pm 0.043
O12	13.020 \pm 0.640	2.481 \pm 0.740	14.103 \pm 0.049
O13	1348.400 \pm 14.564	46.876 \pm 1.506	161.870 \pm 0.966
O14	38.197 \pm 0.828	1.660 \pm 0.389	6.650 \pm 0.058
O15	2.573 \pm 0.554	\leq 2.789	1.062 \pm 0.046

Note. Detections are listed with $\pm 1\sigma$ uncertainties; 3σ upper limits are reported for non-detections.

Table 6
Subsample Correlation Tests

Correlation Test	Subsample Being Tested	Correlation Tests			Correlated?	Linear Regression	
		$P(1)$	$P(2)$	$P(3)$		Intercept	Slope
$L_{63\ \mu\text{m}}$ v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	-3.24 ± 0.18	1.15 ± 0.13
	Transitional Disks Only	0.0%	0.0%	0.1%	Correlated	-4.07 ± 0.21	0.74 ± 0.15
	Full Disks Only	10.0%	0.7%	3.3%	Correlated	-4.42 ± 0.31	0.38 ± 0.18
	Outflow Disks Only	0.2%	0.2%	0.5%	Correlated	-3.01 ± 0.17	0.97 ± 0.15
T_{eff} v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.1%	0.3%	0.4%	Correlated	-7.66 ± 0.93	0.0007 ± 0.0002
	Transitional Disks Only	0.0%	0.1%	0.2%	Correlated	-6.67 ± 0.39	0.0004 ± 0.0001
	Full Disks Only	82.7%	84.7%	87.3%	Not Correlated	-4.68 ± 1.15	-0.0001 ± 0.0003
	Outflow Disks Only	24.7%	29.7%	33.0%	Not Correlated	-6.22 ± 1.70	0.0005 ± 0.0004
T_{eff} v. $L_{63\ \mu\text{m}}$	All Objects	10.9%	0.4%	1%	Not Correlated	-3.09 ± 0.73	0.0004 ± 0.0002
	Transitional Disks Only	0.0%	0.1%	0.2%	Correlated	-3.17 ± 0.30	0.0004 ± 0.0001
	Full Disks Only	10.4%	28.4%	18.0%	Not Correlated	1.81 ± 1.63	-0.0009 ± 0.0004
	Outflow Disks Only	11.8%	14.4%	15.6%	Not Correlated	-3.79 ± 1.45	0.0007 ± 0.0003
L_{bol} v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	-4.61 ± 0.10	1.02 ± 0.19
	Transitional Disks Only	0.1%	0.3%	0.3%	Correlated	-4.96 ± 0.07	0.41 ± 0.11
	Full Disks Only	28.3%	45.7%	38.3%	Not Correlated	-4.94 ± 0.19	0.39 ± 0.50
	Outflow Disks Only	3.0%	7.3%	4.7%	Correlated	-4.04 ± 0.20	0.75 ± 0.35
L_{bol} v. $L_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	-1.23 ± 0.09	0.78 ± 0.15
	Transitional Disks Only	0.0%	0.1%	0.2%	Correlated	-1.23 ± 0.06	0.52 ± 0.09
	Full Disks Only	29.0%	32.9%	36.9%	Not Correlated	-1.60 ± 0.39	0.64 ± 0.99
	Outflow Disks Only	0.2%	0.4%	0.8%	Correlated	-1.08 ± 0.14	0.97 ± 0.25
L_X v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	5.9%	59.0%	80.7%	Not Correlated	-5.59 ± 0.50	-0.18 ± 0.13
	Transitional Disks Only	6.4%	56.9%	53.7%	Not Correlated	-5.62 ± 0.29	-0.14 ± 0.08
	Full Disks Only	65.5%	92.0%	...	Not Correlated	-5.35 ± 0.79	-0.06 ± 0.20
	Outflow Disks Only	9.1%	39.5%	54.8%	Not Correlated	-2.69 ± 0.64	0.40 ± 0.18
L_X v. $L_{63\ \mu\text{m}}$	All Objects	3.0%	69.6%	13.9%	Not Correlated	-2.21 ± 0.36	-0.18 ± 0.10
	Transitional Disks Only	0.3%	37.1%	27.6%	Not Correlated	-2.32 ± 0.34	-0.24 ± 0.09
	Full Disks Only	86.4%	73.2%	80.9%	Not Correlated	-2.03 ± 0.58	-0.03 ± 0.15
	Outflow Disks Only	0.8%	8.9%	18.5%	Not Correlated	0.59 ± 0.53	0.47 ± 0.15
L_{acc} v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	-4.22 ± 0.15	0.66 ± 0.12
	Transitional Disks Only	1.0%	4.7%	4.9%	Correlated	-4.76 ± 0.15	0.28 ± 0.11
	Full Disks Only	3.5%	17.6%	16.1%	Not Correlated	-4.69 ± 0.27	0.34 ± 0.22
	Outflow Disks Only	5.7%	5.2%	8.6%	Not Correlated	-3.88 ± 0.23	0.41 ± 0.22
L_{acc} v. $L_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	-0.97 ± 0.14	0.47 ± 0.10
	Transitional Disks Only	0.3%	4.1%	2.8%	Correlated	-1.06 ± 0.14	0.29 ± 0.09
	Full Disks Only	3.6%	19.9%	18.2%	Not correlated	-1.21 ± 0.61	0.54 ± 0.44
	Outflow Disks Only	0.7%	1.6%	1.2%	Correlated	-0.85 ± 0.19	0.48 ± 0.18
\dot{M} v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.0%	0.1%	0.1%	Correlated	-0.44 ± 0.94	0.55 ± 0.12
	Transitional Disks Only	4.0%	13.5%	12.1%	Not Correlated	-3.03 ± 1.15	0.25 ± 0.14
	Full Disks Only	7.0%	36.8%	25.2%	Not Correlated	-2.68 ± 1.66	0.30 ± 0.20
	Outflow Disks Only	10.8%	1.5%	4.5%	Correlated	-1.91 ± 1.44	0.28 ± 0.19
\dot{M} v. $L_{63\ \mu\text{m}}$	All Objects	0.0%	0.0%	0.0%	Correlated	2.12 ± 0.81	0.44 ± 0.10
	Transitional Disks Only	0.6%	1.2%	2.5%	Correlated	1.24 ± 1.01	0.32 ± 0.12
	Full Disks Only	0.7%	4.4%	4.3%	Correlated	4.50 ± 3.29	0.77 ± 0.40
	Outflow Disks Only	1.2%	1.0%	1.0%	Correlated	1.56 ± 1.21	0.35 ± 0.16
m_{disk} v. $L[\text{O I}]_{63\ \mu\text{m}}$	All Objects	0.0%	26.5%	43.6%	Not Correlated	-5.79 ± 0.25	0.04 ± 0.01
	Transitional Disks Only	0.3%	26.6%	20.0%	Not Correlated	-4.72 ± 0.18	0.03 ± 0.01
	Full Disks Only	1.9%	44.7%	8.2%	Not Correlated	-5.25 ± 0.13	0.01 ± 0.01
	Outflow Disks Only	0.0%	4.4%	2.3%	Correlated	-4.78 ± 0.26	0.03 ± 0.01
m_{disk} v. $L_{63\ \mu\text{m}}$	All Objects	0.0%	5.7%	7.9%	Correlated	-2.14 ± 0.16	0.03 ± 0.01
	Transitional Disks Only	0.6%	7.7%	5.6%	Correlated	-2.13 ± 0.25	0.03 ± 0.01
	Full Disks Only	6.6%	95.1%	86.7%	Not Correlated	-2.05 ± 0.20	0.02 ± 0.01
	Outflow Disks Only	0.0%	4.4%	13.0%	Correlated	-1.71 ± 0.24	0.03 ± 0.01
a_{cavity} v. $L[\text{O I}]_{63\ \mu\text{m}}$	Transitional Disks Only	71.1%	77.0%	69.5%	Not Correlated	-5.10 ± 0.17	-0.0012 ± 0.01
a_{cavity} v. $L_{63\ \mu\text{m}}$	Transitional Disks Only	75.0%	24.6%	26.5%	Not Correlated	-1.46 ± 0.16	0.00 ± 0.01
h_{wall} v. $L[\text{O I}]_{63\ \mu\text{m}}$	Transitional Disks Only	23.6%	78.5%	64.7%	Not Correlated	-5.25 ± 0.20	0.04 ± 0.04
h_{wall} v. $L_{63\ \mu\text{m}}$	Transitional Disks Only	31.0%	23.6%	24.5%	Not Correlated	-1.65 ± 0.18	0.07 ± 0.04

Notes. P is the probability that the correlation between the two listed parameters is obtained by chance; low P values indicate a correlation. The different statistical tests used are (1) Cox Hazard, (2) Kendall Tau, and (3) Spearman Rho. If the average of the three statistical tests is less than 5%, they are listed as “correlated,” in boldface. A linear regression (using the EM method) was performed for all combinations, fitting the log of the quantities listed, where the first parameter listed for each pair is the independent variable. If no correlation is detected, the linear regression may not be significant.




Figure 1. (a) [O I] $63.18 \mu\text{m}$ line luminosity as a function of $63 \mu\text{m}$ continuum luminosity for our sample of transitional disks (red), full disks (blue), and outflow disks (green). upper limits of 3σ are denoted by hollow data points with arrows. Symbols correspond to stellar spectral types: circles are G-type stars (which are included in this plot but neglected in the statistical analysis, for reasons described in the paper), squares are K-type stars, and diamonds are M-type stars. BP Tau (an evolved full disk) is indicated in purple. Targets excluded from statistical tests (for either being a binary that does not meet the criteria in Section 2.1, or being a G-type star) are marked by an asterisk. (b) The ratio of [O I] $63.18 \mu\text{m}$ line luminosity/ $63 \mu\text{m}$ continuum luminosity as a function of $63 \mu\text{m}$ continuum luminosity for our sample of transitional disks (red), full disks (blue), and outflow disks (green). upper limits of 3σ are denoted by hollow data points with arrows. Symbols are as in panel (a). (A color version of this figure is available in the online journal.)

ASURV statistical tests point to indistinguishable line luminosities between the two subsamples (see Tables 7 and 8). However, this difference between full disks and transitional disks becomes clear when we examine the ratio of the [O I] $63.18 \mu\text{m}$ line luminosity to the $63 \mu\text{m}$ continuum luminosity, as shown in Figure 1(b). The ASURV statistical tests indicate that the distribution of line-to-continuum ratios of transitional disks is significantly different from that of full disks (see Tables 7 and 8), with full disks having line-to-continuum ratios larger by a factor of ~ 2 .

Additionally, the best-fit linear regressions in the line luminosity for full disks and transitional disks, as shown in Table 6, are distinct. Transitional disks have steeper best-fit slopes and shallower best-fit intercepts than full disks; both of these effects contribute to larger differences in line luminosity at the relevant continuum luminosities. We checked our ASURV fit results with an alternative Bayesian metric (linmix_err.pro; Kelly 2007) and found similar differences between transitional disks and full disks.

Table 7
Subsample Statistical Difference Tests

Parameter	Subsamples Being Compared	Statistical Difference Tests					Different?
		$P(1)$	$P(2)$	$P(3)$	$P(4)$	$P(5)$	
L [O I] 63 μm	Transitional vs. Full Disks	98.4%	98.4%	78.7%	93.8%	94.1%	Not Different
	Transitional vs. Outflow Disks	0.0%	0.0%	0.0%	0.0%	0.0%	Different
	Full vs. Outflow Disks	0.0%	0.0%	0.0%	0.0%	0.0%	Different
L 63 μm	Transitional vs. Full Disks	9.8%	10.2%	4.8%	9.8%	10.0%	Not Different
	Transitional vs. Outflow Disks	2.1%	1.4%	5.9%	5.9%	...	Different
	Full vs. Outflow Disks	0.1%	0.0%	0.0%	0.1%	0.0%	Different
L [O I] 63 μm / L 63 μm	Transitional vs. Full Disks	2.2%	1.0%	3.0%	3.1%	1.5%	Different
	Transitional vs. Outflow Disks	0.0%	0.0%	0.0%	0.0%	0.0%	Different
	Full vs. Outflow Disks	1.8%	1.7%	4.0%	2.3%	2.0%	Different
T_{eff}	Transitional vs. Full Disks	78.9%	79.1%	50.1%	50.1%	...	Not Different
	Transitional vs. Outflow Disks	30.8%	31.8%	45.2%	45.2%	...	Not Different
	Full vs. Outflow Disks	11.4%	11.5%	10.3%	10.3%	...	Not Different
L_{bol}	Transitional vs. Full Disks	33.7%	34.3%	76.4%	76.4%	...	Not Different
	Transitional vs. Outflow Disks	2.5%	2.8%	1.7%	1.7%	...	Different
	Full vs. Outflow Disks	3.7%	3.4%	0.5%	0.5%	...	Different
L_X	Transitional vs. Full Disks	94.3%	94.4%	68.0%	99.0%	1.9%	Not Different
	Transitional vs. Outflow Disks	74.6%	75.1%	97.1%	73.5%	74.9%	Not Different
	Full vs. Outflow Disks	94.8%	94.8%	48.9%	94.7%	96.3%	Not Different
L_{acc}	Transitional vs. Full Disks	68.8%	69.1%	96.2%	68.9%	69.1%	Not Different
	Transitional vs. Outflow Disks	1.9%	1.0%	5.4%	1.9%	1.3%	Different
	Full vs. Outflow Disks	1.4%	0.9%	8.5%	1.4%	0.9%	Different
\dot{M}	Transitional vs. Full Disks	100.0%	100.0%	90.0%	99.7%	99.7%	Not Different
	Transitional vs. Outflow Disks	1.7%	0.8%	4.1%	1.6%	1.0%	Different
	Full vs. Outflow Disks	1.9%	1.3%	5.2%	2.0%	1.5%	Different
m_{disk}	Transitional vs. Full Disks	59.6%	59.8%	42.6%	53.6%	54.5%	Not Different
	Transitional vs. Outflow Disks	50.7%	50.5%	50.7%	48.3%	48.7%	Different
	Full vs. Outflow Disks	75.4%	75.4%	99.8%	75.4%	75.3%	Different

Notes. P is the probability that the parameter being compared between two subsamples is drawn from the same parent distribution; low P values indicate that two subsamples are different. The different statistical tests are (1) Gehan generalized Wilcoxon test (with permutation variance); (2) Gehan generalized Wilcoxon test (with hypergeometric variance); (3) logrank test; (4) Peto & Peto generalized Wilcoxon test; (5) Peto & Prentice generalized Wilcoxon test. If the average of the five statistical tests is less than 5%, they are listed as “different,” in boldface.

This difference between full disks and transitional disks was previously recognized by Howard et al. (2013) for the GASPS Taurus-Auriga sample only. Our data extends this trend to a much larger sample of transitional disks, suggesting that this lower [O I] 63.18 μm line emission is a characteristic property of transitional disks.

- There is a weak trend for M-type stars to have lower line and continuum luminosities than K-type stars. This trend is most evident in our sample of transitional disks.

Generally, the [O I] 63.18 μm line is spectrally unresolved. Most FWHM are within $\sim 11 \text{ km s}^{-1}$ (the native resolution of PACS) of the expected line width for an unresolved line for PACS ($\sim 98 \text{ km s}^{-1}$ at 60 μm). This result was expected, since [O I] emission originates far out in the disk, $\sim \text{AU}$ from the central star (Woitke et al. 2010). For gas orbiting a Sun-like star, Keplerian velocities go as $V_{\text{Keplerian}} = 30 \text{ km s}^{-1} \cdot (a/\text{AU})^{-1/2}$. Thus, beyond $\sim \text{AU}$ from the central star, we expect line widths on the order of ~ 10 's of km s^{-1} . Even at these distances, Keplerian velocities dominate over thermal velocities ($\sim 1 \text{ km s}^{-1}$, assuming typical O I 63.18 μm gas temperatures of $\sim 100 \text{ K}$; Aresu et al. 2012), or turbulent velocities ($\sim 1 \text{ km s}^{-1}$; Hughes et al. 2011), and are the cause of most of the line broadening. A few objects, all outflow sources, have broader

line widths, as high as 170 km s^{-1} (e.g., RW Aur). In these sources, [O I] 63.18 μm emission is thought to originate from shocks along the jet and/or UV-heated gas in the outflow cavity walls (Podio et al. 2012). Line widths of ~ 100 's of km s^{-1} reflect the similarly large shock velocities. Outflow disks can also have spatially extended [O I] 63.18 μm emission associated with the jet, which is detectable in non-central PACS spaxels (Podio et al. 2012). We verified that [O I] 63.18 μm emission was localized only in the central spaxel for our transitional and full disks. For outflow disks, we only report [O I] 63.18 μm line and 63 μm continuum fluxes from the central spaxel (for more accurate line and continuum fluxes of outflow disks, including neighboring spaxels, see Podio et al. 2012).

Our PACS spectral range fortuitously also includes the considerably fainter o-H₂O 63.32 μm emission line. We confirmed the detection o-H₂O emission in five full disks and outflow disks, previously identified by Riviere-Marichalar et al. (2012). In addition, we report the marginal detections of o-H₂O in IQ Tau, DK Tau, and BP Tau—for which Riviere-Marichalar et al. (2012) previously identified 3σ upper limits. These new detections, from the same observational data, are made possible with our updated version of the Herschel HIPE pipeline and a different line-fitting algorithm. In addition to these objects, we also report the detection of o-H₂O emission from RW Aur (observed

Table 8
Mean Parameter Values

Parameter	Subsample	Kaplan–Meier Estimator Mean \pm Standard Deviation
L [O I] 63 μm	Transitional Disks	$-5.12 \pm 0.07 \log L_{\odot}$
	Full Disks	$-5.09 \pm 0.07 \log L_{\odot}$
	Outflow Disks	$-3.89 \pm 0.22 \log L_{\odot}$
L 63 μm	Transitional Disks	$-1.45 \pm 0.08 \log L_{\odot}$
	Full Disks	$-1.86 \pm 0.17 \log L_{\odot}$
	Outflow Disks	$-0.91 \pm 0.20 \log L_{\odot}$
$\log(L \text{ [O I] } 63 \mu\text{m} / L \text{ 63 } \mu\text{m})$	Transitional Disks	-3.71 ± 0.05
	Full Disks	-3.39 ± 0.08
	Outflow Disks	-2.99 ± 0.10
T_{eff}	Transitional Disks	$4066 \pm 650 \text{ K}$
	Full Disks	$3976 \pm 399 \text{ K}$
	Outflow Disks	$4299 \pm 534 \text{ K}$
L_{bol}	Transitional Disks	$-0.43 \pm 0.12 \log L_{\odot}$
	Full Disks	$-0.34 \pm 0.05 \log L_{\odot}$
	Outflow Disks	$0.13 \pm 0.17 \log L_{\odot}$
L_X	Transitional Disks	$-3.89 \pm 0.18 \log L_{\odot}$
	Full Disks	$-3.91 \pm 0.11 \log L_{\odot}$
	Outflow Disks	$-3.78 \pm 0.21 \log L_{\odot}$
L_{acc}	Transitional Disks	$-1.51 \pm 0.22 \log L_{\odot}$
	Full Disks	$-1.39 \pm 0.16 \log L_{\odot}$
	Outflow Disks	$-0.34 \pm 0.32 \log L_{\odot}$
\dot{M}	Transitional Disks	$-8.60 \pm 0.17 \log M_{\odot}/\text{yr}$
	Full Disks	$-8.42 \pm 0.18 \log M_{\odot}/\text{yr}$
	Outflow Disks	$-7.39 \pm 0.36 \log M_{\odot}/\text{yr}$
m_{disk}	Transitional Disks	$10.38 \pm 3.09 M_{\text{Jupiter}}$
	Full Disks	$12.21 \pm 3.19 M_{\text{Jupiter}}$
	Outflow Disks	$18.10 \pm 6.25 M_{\text{Jupiter}}$

Notes. The Kaplan–Meier estimator provides an estimate of the mean and standard deviation of the quantity measured, while taking data censoring into account. See LaValley et al. (1992) for more.

by the GASPS survey, but not included in Riviere-Marichalar et al. (2012) and the first detection of o-H₂O 63.32 μm emission from a transitional disk: DoAr44 (original to this study). o-H₂O line fluxes and 3σ upper limits for all targets are reported in Table 5.

4. TRENDS WITH OBSERVABLE STELLAR AND DISK PROPERTIES

In this section, we compare our [O I] 63.18 μm line flux results with stellar and disk properties summarized in Section 2.3 in order to identify the origin of the trends described in the previous section. We performed correlation tests between [O I] 63.18 μm and all of these disk/star parameters using the ASURV (LaValley et al. 1992). Tables 6–8 summarize the results of these correlation tests.

4.1. Effective Temperature and Bolometric Luminosity

Since the [O I] 63.18 μm line is generally optically thick (e.g., Aresu et al. 2012), it would be expected that the line flux might increase for increasing stellar effective temperature. Similarly, one might expect that the bolometric luminosity of the host star may affect the line flux. As shown in Table 6, we indeed find a correlation between [O I] 63.18 μm line flux and the effective temperature and bolometric luminosity of the host

star for transitional disks. We also find a correlation between the 63.18 μm continuum flux and the effective temperature and bolometric luminosity. Curiously, we do not find either of these correlations for our sample of full disks. This may be a result of the smaller span of effective temperature and bolometric luminosity covered by full disks, compared to transitional disks.

The observed correlations between bolometric luminosity and effective temperature with [O I] 63.18 μm line flux and 63.18 μm continuum flux in our transitional disk sample are expected on the basis that both the line and continuum emission are expected to be optically thick and thus sensitive primarily to the disk temperature. This explains why the line and continuum emission are correlated with each other, as they both increase with increasing temperature. This correlation (though weaker) was also observed by Meeus et al. (2012), for their smaller sample of Herbig Ae/Be stars. This relationship is visually evident in Figure 1(a), where the symbol of each data point is representative of the star’s spectral type; generally cooler, M-type stars have lower line and continuum fluxes than K-type stars. What is more important, however, is that the effective temperatures and bolometric luminosities between full disks and transitional disks are not statistically different, as illustrated in Tables 7 and 8. This similarity was expected since we attempted to uniformly sample across spectral types within each subsample. This suggests that the effective temperature and bolometric luminosity alone are not enough to explain why full disks have systematically larger [O I] 63.18 μm line fluxes than transitional disks.

4.2. FUV Luminosities and Accretion Rates

Pinte et al. (2010) used disk thermo-chemical models and showed that far-ultraviolet (FUV) radiation can be a significant gas heating mechanism and can promote [O I] 63.18 μm line emission. For low-mass stars, where chromospheric FUV is negligible, most of the FUV luminosity is generated from the infall of disk material onto the central star. This accretion process shocks and superheats the gas, generating FUV emission, which can then heat the surface layers of the surrounding disk. Indeed, we find a correlation between FUV¹³ and both [O I] 63.18 μm line emission and 63.18 μm continuum emission in Table 6. However, for our sample, we find that transitional and full disks have statistically indistinguishable FUV¹³ luminosities, as shown in Tables 7 and 8. Thus, FUV cannot be responsible for the [O I] line flux differences between full and transitional disks.

The literature is not conclusive about any accretion rate difference between transitional disks and full disks. Some studies find that transitional disks have accretion rates an order of magnitude lower than full disks (Najita et al. 2007; Espaillat et al. 2012; Kim et al. 2013). However, when the two samples are drawn from the same spectral type distribution, and the accretion rates are self-consistently derived (and not drawn from the literature), as in our study, no differences are found (Fang et al. 2009). This is illustrated in Tables 7 and 8.

4.3. X-Ray Luminosities

More recent thermo-chemical disk models by Aresu et al. (2011, 2012, 2014) have included the effects of irradiation from stellar X-rays. Aresu et al. (2012) found that X-ray irradiation tends to become a significant driver for the [O I] 63.18 μm

¹³ L_{FUV} is directly related to L_{acc} via Equations (1) and (2), so statistical tests for the two are identical. In all tables, we only list L_{acc} .

line emission only when $L_X > 10^{30}$ erg s⁻¹. Below this limit, FUV irradiation dominates. As shown in Table 6, we find no correlation between L_X and either [O I] 63.18 μ m line luminosity or 63 μ m continuum luminosity. We also find that full disks and transitional disks have statistically indistinguishable X-ray luminosities, as shown in Tables 7 and 8. Furthermore, the observed X-ray luminosities are generally lower than the 10^{30} erg s⁻¹ limit suggested by Aresu et al. (2012), suggesting that X-ray irradiation is not the driving mechanism for the trends between [O I] 63.18 μ m line luminosity in full disks and transitional disks. Newer models by Meijerink et al. (2012) and Aresu et al. (2012) have predicted a correlation between [O I] 63.18 μ m line luminosity and the *sum* of the X-ray luminosity and FUV luminosity, although this trend is not found in either the GASPS Taurus sample (Aresu et al. 2014) or in our larger sample of transitional disks.

While X-rays may not be important for the differences between subsamples, X-ray irradiation may be important for a few of the G-type stars. As noted previously, the G-type stars in our sample tend to have line and continuum fluxes that differ significantly from our other targets. Many of these G-type outliers (e.g., CHX 22, CHX 7, YLW8) have X-ray luminosities at or above the 10^{30} erg s⁻¹ L_X limit of Aresu et al. (2012).

4.4. Disk Structure and Disk Mass

Is the lower [O I] 63.18 μ m line luminosity in transitional disks simply due to the lack of gas in the inner cavity of transitional disks? Kamp et al. (2010) showed for a small number of thermo-chemical disk models that creating an inner cavity (out to 10 AU) completely devoid of gas decreased the [O I] 63.18 μ m line flux by a factor of 1.5. Bruderer (2013) has shown that even though most [O I] emission originates from the outer disk (beyond 10's of AU), depleting gas within the inner cavity of transitional disks can reduce the disk's total [O I] 63.18 μ m luminosity by factors of up to several. Given the large beam of *Herschel*/PACS (~ 1000 's of AU, at the distance of Taurus), a reduction in the [O I] 63.18 μ m line flux by a factor of ~ 2 as we observe would require a depletion of gas in the inner disk by a factor $\gtrsim 100$ (see Figure 18 of Bruderer 2013). These scenarios proposed by Kamp et al. (2010) and Bruderer (2013) seem unlikely in view of our finding that the mass accretion rates of transitional disks in our sample are statistically indistinguishable from those of full disks in Taurus.

One might expect that the heating of the gas, and by extension the luminosity of the [O I] 63.18 μ m line, is affected by the distribution of the dust, hence a correlation between the dust cavity size or wall height and the [O I] 63.18 μ m line. We find no correlation between the cavity sizes or wall heights of transitional disks and either the [O I] 63.18 μ m line luminosities or 63.18 μ m continuum luminosities. This result is shown in Table 6 and graphically in Figure 2. This suggests that either [O I] 63.18 μ m is tracing material well beyond the inner cavity and/or the distribution of gas in the inner cavities of full disks and transitional disks is similar. While this result is suggestive, more work following Kamp et al. (2010) and Bruderer (2013) needs to be done to determine the relationship between the size of transitional disk cavities, the gas-to-dust ratio in these cavities, and [O I] 63.18 μ m emission.

Finally, we find that the disk dust masses for full and transitional disks are statistically indistinguishable and that there is no correlation with the dust mass and either the [O I] 63.18 μ m line luminosities or 63 μ m continuum luminosities. This is

expected, given that both the [O I] 63.18 μ m line and 63 μ m continuum emission are mostly optically thick.

5. TRENDS WITH MODEL-DERIVED DISK PROPERTIES

Thus far, we have been unable to satisfactorily explain the difference in [O I] 63.18 μ m line flux between full and transitional disks. To identify other possible causes for the trends we used the DENT (“Disk Evolution with Neat Theory”) grid of thermo-chemical models by Woitke et al. (2010) to look for correlations between [O I] 63.18 μ m line emission and various disk properties, that are not directly observable. Of the free parameters in the DENT grid (e.g., column density, disk inner/outer radius, grain sizes, inclination, etc.), there are only two parameters that can cause the observed [O I] 63.18 μ m line flux trends: disk flaring, and the disk gas-to-dust ratio.

To illustrate trends within the DENT grid, we have developed a novel approach for analyzing the large suite of DENT disk models (totaling over 300,000 unique disks). Figure 3 illustrates an example of this technique, for the case where we investigate how [O I] 63.18 μ m line and 63.18 μ m continuum emission change as a function of FUV excess luminosity (which is discussed previously in Section 4.2). Using an original MATLAB script, we select a randomized subsample of a few thousand¹⁴ unique disk models from the full DENT grid. Generally, we constrain this randomized subsample to consist of low-mass stars (M and K type), similar to our sample of transitional and full disks. Next, for each of the selected disk models, we search the full DENT grid for all of the disk models that possess identical stellar/disk properties, *except* for the quantity that we are interested in—FUV excess in this case. Since the DENT grid allows for two different FUV excesses (0.001 and 0.1 L_{star}), this results in a few thousand *pairs*¹⁵ of disks.

From this ensemble of disk model pairs, we can perform a number of analyses. Figure 3(a) shows the ensemble of disk models in a plot of [O I] 63.18 μ m line flux versus 63.18 μ m continuum flux,¹⁶ similar to Figure 1(a). In this plot, each of the disk models is represented by a colored point, with the color corresponding to its FUV excess. The vectors connect individual disk pairs. These vectors can be thought of as “evolutionary tracks” which show how one disk would change if the FUV excess changed (in this case, the arrow points in the direction of increasing FUV excess). Due to the extreme number of disk models in the DENT grid, even the randomized subsample in Figure 3(a) is dense and difficult to interpret. To simplify interpretation, Figure 3(b) displays two contour intervals—one for each FUV excess—indicating the region that contains 67% of the disk models for that particular FUV excess. These contours are generated by binning the data in both continuum flux and line flux space (usually with bins 0.25 dex in size). A small number of “evolutionary tracks” are included, to reinforce

¹⁴ In general, our results are not sensitive to the number of disk models selected, as long as it is fairly large ($\gtrsim 100$).

¹⁵ For other stellar/disk parameters where more than two values are possible, we form sets containing the same number of disk models as the number of possible values for that stellar/disk parameter. For example, there are five possible gas-to-dust ratios within the DENT grid; thus, when performing our analysis for gas-to-dust ratios, we form several thousand sets of disk models, each containing five disks that are identical with the exception of their gas-to-dust ratios.

¹⁶ It is not possible to exactly duplicate Figure 1(a) with the DENT grid, as the DENT grid does not include a 63 μ m photometric point. Instead, we used the 65 μ m photometric point as a proxy. For most DENT models, there is not a significant change in the continuum luminosity between 60, 65 or 70 μ m.




Figure 2. [O I] 63.18 μm line luminosity and 63 μm continuum luminosity as a function of gap size and wall height for all of the transitional disks within our sample for which such measurements have been made in the past. 3σ upper limits are denoted by hollow data points with arrows. Symbols are as in Figure 1(a). (An extended, color version of this figure is available in the online journal.)

the concept that we are tracking disk models as a particular quantity is changed. Lastly, Figure 3(c) illustrates the mean “evolutionary track” for all of our disk pairs. To generate this figure, we take each pair and calculate the change (signified by a “ Δ ” in the figure axes) in continuum flux and change in line flux between each pair member. The vector displayed represents the mean change in continuum and line flux for our entire ensemble of disk pairs. The error bars indicate the 1σ variations in this single step-up in FUV excess. This last figure, Figure 3(c) is particularly useful, as it shows information that is easily lost in the large apparent scatter in Figures 3(a) and (b). For example, while it is clear in these other figures that increasing the FUV excess increases the line flux, it is not as obvious how much this line flux changes, and the relative uncertainties. It’s also not obvious in the other figures that the change in continuum flux is so consistent (represented by the very small horizontal error bars) between all the DENT models. In the following sections, we will make use of figures similar to Figures 3(b) and (c) to investigate how changing various stellar/disk parameters affect the observed [O I] 63.18 μm line fluxes and 63.18 μm continuum fluxes.

5.1. Disk Flaring

One of the early predicted trends of the DENT grid was that [O I] 63.18 μm emission may trace the flaring of the disk

(Woitke et al. 2010). In a flared disk, the disk surface is directly illuminated by the central star, causing higher temperatures and stronger [O I] 63.18 μm emission. Within the DENT model grid, geometric flaring of the gas disk is parameterized.¹⁷ By the value of β : the disk scale height, h , as a function of radial distance, r , can be described by

$$h(r) = h_0 \left(\frac{r}{r_0} \right)^\beta, \quad (4)$$

where h_0 is the disc scale height (fixed at 10 AU), and r_0 is a fixed reference distance (fixed at 100 AU). In the DENT grid, there are three possible flaring parameters, $\beta = 0.8$, $\beta = 1.0$, and $\beta = 1.2$. A flaring parameter as low as $\beta = 0.8$ is more appropriate for the late stages of disk evolution (e.g., debris disks) and not for our study of young protoplanetary disks (Kamp et al. 2011). Thus, we have excluded models with $\beta = 0.8$ from our analysis. It is important to note that when we refer to “flaring,” we are referring to the flaring of the *gas* disk.

¹⁷ In principle, the vertical scale height of the gas disk should be self-consistently derived from hydrostatic equilibrium, given the temperature structure of the disk. This is *not* done in the DENT grid. Using parameterized disk structures allows for a wider, unbiased exploration of disk parameter space while still assessing the relative influence of key parameters on observable quantities. See Section 2 of Kamp et al. (2011) for a discussion of the parameterized approach.




Figure 3. DENT grid predictions for how [O I] 63.18 μm line flux and 65 μm continuum flux change, with increasing disk FUV excess (FUV = 0.001 and 0.1), for a random subsample ($N \sim 5000$) of disks around low-mass stars ($\leq 1 M_{\text{Sun}}$). The left panel shows all of the “evolutionary tracks” for this sample of disk models. The middle panel shows the regions that contain 67% of the models as a function of FUV. The “evolutionary tracks” of 10 randomly selected disk models are included for reference. These tracks indicate the path that that particular disk would move if the FUV increased. The right panel shows the mean change (“delta”) in [O I] 63.18 μm line flux and 65 μm continuum flux, with respect to an initially low FUV disk. Arrows point in the direction of increasing FUV. Error bars indicate the 1σ variations in these Δ [O I] 63.18 μm line flux and Δ 65 μm continuum flux during each step in increasing FUV.

(A color version of this figure is available in the online journal.)

In the DENT grid, the dust is either well mixed with the gas or settled. The dust disk scale height, h_{dust} , is parameterized as

$$h_{\text{dust}}(r, a) \propto h(r)a^{-s/2}, \quad (5)$$

where h is the gas scale height (Equation (4)), a is the grain size, and s describes the strength of dust settling: $s = 0$ for a well-mixed disk, and $s = 0.5$ for a settled disk. We find that within the DENT grid, there is no systematic difference between the [O I] 63.18 μm line luminosity of disks with either well-mixed or settled dust disks. Settled dust disks do have systematically lower 63 μm continuum luminosities than well-mixed disks, by ~ 0.7 dex. Instead, we focus on the effects of changing the flaring of the gas disk.

Figure 4 illustrates how the [O I] 63.18 μm line luminosity and 63 μm continuum luminosity change as disks become more flared according to the DENT grid. From Figure 4, we can see that increasing the disk flaring from $\beta = 1.0$ to $\beta = 1.2$ can increase the [O I] 63.18 μm line luminosity by ~ 0.5 dex, while not significantly altering the 63 μm continuum luminosity. This increase in flaring in the DENT grid results in generally warmer gas in the disk surface, resulting in larger [O I] 63.18 μm line luminosities. Since changing the flaring of the disks only changes the [O I] 63.18 μm line luminosity, and not the 63 μm continuum luminosity, this may provide a natural explanation for the decreased [O I] 63.18 μm line luminosity in transitional disks compared to full disks. This would imply that transitional disks are *less* flared than full disks and that their lower [O I] 63.18 μm line luminosities are the result of cooler disk gas surface layers. If the gas in transitional disks is indeed cooler than in full disks,

this might be linked to the reduction or removal of some gas heating mechanism. Aikawa & Nomura (2006) have shown that growth and settling of larger dust grains (~ 10 cm in diameter) leads to decreased photoelectric heating in the disk atmosphere and less disk flaring. However, these large dust grains will quickly settle toward the disk midplane, resulting in reduced far-infrared emission, which we do not see in our sample. An alternative explanation could be that the stellar FUV photons responsible for heating the [O I] emitting disk surface layers (Aresu et al. 2012) are being absorbed at a vertically extended dust inner rim. Future SED modeling may be able to disentangle these two possibilities.

5.2. Disk Gas-to-dust Ratio

A second, though less well-recognized trend in the DENT grid is that [O I] 63.18 μm emission may trace the disk gas-to-dust ratio. From *Herschel*/PACS and millimeters observations combined with dust and gas modeling, Thi et al. (2010) suggested that the transitional disk TW Hya possesses a lower gas-to-dust ratio than the standard interstellar value of 100, though this suggestion has been disputed in recent years (Gorti et al. 2011; Bergin et al. 2013). Meeus et al. (2012) have also suggested, from analysis of the DENT grid, that variations in the [O I] 63.18 μm line luminosities of Ae/Be stars, could be a result of variations in the gas-to-dust ratio, although they do not explore this further.

Figures 5 and 6 illustrate how the [O I] 63.18 μm line luminosity and 63 μm continuum luminosity change as the gas-to-dust ratio changes within the DENT grid. We consider two




Figure 4. DENT grid predictions for how [O I] 63.18 μm line flux and 65 μm continuum flux change, with increasing disk flaring ($\beta = 1.0$ and $\beta = 1.2$), for a random subsample ($N \sim 5000$) of disks around low-mass stars ($\leq 1 M_{\text{Sun}}$). The left panel shows the regions that contain 67% of the models as a function of disk flaring. The “evolutionary tracks” of 10 randomly selected disk models are included for reference. These tracks indicate the path that that particular disk would move if the disk became more flared. The right panel shows the mean change (“delta”) in [O I] 63.18 μm line flux and 65 μm continuum flux, with respect to an initially flatter ($\beta = 1.0$) disk. Arrows point in the direction of increasing disk flaring. Error bars indicate the 1σ variations in these Δ [O I] 63.18 μm line flux and Δ 65 μm continuum flux during each step in increasing flaring.

(A color version of this figure is available in the online journal.)

scenarios: first, the effects of changing the gas-to-dust ratio while holding the dust mass constant, as shown in Figure 5; and second, the effects of changing the gas-to-dust ratio while holding the gas mass constant, as shown in Figure 6. It is necessary to consider these two scenarios independently since identical gas-to-dust ratios can be constructed from different combinations of gas and dust mass.

As shown in Figure 5, increasing the gas-to-dust ratio, while holding the dust mass constant (in other words: we are increasing the gas-to-dust ratio by adding gas), results in increased [O I] 63.18 μm line luminosities. The increase in line luminosity is greatest for low dust masses, where changing the dust to gas ratio from 10^1 to 10^{-3} results in an increase in line luminosity of ~ 2 dex. At higher dust masses, the increase in line luminosity across the same range of gas-to-dust ratio results in an increase in line luminosity of ~ 1 dex. Since the [O I] 63.18 μm line is generally optically thick (e.g., Aresu et al. 2012), the increase in line luminosity with increasing gas mass is likely due to an increased heating rate, perhaps by H_2 photodissociation, collisional de-excitation of H_2^* , or photo-electric heating (e.g., Woitke et al. 2009). From Figure 5, it is also clear that changing the gas-to-dust ratio, while holding the dust mass constant, does not change the 63 μm continuum luminosity. This is not unexpected, since the continuum luminosity is tracing the dust in the disk, which in these cases, remains unchanged.

Figure 6 shows the complicated effects of increasing the gas-to-dust ratio while holding the gas mass constant (in other words, we are increasing the gas-to-dust ratio by removing dust). In general, increasing the gas-to-dust ratio by removing dust significantly decreases the 63 μm continuum luminosity by $0.2 \sim 2$ dex, depending on the gas mass. The behavior of the [O I] 63.18 μm line luminosity as the gas-to-dust ratio changes, while holding the gas mass constant, is even more complicated. For gas masses below $10^{-6} M_{\odot}$, the [O I], line luminosity decreases with increasing gas-to-dust ratio. For gas masses above $10^{-6} M_{\odot}$, the [O I], line luminosity increases with increasing gas-to-dust ratio, although the rate of this

increase decreases with decreasing dust mass. This decrease in [O I] line luminosity with decreasing dust mass may indicate the significance of dust-driven heating processes within the disk, such as polycyclic aromatic hydrocarbon heating and collisional heating (e.g., Woitke et al. 2009).

So, could the lower [O I] 63.18 μm line luminosities of transitional disks be explained by changes in the gas-to-dust ratio? Given the two ways of changing the gas-to-dust ratio, the simplest possible explanation is that transitional disks have lower gas-to-dust ratios, by having less gas mass than full disks. As shown in Figure 5, a decrease of the gas-to-dust ratio of only ~ 0.5 dex would be able to explain the factor of $\sim a$ few lower [O I] 63.18 μm line luminosities in transitional disks, while retaining similar 63 μm continuum luminosities. While there may be specific evolutionary pathways whereby increasing the dust mass can also explain the factor of $\sim a$ few lower [O I] 63.18 μm line luminosities in transitional disks, changes in the dust mass strongly affect the 63 μm continuum luminosities, as shown in Figure 6. Furthermore, our estimates of dust mass from millimeter observations (see Section 4.4) suggest that there is no statistical difference between the dust masses of full and transitional disks.

BP Tau may be an example of a more evolved full disk that is dispersing its gas, and decreasing its gas-to-dust ratio. Dutrey et al. (2003) showed that BP Tau is anomalous in many regards: its CO and dust disk are small and faint; the $^{12}\text{CO } J=2 \rightarrow 1$ transitional is optically thin; and that with respect to the dust, the CO is depleted by a large factor (~ 100). One possible explanation, discussed by Dutrey et al. (2003), is that BP Tau may be depleted in gas with respect to dust and have a lower gas-to-dust ratio than other full disks. As shown in Figure 1(a), BP Tau has an anomalously low [O I] 63.18 μm line luminosity compared to other full disks. This result confirms that BP Tau is indeed different from other full disks. Furthermore, the low [O I] 63.18 μm line luminosity is consistent with the hypothesis of Dutrey et al. (2003), that BP Tau has a lower gas-to-dust ratio than typical full disks by ~ 1 dex.




Figure 5. DENT grid predictions for how [O I] 63.18 μm line flux and 65 μm continuum flux change, with increasing gas-to-dust ratio, while the dust mass remains fixed. Panels (a)–(e) display the effect for different dust masses. Disk models are sampled at random ($N \sim 5000$) and consist of only low-mass stars ($\leq 1 M_{\text{Sun}}$). The panels on the left show the regions that contain 67% of the models as a function of gas-to-dust ratio. The “evolutionary tracks” of 10 randomly selected disk models are included for reference. These tracks indicate the path that that particular disk would move if the gas-to-dust ratio increased. The panels on the right show the mean change (“delta”) in [O I] 63.18 μm line flux and 65 μm continuum flux, with respect to an initially low gas-to-dust disk. Arrows point in the direction of increasing gas-to-dust ratio (corresponding to increasing gas in these figures). Error bars indicate the 1σ variations in these Δ [O I] 63.18 μm line flux and Δ 65 μm continuum flux during each step in increasing gas-to-dust ratio.

(A color version of this figure is available in the online journal.)

6. DISCUSSION

6.1. Implications for Disk Evolution Models

Photoevaporation may be a natural mechanism by which the disk gas-to-dust mass ratio is reduced with time. High-energy stellar photons heat the disk and drive a photoevaporative wind, which primarily removes the gas component from the disk surface. Amongst our sample of transitional disks, CS Cha, TW Hya, T Cha, RXJ1615.3-3255, and YLW8 have been observed with *VLT/VISIR* and present [Ne II] emission lines blueshifted by several km s^{-1} , implying ongoing photoevaporation (Pascucci & Sterzik 2009; Sacco et al. 2012). GM Aur has been observed with *Gemini/TEXES* but with insufficient S/N to precisely determine the line centroid (Najita et al. 2009). While photoevaporation has been detected from these objects, the rate

at which gas is lost via this mechanism is still unknown. If [Ne II] is tracing the very thin EUV irradiated region, the mass loss rate is negligible ($\sim 10^{-10} M_{\odot} \text{ yr}^{-1}$); while, if [Ne II] is tracing the deeper X-ray irradiated layer, the mass loss rate may be significant ($\sim 10^{-8} M_{\odot} \text{ yr}^{-1}$). In the latter case, if we assume that full disks start with a mass of $\sim 22 M_{\text{Jupiter}}$ (the mean value derived from millimeter data; see Section 2.3.3), they could lose half of their gas mass in just 1 Myr via photoevaporation.

Planet-disk interactions may also provide a mechanism for reducing the gas-to-dust ratio in protoplanetary disks (e.g., Espaillat 2013). Rice et al. (2006) showed that pressure gradients at the outer edge of a gap cleared by a giant planet can act as dust filters. In such a scenario, small dust grains and gas flow across the gap and are either lost to the planet or the inner disk (and eventually the host star), while large dust grains remain




Figure 6. DENT grid predictions for how [O I] 63.18 μm line flux and 65 μm continuum flux change, with increasing gas-to-dust ratio, while the gas mass remains fixed. Panels 6(a)–(g) display the effect for different gas masses. Disk models are sampled at random ($N \sim 5000$), and consist of only low-mass stars ($\leq 1 M_{\text{Sun}}$). The panels on the left show the regions that contain 67% of the models as a function of gas-to-dust ratio. The “evolutionary tracks” of 10 randomly selected disk models are included for reference. These tracks indicate the path that that particular disk would move if the gas-to-dust ratio increased. The right panels show the mean change (“delta”) in [O I] 63.18 μm line flux and 65 μm continuum flux, with respect to an initially low gas-to-dust disk. Arrows point in the direction of increasing gas-to-dust ratio (corresponding to decreasing dust in these figures). Error bars indicate the 1σ variations in these Δ [O I] 63.18 μm line flux and Δ 65 μm continuum flux during each step in increasing gas-to-dust ratio.

(A color version of this figure is available in the online journal.)

trapped in the outer disk. This has the effect of removing gas from the outer disk while retaining most of the millimeter- and centimeter-size dust and thus decreasing the gas-to-dust ratio of the outer disk. However, the leak of small, micron-sized dust particles into the inner disk still necessitates some

additional mechanism, such as dust coagulation, to explain the dust cavities in transitional disks (Zhu et al. 2012). Additionally, dust filtration alone is not a realistic mechanism for decreasing the gas-to-dust ratio by 0.5 dex, as suggested by our work. As gas leaves the outer disk and flows into the gap formed by the planet,




Figure 7. *Herschel*/PACS 63 μm spectra. Blue solid lines indicate the best-fit Gaussian line profile for the [O I] 63.18 μm line (as discussed in Section 3). Red dashed lines depict the hypothetical 3σ upper limits. Observations taken by the GASPS team are indicated by the annotation “GASPS Data.” While we re-reduced this data, these observations were previously reported in Howard et al. (2013), Meeus et al. (2012), and Podio et al. (2012).

(An extended, color version of this figure is available in the online journal.)

it will either be accreted onto the planet or will completely cross the gap into the inner disk, where it can then accrete onto the central star. Lubow & D’Angelo (2006) showed that when mass flows across into these gaps formed by giant planets, $\sim 90\%$ of the mass will be accreted onto the planet. Thus, for dust filtration to be the driver of a low gas-to-dust ratio in the outer disk, it is at the expense of putting a large majority of the outer disks’ gas mass directly into planets. If we assume full disks start with a gas mass of $\sim 22 M_{\text{Jupiter}}$ (the mean value derived from millimeter data; see Section 2.3.3), $\sim 7 M_{\text{Jupiter}}$ of gas would need to be lost to planet formation to result in a decrease in the gas-to-dust of 0.5 dex. If instead, we assume that a full protoplanetary disk can be characterized by a minimum mass solar nebula (MMSN; Weidenschilling 1977; Kuchner 2004),

then it would be necessary for the disk to lose even more mass, upward of $\gtrsim 20 M_{\text{Jupiter}}$.¹⁸ These simple calculations also assume that all of the dust in the outer disk is somehow protected, perhaps due to a planet-induced pressure bump. If the loss of dust across the gap is large, these mass estimates would only be

¹⁸ The total disk mass is calculated by integrating the surface mass density from the inner edge of the protoplanetary disk (~ 0.07 AU) to the outer edge (conservatively, ~ 40 AU). Using the MMSN described by Kuchner (2004) ($\Sigma = 4225 \text{ g cm}^{-2} (a/1 \text{ AU})^{-1.78}$) results in a total disk mass of $24 M_{\text{Jupiter}}$. Using the classical MMSN described by Weidenschilling (1977) ($\Sigma = 4200 \text{ g cm}^{-2} (a/1 \text{ AU})^{-1.5}$) results in a total disk mass of $38 M_{\text{Jupiter}}$. A loss of 0.5 dex of the disk mass for these two models correspond to 17 and $26 M_{\text{Jupiter}}$, respectively. Using more liberal estimates of the outer edge of the protoplanetary disk (e.g., 270 AU; Chiang & Goldreich 1997) results in even larger masses.

lower limits. If all of this mass is lost to forming planets, this would suggest the formation of a large number of giant planets at large semimajor axes ($\gtrsim 10$ AU), which does not seem to agree with the current (though still debated) statistics of giant exoplanets (Nielsen et al. 2013; Fressin et al. 2013; Biller et al. 2013). Lastly, while large, Jovian-mass planets can clear gaps and cause global depletions in the gas surface density of disks, they only deplete the surface density of the disk by a factor of a few (e.g., Figure 3 of Lubow & D’Angelo 2006). As discussed in Bruderer (2013) (and in Section 4.4), our observed factor of 2 line flux difference between transitional disks and full disks would require a drop in the surface density by a factor of $\gtrsim 100$.

6.2. Potential Follow-up Observations

Direct measurement of the gas-to-dust ratio in full disks and transitional disks would break our observed degeneracy between gas-to-dust ratio and disk flaring. While the dust mass of protoplanetary disks can be estimated with millimeter observations (e.g., Mohanty et al. 2013), the total gas mass of protoplanetary disks is difficult to directly measure. Combining our observations of the [O I] 63.18 μm line with low J CO rotational lines has been suggested as a possible way to directly measure total disk gas mass. While this method has been implemented for select well-studied disks (e.g., TW Hya, Thi et al. 2010), its reliability is still under discussion (Gorti & Hollenbach 2009; Bergin et al. 2013). Both low J CO and [O I] lines are optically thick, which make them both primarily sensitive to temperature and only weakly dependent on disk mass. Alternatively, observations of isotopologues may provide direct estimates for disk mass. Isotopologues (such as ^{13}C) are minor components within the disk and can be optically thin and directly trace disk mass (modulo the assumed abundances of the relative species). With the significant ($\sim 10\times$) increase in sensitivity allowed by ALMA, detecting emission from minor disk components out to nearby star-forming regions (e.g., Taurus-Auriga) is now possible.

It is difficult to directly measure the flaring of gas in protoplanetary disks. For select nearby and edge-on disks, it may be possible to directly measure the relative vertical distribution of dust (via millimeter emission) and gas (via gas emission lines, such as CO and its isotopologues) with high spatial and spectral resolution observations with ALMA (Rosenfeld et al. 2013). Detailed SED modeling covering the mid-infrared, far-infrared, and millimeter wavelengths may be able to break the degeneracy between disk gas mass and disk scale height. Flared disks intercept more stellar radiation at larger semimajor axes than flatter disks. Emission from these warm outer disk surface layers dominate the SED beyond $\sim 20 \mu\text{m}$ (Chiang & Goldreich 1997).

7. SUMMARY

We obtained *Herschel*/PACS spectra of [O I] 63.18 μm for 21 transitional disks in the Ophiuchus, Chameleon, and Lupus star-forming regions. This survey complements the larger *Herschel* GASPS survey of the Taurus star-forming region (Dent et al. 2013) by quadrupling the number of transitional disks observed with PACS in this wavelength. [O I] 63.18 μm is significant because it traces the cool, outer regions ($\gtrsim 10$ AU) of the protoplanetary disk, where the majority of the disk mass lies. Our primary results can be summarized as follows.

1. Full disks have larger [O I] 63.18 μm line luminosities than transitional disks, while having similar 63.18 μm continuum luminosities. While this result was previously

recognized by Howard et al. (2013) for the GASPS Taurus-Auriga sample, our data extends this trend to a larger sample of transitional disks, suggesting that lower [O I] 63.18 μm line emission is a characteristic property of transitional disks.

2. For all of our targets, we self-consistently derived stellar and disk parameters that have been previously shown to affect [O I] 63.18 μm emission. While [O I] 63.18 μm can correlate with these parameters, we found that transitional disks and full disks have statistically indistinguishable effective temperatures, bolometric luminosities, FUV luminosities, accretion rates, and X-ray luminosities. Thus, these properties cannot be responsible for the lower [O I] 63.18 μm line luminosities of transitional disks.
3. We found no correlation between the [O I] 63.18 μm line luminosities of transitional disks and either their disk masses (as inferred from millimeter photometry), dust cavity sizes, or wall heights (as inferred from SED and interferometric image modeling). This suggests that the decrease in [O I] 63.18 μm emission is not simply due to a lack of material in the inner cavity of transitional disks, though more modeling is needed to confirm this result (e.g., Bruderer 2013).
4. Using the DENT grid of thermo-chemical protoplanetary disk models (Woitke et al. 2010), we determined that the lower [O I] 63.18 μm line luminosities in transitional disks could result from either a decrease in disk flaring or a decrease in gas-to-dust ratio via a global depletion of gas mass. Decreasing the disk flaring results in less stellar irradiation impinging on the surface of the outer disk, thus decreasing the disk temperature and reducing [O I] 63.18 μm emission. Decreasing the gas-to-dust ratio by removing gas mass results in a decrease in the amount of heating from H_2 photo-dissociation, collisional de-excitation of H_2^+ , and/or photo-electric heating (e.g., Woitke et al. 2009). Both photoevaporation and planet formation can result in a decrease in gas mass, although their efficiencies are still not well constrained. While additional observations are needed to disentangle the effects of disk flaring and gas-to-dust ratio, our results show that transitional disks are more evolved than their full disk counterparts, possibly even at large radii.

I.P., J.T.K., C.E., and S.A. acknowledge NASA/JPL for funding support. J.T.K. and I.P. thank Elisabetta Rigliaco for helpful discussions on mass accretion rate estimates. The authors would also like to thank the referee, Kees Dullemond, for a very constructive review.

APPENDIX

Figure 7 includes all of the reduced *Herschel*/PACS [O I] 63.18 μm spectra used in this work, and is only provided in the online version of the article. Figure 2 displays null correlations of various stellar and disk parameters with the [O I] 63.18 μm line luminosity and nearby continuum luminosity and are only provided in the online version of the article.

REFERENCES

- Aikawa, Y., & Nomura, H. 2006, *ApJ*, 642, 1152
 Alcalá, J. M., Covino, E., Sterzik, M. F., et al. 2000, *A&A*, 355, 629
 Alexander, R. D., Clarke, C. J., & Pringle, J. E. 2006, *MNRAS*, 369, 229

- Alexander, R. D., Pascucci, I., Andrews, S., Armitage, P., & Cieza, L. 2014, in *Protostars and Planets VI*, ed. H. Beuther, C. Dullemond, T. Henning, & R. Klessen, in press (arXiv:1311.1819)
- Andrews, S. M., Rosenfeld, K. A., Wilner, D. J., & Bremer, M. 2011, *ApJ*, **732**, 42
- Andrews, S. M., & Williams, J. P. 2005, *ApJ*, **631**, 1134
- Andrews, S. M., Wilner, D. J., Hughes, A. M., Qi, C., & Dullemond, C. P. 2009, *ApJ*, **700**, 1502
- Antoniucci, S., Garcí López, R., Nisini, B., et al. 2011, *A&A*, **534**, 32
- Aresu, G., Kamp, I., Meijerink, R., et al. 2011, *A&A*, **526**, 163
- Aresu, G., Kamp, I., Meijerink, R., et al. 2014, *A&A*, in press (arXiv:1402.2488)
- Aresu, G., Meijerink, R., Kamp, I., et al. 2012, *A&A*, **547**, 69
- Bary, J. S., Weintraub, D. A., Shukla, S. J., Leisenring, J. M., & Kastner, J. H. 2008, *ApJ*, **678**, 1088
- Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Guesten, R. 1990, *AJ*, **99**, 924
- Bergin, E. A., Cleeves, L. I., Gorti, U., et al. 2013, *Natur*, **493**, 644
- Biazzo, K., Alcalá, J. M., Covino, E. L., et al. 2012, *A&A*, **547**, 104
- Biller, B. A., Liu, M. C., Wahhaj, Z., et al. 2013, *ApJ*, **777**, 160
- Bouvier, J., & Appenzeller, I. 1992, *A&AS*, **92**, 481
- Bouvier, J., Cabrit, S., Fernandez, M., Martin, E. L., & Matthews, J. M. 1993, *A&AS*, **101**, 485
- Brandner, W., Alcalá, J. M., Kunkel, M., Moneti, A., & Zinnecker, H. 1996, *A&A*, **307**, 121
- Briceno, C., Calvet, N., Kenyon, S., & Hartmann, L. 1999, *AJ*, **118**, 1354
- Brown, J. M., Blake, G. A., Dullemond, C. P., et al. 2007, *ApJ*, **664**, 107
- Brown, J. M., Blake, G. A., Qi, C., et al. 2009, *ApJ*, **704**, 496
- Bruderer, S. 2013, *A&A*, **559**, A46
- Cabrit, S., Edwards, S., Strom, S. E., & Strom, K. M. 1990, *ApJ*, **354**, 687
- Calvet, N., D'Alessio, P., Hartmann, L., et al. 2002, *ApJ*, **658**, 1008
- Carkner, L., Kozak, J. A., & Feigelson, E. D. 1998, *AJ*, **116**, 1933
- Chen, H., Myers, P. C., Ladd, E. F., & Wood, D. O. S. 1995, *ApJ*, **445**, 377
- Chiang, E. I., & Goldreich, P. 1997, *ApJ*, **490**, 368
- Cohen, M., & Kuhl, L. V. 1979, *ApJS*, **41**, 743
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *yCat*, **2246**, 0
- Daemgen, S., Petr-Gotzens, M., Correia, S., et al. 2013, *A&A*, **554**, 43
- Dahm, S. E. 2008, *AJ*, **136**, 521
- D'Alessio, P. D., Calvet, N., Hartmann, L., Franco-Hernández, R., & Servín, H. 2006, *ApJ*, **638**, 314
- Damiani, F., Micela, G., Sciortino, S., & Harnden, F. R., Jr. 1995, *ApJ*, **446**, 331
- DENIS Consortium 2005, *yCat*, **2263**, 0
- Dent, W. R. F., Thi, W. F., Kamp, I., et al. 2013, *PASP*, **125**, 477
- Dullemond, C. P., & Dominik, C. 2005, *A&A*, **434**, 971
- Dutrey, A., Guilloteau, S., & Simon, M. 2003, *A&A*, **402**, 1003
- Edwards, S., Cabrit, S., Strom, S. E., et al. 1987, *ApJ*, **321**, 473
- Edwards, S., Hartigan, P., Ghandour, L., & Andrulis, C. 1994, *AJ*, **108**, 1056
- Españallat, C. 2013, in *IAU Symp. 299, Exploring the Formation and Evolution of Planetary Systems*, ed. M. Booth, B. C. Matthews, & J. R. Graham (Cambridge: Cambridge Univ. Press), 140
- Españallat, C., Calvet, N., D'Alessio, P., et al. 2007, *ApJL*, **670**, L135
- Españallat, C., D'Alessio, P., Hernández, J., et al. 2010, *ApJ*, **717**, 441
- Españallat, C., Furlan, E., D'Alessio, P., et al. 2011, *ApJ*, **728**, 49
- Españallat, C., Ingleby, L., Hernández, J., et al. 2012, *ApJ*, **747**, 103
- Españallat, C., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, C. Dullemond, T. Henning, & R. Klessen, in press (arXiv:1402.7103)
- Fang, M., van Boekel, R., Wang, W., et al. 2009, *A&A*, **504**, 461
- Feigelson, E. D., Casanova, S., Montmerle, T., & Guibert, J. 1993, *ApJ*, **416**, 623
- Feigelson, E. D., & Kriss, G. A. 1989, *ApJ*, **338**, 262
- Fernandez, M. 1995, *A&AS*, **113**, 473
- Fressin, F., Torres, G., Charbonneau, D., et al. 2013, *ApJ*, **766**, 81
- Furlan, E., Watson, D. M., McClure, M. K., et al. 2009, *ApJ*, **703**, 1964
- Gauvin, L. S., & Strom, K. M. 1992, *ApJ*, **385**, 217
- Gorti, U., & Hollenbach, D. 2009, *ApJ*, **690**, 1539
- Gorti, U., Hollenbach, D., Najita, J., & Pascucci, I. 2011, *ApJ*, **735**, 90
- Güdel, M., Briggs, K. R., Arzner, K., et al. 2007, *A&A*, **468**, 353
- Guenther, E. W., & Emerson, J. P. 1997, *A&A*, **321**, 803
- Guenther, E. W., Esposito, M., & Mundt, R. 2007, *A&A*, **467**, 1147
- Guilloteau, S., Dutrey, A., Piétu, V., & Boehler, Y. 2011, *A&A*, **529**, 105
- Gullbring, E., Hartmann, L., Briceno, C., & Calvet, N. 1998, *ApJ*, **492**, 323
- Hartigan, P., Edwards, S., & Ghandour, L. 1995, *ApJ*, **452**, 736
- Hartmann, L., Calvet, N., Gullbring, E., & D'Alessio, P. 1998, *ApJ*, **495**, 385
- Henning, T., Pfau, W., Zinnecker, H., & Prusti, T. 1993, *A&A*, **276**, 129
- Herbst, W., Herbst, D. K., Grossman, E. J., & Weinstein, D. 1994, *AJ*, **108**, 1906
- Herczeg, G. J., & Hillenbrand, L. A. 2008, *ApJ*, **681**, 594
- Herczeg, G. J., Najita, J. R., Hillenbrand, L. A., & Pascucci, I. 2007, *ApJ*, **670**, 509
- Howard, C. D., Sandell, G., Vacca, W. D., et al. 2013, *ApJ*, **776**, 21
- Hughes, A. M., Wilner, D. J., Andrews, S. M., Qi, C., & Hogerheijde, M. R. 2011, *ApJ*, **727**, 85
- Hughes, J., Hartigan, P., Krautter, J., & Kelemen, J. 1994, *AJ*, **108**, 1071
- Ireland, M. J., & Kraus, A. L. 2008, *ApJ*, **678**, 59
- Kamp, I., Tilling, I., Woitke, P., Thi, W.-F., & Hogerheijde, M. 2010, *A&A*, **510**, 18
- Kamp, I., Woitke, P., Pinte, C., et al. 2011, *A&A*, **532**, 85
- Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., & Weintraub, D. A. 1999, *ApJ*, **525**, 837
- Kelly, B. C. 2007, *ApJ*, **665**, 1489
- Kenyon, S. J., Brown, D. L., Tout, C. A., & Berline, P. 1998, *AJ*, **115**, 2491
- Kenyon, S. J., Gomez, M., & Whitney, B. A. 2008, in *Handbook of Star Forming Regions*, Vol. 1, ed. B. Reipurth (San Francisco, CA: ASP), 405
- Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, **101**, 117
- Kim, K. H., Watson, D. M., Manoj, P., et al. 2009, *ApJ*, **700**, 1017
- Kim, K. H., Watson, D. M., Manoj, P., et al. 2013, *ApJ*, **769**, 149
- Kraus, A. L., & Hillenbrand, L. A. 2009, *ApJ*, **704**, 531
- Kraus, A. L., Ireland, M. J., Hillenbrand, L. A., & Martinache, F. 2012, *ApJ*, **745**, 19
- Kuchner, M. J. 2004, *ApJ*, **612**, 1147
- LaValley, M., Isobe, T., & Feigelson, E. 1992, in *ASP Conf. Ser. 25, Astronomical Data Analysis Software and Systems I*, ed. D. M. Worrall, C. Biemesderfer, & J. Barnes (San Francisco, CA: ASP), 245
- Lubow, S. H., & D'Angelo, G. 2006, *ApJL*, **641**, L526
- Lubow, S. H., Seibert, M., & Artymowicz, P. 1999, *ApJ*, **526**, 1001
- Luhman, K. L. 1999, *ApJ*, **525**, 466
- Luhman, K. L. 2000, *ApJ*, **544**, 1044
- Luhman, K. L. 2004, *ApJ*, **602**, 816
- Luhman, K. L. 2007, *ApJS*, **173**, 104
- Magazzu, A., Rebolo, R., & Pavlenko, I. V. 1992, *ApJ*, **392**, 159
- Mathis, J. S. 1990, *ARA&A*, **28**, 37
- McCabe, C., Ghez, A. M., & Prato, L. 2006, *ApJ*, **636**, 932
- McClure, M. K., Furlan, E., Manoj, P., et al. 2010, *ApJS*, **188**, 75
- Meeus, G., Montesinos, B., Mendigutía, I., et al. 2012, *A&A*, **544**, 78
- Meijerink, R., Aresu, G., Kamp, I., et al. 2012, *A&A*, **547**, 68
- Merín, B., Brown, J. M., Oliveira, I., et al. 2010, *ApJ*, **718**, 1200
- Merín, B., Jørgensen, J., Spezzi, L., et al. 2008, *ApJS*, **177**, 551
- Mohanty, S., Greaves, J., Mortlock, D., et al. 2013, *ApJ*, **773**, 168
- Monet, D. G., Levine, S. E., Canzian, B., et al. 2003, *AJ*, **125**, 984
- Muzerolle, J., Allen, L. E., et al. 2010, *ApJ*, **708**, 1107
- Muzerolle, J., Calvet, N., & Hartmann, L. 2001, *ApJ*, **550**, 944
- Najita, J. R., Doppmann, G. W., Bitner, M. A., et al. 2009, *ApJ*, **697**, 957
- Najita, J. R., Strom, S. E., & Muzerolle, J. 2007, *MNRAS*, **378**, 369
- Neuhaeuser, R., Sterzik, M. F., Schmitt, J. H. M. M., Wichmann, R., & Krautter, J. 1995, *A&A*, **297**, 391
- Nielsen, E. L., Liu, M. C., Wahhaj, Z., et al. 2013, *ApJ*, **776**, 4
- Nuernberger, D., Chini, R., & Zinnecker, H. 1997, *A&A*, **324**, 1036
- Osterloh, M., & Beckwith, S. V. W. 1995, *ApJ*, **539**, 288
- Ott, S. 2010, in *ASP Conf. Ser. 434, Astronomical Data Analysis Software and Systems XIX*, ed. Y. Mizumoto, K.-I. Morita, & M. Ohishi (San Francisco, CA: ASP), 139
- Pascucci, I., Apai, D., Hardegree-Ullman, E. E., et al. 2008, *ApJ*, **673**, 477
- Pascucci, I., & Sterzik, M. 2009, *ApJ*, **702**, 724
- Pascucci, I., & Tachibana, S. 2010, in *Protoplanetary Dust: Astrophysical and Cosmochemical Perspectives*, ed. D. Apai & D. S. Lauretta (Cambridge: Cambridge Univ. Press), 263
- Pinte, C., Woitke, P., Ménard, F., et al. 2010, *A&A*, **518**, L126
- Podio, L., Kamp, I., Flower, D., et al. 2012, *A&A*, **545**, 44
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, **518**, L2
- Prato, L., Greene, T. P., & Simon, M. 2003, *ApJ*, **584**, 853
- Rebull, L. M., Padgett, D. L., & McCabe, C.-E. 2010, *ApJS*, **186**, 259
- Reipurth, B. 2008a, *Handbook of Star-forming Regions: Volume I, The Northern Sky* (San Francisco, CA: ASP)
- Reipurth, B. 2008b, *Handbook of Star-forming Regions: Volume II, The Southern Sky* (San Francisco, CA: ASP)
- Riaz, B., Gizis, J. E., & Hmiel, A. 2006, *ApJ*, **639**, 79
- Rice, W. K. M., Armitage, P. J., Wood, K., & Lodato, G. 2006, *MNRAS*, **372**, L9
- Rigliaco, E., Natta, A., Randich, S., et al. 2011, *A&A*, **526**, L6
- Rigliaco, E., Natta, A., Testi, L., et al. 2012, *A&A*, **548**, 56
- Riviere-Marichalar, P., Ménard, F., Thi, W. F., et al. 2012, *A&A*, **538**, L3
- Rosenfeld, K. A., Andrews, S. M., Wilner, D. J., Kastner, J. H., & McClure, M. K. 2013, *ApJ*, **775**, 136

- Sacco, G. G., Flaccomio, E., Pascucci, I., et al. 2012, *ApJ*, 747, 142
- Salyk, C., Blake, G. A., Boogert, A. C. A., & Brown, J. M. 2009, *ApJ*, 699, 330
- Schisano, E., Covino, E., Alcalá, J. M., et al. 2009, *A&A*, 501, 1013
- Shu, F., Najita, J., Ostriker, E., et al. 1994, *ApJ*, 429, 781
- Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 593
- Spezzi, L., Alcalá, J. M., Frasca, A., Covino, E., & Gandolfi, D. 2007, *A&A*, 470, 281
- Strom, K. M., Strom, S. E., Edwards, S., Cabrit, S., & Skrutskie, M. F. 1989, *AJ*, 97, 1451
- Taguchi, Y., Itoh, Y., & Mukai, T. 2009, *PASJ*, 61, 251
- Thi, W.-F., Mathews, G., Ménard, F., et al. 2010, *A&A*, 518, L125
- Weidenschilling, S. J. 1977, *Ap&SS*, 51, 153
- White, N. E., Giommi, P., & Angelini, L. 2000, *yCat*, 9031, 0
- White, R. J., & Basri, G. 2003, *ApJ*, 582, 1109
- White, R. J., & Ghez, A. M. 2001, *ApJ*, 556, 265
- Wilking, B. A., Meyer, M. R., Robinson, J. G., & Greene, T. P. 2005, *AJ*, 130, 1733
- Williams, J. P., & Cieza, L. A. 2011, *ARA&A*, 49, 67
- Woitke, P., Kamp, I., & Thi, W.-F. 2009, *A&A*, 501, 383
- Woitke, P., Pinte, C., Tilling, I., et al. 2010, *A&A*, 510, 18
- Yang, H., Herczeg, G. J., Linsky, J. L., et al. 2012, *ApJ*, 744, 121
- Zhu, Z., Nelson, R. P., Dong, R., Espaillat, C., & Hartmann, L. 2012, *ApJ*, 755, 6