



CrossMark
click for updates

Cite this: *RSC Adv.*, 2015, 5, 19379

Received 21st November 2014
Accepted 9th February 2015

DOI: 10.1039/c4ra15004j

www.rsc.org/advances

Unexpected rearrangement during the gas-phase dehalogenation approach to benzodithiophenes†


R. Alan Aitken* and Adebayo O. Oyewale‡

A range of halogenated methylthiophenes undergo dehalogenative condensation upon FVP over magnesium or calcium to give products including benzodithiophenes; in some cases this involves a novel rearrangement by equilibration of alkenylthienyl radicals *via* thiophene-fused cyclopropylmethyl intermediates.

Some time ago we described the use of flash vacuum pyrolysis (FVP) over a bed of freshly sublimed magnesium as an effective and general method for dehalogenative coupling of benzyl halides, mono and dihaloalkanes and related compounds, with various pieces of evidence favouring the intermediacy of surface-adsorbed organometallic species rather than free gas-phase radicals.^{1,2} Although the six possible benzodithiophenes were prepared by conventional multi-step synthesis many years ago,³ related compounds such as a dialkylbenzodithiophene⁴ and naphthodithiophenes^{5,6} have been of recent interest for their electronic properties and this inspired us to apply the gas phase dehalogenation method to their synthesis.

The required dichloromethylthiophenes were prepared by reaction of the corresponding aldehydes with PCl₅ and in common with previous workers,⁷ we found them to be highly unstable so they were used immediately for the pyrolysis studies. Their ¹³C NMR spectra, not previously reported, all showed a distinctive signal at 62–68 ppm for the CHCl₂ carbon. We previously described briefly the reaction of a range of substituted benzylidene chlorides ArCHCl₂ over magnesium under FVP conditions which resulted in complete dehalogenation to give mainly the corresponding stilbenes,¹ and this pattern was repeated in the thiophene series. Thus, FVP of 2-dichloromethylthiophene **1** at 600 °C through glass wool

coated with freshly sublimed magnesium gave mainly the (*E*)- and (*Z*)-dithienylethenes along with a little 2-methylthiophene and the cyclisation product benzo[1,2-*b*:4,3-*b'*]dithiophene **2** identified by comparison with an authentic sample³ (Scheme 1). Likewise the isomeric 3-dichloromethylthiophene **3** gave the (*E*)- and (*Z*)-alkenes, the methylthiophene and





Scheme 1

EaStCHEM School of Chemistry, University of St Andrews, North Haugh, St Andrews, Fife, KY16 9ST, UK. E-mail: raa@st-and.ac.uk; Fax: +44 (0)1334 463808; Tel: +44 (0) 1334 463865

† Electronic supplementary information (ESI) available: Full experimental details for synthetic and pyrolytic studies. See DOI: 10.1039/c4ra15004j

‡ Current address: Department of Chemistry, Ahmadu Bello University, Zaria, Nigeria, E-mail: aoyewale@abu.edu.ng






benzo[2,1-*b*:3,4-*b'*]dithiophene **4**. The methyl compound **5** gave mainly the known⁸ crystalline (*E*)-alkene but this was accompanied by an insoluble polymer assumed to be poly(2,5-thienyleneethylene) **7** formed from the quinonoid dimethylene compound **6**. In the previous work,¹ cyclisation was promoted by an *ortho* halogen substituent and 2-chlorobenzylidene chloride gave phenanthrene in 47% yield. It was therefore expected that benzodithiophene formation would be increased with thiophenes bearing adjacent dichloromethyl and halogen substituents. The two new compounds **8** and **15** were therefore prepared from the corresponding aldehydes^{9,10} by reaction with PCl_5 and subjected to FVP over magnesium.

With compound **8** at 600 °C there was only partial reaction to give mainly products **9–13** still containing halogen (Scheme 2, Table 1) so the temperature was increased to 650 °C. This had the desired effect in reducing halogen-containing products and increasing the benzodithiophene yield. However, most surprisingly, not only the expected isomer **2**, but also **4** and **14** were observed. The volatility of magnesium places an upper limit of around 650 °C on the temperature of operation, so we substituted it with calcium for which 700 °C is easily accessible and, under these conditions, there was even more complete dehalogenation to give as the main product a mixture of the three isomeric benzodithiophenes **2**, **4** and **14**.

With the pseudoisomeric system **15** similar results were obtained: at 600 °C the main products were **16–20** along with




not only the expected benzodithiophene **4** but also the isomer **14**. Either moving to 650 °C with magnesium or 700 °C with calcium served to increase the yield of the latter two compounds with the rearranged product **14** actually predominating over **4** (Scheme 3, Table 1).

It therefore appears that, under the conditions used, an unusual isomerisation process is occurring. Authentic samples of both **2**³ and **14**¹¹ were prepared and resubjected to the pyrolysis conditions but were recovered completely unchanged. The key evidence as to the mechanism of the rearrangement was obtained by synthesis³ of the expected intermediates **21** and **22** formed by dehalogenative coupling of the dichloromethyl groups

Table 1 Products from pyrolysis of **8** and **15**

Starting material	Metal	Temperature °C	Products (%)									
			9/16	10/17	11	12	13/18	19	20	2	4	14
8	Mg	600	25	4	13	6	3				Trace	
8	Mg	650	14	4	10	12	Trace				8	10
8	Ca	700	Trace	Trace		8					16	16
15	Mg	600	28	10			6	18	10			2
15	Mg	650	24	8								2
15	Ca	700	21	3			12	9	3			3





Scheme 5

but with the ring halogens still in place. Subjecting either of these compounds to FVP over magnesium at 650 °C gave a mixture of the three isomeric benzodithiophenes **2**, **4**, and **14** (Scheme 4). In both cases the main product was the one expected without rearrangement but the formation of the other isomers and particularly a significant amount of **14** from **21** as well as from **22** supports the occurrence of the rearrangement process at this stage in the dehalogenation.

In view of all the evidence we propose that this occurs as shown in Scheme 5 by way of a cyclopropenylmethyl radical (or its metal-adsorbed equivalent) which can be formed either

from the 2-alkenyl-3-thienyl or 3-alkenyl-2-thienyl radicals and this provides a mechanism for interconversion between them leading to the two different modes of thiophene ring fusion. Further applications of FVP over reactive metals will be reported shortly.

Notes and references

- 1 R. A. Aitken, P. K. G. Hodgson, A. O. Oyewale and J. J. Morrison, *Chem. Commun.*, 1997, 1163–1164.
- 2 R. A. Aitken, P. K. G. Hodgson, J. J. Morrison and A. O. Oyewale, *J. Chem. Soc., Perkin Trans. 1*, 2002, 402–415.
- 3 S. Gronowitz and T. Dahlgren, *Chem. Scr.*, 1977, **12**, 57–67.
- 4 T. M. Pappenfus, D. T. Seidenkranz and E. W. Reinheimer, *Heterocycles*, 2012, **85**, 355–364.
- 5 R. Umeda, H. Fukuda, K. Miki, S. M. A. Rahman, M. Sonoda and Y. Tobe, *C. R. Chimie*, 2009, **12**, 378–384.
- 6 S.-W. Cheng, D.-Y. Chiou, Y.-Y. Lai, R.-H. Yu, C.-H. Lee and Y.-J. Cheng, *Org. Lett.*, 2013, **15**, 5338–5341.
- 7 M. S. Baird and I. Bruce, *J. Chem. Res., Synop.*, 1990, 134–135; *J. Chem. Res., Miniprint*, 1990, 946–965.
- 8 R. V. Hoffman, G. G. Orphanides and H. Shechter, *J. Am. Chem. Soc.*, 1978, **100**, 7927–7933.
- 9 S. Gronowitz, P. Moses, A.-B. Hörnfeldt and R. Håkansson, *Ark. Kemi*, 1961, **17**, 165–177.
- 10 B. Yom-Tov and S. Gronowitz, *Chem. Scr.*, 1973, **3**, 165–170.
- 11 R. M. Kellogg, M. B. Groen and H. Wynberg, *J. Org. Chem.*, 1967, **32**, 3093–3100.

