

Proton Transfer Reactions of N-Aryl Triazolium Salts: Unusual *Ortho*-Substituent Effects

David E. Tucker,^a Peter Quinn,^a Richard S. Massey,^a Christopher J. Collett,^b David J. Jasiewicz,^a Christopher R. Bramley,^a Andrew D. Smith^{b,*} and AnnMarie C. O'Donoghue^{a,*}

^a*Department of Chemistry, Durham University, South Road, Durham DH1 3LE, UK.*


^b*EaStCHEM, School of Chemistry, University of St Andrews, North Haugh, St Andrews, Fife KY16 9ST, UK*

EMAIL: annmarie.odonoghue@durham.ac.uk, ads10@st-andrews.ac.uk

ABSTRACT:


INTRODUCTION

Triazol-3-ylidenes **1** form a class of stable N-heterocyclic carbenes (NHC) that are frequently utilized as efficient, selective organic catalysts in a broad range of transformations.^[1-7] Closely related to other stable NHCs including imidazole-2-ylidenes **2**, imidazolin-2-ylidenes **3**, thiazol-2-ylidenes **4** and trihydropyrimidin-2-ylidenes **5**, these carbenes have seen application in diverse areas of chemistry in addition to organic catalysis.^[8-17] Common to most applications of NHCs of this type is the *in situ* generation of the carbene from the conjugate acid azolium salt precursor by use of an appropriate base. We and others have reported the kinetic acidities towards hydroxide ion and estimates of the aqueous pK_a values of the conjugate acid precursors to NHCs **1-5**.^[18-23]



In particular, we showed that the pD rate profiles for the deuterium exchange reactions of triazolium salt precursors **6** to triazol-3-ylidenes **7** reveal distinct differences from analogous data for NHCs **2-5**.^[21] The presence of the additional ring nitrogen in triazolium ions **6** allows for alternative deuterium exchange mechanisms under more acidic conditions. Deuteration at N(1) can occur to give dicationic triazolium ions **8**, which are precursors to monocationic N-heterocyclic carbenes **9**. At higher pD values, $\log k_{ex}$ values were observed to increase linearly with pD consistent with a first order dependence on deuteroxide ion and exchange via triazolium salt **7** (Path A, Scheme 1). At lower pD values, upward deviations were observed from the line of unit slope through the $\log k_{ex} - pD$ data consistent with a change in mechanism for deuterium exchange.

Scheme 1 Potential Mechanistic Pathways for Deuterium Exchange at C(3)-H of Triazolium Ions **6**.




Of the large series of triazolium salts studied in aqueous solution, the altered dependence of $\log k_{\text{ex}}$ values on pD under more acidic conditions was most prevalent, and occurred at significantly higher pD values, for N-pentafluorophenyl triazolium salt **6a** (Ar = C_6F_5).^[21] For the structurally homologous series of triazolium salts **6**, the pD value for onset of the change in slope decreased in the order **6a** > **6b** > **6c** > **6d** ~ **6e** ~ **6f** (Ar = C_6F_5 (a); 4-CN-Ph (b); 4-F-Ph (c); H (d); 2,4,6-(Me₃)-Ph (e); 4-MeO-Ph (f)). Only in the case of triazolium salt **6a**, the initial decreased dependence on pD was further followed by a downward break at the lowest pD s studied. Data for **6a** could be fit by an equation describing either of two kinetically equivalent mechanistic options (Paths B or C, Scheme 1), which *both* require protonation at N1 of the triazolium ring. Path B involves a solvent promoted deuterium exchange reaction of the cationic triazolium salt **6a** with removal of active species *via* N-protonation to a dicationic triazolium ion **8a** to allow for the observed continued decrease of rate

constants with pD . Path C entails no solvent reaction of the monocationic salt **6a**, but instead the initial N-protonation at N(1) of salt **6a** followed by deuteroxide-catalyzed exchange on the dicationic triazolium salt **8a**. Based on an analysis of rate constants obtained through assumption of either pathway, we suggested that Path C was the more likely option.


In order to explain the differences in the observed kinetic data for **6a-f**, we postulated that the presence of an *ortho*-heteroatom, *e.g.* fluorine, could favour protonation at N(1) resulting in an increased prevalence of dicationic triazolium salts **8** in the normal pD range.^[21] Of the substrates studied, only the N-pentafluorophenyltriazolium salt **6a** had *ortho*-heteroatoms rather than hydrogen or methyl groups on the N-aryl ring. A higher pK_a (N1) for salt **6a** seems counter-intuitive based on the electron-withdrawing, through-bond, inductive substituent effect of fluorine, which would be expected to decrease basicity at N(1). However, protonation at N(1) may be favoured in order to suppress unfavourable electrostatic interactions between this nitrogen and spatially proximal N-aryl *ortho*-fluorines, or, as a result of stabilizing $N^+ \cdots H \cdots ortho-F$ interactions in the N-protonated salt.

In related mechanistic studies of triazolium catalysis of benzoin and Stetter reactions by triazolium salts, we have also observed unexpected substituent effects in the presence of *ortho*-heteroatom substituents on the N-aryl ring of catalyst.^[24] The common first step of both of these reactions involves the reaction of an aryl aldehyde and triazolium salt in the presence of base to give a hydroxy aryl intermediate **10** (Scheme 2, shown for benzoin condensation). We have observed that N-aryl *ortho*-X-heteroatom substituents significantly increase rate and equilibrium constants for formation of adducts **10** relative to *para*-substituted and *ortho*-alkyl analogues. One possible explanation of these observations is the presence of an O-H...X interaction in the product hydroxyaryl adduct similar to the $N^+ \cdots H \cdots ortho-F$ interaction proposed above.

Scheme 2 Triazolium Catalysis of the Benzoin Condensation.



In order to further probe these *ortho*-substituent effects, we have studied the deuterium exchange reactions of a number of additional triazolium salts **11-14** with heteroatoms or heteroatom substituents in the 2- and/or 6-positions of the N-aryl ring. The data for these substrates also provides further evidence that Path C, and not Path B, is accountable for the altered dependence of $\log k_{\text{ex}}$ values on pD under more acidic conditions. In addition, we observed that the N-pyridyl triazolium salt **13** displays a different N-aryl substituent effect to all other triazolium substrates. Formal acid catalysis of deuterium exchange was uniquely observed for N-pyridyltriazolium salt **13** under more acidic condition providing evidence for a possible intramolecular deprotonation reaction involving the pyridyl substituent.



EXPERIMENTAL

The syntheses of triazolium salts **11-14**, the preparation of solutions, the determination of pD and NMR methods are described in the Supporting Information.

Kinetic measurements

The kinetic procedures for the measurement of rate constants for deuterium exchange for triazolium salts **11-14** were identical to those previously reported for the study of analogous salts **6a-f**.^[21] Due to the lability of the triazolium salts towards C(3)-H/D-exchange in unbuffered D₂O solvent, reactions were initiated by addition of a solution, containing internal standard (tetramethylammonium deuteriosulfate) and buffer or DCl, directly to the rigorously dried triazolium salt. The final substrate and internal standard concentrations in the D₂O reaction solutions were 5 mM and 1 mM, respectively. Reaction solutions in NMR tubes were incubated at 25 °C in a thermostatted water bath. pD values were recorded at the beginning and end of each reaction and were found to be constant within error (± 0.03). The progress of the C(3)-H/D deuterium exchange reaction was followed by ¹H NMR spectroscopy during the disappearance of 75-90% of the C(3)-H signal of each substrate. There was no change in the integrated areas of signals due to all other protons of triazolium salts **11-13** during this period, and no appearance of new signals, consistent with the absence of any parallel decomposition or hydrolysis reactions under the reaction conditions. For N-pyrimidinyl salt **14** a competing reaction was observed to occur to a small extent accounting for $\leq 6\%$ of total products. Based on the NMR signals due to the products formed, this reaction was presumed to be a nucleophilic aromatic substitution reaction of DO⁻ or D₂O at C(2) of the pyrimidinyl ring.

The observed pseudo first order rate constants for exchange of the C(3)-proton for deuterium, k_{ex} (s⁻¹), were obtained from non-linear least square fitting of reaction progress against time to a first order exponential decay function. Reaction progress was defined by values of $f(s)$, the fraction of remaining unexchanged substrate, which were calculated from Eqn. (1), where $A_{\text{C}_3\text{H}}$ and A_{std} are the integrated areas of the singlet due to the C(3)-H of the triazolium salt and the broad triplet at 3.3 ppm due to the methyl hydrogens of internal standard, tetramethylammonium deuteriosulfate. In the case of triazolium salt **14**, small corrections were made for the parallel S_NAr reaction of substrate such that the corrected $f(s)$ values represented reaction due to deuterium exchange only.

$$f(s) = \frac{(A_{\text{C}_3\text{H}} / A_{\text{std}})_t}{(A_{\text{C}_3\text{H}} / A_{\text{std}})_0} \quad (1)$$

Representative NMR spectral overlays of the deuterium exchange reactions, first order kinetic plots, tabulated k_{ex} data and $\log k_{\text{ex}} - \text{pD}$ rate profiles are included in

the Supporting Information for each triazolium salt **11-14** (Figures S1-S16, Tables S1-S4).

RESULTS AND DISCUSSION

The C(3)-H/D deuterium exchange reactions of triazolium salts **11-14** were performed in aqueous acetic acid buffer or DCl solutions at a range of pD values and at constant ionic strength, $I = 1.0$ (KCl). For these substrates, deuterium exchange was too fast to monitor above pD 4 at 25 °C. Buffer catalysis of deuterium exchange was found to be insignificant in all previous studies of azolium ion conjugate acids of N-heterocyclic carbenes including representative triazolium salts.^[18, 20-21] Hence, it was assumed that buffer catalysis of exchange was not significant and the observed pseudo first order rate constants for exchange, k_{ex} (s^{-1}) were used directly on pD - rate profiles of deuterium exchange.

As in the previous study of a large series of triazolium salts, values of $\log k_{\text{ex}}$ for **11** and **12** (Figure 1, ▲ and ▼) increase with pD in the region from pD = 0 to 4.5. Figure 1 shows the pD rate profiles for these salts in comparison with those for two previously studied N-pentafluorophenyl and N-phenyl triazolium salts **6a** and **6d**, respectively. Figure 2 shows the pD rate profiles for N-pyridyl and N-pyrimidinyl salts **13** and **14** also in comparison with previous data for **6a** and **6d**, respectively. The data for N-pyridyl triazolium salt **13** (Figure 2, ◆) was distinctly different from all other triazolium salts studied including N-pentafluorophenyl substrate **6a**. Values of $\log k_{\text{ex}}$ for **13** decrease with pD in the region from pD = 0 to 1.3 and increase with pD in the region 1.3-4.5. By contrast, the $\log k_{\text{ex}} - \text{pD}$ profile for N-pyrimidinyl salt **14** (Figure 2, ◇) is comparable to those for **11** and **12** displaying an increase of rate constants for exchange with pD in the whole region studied.

*Deuterium Exchange Reactions of N-2,4,6-Tribromophenyl- **11** and N-2,6-Dichlorophenyl- **12** Triazolium Tetrafluoroborates*

Deuterium exchange kinetic data for N-2,4,6-tribromophenyl- and N-2,6-dichlorophenyl triazolium salts **11** and **12** show the same dependencies on pD as observed for N-pentafluorophenyl salt **6a**. In particular, there is a marked change in the dependence of $\log k_{\text{ex}}$ values on pD under more acidic conditions closely similar

to data for **6a** and more significant than for **6b-6f**. At $pD_s > 1.5$, values of $\log k_{ex}$ for **11** and **12** increase linearly with pD and the data may be fit by a line of unit slope indicating a first order dependence on deuteroxide ion concentration in this region. This is consistent with a mechanism involving deuteroxide-catalyzed C3-H/D exchange of the monocationic triazolium ion substrate (Path A, Scheme 1). At $pD_s < 1.5$, the dependencies of $\log k_{ex}$ on pD decrease and data points in this region deviate upwards from the line of unit slope that fits the remaining data at higher pD_s . There is also the beginning of a further downward break at the lowest pD values as observed previously for N-pentafluorophenyl triazolium salt **6a**. By comparison, the profile for unsubstituted N-phenyl salt **6d** (Figure 2, ■) is essentially linear with slope unity for all data points except at $pD_s < 0.2$.

As in our previous study for **6a**, the data for salts **11** and **12** fit well to a kinetic scheme allowing for the occurrence of either Paths B or C at lower pD_s in conjunction with Path A at higher pD_s . The $\log k_{ex} - pD$ data fits well to Eqns. (2) or (3) which allow for Paths A and B or Paths A and C, respectively. In these equations, k_{DO} ($M^{-1}s^{-1}$) is the second order rate constant for deprotonation of monocationic triazolium ion (*c.f.* **6**, Scheme 1) by deuteroxide, $K_w = 10^{-14.87}$ is the ion product of D_2O at 25 °C, $\gamma_{DO} = 0.73$ is the activity coefficient for deuteroxide ion under our experimental conditions, K_a^N is the acidity constant for ionization at N, k_{D_2O} (s^{-1}) is the first order rate constant for deprotonation of monocationic triazolium ion at C(3) by solvent D_2O and k_{DO}' ($M^{-1}s^{-1}$) is the second order rate constant for deprotonation of dicationic triazolium ion (*c.f.* **8**, Scheme 1) by deuteroxide ion. Fitting to either equation yields identical values for k_{DO} and K_a^N (Table 1), whereas values for k_{D_2O} or k_{DO}' are obtained by fitting to Eqn. (2) or Eqn. (3), respectively (Table 2). For comparison, previously reported data^[21] for N-pentafluorophenyl and phenyl salts **6a** and **6d** are also included in Tables 1 and 2.

$$\log k_{ex} = \log \frac{K_a^{N1} \left(\frac{k_{DO} K_w}{\gamma_{DO}} 10^{pD} + (K_a^{N1} k_{D_2O}) \right)}{(K_a^{N1} + 10^{-pD})} \quad (2)$$


$$\log k_{\text{ex}} = \log \frac{k_{\text{DO}} K_{\text{a}}^{\text{N}} + 10^{-\text{pD}}}{(K_{\text{a}}^{\text{N}} + 10^{-\text{pD}})} \quad (3)$$

Values of the second order rate constants, k_{DO} , for C(3)-deprotonation by deuterioxide ion of triazolium salts **6a**, **6d**, **11** and **12** decrease in the order **6a** > **11** > **12** > **6d** although the difference across this series is only 10-fold (Table 1). Similarly small N-aryl substituent effects were observed for the twenty triazolium salts previously studied with k_{DO} values only varying by a maximum of 37-fold across this large series. The order of reactivity of triazolium salts corresponds to an increase in rate constant for deprotonation at C(3) with more electron-withdrawing N-aryl substituents.

The acidity constants for protonation at nitrogen, K_{a}^{N} , increase in the order **6a** < **11** < **12** and correspond to $\text{p}K_{\text{a}}^{\text{N}}$ values of 0.18, 0.04, 0.01, respectively (Table 1). The relatively large fitting errors associated with these K_{a}^{N} values are expected as only ~50% N-protonation has occurred at $\text{pD} = 0$. The values for $\text{p}K_{\text{a}}^{\text{N}}$ are very similar within error, and the data shows that there is a small increase in the degree of N-protonation within the normal pH range in the order **12** < **11** < **6a**. Importantly, as observed previously for **6a**, the $\text{p}K_{\text{a}}^{\text{N}}$ values for **11** and **12** must be substantially higher than for N-phenyl salt **6d** (Figure 2) and other N-aryl salts **6b-f**, as the pD profiles for the latter suggests no substantial protonation at N1 occurs in the normal pD range implying lower $\text{p}K_{\text{a}}^{\text{N}}$ values. In particular, the profiles for N-4-cyanophenyl **6b** and 2,6-dichlorophenyl **12** substrates are almost superimposable at $\text{pDs} > 1.5$ yielding closely similar k_{DO} values (Figure S17), however, the altered dependence on pD is more marked for the latter under more acidic conditions. As the electronic substituent effect on k_{DO} is similar in both cases, this supports the existence of an additional *ortho*-chloro substituent effect to explain the observed differences in $\text{p}K_{\text{a}}^{\text{N}}$. This small N-aryl net *donor* effect on $\text{p}K_{\text{a}}^{\text{N}}$ in these cases is opposite to the normal inductive through-bond electron accepting substituent effect of these substituents, which would favour an increase in acidity as observed at C(3). By contrast, we observe an increase

in the basicity of N(1) in the presence of *ortho*-fluoro, bromo and chloro substituents relative to other N-aryl triazolium ions **6b-f**.

Previously, we suggested that Path C rather than B was the most likely mechanism under more acidic conditions to explain the altered dependence of $\log k_{\text{ex}}$ values on pD . This was mainly based on a comparison of rate constants for deuterium exchange for triazolium ion **6a** with analogous literature data for thiazolium ions **15**.^[23, 25] There is no additional site for protonation within the thiazolium ring of **15** unlike for the triazolium analogue **6a**. Washabaugh and Jencks did observe the onset of a true pD -independent solvent reaction for four thiazolium salts **15** in concentrated DCl solutions at $pDs \ll 0$ yielding rate constants, k_{D_2O} , that range from $1.6 \times 10^{-9} \text{ s}^{-1}$ up to $9.4 \times 10^{-8} \text{ s}^{-1}$. In a comparison of data for triazolium and thiazolium salts, we previously noted that the difference in k_{DO} values is significantly smaller than for k_{D_2O} values. As an example, $k_{DO} = 4.67 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$ for the N-cyanomethyl thiazolium salt **15** ($R_1 = \text{CN}$, $R_2 = \text{H}$), which is 14-fold lower than $k_{DO} = 6.82 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$ for N-pentafluorophenyl triazolium salt **6a** (Table 1), whereas there is a much greater 650-fold difference for the corresponding k_{D_2O} values [$k_{D_2O} = 9.4 \times 10^{-8} \text{ s}^{-1}$ for **15** ($R_1 = \text{CN}$, $R_2 = \text{H}$) versus $k_{D_2O} = 6.1 \times 10^{-5} \text{ s}^{-1}$ for **6a** (Table 2)]. It is difficult to explain a substantially larger ring effect on the solvent compared with the deuterioxide-catalyzed deuterium exchange reactions, and, we suggested that this provides evidence that Path B is likely not occurring under our conditions. Further, the N-protonated dicationic substrate would be expected to be more acidic at C(3) than the monocationic analogue, and it seems logical (due to the requirement for N-protonation to explain the observed $\log k_{\text{ex}} - pD$ data for **6a**) that a deuterioxide reaction of the dication would be more likely than a water reaction on the monocation given the greater reactivity of *both* the substrate and the base in the former case albeit at very low concentrations of DO^- .



The new data in Table 2 for triazolium salts **11** and **12** supports these conclusions. The k_{D_2O} values calculated for **11** and **12** using Eqn. (2) are 85 - 12,500

fold larger than observed for the true pD -independent solvent reactions of corresponding thiazolium ions, whereas there is a much smaller ring effect on k_{DO} values, again suggesting that Path B does not occur for the former triazolium ions. Estimates for k_{DO}' calculated for C(3)-deprotonation of the dicationic conjugate acids of **11** and **12** (*c.f.* **8** in Scheme 2), for reaction *via* Path C, are all at the diffusional limit (Table 2). This is logical given that k_{DO} values for the less reactive monocationic salts are already as high as $\sim 10^8 \text{ M}^{-1}\text{s}^{-1}$. Further, as will be seen below, the observed acid catalysis in the case of N-pyridyl salt **13** is most logically accounted for in tandem with Path C for the other triazolium salts at lower pD s.

Deuterium Exchange Reactions of N-Pyridyltriazolium 13 and N-2,6-Pyrimidinyl- 14 Triazolium Tetrafluoroborates

As for triazolium salts **11-12**, values of $\log k_{ex}$ for N-pyridyl salt **11** increase linearly at pD s > 1.5 (Figure 2, \blacklozenge), and the data may be fit by a line of unit slope consistent with a first order dependence on deuterioxide ion concentration in this region and deuterium exchange via Path A (Scheme 1). In contrast with data for the other triazolium salts in Figure 1, values of $\log k_{ex}$ for **13** *decrease* with pD in the region from $pD = 0$ to 1.3. This observed acid catalysis of deuterium exchange *requires* protonation of substrate and a subsequent pD -independent C(3)-deprotonation of the resulting dicationic substrate.


The $\log k_{ex} - pD$ data for N-pyridyl salt **13** fits well to Eqn. (4), which allows for both deuterioxide-catalyzed exchange on the monocationic triazolium salt (Path A, Scheme 1) and, additionally, a pD -independent C(3)-deprotonation reaction on the dicationic substrate. In Eqn. (4), k_{DO} , K_w and γ_{DO} are as defined above. The first order rate constant k' (s^{-1}) refers to pD -independent deprotonation of dicationic substrate and mechanistic options for this process are discussed below. In this case, either the pyridyl nitrogen or N1 of the triazolium ring could potentially be protonated to give a dicationic species. The $pK_a^N = -0.05$ calculated for N-pyridyl substrate **13** (Table 1) is similar to pK_a^N values for triazolium salts **6a**, **11** and **12** for which additional protonation can only occur at N(1) of the triazolium ring. However, $pK_a^N = -0.05$ may also be consistent with the acidity constant for the pyridinium nitrogen. Numerous solution studies establish the pK_a of the N-protonated pyridinium ion at ~ 5 ^[ref] and this

would be expected to substantially decrease in the presence of a monocationic triazolium substituent.

$$\log k_{\text{ex}} = \log \frac{k_{\text{DO}} K_{\text{a}}^{-1} (10^{-pD})}{(K_{\text{a}}^{\text{N}} + 10^{-pD})} + k' (10^{-pD}) \quad (4)$$

Possible mechanisms formally consistent with the observed acid catalysis of deuterium exchange for N-pyridyl salt **13** at lower pD s are shown in Scheme 3. In analyzing these options, a key consideration is whether these mechanisms can explain why a pD -independent reaction of N-protonated salt is possible for **13** but has not been observed for any other triazolium ion. Option D1 (Scheme 3) involves initial protonation on the N1 of the triazolium ring followed by pD -independent deprotonation by D_2O *without* direct involvement of the pyridyl ring. This mechanism may be discounted as acid catalysis of deuterium exchange has not been observed for any other triazolium ion studied to date, and a remote pyridyl substituent would not be expected to drastically increase the rate of deprotonation of dicationic substrate by solvent, especially as k_{DO} values for exchange *via* Path A (Scheme 1) are similar for **13** and other triazolium salts studied. Option D2 (Scheme 3) involves initial protonation on the pyridyl nitrogen, rather than N1 of the triazolium ring, followed by pD -independent deprotonation by water. Remote protonation on the adjacent pyridyl ring would not significantly increase the rate of deprotonation at C(3) by solvent to enable competition with Path C (Scheme 1). The latter would be expected to be faster due to N1-protonation on the more proximal triazole ring with subsequent C(3) deprotonation by more basic deuterioxide ion. N-protonation of the pyridyl substituent could increase the rate of deprotonation by water at C(3) to compete with a water reaction of the monocationic triazolium salt (Path B, Scheme 1)). However, as the data for all other triazolium salts requires protonation at N1, this latter option is self-contradictory. Protonation of the more proximal triazolium ring should then also result in a competing water reaction like D1 above and acid catalysis would also be expected for the other triazolium salts.

Scheme 3 Potential Mechanistic Options for pD -independent deprotonation of a dicationic N-deuterated N-pyridyl triazolium salt.



Another possibility is that a shared hydrogen bond forms between N1 on the triazole and the pyridyl nitrogen thereby accelerating the rate of deprotonation at C(3) by solvent (Option D3, Scheme 3). If the pK_a s of the two hydrogens are closely matched, then a stronger hydrogen bond could result depending on spatial geometry around the two nitrogens. It is difficult to envisage why this would occur only for a solvent deprotonation reaction and not for deprotonation by deuteroxide ion. A final option (D4, Scheme 3) is compatible with the occurrence of acid catalysis in the case of the N-pyridyl salt only, and also supports Path C as the main mechanism for deuterium exchange at lower pD s for the other triazolium salts **6a**, **11** and **12** rather than Path B. In this mechanism, observed acid catalysis can be explained by N-protonation on the triazolium ring accompanied by intramolecular deprotonation at C(3) by the pyridyl nitrogen. This intramolecular deprotonation could be direct or may involve one or more solvent molecules. As this option is not available to the other substrates **6a**, **11** and **12**, it can provide an explanation for the singular occurrence of acid catalysis in the case of N-pyridyl substrate **13** only.

The $\log k_{\text{ex}} - pD$ profile for N-pyrimidinyl salt **14** (Figure 2, \diamond) is comparable to those for **11** and **12** displaying an increase of rate constants for exchange with pD in the whole region studied and no acid catalysis of deuterium exchange is observed at lower pD s. This suggests that intramolecular catalysis, through C(3) deprotonation


by the adjacent pyrimidine ring, is not occurring in this case, which is logical given the much decreased basicity of a simple monocyclic pyrimidine relative to a pyridine nitrogen (pK_a s of 5.1 and 1.1 for N-protonated pyridinium and pyrimidinium ions, respectively).^{REFS} The data for N-pyrimidinyl salt **14** fits well to Eqns. (2) or (3), and the resulting values for k_{DO} , K_a^N , and, k_{D2O} or k_{DO}' are shown in Tables 1 and 2. Significantly, the observed pK_a^N for the pyrimidinyl substrate **14** is very similar to that observed for N-pyridyl substrate **13**, and provides further evidence that protonation occurs on the triazolium N(1) rather than the pyridine or pyrimidinium rings, which would be expected to yield more different pK_a s. This lends further support to reaction *via* Path C as the dominant mechanism for all salts, except N-pyridyl triazolium ion **13**, at lower pD s.

Estimation of Carbon acid pK_a Values

The carbon acid pK_a values for deprotonation at C(3) for monocationic triazolium salts **11-14** may be determined using Eqn. (5), which is derived for Scheme 4.^[18, 20-22, 26-30] In this equation, k_{HO} ($M^{-1}s^{-1}$) is the second order rate constant for deprotonation at C(3) by hydroxide ion, which may be calculated from the corresponding k_{DO} value using a value of $k_{HO}/k_{DO} = 2.4$ ^[31] for the secondary solvent isotope effect on the basicity of HO^- in H_2O versus DO^- in D_2O . As discussed previously,^[18, 20-21] the absence of significant general base catalysis of exchange provides evidence that the reverse protonation of the triazol-3-ylidene **7** by water is equal or close to the limiting rate constant for the physical process of dielectric relaxation of solvent ($k_{HOH} \leq k_{reorg} = 10^{11} s^{-1}$ ^[32-33]).

$$pK_a = pK_w + \log \frac{k_{HOH}}{k_{HO}} \quad (5)$$

Scheme 4 Equilibrium for deprotonation by hydroxide of triazolium ions **6** at C(3).



The C(3)-H pK_a values for triazolium salts **11-14** are similar and range from 16.7-17.3. The values are comparable to those estimated for related thiazolium ions

(*c.f.* **15**), however, are substantially lower than our previously published values for the conjugate acids of imidazole-2-ylidenes **2**, imidazolin-2-ylidenes **3**, and trihydropyrimidin-2-ylidenes. This is due to the presence of the additional electron withdrawing ring nitrogen, which increases the stability of the formally neutral NHC **7** relatively to the cationic conjugate acid **6**. This large in-plane electron withdrawing effect of nitrogen is relatively common in heterocyclic systems. As mentioned earlier, the pK_a of N-protonated pyrimidine is 4 units lower than for the pyridinium ion due to the additional ring nitrogen atom in the former system.

The C(3)-carbon acid pK_a values for N(1)-protonated dicationic triazolium salts (*c.f.* **8**, Scheme 1) are expected to be substantially lower than for monocationic triazolium ions **6**. The k_{DO}' values for N-protonated **11-14** (Table 2) are at the diffusional limit and it is predicted that the reverse protonation of the monocationic NHCs (*c.f.* **9**) will fall below the upper limiting rate constant for protonation by solvent ($k_{reorg} = 10^{11} \text{ s}^{-1}$). Using an average value of $k_{DO}' = 5 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$ together with $k_{HOH} \leq 10^{11} \text{ s}^{-1}$ in Eqn. (5) yields an upper limit estimate of $pK_a \leq 15.3$ for the C(3)-carbon acidity of a dicationic triazolium ions. These pK_a s establish the conjugate acids **8** of monocationic carbenes **9** as the most acidic of all NHC families studied to date in aqueous solution. Furthermore, to our knowledge, a monocationic NHC **9** or an N1-alkylated analogue has not been directly isolated to date, and these results provide evidence for the transient formation of these species in aqueous solution under acidic conditions.^[34]

CONCLUSIONS

Our studies of the deuterium exchange reactions of *ortho*-disubstituted triazolium salts **11-12** provides additional evidence for formal 2-heteroatom donor effects on pK_a values at N(1). In particular, there is a marked change in the dependence of $\log k_{ex}$ values on pD under more acidic conditions closely similar to data for **6a** and more significant than for **6b-6f** where the latter have only hydrogens or methyl groups rather than halogens in the 2-position. Although the pD profiles for all the triazolium ions **6a-f** and **11-12** are very similar above pD 1.5 yielding k_{DO} values within ~10-fold of each other, the altered dependence on pD under more acidic conditions is more dominant for 2-halo-substituted ions **6a**, **11** and **12** than for **6b-6f**. The calculated pK_a^N values decrease in the order $F > Br > Cl$ although the difference is very small

across the series. By contrast, the corresponding C(3)-carbon acid pK_a s increase across this series as a result of a normal electron-withdrawing inductive effect of these substituents. The apparent donor effect on pK_a^N , which favours protonation at N(1), may be to suppress unfavourable electrostatic interactions between N(1) and spatially proximal N-aryl *ortho*-halogens, or, could be a result of stabilizing $N^+H\dots ortho-X$ interactions in the N-protonated salt.

The present study also reveals unique substituent effects for the N-(2-pyridyl)-triazolium system **13**, which shows distinct acid catalysis of deuterium exchange at lower pD values. This data is best explained by invoking an intramolecular general base role for the N-(2-pyridyl) substituent in conjunction with N1-protonation on the triazolium ring.

Overall, these results highlight the varying roles of *ortho*-heteroatoms in influencing the chemical behavior of widely-used triazolium salt organic catalysts.

ACKNOWLEDGEMENTS

We thank the EPSRC (DET, RSM and CJC) and Durham University (PQ, DJJ and CRB) for funding of this project.

REFERENCES

- [1] A. T. Biju, N. Kuhl, F. Glorius, *Acc. Chem. Res.* **2011**, *44*, 1182.
- [2] X. Bugaut, F. Glorius, *Chem. Soc. Rev.* **2012**, *41*, 3511.
- [3] D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.* **2007**, *107*, 5606.
- [4] J. Mahatthananchai, J. W. Bode, *Chem. Sci.* **2012**, *3*, 192.
- [5] N. Marion, S. Díez-González, S. P. Nolan, *Angew. Chem. Int. Ed.* **2007**, *46*, 2988.
- [6] C. A. Rose, S. Gundala, C.-L. Fagan, J. F. Franz, S. J. Connon, K. Zeitler, *Chem. Sci.* **2012**, *3*, 735.
- [7] S. J. Ryan, L. Candish, D. W. Lupton, *Chem. Soc. Rev.* **2013**, *42*, 4906.
- [8] R. W. Alder, M. E. Blake, L. Chaker, J. N. Harvey, F. Paolini, J. Schutz, *Angew. Chem. Int. Ed.* **2004**, *43*, 5896.
- [9] R. H. Crabtree, *Coord. Chem. Rev.* **2013**, *257*, 755.
- [10] S. J. Hock, L.-A. Schaper, W. A. Herrmann, F. E. Kuehn, *Chem. Soc. Rev.* **2013**, *42*, 5073.
- [11] W. Liu, R. Gust, *Chem. Soc. Rev.* **2013**, *42*, 755.
- [12] O. R. Luca, R. H. Crabtree, *Chem. Soc. Rev.* **2013**, *42*, 1440.
- [13] C. D. Martin, M. Soleilhavoup, G. Bertrand, *Chem. Sci.* **2013**, *4*, 3020.
- [14] D. Martin, M. Melaimi, M. Soleilhavoup, G. Bertrand, *Organometallics* **2011**, *30*, 5304.

- [15] M. Melaimi, M. Soleilhavoup, G. Bertrand, *Ang. Chem. Int. Ed.* **2010**, *49*, 8810.
- [16] B. M. Neilson, A. G. Tennyson, C. W. Bielawski, *J. Phys. Org. Chem.* **2012**, *25*, 531.
- [17] V. P. Reddy, A. Vadapalli, E. Sinn, N. Hosmane, *J. Organomet. Chem.* **2013**, *747*, 43.
- [18] T. L. Amyes, S. T. Diver, J. P. Richard, F. M. Rivas, K. Toth, *J. Am. Chem. Soc.* **2004**, *126*, 4366.
- [19] J. P. Guthrie, R. A. M. O'Ferrall, A. C. O'Donoghue, W. E. Waghorne, I. Zrinski, *J. Phys. Org. Chem.* **2003**, *16*, 582.
- [20] E. M. Higgins, J. A. Sherwood, A. G. Lindsay, J. Armstrong, R. S. Massey, R. W. Alder, A. C. O'Donoghue, *Chem. Commun.* **2011**, *47*, 1559.
- [21] R. S. Massey, C. J. Collett, A. G. Lindsay, A. D. Smith, A. C. O'Donoghue, *J. Am. Chem. Soc.* **2012**, *134*, 20421.
- [22] A. C. O'Donoghue, R. S. Massey, in *Contemporary Carbene Chemistry, Vol. 7* (Eds.: R. A. Moss, M. P. Doyle), Wiley, Hoboken, New Jersey, **2014**, pp. 75.
- [23] M. W. Washabaugh, W. P. Jencks, *Biochemistry-U.S.* **1988**, *27*, 5044.
- [24] C. J. Collett, R. S. Massey, O. R. Maguire, A. S. Batsanov, A. C. O'Donoghue, A. D. Smith, *Chem. Sci.* **2013**, *4*, 1514.
- [25] M. W. Washabaugh, W. P. Jencks, *J. Am. Chem. Soc.* **1989**, *111*, 683.
- [26] T. L. Amyes, J. P. Richard, *J. Am. Chem. Soc.* **1992**, *114*, 10297.
- [27] T. L. Amyes, J. P. Richard, *J. Am. Chem. Soc.* **1996**, *118*, 3129.
- [28] R. J. Delley, S. Bandyopadhyay, M. A. Fox, C. Schliehe, D. R. W. Hodgson, F. Hollfelder, A. J. Kirby, A. C. O'Donoghue, *Org. Biomol. Chem.* **2012**, *10*, 590.
- [29] J. P. Richard, G. Williams, J. L. Gao, *J. Am. Chem. Soc.* **1999**, *121*, 715.
- [30] J. P. Richard, G. Williams, A. C. O'Donoghue, T. L. Amyes, *J. Am. Chem. Soc.* **2002**, *124*, 2957.
- [31] V. Gold, S. Grist, *J. Chem. Soc. Perkin Trans. 2* **1972**, 89.
- [32] U. Kaatze, *J. Chem. Eng. Data* **1989**, *34*, 371.
- [33] U. Kaatze, R. Pottel, A. Schumacher, *J. Phys. Chem.* **1992**, *96*, 6017.
- [34] Rate constants k_{DO} are within error of typical bimolecular values for diffusion of small molecules in solution ($k_{\text{d}} = 5 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$). This supports a stepwise, rather than concerted, C(3)-H/D exchange mechanism on dicationic salt **8** via a distinct monocationic NHC intermediate.




Figure 1. pD -rate profiles for the deuterium exchange reactions of the C(3)-proton of triazolium salts **11** (\blacktriangle) and **12** (\blacktriangledown) in D_2O at 25 °C and $I = 1.0$ (KCl). Also plotted for comparison are $\log k_{\text{ex}} - pD$ data taken from R. S. Massey *et al*^[21] for the deuterium exchange reactions of triazolium salts **6a** (\bullet) and **6d** (\blacksquare). The solid lines show the fits of the data to eq 3. For triazolium salt **11** and **12**, extra datapoints (\triangle and \triangledown) were obtained in 2 M DCl (shown as open symbols on the same profiles as for **6a**).




Figure 2. pD -rate profiles for the deuterium exchange reactions of the C(3)-proton of triazolium salts **13** (\blacklozenge) and **14** (\diamond) in D_2O at 25 °C and $I = 1.0$ (KCl). Also plotted for comparison are $\log k_{\text{ex}} - pD$ data taken from R. S. Massey *et al*^[21] for the deuterium exchange reactions of triazolium salts **6a** (\bullet) and **6d** (\blacksquare). The solid lines show the fits of the data to eq 3.

Table 1 Second order rate constants for deuterioxide-catalysed hydrogen-deuterium exchange at C(3) (k_{DO} , $\text{M}^{-1}\text{s}^{-1}$), carbon acid $\text{p}K_{\text{a}}^{\text{C3}}$ and $\text{p}K_{\text{a}}^{\text{N1}}$ values in aqueous solution at 25°C and ionic strength, $I = 1.0$ (KCl).

Salt	$k_{\text{DO}} (\text{M}^{-1}\text{s}^{-1})^{\text{a}}$	$\text{p}K_{\text{a}}^{\text{C3c}}$	$K_{\text{a}}^{\text{N1}} (\text{M})^{\text{d}}$	$\text{p}K_{\text{a}}^{\text{N1}}$
11	$4.29 (\pm 0.13) \times 10^8^{\text{a}}$	16.7	$1.1 (\pm 0.3)$	0.04
12	$2.71 (\pm 0.12) \times 10^8^{\text{a}}$	16.9	$1.3 (\pm 1.3)$	0.11
13	$1.17 (\pm 0.06) \times 10^{8\text{a}}$	17.3	$0.9 (\pm 0.4)$	-0.05
14	$1.07 (\pm 0.03) \times 10^{8\text{a}}$	17.3	$2.3 (\pm 1.6)$	0.36
6a	$6.82 (\pm 0.25) \times 10^8^{\text{b}}$	16.5^{b}	$1.5 (\pm 0.4)^{\text{b}}$	0.18^{b}
6d	$6.82 (\pm 0.13) \times 10^7^{\text{b}}$	17.5^{b}	—	—

^aValues of k_{DO} ($\text{M}^{-1}\text{s}^{-1}$) obtained by fitting $\log k_{\text{ex}} - \text{p}D$ data to Eqn. (2) or (3). ^bTaken from R. S. Massey *et al.*^[21] ^c $\text{p}K_{\text{a}}^{\text{C3}}$ values obtained by application of Eqn. (5). ^dValues of K_{a}^{N1} (M) obtained by fitting $\log k_{\text{ex}} - \text{p}D$ data to Eqn. (2) or (3).

Table 2 Predicted rate constants for k_{D2O} or k_{DO}' based on kinetic fitting to Eqn. (2) or Eqn. (3).

Salt	$k_{\text{D2O}} (\text{s}^{-1})^{\text{a}}$	$k_{\text{DO}}' (\text{M}^{-1}\text{s}^{-1})^{\text{d}}$
11	$2.0 \times 10^{-5} (\pm 3.8 \times 10^{-6})$	$1.5 \times 10^{10} (\pm 2.8 \times 10^9)$
12	$8.0 \times 10^{-6} (\pm 6.6 \times 10^{-6})$	$5.7 \times 10^9 (\pm 4.7 \times 10^9)$
14	$4.7 \times 10^{-6} (\pm 2.6 \times 10^{-6})$	$5.9 \times 10^9 (\pm 3.3 \times 10^9)$
6a	$6.1 \times 10^{-5} (\pm 3.6 \times 10^{-7})^{\text{b}}$	$3.3 \times 10^{10} (\pm 2.0 \times 10^9)^{\text{b}}$
13	$1.4 \times 10^{-4} (\pm 4.3 \times 10^{-5})^{\text{c}}$	—

^aValues of k_{D2O} (s^{-1}) obtained by fitting $\log k_{\text{ex}} - \text{p}D$ data to Eqn. (2). ^bTaken from R. S. Massey *et al.*^[21] ^cValue of k' (s^{-1}) obtained by fitting $\log k_{\text{ex}} - \text{p}D$ data to Eqn. (4). ^dValues of k_{DO}' ($\text{M}^{-1}\text{s}^{-1}$) obtained by fitting $\log k_{\text{ex}} - \text{p}D$ data to Eqn. (3).