

Cosmological implications of baryon acoustic oscillation measurements

Éric Aubourg,¹ Stephen Bailey,² Julian E. Bautista,¹ Florian Beutler,² Vaishali Bhardwaj,^{3,2} Dmitry Bizyaev,⁴ Michael Blanton,⁵ Michael Blomqvist,⁶ Adam S. Bolton,⁷ Jo Bovy,⁸ Howard Brewington,⁴ J. Brinkmann,⁴ Joel R. Brownstein,⁷ Angela Burden,⁹ Nicolás G. Busca,^{1,10,11} William Carithers,² Chia-Hsun Chuang,¹² Johan Comparat,¹² Rupert A. C. Croft,^{13,14} Antonio J. Cuesta,^{15,16} Kyle S. Dawson,⁷ Timothée Delubac,¹⁷ Daniel J. Eisenstein,¹⁸ Andreu Font-Ribera,² Jian Ge,¹⁹ J.-M. Le Goff,²⁰ Satya Gontcho A. Gontcho,¹⁶ J. Richard Gott III,²¹ James E. Gunn,²¹ Hong Guo,^{22,7} Julien Guy,^{23,2} Jean-Christophe Hamilton,¹ Shirley Ho,¹³ Klaus Honscheid,^{24,25} Cullan Howlett,⁹ David Kirkby,⁶ Francisco S. Kitaura,²⁶ Jean-Paul Kneib,^{17,27} Khee-Gan Lee,²⁸ Dan Long,⁴ Robert H. Lupton,²¹ Mariana Vargas Magaña,¹ Viktor Malanushenko,⁴ Elena Malanushenko,⁴ Marc Manera,^{9,29} Claudia Maraston,⁹ Daniel Margala,⁶ Cameron K. McBride,¹⁸ Jordi Miralda-Escudé,^{30,16} Adam D. Myers,³¹ Robert C. Nichol,⁹ Pasquier Noterdaeme,³² Sebastián E. Nuza,²⁶ Matthew D. Olmstead,⁷ Daniel Oravetz,⁴ Isabelle Pâris,³³ Nikhil Padmanabhan,¹⁵ Nathalie Palanque-Delabrouille,^{2,20} Kaike Pan,⁴ Marcos Pellejero-Ibanez,^{34,35} Will J. Percival,⁹ Patrick Petitjean,³² Matthew M. Pieri,³⁶ Francisco Prada,^{12,37,38} Beth Reid,^{2,39} James Rich,²⁰ Natalie A. Roe,² Ashley J. Ross,^{9,25} Nicholas P. Ross,⁴⁰ Graziano Rossi,^{41,20} Jose Alberto Rubiño-Martín,^{34,35} Ariel G. Sánchez,⁴² Lado Samushia,^{43,44} Ricardo Tanausú Génova Santos,³⁴ Claudia G. Scóccola,^{12,34,45} David J. Schlegel,² Donald P. Schneider,^{46,47} Hee-Jong Seo,^{25,48} Erin Sheldon,⁴⁹ Audrey Simmons,⁴ Ramin A. Skibba,⁵⁰ Anže Slosar,⁴⁹ Michael A. Strauss,²¹ Daniel Thomas,⁹ Jeremy L. Tinker,⁵ Rita Tojeiro,⁹ Jose Alberto Vazquez,⁴⁹ Matteo Viel,^{33,51} David A. Wake,^{52,53} Benjamin A. Weaver,⁵ David H. Weinberg,^{25,54} W. M. Wood-Vasey,⁵⁵ Christophe Yèche,²⁰ Idit Zehavi,⁵⁶ and Gong-Bo Zhao^{57,9}

(BOSS Collaboration)

¹*APC, Astroparticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris Cité, 10, rue Alice Domon & Léonie Duquet, 75205 Paris Cedex 13, France*

²*Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720, USA*

³*Department of Astronomy, University of Washington, Box 351580, Seattle, Washington 98195, USA*

⁴*Apache Point Observatory, P.O. Box 59, Sunspot, New Mexico 88349-0059, USA*

⁵*Center for Cosmology and Particle Physics, New York University, New York, New York 10003, USA*

⁶*Department of Physics and Astronomy, UC Irvine, 4129 Frederick Reines Hall, Irvine, California 92697, USA*

⁷*Department Physics and Astronomy, University of Utah, Utah 84112, USA*

⁸*Institute for Advanced Study, Einstein Drive, Princeton, New Jersey 08540, USA*

⁹*Institute of Cosmology & Gravitation, Dennis Sciama Building, University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom*

¹⁰*Observatório Nacional, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil*

¹¹*Laboratório Interinstitucional de e-Astronomia, - LIneA, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil*

¹²*Instituto de Física Teórica (UAM/CSIC), Universidad Autonoma de Madrid, Cantoblanco, E-28049 Madrid, Spain*

¹³*Department of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, Pennsylvania 15213, USA*

¹⁴*Astrophysics, University of Oxford, Keble Road, Oxford OX13RH, United Kingdom*

¹⁵*Department of Physics, Yale University, 260 Whitney Ave, New Haven, Connecticut 06520, USA*

¹⁶*Institut de Ciències del Cosmos, Universitat de Barcelona, IEEC-UB, Martí i Franquès 1, E08028 Barcelona, Spain*

¹⁷*Laboratoire d'astrophysique, Ecole Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, CH-1290 Versoix, Switzerland*

¹⁸*Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, Massachusetts 02138, USA*

¹⁹*Department of Astronomy, University of Florida, Gainesville, Florida 32611, USA*

²⁰*CEA, Centre de Saclay, IRFU, 91191 Gif-sur-Yvette, France*

²¹*Department of Astrophysical Sciences, Princeton University, Ivy Lane, Princeton, New Jersey 08544, USA*

²²*Key Laboratory for Research in Galaxies and Cosmology of Chinese Academy of Sciences, Shanghai Astronomical Observatory, Shanghai 200030, China*

²³*LPNHE, CNRS/IN2P3, Université Pierre et Marie Curie Paris 6, Université Denis Diderot Paris 7, 4 place Jussieu, 75252 Paris CEDEX, France*

²⁴*Department of Physics, The Ohio State University, Columbus, Ohio 43210, USA*

²⁵*Center for Cosmology and Astro-Particle Physics, Ohio State University, Columbus, Ohio 43210, USA*

- ²⁶*Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany*
- ²⁷*CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France*
- ²⁸*Max-Planck-Institut für Astronomie, Königstuhl 17, D69117 Heidelberg, Germany*
- ²⁹*University College London, Gower Street, London WC1E 6BT, United Kingdom*
- ³⁰*Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain*
- ³¹*Department of Physics and Astronomy, University of Wyoming, Laramie, Wyoming 82071, USA*
- ³²*Institut d'Astrophysique de Paris, UPMC-CNRS, UMR7095, 98bis boulevard Arago, 75014 Paris, France*
- ³³*INAF, Osservatorio Astronomico di Trieste, Via G. B. Tiepolo 11, 34131 Trieste, Italy*
- ³⁴*Instituto de Astrofísica de Canarias (IAC), C/Vía Láctea, s/n, E-38200, La Laguna, Tenerife, Spain*
- ³⁵*Departamento Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain*
- ³⁶*A*MIDEX, Aix-Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, Marseille, France*
- ³⁷*Campus of International Excellence UAM+CSIC, Cantoblanco, E-28049 Madrid, Spain*
- ³⁸*Instituto de Astrofísica de Andalucía (CSIC), E-18080 Granada, Spain*
- ³⁹*Department of Physics, University of California, 366 LeConte Hall, Berkeley, California 94720, USA*
- ⁴⁰*Department of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, Pennsylvania 19104, USA*
- ⁴¹*Department of Astronomy and Space Science, Sejong University, Seoul, 143-747, Korea*
- ⁴²*Max-Planck-Institut für extraterrestrische Physik, Postfach 1312, Giessenbachstrasse, 85748 Garching, Germany*
- ⁴³*Department of Physics, Kansas State University, 116 Cardwell Hall, Manhattan, Kansas 666506, USA*
- ⁴⁴*National Abastumani Astrophysical Observatory, Iliia State University, 2A Kazbegi Avenue GE-1060 Tbilisi, Georgia*
- ⁴⁵*Departamento de Física Teórica, Universidad Autónoma de Madrid, E-28049 Cantoblanco, Madrid, Spain*
- ⁴⁶*Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*
- ⁴⁷*Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*
- ⁴⁸*Department of Physics and Astronomy, Ohio University, 251B Clippinger Labs, Athens, Ohio 45701*
- ⁴⁹*Brookhaven National Laboratory, 2 Center Road, Upton, New York 11973, USA*
- ⁵⁰*Center for Astrophysics and Space Sciences, Department of Physics, University of California, 9500 Gilman Dr., San Diego, California 92093 USA*
- ⁵¹*INFN/National Institute for Nuclear Physics, Via Valerio 2, 34127 Trieste, Italy*
- ⁵²*Department of Astronomy, University of Wisconsin-Madison, 475 N. Charter Street, Madison, Wisconsin, 53706, USA*
- ⁵³*Department of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, United Kingdom*
- ⁵⁴*Department of Astronomy, Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁵*PITT PACC, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁵⁶*Department of Astronomy, Case Western Reserve University, Cleveland, Ohio 44106, USA*
- ⁵⁷*National Astronomy Observatories, Chinese Academy of Science, Beijing 100012, Peoples Republic of China*

(Received 18 November 2014; published 14 December 2015)

We derive constraints on cosmological parameters and tests of dark energy models from the combination of baryon acoustic oscillation (BAO) measurements with cosmic microwave background (CMB) data and a recent reanalysis of Type Ia supernova (SN) data. In particular, we take advantage of high-precision BAO measurements from galaxy clustering and the Lyman- α forest (LyaF) in the SDSS-III Baryon Oscillation Spectroscopic Survey (BOSS). Treating the BAO scale as an uncalibrated standard ruler, BAO data alone yield a high confidence detection of dark energy; in combination with the CMB angular acoustic scale they further imply a nearly flat universe. Adding the CMB-calibrated physical scale of the sound horizon, the combination of BAO and SN data into an “inverse distance ladder” yields a measurement of $H_0 = 67.3 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$, with 1.7% precision. This measurement assumes standard prerecombination physics but is insensitive to assumptions about dark energy or space curvature, so agreement with CMB-based estimates that assume a flat Λ CDM cosmology is an important corroboration of this minimal cosmological model. For constant dark energy (Λ), our BAO + SN + CMB combination yields matter density $\Omega_m = 0.301 \pm 0.008$ and curvature $\Omega_k = -0.003 \pm 0.003$. When we allow more general forms of evolving dark energy, the BAO + SN + CMB parameter constraints are always consistent with flat Λ CDM

values at $\approx 1\sigma$. While the overall χ^2 of model fits is satisfactory, the Ly α BAO measurements are in moderate ($2\text{--}2.5\sigma$) tension with model predictions. Models with early dark energy that tracks the dominant energy component at high redshift remain consistent with our expansion history constraints, and they yield a higher H_0 and lower matter clustering amplitude, improving agreement with some low redshift observations. Expansion history alone yields an upper limit on the summed mass of neutrino species, $\sum m_\nu < 0.56$ eV (95% confidence), improving to $\sum m_\nu < 0.25$ eV if we include the lensing signal in the Planck CMB power spectrum. In a flat Λ CDM model that allows extra relativistic species, our data combination yields $N_{\text{eff}} = 3.43 \pm 0.26$; while the Ly α BAO data prefer higher N_{eff} when excluding galaxy BAO, the galaxy BAO alone favor $N_{\text{eff}} \approx 3$. When structure growth is extrapolated forward from the CMB to low redshift, standard dark energy models constrained by our data predict a level of matter clustering that is high compared to most, but not all, observational estimates.

DOI: [10.1103/PhysRevD.92.123516](https://doi.org/10.1103/PhysRevD.92.123516)

PACS numbers: 98.80.-k, 95.36.+x

I. INTRODUCTION

Acoustic oscillations that propagate in the pre-recombination universe imprint a characteristic scale in the clustering of matter, providing a cosmological ‘‘standard ruler’’ that can be measured in the power spectrum of cosmic microwave background (CMB) anisotropies and in maps of large-scale structure at lower redshifts [1–5]. While distance scale measurements with Type Ia supernovae (SNIa) are calibrated against systems in the local Hubble flow [6–8], the baryon acoustic oscillation (BAO) scale is computed from first principles, using physical parameters (such as the radiation, matter, and baryon densities) that are well constrained by CMB data. The difference between absolute and relative measurements, the sharpening of BAO precision with increasing redshift, and the entirely independent systematic uncertainties make BAO and SNe highly complementary tools for measuring the cosmic expansion history and testing dark energy models. In spectroscopic surveys, BAO measurements in the line-of-sight dimension allow direct determinations of the expansion rate $H(z)$, in addition to the constraints from transverse clustering on the comoving angular diameter distance $D_M(z) \propto \int_0^z cH^{-1}(z)dz$ in a flat spatial metric.¹

The first clear detections of low-redshift BAO [10,11] came from galaxy clustering analyses of the Two Degree Field Galaxy Redshift Survey (2dFGRS, [12]) and the luminous red galaxy (LRG, [13]) sample of the Sloan Digital Sky Survey (SDSS, [14]). Analyses of the final 2dFGRS and SDSS-II redshift surveys yielded BAO distance measurements with aggregate precision of 2.7% at $z \approx 0.275$ [15], subsequently sharpened to 1.9% [16] by application of reconstruction methods [17] that suppress nonlinear degradation of the BAO feature. The WiggleZ survey [18] pushed BAO measurements to higher redshifts, achieving 3.8% aggregate precision from galaxies in the

redshift range $0.4 < z < 1.0$ [19]. The Six Degree Field Galaxy Survey (6dFGS, [20]) took advantage of its 17,000 deg² sky coverage to provide a BAO measurement at low redshift, achieving 4.5% precision at $z = 0.1$ [21]. A recent reanalysis that applies reconstruction to the main galaxy sample [22] of SDSS-II obtained 3.8% precision at $z = 0.15$ [23], in a sky area that has minimal (< 3%) overlap with 6dFGS.

The Baryon Oscillation Spectroscopic Survey (BOSS, [24]) of SDSS-III [25] has two defining objectives: to measure the BAO distance scale with 1% precision from a redshift survey of 1.5 million luminous galaxies at $z = 0.2\text{--}0.7$, and to make the first BAO measurement at $z > 2$ using 3-dimensional structure in the Ly α forest absorption towards a dense grid of 160,000 high-redshift quasars. This paper explores the cosmological implications of BAO measurements from the BOSS Data Release 11 (DR11) data sample, in combination with a variety of other cosmological data. The measurements themselves, including detailed discussion of statistical uncertainties and extensive tests for systematic errors, have been presented in previous papers. For the galaxy survey, the authors of [26] report a 1.4% measurement of $D_M(z)$ and a 3.5% measurement of $H(z)$ at $z = 0.57$ (1σ uncertainties, with a correlation coefficient of -0.52), and a 2.0% measurement of $D_V(z) \equiv [D_M^2(z) \times cz/H(z)]^{1/3}$ from lower redshift BOSS galaxies at $z = 0.32$. The $D_V(z)$ precision at $z = 0.57$ is 1.0%. For the Ly α forest (often abbreviated as Ly α F below), we combine constraints from the autocorrelation function, with 2.6% precision on $H(z)$ and 5.4% precision on $D_M(z)$ [27], and the quasar-forest cross correlation, with precision of 3.3% on $H(z)$ and 3.7% on $D_M(z)$ [28], both at an effective redshift $z \approx 2.34$. While some cosmological analysis appears in these papers, the combination of galaxy and Ly α forest BAO measurements and the addition of other data allow us to constrain broader classes of cosmological models and to search for deviations from standard assumptions.

The combination of BAO measurements with precise CMB measurements from the Planck and WMAP satellites

¹We use the notation $D_M(z)$ to refer to the comoving angular diameter distance, which is also referred to in the literature as the proper motion distance [9]. This notation avoids confusion with the proper angular diameter distance $D_A(z) = D_M(z)/(1+z)$.

already yields tight constraints on the parameters of the Λ CDM cosmological model (inflationary cold dark matter with a cosmological constant and zero space curvature²) and on one-parameter extensions of this model that allow, e.g., nonzero curvature, an evolving dark energy density, or a cosmologically significant neutrino mass [29]. We also take advantage of another major recent advance, a comprehensive reassessment of the SNIa distance scale by [30] using data from the 3-year Supernova Legacy Survey [31] and SDSS-II Supernova Survey [32,33] samples and additional data at low and high redshifts. We will examine the consistency of the BAO and SNIa results for relative distances and the constraints on H_0 that emerge from an “inverse distance ladder” that combines the two data sets, in essence using SNIa to transfer the absolute calibration of the BAO scale from the intermediate redshifts where it is precisely measured down to $z = 0$. Our primary focus will be on the cosmological parameter constraints and model tests that come from combining the BAO and SNIa data with Planck CMB data.³ When fitting models to these data, we will also examine their predictions for observable measures of structure growth and compare the results to inferences from weak lensing, clusters, redshift-space distortions, and the 1-d Ly α power spectrum. The interplay of BAO, CMB, and SNIa constraints, and the more general interplay between measurements of expansion history and structure growth, are reviewed at length by [35], along with detailed introductions to the methods themselves. In particular, Sec. IV of [35] provides a thorough description of the BAO method and its motivation.

Section II describes the basic methodology of our analysis, including the relevant underlying equations, and reviews the BAO, CMB, and SN measurements that we adopt for our constraints, concluding with variants of “BAO Hubble diagrams” that illustrate our qualitative results. Section III presents the constraints obtainable by assuming that the BAO scale is a standard ruler independent of redshift without computing its physical scale; in particular, we demonstrate that galaxy and Ly α BAO alone yield a convincing detection of dark energy and that addition of the angular scale of the CMB acoustic peaks requires a nearly flat universe if dark energy is a cosmological constant. Section IV presents our inverse distance

ladder determination of H_0 , which assumes standard recombination physics but does not assume a specific dark energy model or a flat universe. Section V describes our constraints on the parameters of standard dark energy models, while Sec. VI considers models that allow early dark energy, decaying dark matter, cosmologically significant neutrino mass, or extra relativistic species. We compare the predictions of our BAO + SN + CMB constrained models to observational estimates of matter clustering in Sec. VII and summarize our overall conclusions in Sec. VIII.

II. METHODOLOGY, MODELS, AND DATA SETS

A. Methodology

A homogenous and isotropic cosmological model is specified by the curvature parameter k entering the Friedman-Robertson-Walker metric

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right], \quad (1)$$

which governs conversion between radial and transverse distances, and by the evolution of $a(t) = (1 + z)^{-1}$. In general relativity (GR), this evolution is governed by the Friedmann equation [36], which can be written in the form

$$\frac{H^2(a)}{H_0^2} = \frac{\rho(a)}{\rho_0} + \Omega_k a^{-2}, \quad (2)$$

where $H \equiv \dot{a}/a$ is the Hubble parameter, $\rho(a)$ is the total energy density (radiation + matter + dark energy), and the subscript 0 denotes the present day ($a = 1$). We define the density parameter of component x by the ratio

$$\Omega_x = \frac{\rho_x}{\rho_{\text{crit}}} = \frac{8\pi G}{3H^2} \rho_x \quad (3)$$

and the curvature parameter

$$\Omega_k = 1 - \sum \Omega_x, \quad (4)$$

where the sum is over all matter and energy components and $\Omega_k = 0$ for a flat ($k = 0$) universe. Density parameters and ρ_{crit} always refer to values at $z = 0$ unless a dependence on a or z is written explicitly, e.g., $\Omega_x(z)$. We will frequently refer to the Hubble constant H_0 through the dimensionless ratio $h \equiv H_0/100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The dimensionless quantity $\omega_x \equiv \Omega_x h^2$ is proportional to the physical density of component x at the present day.

Given the curvature parameter and $H(z)$ from the Friedmann equation, the comoving angular diameter distance can be computed as

²Throughout the paper, the notation Λ CDM refers to spatially flat models; cosmological constant models allowing nonzero curvature are denoted $o\Lambda$ CDM.

³We use the Planck 2013 data, which were publicly available at the time of our analysis and paper submission. Because best-fit parameter values from the Planck 2015 data are similar to those from the Planck 2013 data [34], we expect that using the 2015 data would make little difference to our results, though with some modest improvements in parameter uncertainties. We will present analyses that use the Planck 2015 data in concert with BOSS DR12 BAO measurements in future work.

$$D_M(z) = \frac{c}{H_0} S_k \left(\frac{D_C(z)}{c/H_0} \right) \approx D_C(z) \left[1 + \frac{1}{6} \Omega_k \left(\frac{D_C(z)}{c/H_0} \right)^2 \right], \quad (5)$$

where the line-of-sight comoving distance is

$$D_C(z) = \frac{c}{H_0} \int_0^z dz' \frac{H_0}{H(z')} \quad (6)$$

and

$$S_k(x) = \begin{cases} \sin(\sqrt{-\Omega_k}x)/\sqrt{-\Omega_k} & \Omega_k < 0, \\ \sinh(\sqrt{\Omega_k}x)/\sqrt{\Omega_k} & \Omega_k > 0, \\ x & \Omega_k = 0. \end{cases} \quad (7)$$

Positive k corresponds to negative Ω_k . We do not use the small Ω_k approximation of Eq. (5) in our calculations, but we provide it here to illustrate that for small nonzero curvature the change in distance is linear in Ω_k and quadratic in $D_C(z)$.

Curvature affects $D_M(z)$ both through its influence on $H(z)$ and through the geometrical factor in Eq. (5). The luminosity distance (relevant to supernovae) is $D_L = D_M(1+z)$.

The energy components considered in our models are pressureless (cold) dark matter, baryons, radiation, neutrinos, and dark energy. The densities of CDM and baryons scale as a^{-3} ; we refer to the density parameter of these two components together as Ω_{cb} . The energy density of neutrinos with nonzero mass scales like radiation at early times and like matter at late times, with

$$\frac{\rho_{\nu+r}(a)}{\rho_{\text{crit}}} = \frac{8\pi^3 k_B^4 G}{45\hbar^3 c^5 H_0^2} \times \left[T_{\text{CMB}}(a)^4 + T_\nu(a)^4 \sum_i I(m_i c^2 / k_B T_\nu(a)) \right], \quad (8)$$

where both CMB temperature T_{CMB} and neutrino temperature scale inversely with scale factor, and the neutrino temperature is given by $T_\nu = T_{\text{CMB}} (\frac{4}{11})^{1/3} g_c$, where $g_c = (3.046/3)^{1/4}$ accounts for a small amount of heating of neutrinos due to electron-positron annihilation. The sum in the above expression is over neutrino species with masses m_i . The integral I is given by

$$I(r) = \frac{15}{\pi^4} \int_0^\infty \frac{\sqrt{x^2 + r^2}}{e^x + 1} x^2 dx \quad (9)$$

and must be evaluated numerically. For massless neutrinos $I(0) = \frac{7}{8}$, while in the limit of very massive neutrinos

$I(r) \sim 45\zeta(3)(2\pi^4)^{-1}r$ [for $r \gg 1$; here $\zeta(3)$ is the Riemann function], i.e., scaling proportionally with a so that neutrinos behave like pressureless matter. When we refer to the $z=0$ matter density parameter Ω_m , we include the contributions of radiation (which is small compared to the uncertainties in Ω_m) and neutrinos (which are nonrelativistic at $z=0$), so that $\Omega_m + \Omega_{\text{de}} + \Omega_k \equiv 1$. Following the Planck Collaboration [29], we adopt $\sum m_\nu = 0.06$ eV with one massive and two massless neutrino species in all models except the one referred to as νCDM , where it is a free parameter. The default implies $\omega_\nu = 6.57 \times 10^{-4}$ including massless species and $\omega_\nu = 6.45 \times 10^{-4}$ excluding them. The effect of finite neutrino temperature at $z=0$ is a very small 10^{-4} relative effect. The adopted values are close to the minimum value allowed by neutrino oscillation experiments.

We consider a variety of models for the evolution of the energy density or equation-of-state parameter $w = p_{\text{de}}/\rho_{\text{de}}$. Table I summarizes the primary models discussed in the paper, though we consider some additional special cases in Sec. VI. ΛCDM represents a flat universe with a cosmological constant ($w = -1$). $o\Lambda\text{CDM}$ extends this model to allow nonzero Ω_k . $w\text{CDM}$ adopts a flat universe and constant w , and $ow\text{CDM}$ generalizes to nonzero Ω_k . $w_0 w_a \text{CDM}$ and $ow_0 w_a \text{CDM}$ allow $w(a)$ to evolve linearly with $a(t)$, $w(a) = w_0 + w_a(1-a)$. PolyCDM adopts a quadratic polynomial form for $\rho_{\text{de}}(a)$ and allows nonzero space curvature, to provide a highly flexible description of the effects of dark energy at low redshift. Finally, slow roll dark energy is an example of a one-parameter evolving- w model, based on a quadratic dark energy potential.

We focus in this paper on parameter constraints and model tests from measurements of cosmic distances and expansion rates, which we refer to collectively as ‘‘expansion history’’ or ‘‘geometric’’ constraints. We briefly consider comparisons to measurements of low-redshift matter clustering in Sec. VII. In this framework, the crucial roles of CMB anisotropy measurements are to constrain the parameters (mainly ω_m and ω_b) that determine the BAO scale and to determine the angular diameter distance to the redshift of recombination. For most of our analyses, this approach allows us to use a highly compressed summary of CMB constraints, discussed in Sec. II C below, and to compute parameter constraints with a simple and fast Markov chain Monte Carlo (MCMC) code that computes expansion rates and distances from the Friedmann equation. The code is publicly available [37] with data used in this paper.

B. BAO data

The BAO data in this work are summarized in Table II and more extensively discussed below.

The robustness of BAO measurements arises from the fact that a sharp feature in the correlation function (or an oscillatory feature in the power spectrum) cannot be readily

TABLE I. Models considered in the paper and section in the paper where they are discussed. The top section is the minimal cosmological model (with and without curvature) and various extensions in the dark energy sector. PolyCDM is a model used to mimic nonparametric methods (i.e., flexible models where the only *de facto* assumption is smoothness of the expansion history). The bottom group are various extensions of the minimal model to which we are sensitive only in conjunction with the CMB data. Throughout, Ω_{cb} is the $z = 0$ density parameter of baryons + CDM and $\rho_{\nu+r}(z)$ is the energy density of radiation + massive neutrinos. All models except ν CDM and $\Delta N_{\text{eff}}\Lambda$ CDM adopt $\sum m_\nu = 0.06$ eV and the standard radiation content $N_{\text{eff}} = 3.046$.

Name	Friedmann equation (H^2/H_0^2)	Curvature	Section(s)
Λ CDM	$\Omega_{cb}a^{-3} + \Omega_\Lambda + \rho_{\nu+r}(z)/\rho_{\text{crit}}$	no	III-V
$o\Lambda$ CDM	$\Omega_{cb}a^{-3} + \Omega_\Lambda + \rho_{\nu+r}(z)/\rho_{\text{crit}} + \Omega_k a^{-2}$	yes	III-V
w CDM	$\Omega_{cb}a^{-3} + \Omega_{\text{de}}a^{-3(1+w)} + \rho_{\nu+r}(z)/\rho_{\text{crit}}$	no	V
ow CDM	$\Omega_{cb}a^{-3} + \Omega_{\text{de}}a^{-3(1+w)} + \rho_{\nu+r}(z)/\rho_{\text{crit}} + \Omega_k a^{-2}$	yes	V
w_0w_a CDM	$\Omega_{cb}a^{-3} + \Omega_{\text{de}}a^{-3(1+w_0+w_a)} \exp[-3w_a(1-a)] + \rho_{\nu+r}(z)/\rho_{\text{crit}}$	no	V
Slow roll dark energy	$\Omega_{cb}a^{-3} + \rho_{\nu+r}(z)/\rho_{\text{crit}} + \Omega_{\text{DE}}[a^{-3}/(\Omega_m a^{-3} + \Omega_{\text{DE}})]^{\delta w_0/\Omega_{\text{DE}}}$	no	V
ow_0w_a CDM	$\Omega_{cb}a^{-3} + \Omega_{\text{de}}a^{-3(1+w_0+w_a)} \exp[-3w_a(1-a)] + \rho_{\nu+r}(z)/\rho_{\text{crit}} + \Omega_k a^{-2}$	yes	IV-V
PolyCDM	$\Omega_{cb}a^{-3} + (\Omega_1 + \Omega_k)a^{-2} + \Omega_2 a^{-1} + (1 - \Omega_{cb} - \Omega_k - \Omega_1 - \Omega_2)$	yes ^a	IV
Early dark energy	See relevant section.	no	VI A
Decaying dark matter	See relevant section.	no	VIB
ν CDM	free neutrino mass ($\sum m_\nu < 1$ eV)	no	VIC
$\Delta N_{\text{eff}}\Lambda$ CDM	nonstandard radiation component ($2 < N_{\text{eff}} < 5$)	no	VID
Tuned oscillation	See relevant section.	no	VIE

^awith Gaussian prior $\Omega_k = 0 \pm 0.1$.

mimicked by systematics, whether observational or astrophysical, as these should be agnostic about the BAO scale and hence smooth over the relevant part of the correlation function (or power spectrum). In most current analyses, the BAO scale is determined by adopting a fiducial cosmological model that translates angular and redshift separations to comoving distances but allowing the location of the BAO feature itself to shift relative to the fiducial model expectation. One then determines the likelihood of obtaining the observed two-point correlation function or power spectrum as a function of the BAO offsets, while marginalizing over nuisance parameters. These nuisance parameters characterize “broad-band” physical or observational effects that smoothly change the shape or amplitude of the underlying correlation function or power spectrum, such as scale-dependent bias of galaxies or the Ly α F, or distortions caused by continuum fitting or by variations in

star-galaxy separation. In an isotropic fit, the measurement is encoded in the α parameter, the ratio of the measured BAO scale to that predicted by the fiducial model. In an anisotropic analysis, one separately constrains α_\perp and α_\parallel , the ratios perpendicular and parallel to the line of sight. In real surveys the errors on α_\perp and α_\parallel are significantly correlated for a given redshift slice, but they are typically uncorrelated across different redshift slices. While the values of α are referred to a specified fiducial model, the corresponding physical BAO scales are insensitive to the choice of fiducial model within a reasonable range.

The BAO scale is set by the radius of the sound horizon at the drag epoch z_d when photons and baryons decouple,

$$r_d = \int_{z_d}^{\infty} \frac{c_s(z)}{H(z)} dz, \quad (10)$$

TABLE II. BAO constraints used in this work. These values are taken from [21] (6dFGS), [23] (MGS), [26] (BOSS galaxies), [27] (BOSS Ly α F auto-correlation), and [28] (BOSS Ly α F cross correlation). For our likelihood calculations, we adopt Gaussian approximations for 6dFGS and LOWZ (with 6dFGS truncated at $\Delta\chi^2 = 4$), while for others we use the full χ^2 look-up tables. The Ly α F autocorrelation and cross-correlation results are used directly; the combined Ly α F numbers are provided here for convenience.

Name	Redshift	D_V/r_d	D_M/r_d	D_H/r_d	r_{off}
6dFGS	0.106	3.047 ± 0.137
MGS	0.15	4.480 ± 0.168
BOSS LOWZ Sample	0.32	8.467 ± 0.167
BOSS CMASS Sample	0.57	...	14.945 ± 0.210	20.75 ± 0.73	-0.52
Ly α F auto-correlation	2.34	...	37.675 ± 2.171	9.18 ± 0.28	-0.43
Ly α F-QSO cross correlation	2.36	...	36.288 ± 1.344	9.00 ± 0.30	-0.39
Combined Ly α F	2.34	...	36.489 ± 1.152	9.145 ± 0.204	-0.48

where the sound speed in the photon-baryon fluid is $c_s(z) = 3^{-1/2}c[1 + \frac{3}{4}\rho_b(z)/\rho_\gamma(z)]^{-1/2}$. A precise prediction of the BAO signal requires a full Boltzmann code computation, but for reasonable variations about a fiducial model the *ratio* of BAO scales is given accurately by the ratio of r_d values computed from the integral (10). Thus, a measurement of α_\perp from clustering at redshift z constrains the ratio of the comoving angular diameter distance to the sound horizon:

$$D_M(z)/r_d = \alpha_\perp D_{M,\text{fid}}(z)/r_{d,\text{fid}}. \quad (11)$$

A measurement of α_\parallel constrains the Hubble parameter $H(z)$, which we convert to an analogous quantity:

$$D_H(z) = c/H(z), \quad (12)$$

with

$$D_H(z)/r_d = \alpha_\parallel D_{H,\text{fid}}(z)/r_{d,\text{fid}}. \quad (13)$$

An isotropic BAO analysis measures some effective combination of these two distances. If redshift-space distortions are weak, which is a good approximation for luminous galaxy surveys after reconstruction but not for the Ly α F, then the constrained quantity is the volume averaged distance

$$D_V(z) = [zD_H(z)D_M^2(z)]^{1/3}, \quad (14)$$

with

$$D_V(z)/r_d = \alpha D_{V,\text{fid}}(z)/r_{d,\text{fid}}. \quad (15)$$

There are different conventions in use for defining r_d , which differ at the 1%–2% level, but ratios of r_d for different cosmologies are independent of the convention provided one is consistent throughout. In this work we adopt the CAMB convention for r_d , i.e., the value that is reported by the linear perturbations code CAMB [38]. In practice, we use the numerically calibrated approximation

$$r_d \approx \frac{55.154 \exp[-72.3(\omega_\nu + 0.0006)^2]}{\omega_{cb}^{0.25351} \omega_b^{0.12807}} \text{ Mpc}. \quad (16)$$

This approximation is accurate to 0.021% for a standard radiation background with $N_{\text{eff}} = 3.046$, $\sum m_\nu < 0.6$ eV, and values of ω_b and ω_{cb} within 3σ of values derived by Planck. It supersedes a somewhat less accurate (but still sufficiently accurate) approximation from [26] [their Eq. (55)]. Note that $\omega_\nu = 0.0107(\sum m_\nu/1.0 \text{ eV})$, and a 0.5 (1.0) eV neutrino mass changes r_d by -0.26% (-0.92%) for fixed ω_{cb} . For neutrino masses in the range allowed by current cosmological constraints, the CMB constrains ω_{cb} rather than $\omega_{cb} + \omega_\nu$ because neutrinos

remain relativistic at recombination, even though they are nonrelativistic at $z = 0$. For the case of extra relativistic species, a useful fitting formula is

$$r_d \approx \frac{56.067 \exp[-49.7(\omega_\nu + 0.002)^2]}{\omega_{cb}^{0.2436} \omega_b^{0.128876} [1 + (N_{\text{eff}} - 3.046)/30.60]} \text{ Mpc}, \quad (17)$$

which is accurate to 0.119% if we restrict to neutrino masses in the range $0 < \sum m_\nu < 0.6$ eV and $3 < N_{\text{eff}} < 5$. Increasing N_{eff} by unity decreases r_d by about 3.2%.

For Λ CDM models (with $\sum m_\nu = 0.06$ eV, $N_{\text{eff}} = 3.046$) constrained by Planck, $r_d = 147.49 \pm 0.59$ Mpc. This 0.4% uncertainty is only slightly larger for $o\Lambda$ CDM, ow CDM, or even ow_0w_d CDM (see Table I for cosmological model definitions), because the relevant quantities ω_{cb} and ω_b are constrained by the relative heights of the acoustic peaks, not by their angular locations. The inference of matter energy densities from peak heights thus depends on correct understanding of physics in the prerecombination epoch, where curvature and dark energy are negligible in any of these models.

BAO measurements constrain cosmological parameters through their influence on r_d , their influence on $D_H(z)$ via the Friedmann equation, and their influence on $D_M(z)$ via Eq. (5). For standard models, the 0.4% error on r_d from Planck is small compared to current BAO measurement errors, so the constraints come mainly through D_H and D_M . From the Friedmann equation, we see that $D_H(z)$ is directly sensitive to the total energy density at redshift z , while $D_M(z)$ constrains an effective average of the energy density and is also sensitive to curvature.

Measurements in Table II are reported in terms of D_V/r_d , D_M/r_d , and D_H/r_d , using the r_d convention of Eq. (16). Expressed in these terms, the results are independent of the fiducial cosmologies assumed in the individual analysis papers. Note that some of the referenced papers quote values of D_A/r_d rather than D_M/r_d , differing by a factor $(1+z)$. An anisotropic analysis yields anticorrelated errors on D_M and D_H , and the correlation coefficients are reported in Table II. Each sample spans a range of redshift, and the quoted effective redshift is usually weighted by statistical contribution to the BAO measurement. Because redshift-space positions are scaled to comoving coordinates based on a fiducial cosmological model, and BAO measurements are obtained as ratios relative to that fiducial model, the values of the effective redshift in Table II can be treated as exact, e.g., one should compare the BOSS CMASS numbers to model predictions computed at $z = 0.5700$.

1. Galaxy BAO measurements

The most precise BAO measurements to date come from analyses of the BOSS DR11 galaxy sample by [26]. BOSS uses the same telescope [39] as the original SDSS, with spectrographs [40] that were substantially upgraded to improve throughput and increase multiplexing (from 640

fibers per plate to 1000). Redshift completeness for the primary BOSS sample is nearly 99%, with typical redshift uncertainty of a few tens of km s^{-1} [41]. The DR11 sample has a footprint of 8377 deg^2 , compared to $10,500 \text{ deg}^2$ expected for the final BOSS galaxy sample to appear in DR12.

BOSS targets two distinct samples of luminous galaxies selected by different flux and color cuts [24]: CMASS, designed to approximate a constant threshold in galaxy stellar mass in the range $0.43 < z < 0.7$, and LOWZ, which provides roughly three times the density of the SDSS-II LRG sample in the range $0.15 < z < 0.43$. Analysis of both samples incorporates reconstruction [16,17] to sharpen the BAO peak by partly reversing nonlinear effects, thus improving measurement precision. For CMASS, we use results of the anisotropic analysis by [26], which yields $D_M/r_d = 14.945 \pm 0.210$ (1.4% precision) and $D_H/r_d = 20.75 \pm 0.73$ (3.5% precision) with anticorrelated errors ($r = -0.52$).

The LOWZ sample does not have sufficient statistical power for a robust anisotropic analysis, so we use the measurement of $D_V = 8.467 \pm 0.167$ at $z = 0.320$ (discussed in detail by [42]). We have not included results from the SDSS-II LRG or WiggleZ surveys cited in the Introduction because these partly overlap the BOSS volume and are not statistically independent. We do include results of a new analysis [23] that uses reconstruction to achieve a 3.8% D_V/r_d measurement from the SDSS main galaxy sample (MGS) at effective redshift $z = 0.150$, which should be nearly independent of the BOSS LOWZ measurement. We also include the 6dFGS measurement of $D_V/r_d = 3.047 \pm 0.137$ (4.5% precision) at $z = 0.106$. Because the 6dFGS BAO detection is of moderate statistical significance and we do not have a full χ^2 surface for it, we truncate its χ^2 contributions at $\Delta\chi^2 = 4$ to guard against non-Gaussian tails of the error distribution. In practice the 6dFGS measurement carries little statistical weight in our constraints. These galaxy BAO measurements are listed in the first four lines of Table II.

2. BOSS Ly α forest BAO measurements

The BAO scale was first measured at higher redshift ($z \sim 2.4$) from the autocorrelation of the Ly α forest fluctuations in the spectra of high-redshift quasars from BOSS DR9 ([43–45]) following the pioneering work of measuring 3D fluctuations in the forest [46]. Here we use the results from [27], who present an improved measurement using roughly twice as many quasar spectra from BOSS DR11. The DR11 quasar catalog was made publicly available simultaneously with the DR12 catalog in 2015. The catalog construction is similar to that of the DR10 quasar catalog described by [47]. The BOSS quasar selection criteria are described by [48] and the background methodology papers [49–52].

The measurement of Ly α BAO peak positions is marginalized over parameters describing broad-band distortion of the correlation function using the methodology of [45]. Because of the low effective bias factor of the Ly α , redshift-space distortion strongly enhances the BAO peak in the line-of-sight direction. The measurement of D_H/r_d is therefore more precise (3.1%) than that of D_M/r_d (5.8%), as seen in line 5 of Table II. The errors of these two measurements are anticorrelated, with $r_{\text{off}} = -0.43$, and the optimally measured combination $D_H^{0.7} D_M^{0.3}$ is determined with a precision of $\sim 2\%$. While the overall signal-to-noise ratio of the BAO measurement is high, the detection significance for transverse separations ($\mu < 0.5$) is only moderate, as one can see in Fig. 3 of [27].

At the same redshift, BAO have also been measured in the cross correlation of the Ly α forest with the density of quasars in BOSS DR11 [28]. While the number of quasar-pixel pairs is much lower than the number of pixel-pixel pairs in the autocorrelation function, the clustering signal is much stronger because of the high bias factor of quasars. For the same reason, redshift-space distortion is much weaker in the cross correlation, and in this case the measurement precision is comparable for D_M/r_d (3.7%) and for D_H/r_d (3.3%) (line 6 of Table II). The higher precision of the transverse measurement makes the cross-correlation measurement an especially valuable complement to the autocorrelation measurement.

Even though these results are derived from the same volume, we can consider them as independent because their uncertainties are not dominated by cosmic variance. They are dominated instead by the combination of noise in the spectra and sparse sampling of the structure in the survey volume, both of which affect the autocorrelation and cross correlation almost independently. A number of tests using mock catalogs and several analysis procedures are presented in [27], finding good agreement between error estimates from the likelihood function and from the variance in mock catalogs. This independence allows us to add the χ^2 surfaces from both publications, which are publicly available [53]. While we use these two χ^2 surfaces separately, the last line of Table II lists the D_M/r_d and D_H/r_d constraints from the combined measurement, with respective precision of 3.2% and 2.2% and a correlation coefficient $r_{\text{off}} = -0.48$.

We caution that BAO measurement from Ly α data is a relatively new field, pioneered entirely by BOSS, in contrast to the now mature subject of galaxy BAO measurement, which has been studied both observationally and theoretically by many groups. The authors of [27] present numerous tests using mock catalogs and different analysis procedures, finding good agreement between error estimates from the likelihood surface and from mock catalog variance and identifying no systematic effects that are comparable to the statistical errors. However, the analysis uses only 100 mock catalogs, limiting the external

tests of the tails of the error distribution. The systematics and error estimation of the cross-correlation measurement have also been less thoroughly examined than those of the autocorrelation measurement, though continuing investigations within the BOSS Collaboration find good agreement with the errors and covariances reported in the publications above. On the theoretical side, [54] and [55] have examined the potential impact of UV background fluctuations on LyaF BAO measurement, finding effects that are much smaller than the current statistical errors.

We anticipate significant improvements in the LyaF analyses of the Data Release 12 sample, thanks to the larger data set and ongoing work on broadband distortion modeling, larger mock catalog samples, and spectrophotometric calibration. For the current paper, we adopt the BAO likelihood surfaces as reported in [28] and [27].

C. Cosmic microwave background data

In this paper we focus on constraints on the expansion history of the homogeneous cosmological model. For this purpose, we compress the cosmic microwave background (CMB) measurements to the variables governing this expansion history. This approach greatly simplifies the required computations, allowing us to fit complex models that have a simple solution to the Friedmann equation without the need to numerically solve for the evolution of perturbations. It is also physically illuminating, making clear what relevant quantities the CMB determines and distinguishing expansion history constraints from those that depend on the evolution of clustering. For some models or special cases we use more complete CMB results obtained by running the industry-standard COSMOMC [56] or by relying on the publicly available Planck MCMC chains [57]. Note that we use the 2013 Planck results not the 2015 results, which were not available at the time of our analysis.

The CMB plays two distinct but important roles in our analysis. First, we treat the CMB as a “BAO experiment” at redshift $z_\star = 1090$, measuring the angular scale of the sound horizon at very high redshift. Here we ignore the small dependence of the last-scattering redshift z_\star on cosmological parameters and the fact that the relevant scale for the CMB is r_\star rather than the drag redshift r_d that sets the BAO scale in low-redshift structure. We have checked that both approximations are valid to around 0.1σ for the case of BAO and Planck data and the Λ CDM model. In its second important role, the CMB calibrates the absolute length of the BAO ruler through its determination of ω_b and ω_{cb} .

Inspired by the existence of well-known degeneracies in CMB data [58–60], we compress the CMB measurements into three variables: ω_b , ω_{cb} and $D_M(1090)/r_d$. The mean vector and the 3×3 covariance matrix are used to describe the CMB constraints by a simple Gaussian likelihood. In order to calibrate these variables, we rely on the publicly

available Planck chains. In particular, we use the BASE_ALENS chains with the PLANCK_LOWL_LOWLIKE data set corresponding to the Planck data set with low- ℓ WMAP polarization (referred to in this paper as Planck + WP). We find that the data vector

$$\mathbf{v} = \begin{pmatrix} \omega_b \\ \omega_{cb} \\ D_M(1090)/r_d \end{pmatrix} \quad (18)$$

can be described by a Gaussian likelihood with mean

$$\mu_{\mathbf{v}} = \begin{pmatrix} 0.02245 \\ 0.1386 \\ 94.33 \end{pmatrix} \quad (19)$$

and covariance

$$C_{\mathbf{v}} = \begin{pmatrix} 1.286 \times 10^{-7} & -6.033 \times 10^{-7} & 1.443 \times 10^{-5} \\ -6.033 \times 10^{-7} & 7.542 \times 10^{-6} & -3.605 \times 10^{-5} \\ 1.443 \times 10^{-5} & -3.605 \times 10^{-5} & 0.004264 \end{pmatrix}. \quad (20)$$

The fractional diagonal errors on ω_b , ω_{cb} , and $D_M(1090)/r_d$ are 1.5%, 1.9%, and 0.06%, respectively. We similarly compress the WMAP 9-year data into

$$\mu_{\mathbf{v}} = \begin{pmatrix} 0.02259 \\ 0.1354 \\ 94.51 \end{pmatrix} \quad (21)$$

and covariance

$$C_{\mathbf{v}} = \begin{pmatrix} 2.864 \times 10^{-7} & -4.809 \times 10^{-7} & -1.111 \times 10^{-5} \\ -4.809 \times 10^{-7} & 1.908 \times 10^{-5} & -7.495 \times 10^{-6} \\ -1.111 \times 10^{-5} & -7.495 \times 10^{-6} & 0.02542 \end{pmatrix}. \quad (22)$$

For reference in interpreting the cosmological constraints from CMB + BAO data, especially in Sec. VI below, note that the contributions to $D_M(1090)$ accumulate over a wide range of redshift, with 14%, 25%, 38%, 47%, 69%, 88%, and 99% of the integral in Eq. (6) coming from redshifts $z < 0.5, 1.0, 2.0, 3.0, 10, 50,$ and 640 , respectively.

The BASE_ALENS model corresponds to the basic flat Λ CDM cosmology with explicit marginalization over the foreground lensing potential. Our decision to use the flat model was intentional, since we found that in curved models there is significant non-Gaussian correlation of ω_b and ω_m with curvature. Because our BAO data inevitably collapse more complex models to nearly flat ones, use of the flat data is more appropriate. We have tested the data compression in a couple of simple cases by comparing results of BAO + CMB data to COSMOMC chains and found

less than 0.5σ differences in best-fit parameter values between using compressed and full chains. The residual differences are driven by the fact that our compressed likelihood attempts to extract purely geometric information from the CMB data (for example, values of ω_b and ω_m are different at roughly the same level between chains that marginalize over lensing potential and those that do not). For BAO-only data combinations the results are completely consistent.

Throughout the paper, we refer to the constraints represented by Eqs. (18)–(20) simply as “Planck” (although they also include information from WMAP polarization measurements). In Sec. III we treat the CMB as a BAO experiment measuring $D_M(1090)/r_d$, but we eliminate its calibration of the absolute BAO scale by artificially blowing up the errors on ω_b and ω_{cb} ; we denote this case as “+Planck D_M ”. Conversely, in Sec. IV we use the CMB information on ω_b and ω_{cb} to set the size of our standard ruler r_d but omit the $D_M(1090)/r_d$ information by artificially inflating its errors; we denote this case as “+ r_d ”. When we use a full Planck chain instead of the compressed information, we adopt the notation “Planck (full)” and specify what additional parameters (such as A_{lens} , N_{eff} , or tensor-to-scalar ratio r) are being varied in the chain.

If one assumes a flat universe, a cosmological constant ($w = -1$), standard relativistic background ($N_{\text{eff}} = 3.046$), and minimal neutrino mass ($\sum m_\nu = 0.06$ eV), then the CMB data summarized by Eqs. (18)–(20) also provide a precise constraint on the Hubble parameter h , and thus on Ω_{cb} , Ω_b , and Ω_Λ . At various points in the paper we refer to a “fiducial” Planck Λ CDM model for which we adopt $\Omega_b h^2 = 0.022032$, $\Omega_m = 0.3183$, and $h = 0.6704$, which are the best fit parameters for “Planck + WP” combination as cited in the Table 2 of [29]. The CMB constraints on h and Ω_m become much weaker if one allows $w \neq -1$ or $\Omega_k \neq 0$, so for more general models BAO data or other constraints are needed to restore high precision on cosmological parameters.

D. Supernova data

A comprehensive set of relative luminosity distances of 740 SNIa was presented in [30], based on a joint calibration and training set of the SDSS-II Supernova Survey [33] and the Supernova Legacy Survey (SNLS) 3-year data set [31]. The 374 supernovae from SDSS-II and 239 from SNLS were combined with 118 nearby supernovae from [6,61–65] and nine high-redshift supernovae discovered and studied by HST [66]. We use this set, dubbed Joint Light-curve Analysis (JLA), rather than the Union 2.1 compilation of [67] because of the demonstrated improvement in calibration and corresponding reduction in systematic uncertainties presented in [30].

While [30] also provide a full COSMOMC module and a covariance matrix in relevant parameters, we here instead use their compressed representation of relative distance

constraints due to conceptual simplicity and a drastic increase in computational speed when combining with other cosmology probes. The compressed information consists of a piece-wise linear function fit over 30 bins (leading to 31 nodes) spaced evenly in $\log z$ (to minimum $z \sim 0.01$) with a 31×31 covariance matrix that *includes all of the systematics* from the original analysis. SNIa constrain *relative* distances, so the remaining marginalization required to use this compressed representation in a cosmological analysis is over the fiducial absolute magnitude of a SNIa, M_B . In Sec. IV we also utilize a similar compression of the Union 2.1 SN data set, which we have constructed in analogous fashion.

E. Visualizing the BAO constraints

Figure 1 shows the “Hubble diagram” (distance vs redshift) from a variety of recent BAO measurements of D_V/r_d , D_M/r_d , or zD_H/r_d ; these three quantities converge at low redshift. In addition to the data listed in Table II, we show measurements from the DR7 data set of SDSS-II by [15] and from the WiggleZ survey by [19], which are not included in our cosmological analysis because they are not independent of the (more precise) BOSS measurements in similar redshift ranges. Curves represent the predictions of the fiducial Planck Λ CDM model, whose parameters are determined independently of the BAO measurements but depend on the assumptions of a flat universe and a




FIG. 1 (color online). The BAO “Hubble diagram” from a world collection of detections. Blue, red, and green points show BAO measurements of D_V/r_d , D_M/r_d , and zD_H/r_d , respectively, from the sources indicated in the legend. These can be compared to the correspondingly colored lines, which represent predictions of the fiducial Planck Λ CDM model (with $\Omega_m = 0.3183$, $h = 0.6704$; see Sec. II C). The scaling by \sqrt{z} is arbitrary, chosen to compress the dynamic range sufficiently to make error bars visible on the plot. Filled points represent BOSS data, which yield the most precise BAO measurements at $z < 0.7$ and the only measurements at $z > 2$. For visual clarity, the Ly α cross-correlation points have been shifted slightly in redshift; auto-correlation points are plotted at the correct effective redshift.

cosmological constant. Overall, there is impressively good agreement between the CMB-constrained Λ CDM model and the BAO measurements, especially as no parameters have been adjusted in light of the BAO data. However, there is noticeable tension between the Planck Λ CDM model and the LyaF BAO measurements.

Figure 2 displays a subset of these BAO measurements with scalings that elucidate their physical content. In the upper panel, we plot $H(z)/(1+z)$, which is the proper




FIG. 2 (color online). BAO measurements and model predictions of $H(z)$ and $D_M(z)$ as a function of redshift, with physically informative scalings. The top panel shows $H(z)/(1+z)$, the proper velocity between two objects 1 comoving Mpc apart. The bottom panel shows $c \ln(1+z)/D_M(z)$, a scaling that matches a constant line $H(z) = (1+z)H_0$ in the top panel to the same constant line in the bottom panel for a flat universe. Filled circles and squares show the BOSS CMSSZ and LyaF measurements of $H(z)$ and $D_M(z)$, respectively; we show the LyaF-quasar cross correlation as crosses to distinguish from the LyaF autocorrelation measurements. Filled triangles in the bottom panel show the BOSS LOWZ and MGS measurements of $D_V(z)$ converted to $D_M(z)$. Open squares show the value of $H_0 = 67.3 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$ determined from the combination of BAO and SNIa data described in Sec. IV. The grey swath in both panels is the prediction from the Planck Λ CDM cosmology including 1σ parameter errors; in the top panel, one can easily see the model transition from deceleration to acceleration at $z \approx 0.6$. The dashed line shows the Λ CDM prediction using the best-fit WMAP parameters, which has lower $\Omega_m h^2$. Dotted curves show models that match the best-fit Planck values of ω_{cb} , ω_b , and $D_M(1090)/r_d$ but have $\Omega_k = 0.01$ (blue), $w = -0.7$ (green), or $w = -1.3$ (red). The x axis is set to $\sqrt{1+z}$ both for display purposes and so that a pure matter universe ($\Omega_m = 1$) appears as a decreasing straight line on the top panel.

velocity between two objects with a constant comoving separation of 1 Mpc. This quantity is declining in a decelerating universe and increasing in an accelerating universe. We set the x axis to be $\sqrt{1+z}$, which makes $H(z)/(1+z)$ a straight line of slope H_0 in an Einstein-de Sitter ($\Omega_m = 1$) model. For the transverse BAO measurements in the lower panel, we plot $c \ln(1+z)/D_M(z)$, chosen so that a constant (horizontal) line in the $H(z)/(1+z)$ plot would produce the same constant line in this panel, assuming a flat universe. This quantity would decrease monotonically in a nonaccelerating flat cosmology. The quantities in both the upper and lower panels approach H_0 as z approaches zero, independent of other cosmological parameters. We convert the BOSS LOWZ and MGS measurements of $D_V(z)$ to $D_M(z)$ in the lower panel assuming the fiducial Planck Λ CDM parameters; this is a robust approximation because all acceptable cosmologies produce similar scaling at these low redshifts. Note that the $H(z)$ and $D_M(z)$ measurements from a given data set (i.e., at a particular redshift) are covariant, in the sense that the points on these panels are anticorrelated (see Table II). For example, if $H(z)$ at $z = 2.34$ were scattered upward by a statistical fluctuation, then the $z = 2.34$ point in the lower panel would be scattered downward.

As discussed below in Sec. IV, the galaxy BAO and JLA supernova data can be combined to yield an “inverse distance ladder” measurement of H_0 , which utilizes the CMB measurements of ω_{cb} and ω_b but no other CMB information. This value of H_0 is robust to a wide range of assumptions about dark energy evolution and space curvature, although it does assume a standard radiation background for the calculation of r_d . We plot the resulting determination of $H_0 = 67.3 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$ as the open square in both panels.

The grey swath in both panels of Fig. 2 represents the 1σ region for the fiducial Planck Λ CDM model, with the top panel clearly showing the transition from deceleration to acceleration at $z \approx 0.6$. Formally, we are scaling both panels by $(r_d/r_{d,\text{fid}})$, so that the comparison of the BAO data points to the CMB prediction is invariant to changes in the sound horizon. The galaxy BAO measurements of $D_M(z)$ from BOSS and MGS are in excellent agreement with the predictions of this model (as are the other measurements shown previously in Fig. 1), and the combination of BAO and SNe yields an H_0 value in excellent agreement with this model’s prediction. The expansion rate $H(0.57)$ from CMSSZ is high compared to the model prediction, at moderate significance. Compared to Planck, the best-fit value of $\Omega_m h^2$ from the 9-year WMAP analysis [68] is lower, 0.143 vs 0.137, implying lower Ω_m and slightly higher h for a Λ CDM model. The model using these best-fit parameters, shown by the dashed lines, agrees better with the CMSSZ $H(z)$ measurement but is in tension with the distance data, especially the CMSSZ value of $D_M(0.57)$.

The Ly α forest measurements are much more difficult to reconcile with the Λ CDM model: compared to the Planck curve, the Ly α F BAO $H(z)$ is low and $[D_M(z)]^{-1}$ is high. It is important to keep the error anticorrelation in mind when assessing significance—if $H(z)$ fluctuates up then $1/D_M(z)$ will fluctuate down, which tends to reduce the tension relative to the CMB. However, our subsequent analyses (and those already reported by [27]) will show that the discrepancy is significant at the 2–2.5 σ level. The dotted curves show predictions of cosmological models with $\Omega_k = 0.01$ or $1 + w = \pm 0.3$. While changing curvature or the dark energy equation of state can improve agreement with some of the data points, it worsens agreement with other data points, and on the whole (as demonstrated quantitatively in Sec. V) such variations do not noticeably improve the fit to the combined CMB, BAO, and SN data.

Not plotted in Fig. 2 is the value of $D_M(1090)$ that comes from the angular acoustic scale in the CMB. Connecting the acoustic scale measured in CMB anisotropy to that measured in large-scale structure does require model assumptions about structure formation at the recombination epoch. However, it would be difficult to move the relative calibration significantly without making substantial changes to the CMB damping tail, which is already well constrained by observations. Using the ratio of $D_M(1090)/r_d$ in Eq. (19) and $r_d = 147.49$ Mpc, we find $c \ln(1+z)/D_M(z) = 151 \text{ km s}^{-1} \text{ Mpc}^{-1}$ at $z = 1090$ with percent level accuracy, a factor of 2 larger than any of the low-redshift values in Fig. 2. On their own, the BAO data in Fig. 2 clearly favor a universe that transitions from deceleration at $z > 1$ to acceleration at low redshifts, and this evidence becomes overwhelming if one imagines the corresponding CMB measurements off the far left of the plot. We quantify these points in the following section.

It is tempting to consider a flat cosmology with a constant $H/(1+z)$ as an alternative model of these data [69]. Note that although this form of $H(z)$ occurs in coasting (empty) cosmologies in general relativity, those models have open curvature and hence a sharply different $D_M(z)$. But even for the flat model, the data are not consistent with a constant $H(z)/(1+z)$, first because the increase in $c \ln(1+z)/D_M(z)$ from $z = 0.57$ to $z = 0.0$ is statistically significant, and second because of the factor of 2 change of this quantity relative to that inferred from the CMB angular acoustic scale. The change from $z = 0.57$ to $z = 0$ is more significant than the plot indicates because the data points are correlated; this occurs because the H_0 value results from normalizing the SNe distances with the BAO measurements. We measure the ratio of the values, $H_0 D_M(0.57)/c \ln(1.57)$, to be 1.080 ± 0.014 from the combination of BAO and SNe data sets, a 5.5 σ rejection of a constant hypothesis and an indication of the strength of the SNe data in detecting the low-redshift accelerating expansion.

III. BAO AS AN UNCALIBRATED RULER

A. Convincing detection of dark energy from BAO data alone

For quantitative constraints, we start by considering BAO data alone with the simple assumption that the BAO scale is a standard comoving ruler, whose length is independent of redshift and orientation but is *not* necessarily the value computed using CMB parameter constraints. A similar analysis has been presented in [70]. In this case, a simple dimensional analysis shows that in addition to fractional densities in cosmic components, one can constrain the dimensionless quantity $P = c/(H_0 r_d)$.

Figure 3 presents constraints on relevant quantities in Λ CDM models, which assume that dark energy is a cosmological constant but allow $\Omega_\Lambda = 0$ and arbitrary Ω_k . The combination of galaxy and Ly α F BAO measurements yields a marginalized constraint of $\Omega_\Lambda = 0.73_{-0.68}^{+0.25}$ at 99.7% confidence, implying a $> 3\sigma$ detection of dark energy from BAO alone *without* CMB data.

These constraints become much tighter if we assume that the CMB is measuring the *same* acoustic scale, functioning as an additional BAO experiment at a much higher redshift. As discussed in Sec. II C, we implement this case by retaining the high-precision CMB measurement of $D_M(1090)/r_d$ but drastically inflating (by a factor of 100) the CMB errors on ω_{cb} and ω_b , so that the value of r_d itself remains effectively unknown. Combining the CMB measurement with galaxy or Ly α F BAO alone yields a strong detection of nonzero Ω_Λ , but with different central values reflecting the tensions already discussed in Sec. II E and examined further in Secs. V–VI. Combining all three measurements yields a marginalized $\Omega_\Lambda = 0.72_{-0.034}^{+0.030}$ (at 68% confidence, with reasonably Gaussian errors), implying $> 20\sigma$ preference for a low-density universe dominated by dark energy. The dimensionless quantity $P = c/(H_0 r_d) = 29.63_{-0.45}^{+0.48}$ is determined with 1.6% precision. Most importantly, this data combination also requires a nearly flat universe, with a total density $\Omega_m + \Omega_\Lambda = 1.011_{-0.016}^{+0.014}$ determined to 1.5% and consistent with the critical density. Thus, with the minimal assumption that the BAO scale is a standard ruler, these data provide strong support for the standard cosmological model.

B. External calibration of r_d

We proceed further by computing the sound horizon scale r_d from the standard physics of the pre-recombination universe but adopting empirical constraints external to the CMB. In particular, we adopt a prior on the baryon density of $\omega_b = 0.02202 \pm 0.00046$ determined from big bang nucleosynthesis (BBN) and the observed primordial deuterium abundance [71], and we assume a standard relativistic background ($N_{\text{eff}} = 3.046$, $\omega_\nu \approx 0$). For any values of Ω_m and the Hubble parameter h that arise in our MCMC




FIG. 3 (color online). Constraints from BAO on the parameters of Λ CDM models, treating the BAO scale as a redshift-independent standard ruler of unknown length. Green curves/contours in each panel show the combined constraints from galaxy and Ly α F BAO, with no CMB information. Black curves/contours include the measurement of $D_M(1090)/r_d$ from the CMB acoustic scale, again with no assumption about the value of r_d except that it is the same scale as the lower redshift measurements. This combination of BAO measurements yields precise constraints on Ω_Λ (top panel) and the dimensionless quantity $c/(H_0 r_d)$ (bottom panel), and it requires a low density ($\Omega_m \approx 0.29$), nearly flat universe (middle panel). Blue and red curves in the top and bottom panels show the result of combining the CMB BAO measurement with either the galaxy or Ly α F BAO measurement separately. The dotted line in the middle panel marks $\Omega_m + \Omega_\Lambda = 1$.




FIG. 4 (color online). Constraints on Ω_m and h in a flat Λ CDM model from galaxy BAO (red), Ly α F BAO (blue), and the combination of the two (green), using a BBN prior on ω_b and standard physics to compute the sound horizon r_d but incorporating no CMB information. Contours are plotted at 68%, 95%, and 99.7% confidence (the interior white region of the green “donut” is 68%). Black contours show the entirely independent constraints on Ω_m and h in Λ CDM from full Planck CMB chains.

chain, we can then compute the value of r_d from Eq. (16). Compared to the previous section, the addition of the physical scale allows us to convert the measured value of $c/(H_0 r_d)$ into a measurement of the dimensional parameter H_0 . In practice, we derive constraints in a separate MCMC run where, instead of a flat prior on P , we have a flat prior on h and the above prior on ω_b . We also fix the curvature parameter Ω_k to zero. Results are presented in Fig. 4. The red (galaxy BAO) and blue (Ly α F BAO) contours in this figure use no CMB information at all, but they *do* assume a spatially flat universe in contrast to Fig. 3.

The point of this exercise is the following. The homogeneous part of the minimal Λ CDM model has just two adjustable parameters, Ω_m and h , which matches the 2 degrees of freedom offered by a measurement of anisotropic BAO at a single redshift. (The weak BBN prior is required to fix the magnitude of r_d , but it does not affect the expansion history.) One can therefore get meaningful constraints from either galaxy BAO or Ly α F BAO alone, though this is no longer true if one allows nonzero curvature and therefore introduces a third parameter. There is substantial $\Omega_m - h$ degeneracy for either measurement individually, but both are generally compatible with standard values of these parameters. The tension of the Ly α F BAO with the Planck Λ CDM model manifests itself here as a best fit at relatively low matter density and high Hubble parameter. Combining the galaxy and Ly α F measurements produces a precise measurement of both Ω_m and the Hubble parameter *coming from BAO alone*, independent of CMB data. In combination, we find $h = 0.67 \pm 0.013$ and $\Omega_m = 0.29 \pm 0.02$ (68% confidence). The small black ellipse in Fig. 4 shows the Planck constraints for Λ CDM, computed from full Planck chains, which are in

excellent agreement with the region allowed by the joint BAO measurements.

IV. BAO, SNIA, AND THE INVERSE DISTANCE LADDER

The traditional route to measuring the Hubble constant H_0 is built on a distance ladder anchored in the nearby Universe: stellar distances to galaxies within ~ 20 Mpc are used to calibrate secondary indicators, and these in turn are used to measure distances to galaxies “in the Hubble flow,” i.e., far enough away that peculiar velocities are a subdominant source of uncertainty when inferring $H_0 = v/d$ [72]. The most powerful implementations of this program in recent years have used Cepheid variables—calibrated by direct parallax, by distance estimates to the LMC, or by the maser distance to NGC 4258—to determine distances to host galaxies of SNIa, which are the most precise of the available secondary distance indicators [73–75].

Because the BAO scale can be computed in absolute units from basic underlying physics, the combination of BAO with SNIa allows a measurement of H_0 via an “inverse distance ladder,” anchored at intermediate redshift. The BOSS BAO data provide absolute values of D_V at $z = 0.32$ and D_M at $z = 0.57$ with precision of 2.0% and 1.4%, respectively. The JLA SNIa sample provides a high-precision relative distance scale, which transfers the BAO measurement down to low redshift, where H_0 is simply the slope of the distance-redshift relation. Equivalently, this procedure calibrates the absolute magnitude scale of SNIa using BAO distances instead of the Cepheid distance scale. Although the extrapolation from the BAO redshifts to low redshifts depends on the dark energy model, the SNIa relative distance scale is precisely measured over a well sampled redshift interval which includes the BAO redshifts, so this extrapolation introduces practically no uncertainty even when the dark energy model is extremely flexible. CMB data enter the inverse distance ladder by constraining the values of ω_m and ω_b and thus allowing computation of the sound horizon scale r_d .

Figure 5 provides a conceptual illustration of this approach, zeroing in on the $z < 1$ portion of the Hubble diagram. Filled points show $c \ln(1+z)/D_M(z)$ from the CMASS, LOWZ, MGS, and 6dFGS BAO measurements, where for illustrative purposes only we have converted the latter three measurements from $D_V(z)$ to $D_M(z)$ using Planck Λ CDM parameters. The error bars on these points include the 0.4% uncertainty in r_d arising from the uncertainties in the Planck determination of ω_m and ω_b , but this is a small contribution to the error budget. Crosses show the binned SNIa distance measurements, with the best absolute magnitude calibration from the joint BAO + SNIa fit. We caution that systematic effects introduce error correlations across redshift bins in the SNIa data, which are accounted for in our full analysis. To allow flexibility in the dark energy model, we adopt the PolyCDM




FIG. 5 (color online). Determination of H_0 by the “inverse distance ladder” combining BAO absolute distance measurements and SNIa relative distance measurements, with CMB data used to calibrate the sound horizon scale r_d . The quantity $c \ln(1+z)/D_M(z)$ converges to H_0 at $z = 0$. Filled circles show the four BAO measurements, normalized with $r_d = 147.49$ Mpc; for the three lower redshift points, D_V has been converted to D_M assuming Λ CDM. Crosses show the SNIa measurements, with error bars representing diagonal elements of the covariance matrix. Because the absolute luminosity of SNIa is not known *a priori*, the SNIa points are free to shift vertically by a constant factor, which is chosen here to produce the best joint fit with the BAO data. The red square and error bar shows the value $H_0 = (67.3 \pm 1.1) \text{ km s}^{-1} \text{ Mpc}^{-1}$ determined by the full inverse distance ladder procedure described in the text. The black curve shows the prediction for a Λ CDM model with $\Omega_m = 0.3$ and the best-fit H_0 , and green curves show ten PolyCDM models randomly selected from our MCMC chain that have $\Delta\chi^2 < 4$ relative to the best-fit PolyCDM model. This H_0 determination assumes standard prerecombination physics to evaluate r_d . For nonstandard energy backgrounds (e.g., extra relativistic species or early dark energy) the more general result is described by Eq. (23).

parametrization described in Sec. II A, imposing a loose Gaussian prior $\Omega_k = 0 \pm 0.1$ to suppress high curvature models that are clearly inconsistent with the CMB. Thin green curves in Fig. 5 show $c \ln(1+z)/D_M(z)$ for ten PolyCDM models that have $\Delta\chi^2 < 4$ relative to the best-fit model, selected from the MCMC chains described below. The intercept of these curves at $z = 0$ is the value of H_0 . While low-redshift BAO measurements like those of 6dFGS and MGS incur minimal uncertainty from the extrapolation to $z = 0$, the statistical error is necessarily large because of the limited volume at low z . It is evident from Fig. 5 that using SNIa to transfer intermediate redshift BAO measurements to the local Universe yields a much more precise determination of H_0 than using only low-redshift BAO measurements, even allowing for great flexibility in the dark energy model.

To compute our H_0 constraints, we adopt the $D_M(z)$ and $H(z)$ constraints from CMASS BAO (including covariance), the $D_V(z)$ constraints from LOWZ, MGS, and 6dFGS BAO, the compressed JLA SNIa data set with its full 31×31 covariance matrix, and an r_d constraint from Planck (see Sec. II C). Marginalizing over the PolyCDM parameters yields $H_0 = 67.3 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$, a 1.7% measurement. Even if we include the CMB angular diameter distance at its full precision, our central value and error bar on H_0 change negligibly because the flexibility of the PolyCDM model effectively decouples low- and high-redshift information.

As a byproduct of our H_0 measurement, we determine the absolute luminosity of a fiducial SNIa to be $M_B = -19.14 \pm 0.042 \text{ mag}$. Here we define a fiducial SNIa as having SALT2 (as retrained in [30]) light-curve width and color parameters $x_1 = 0$ and $C = 0$ and having exploded in a galaxy with a stellar mass $< 10^{10} M_\odot$.

Our best-fit H_0 and its 1σ uncertainty are shown by the open square and error bar in Figs. 2 and 5. To characterize the sources of error, we have repeated our analyses after multiplying either the CMB, SN, or BAO covariance matrix by a factor of 10 (and thus reducing errors by $\sqrt{10}$). Reducing the CMB errors, so that they yield an essentially perfect determination of r_d , makes almost no difference to our H_0 error, because the 0.4% uncertainty in r_d is already small. Reducing either the SNIa or BAO errors shrinks the H_0 error by approximately a factor of 2, indicating that the BAO measurement uncertainties and the SNIa measurement uncertainties make comparable contributions to our error budget; the errors add (roughly) linearly rather than in quadrature because both measurements constrain the redshift evolution in our joint fit. If we replace PolyCDM with ow_0w_a CDM in our analysis, substituting a different but still highly flexible dark energy model, the derived value of H_0 drops by less than 0.2σ and the error bar is essentially unchanged. If we instead fix the dark energy model to Λ CDM, the central value and error bar are again nearly unchanged, because with the dense sampling provided by SNe the extrapolation from the BAO redshifts down to $z = 0$ is also only a small source of uncertainty. To test sensitivity to the SN data set, we constructed a compressed description of the Union 2.1 compilation [67] analogous to that of the JLA compilation; substituting Union 2.1 for JLA makes negligible difference to our best-fit H_0 while increasing the error bar by about 30% (see Table III). Finally, if we substitute the WMAP9 constraints on ω_m and ω_b for the Planck constraints, the central H_0 decreases by 0.5% (to $66.9 \text{ km s}^{-1} \text{ Mpc}^{-1}$) and the error bar grows by 8% (to $1.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

To summarize, this 1.7% determination of H_0 is robust to details of our analysis, with the error dominated by the BAO and SNIa measurement uncertainties. The key assumptions behind this method are (a) standard matter and radiation content, with three species of light neutrinos,

TABLE III. Constraints on H_0 (in $\text{km s}^{-1} \text{ Mpc}^{-1}$) from the inverse distance ladder, assuming $r_d = 147.49 \pm 0.59 \text{ Mpc}$ as inferred from Planck with a standard radiation background. The bottom two lines substitute the Union 2.1 SN data set for the JLA data set. Error bars are 1σ .

Combination	Model	H_0
Galaxy BAO + SN + r_d	PolyCDM	67.3 ± 1.1
Galaxy BAO + SN + r_d	ow_0w_a CDM	67.1 ± 1.1
Galaxy BAO + Union SN + r_d	PolyCDM	67.3 ± 1.5
Galaxy BAO + Union SN + r_d	ow_0w_a CDM	67.2 ± 1.5

and (b) no unrecognized systematics at the level of our statistical errors in the CMB determinations of ω_m and ω_b , in the BAO measurements, or in the SNIa measurements used to tie them to $z = 0$. Note that the SNIa covariance matrix already incorporates the detailed systematic error budget of [30]. The measurement systematics are arguably smaller than those that affect the traditional distance ladder. Thus, with the caveat that it assumes a standard matter and radiation content, this measurement of H_0 is more precise and probably more robust than current distance-ladder measurements.

Nonstandard radiation backgrounds remain a topic of intense cosmological investigation, and a convincing mismatch between H_0 determinations from the forward and inverse distance ladders could be a distinctive signature of nonstandard physics that alters r_d . We can express our constraint in a more model-independent form as

$$H_0 = (67.3 \pm 1.1) \times (147.49 \text{ Mpc}/r_d) \text{ km s}^{-1} \text{ Mpc}^{-1}. \quad (23)$$

Raising N_{eff} from 3.046 to 4.0 would increase our central value of H_0 to $69.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [Eq. (17), but see further discussion in Sec. VID].

Figure 6 compares our H_0 determination to several other values from the literature. The lower two points show our results using either the PolyCDM model or the ow_0w_a CDM model. The top three points show recent distance-ladder determinations from Riess *et al.* [73], Freedman *et al.* [74], and a reanalysis of the Riess *et al.* data set by [76]. There is mild ($\approx 2\sigma$) tension between these determinations and our value. The central two points show the values of H_0 inferred from Planck or WMAP CMB data *assuming a flat Λ CDM model*, with values and uncertainties taken from the MCMC chains provided by the Planck Collaboration. These inferences of H_0 are much more model dependent than our inverse distance ladder measurement; with the ow_0w_a CDM or PolyCDM dark energy models the errors on H_0 from CMB data alone increase by more than order of magnitude because of the CMB geometric degeneracy. Consistency of these H_0 values is therefore a consistency test for the Λ CDM model, which it passes here with flying colors.




FIG. 6 (color online). Constraints on the Hubble constant H_0 from this paper’s inverse distance ladder analysis (blue, at bottom), from three direct distance ladder estimates (red, at top), and from Planck or WMAP CMB data assuming Λ CDM (green, middle). All error bars are 1σ . The inverse distance ladder estimates assume $r_d = 147.49 \pm 0.59$ Mpc, based on Planck constraints for a standard radiation background, while the green points make the much stronger assumptions of a flat universe with a cosmological constant.

Our results can be compared to those of several other recent analyses. The authors of [77] determine H_0 from a collection of BAO data sets using the Planck-calibrated value of r_d . They do not incorporate SNIa, but they assume a flat Λ CDM model, which allows them to obtain a tight constraint $H_0 = 68.11 \pm 0.86 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The authors of [78] carry out a more directly comparable inverse distance ladder measurement with essentially the same data sets but cosmological models that are 1-parameter extensions of Λ CDM, finding $H_0 = 68.0 \pm 1.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$ for either $o\Lambda$ CDM or w CDM. The authors of [79] carry out a rather different analysis that uses age measurements for early-type galaxies to provide an absolute time scale. In combination with BAO and SNIa, they then constrain the acoustic oscillation scale $r_d = 101.9 \pm 1.9 h^{-1} \text{ Mpc}$ independent of CMB data or early universe physics. Their result, which assumes an $o\Lambda$ CDM cosmology, can be cast in a form similar to ours, $H_0 = (69.9 \pm 1.3) \times (147.49 \text{ Mpc}/r_d) \text{ km s}^{-1} \text{ Mpc}^{-1}$; the agreement implies that their stellar evolution age scale is consistent with the scale implied by early-universe BAO physics. As an H_0 determination, our analysis makes much more general assumptions about dark energy than these other analyses, but it yields a consistent result. It is also notable that our value of H_0 agrees with the value of $67 \pm 2 \text{ km s}^{-1} \text{ Mpc}^{-1}$ inferred from a median-statistics analysis of *direct* distance ladder estimates *circa* 2001 ([80], see [81] for a 2011 update).

From Fig. 5, it is visually evident that the relative distance scales implied by our BAO and SN are in fairly good agreement. We have converted SN luminosity distances to comoving angular diameter distances with

$D_M(z) = D_L(z)/(1+z)$, a relation that holds in any metric theory of gravity (see Sec. IV.2 of [82] and references therein). As a quantitative consistency test, we refit the PolyCDM model with an additional free parameter that artificially modifies the luminosity distance by $D_L(z) \rightarrow D_L(1+z)^\beta$, finding $\beta = 0.13 \pm 0.063$. This result is consistent with the expected $\beta = 0$ at 2σ , but there is a mild tension because the SN data are in good agreement with Λ CDM predictions while the ratio of $D_M(z)$ between the CMASS and LOWZ samples is somewhat higher than expected in Λ CDM.

V. CONSTRAINTS ON DARK ENERGY MODELS

We now turn to constraints on dark energy and space curvature from the combination of BAO, CMB, and SNIa data. In this section, we consider models with standard matter and radiation content, including three neutrino species with the minimal allowed mass $\sum m_\nu = 0.06 \text{ eV}$ (although the cosmological differences between 0.06 and 0 eV are negligible relative to current measurement errors). In Sec. VI, we will consider models that allow dynamically significant neutrino mass, extra relativistic species, dark matter that decays into radiation, or “early” dark energy that is dynamically non-negligible even at high redshifts.

To set the scene, Fig. 7 compares the predictions of models constrained by CMB data to the BOSS BAO constraints on D_M and D_H at $z = 0.57$ and $z = 2.34$, from CMASS galaxies and the Ly α F, respectively. Black dots mark best-fit values of (D_M, D_H) , and contours are shown at $\Delta\chi^2 = 2.30, 6.18, \text{ and } 11.83$ (coverage fractions of 68%, 95%, and 99.7% for a 2-d Gaussian). The top row shows results for $o\Lambda$ CDM models, which assume a constant dark energy density but allow nonzero space curvature. Here we have taken models from the Planck Collaboration MCMC chains, based on the combination of Planck, WMAP polarization, and ACT/SPT data. The upper right panel shows the one-dimensional PDF for the curvature parameter Ω_k based on the CMB data alone. Each point in the left and middle panels represents a model from the chains, color-coded by the value of Ω_k on the scale in the right panel. The green cross-hairs mark the predicted (D_M, D_H) from the flat Λ CDM model that best fits the CMB data alone. This model lies just outside the 68% contour for CMASS, but it is discrepant at $> 95\%$ with the Ly α F measurements, as remarked already by [27]. When the flatness assumption is dropped, both the galaxy and Ly α F BAO data strongly prefer Ω_k close to zero, firmly ruling out the slightly closed ($\Omega_k \sim -0.05$) models that are allowed by the CMB alone.

The bottom row shows results for w CDM models, which assume a flat universe but allow a constant equation-of-state parameter $w \equiv p/\rho \neq -1$ for dark energy. The CMB data alone are consistent with a wide range of w values, and they are generally better fit with $w < -1$. However, the combination with CMASS BAO data sharply




FIG. 7 (color online). BAO constraints in the $D_M - D_H$ planes at $z = 0.57$ (left) and $z = 2.34$ (middle) compared to predictions of Λ CDM (top row) or w CDM (bottom row) models constrained by CMB data. Black curves show 68%, 95%, and 99.7% likelihood contours from the CMASS and LyaF BAO measurements, relative to the best-fit values (black dots). Colored points represent individual models from Planck + WP + ACT/SPT MCMC chains, which are color-coded by the value of Ω_k (top row) or w (bottom row) as illustrated in the right panels. Green cross-hairs mark the predictions of the flat Λ CDM model that best fits the CMB data. White curves show 68% and 95% likelihood contours for the CMB data alone.

limits the acceptable range of w , favoring values close to -1.0 (a cosmological constant). The fit to the LyaF BAO results could be significantly improved by going to $w \leq -1.3$, but this change would be inconsistent with the CMASS measurements. This example illustrates a general theme of our results: parameter changes that improve agreement with the LyaF BAO measurements usually run afoul of the galaxy BAO measurements.

More quantitative constraints appear in Fig. 8 and Table IV. We begin with Λ CDM and continue to the progressively more flexible models described in Table I. For CMB data, we now use the compression of Planck or WMAP9 constraints described in Sec. II C.

We include all of the BAO data listed in Table II. Omitting the LyaF BAO data makes almost no difference to the central values or error bars on model parameters, though it has a significant impact on goodness-of-fit as we discuss later.

For Λ CDM and Λ CDM, the combination of Planck CMB constraints and BAO is remarkably powerful, a point already emphasized by [29]. Adding SN data makes

negligible difference to the parameter constraints of these models; SN + Planck constraints have nearly identical central values to BAO + Planck, but larger errors. In Λ CDM, substituting BAO + SN + WMAP9 for BAO + SN + Planck has a tiny effect, shifting $\Omega_m = 0.302 \pm 0.008$ to $\Omega_m = 0.300 \pm 0.008$ with a small compensating shift in h . Figure 8 illustrates the extremely tight curvature constraint that comes from combining CMB and BAO data: for Λ CDM we find $\Omega_k = -0.003 \pm 0.003$ using Planck CMB or $\Omega_k = -0.004 \pm 0.004$ using WMAP9.

Supernovae play a much more important role in models that allow $w \neq -1$, as their high precision relative distance measurements provide strong constraints on low-redshift acceleration. For both w CDM and Λ CDM, SN + Planck and BAO + Planck constraints are perfectly consistent but complementary, and the combination of all three data sets provides much tighter error bars than any pairwise combination. For w CDM we find $w = -0.97 \pm 0.05$. For Λ CDM the curvature constraint from BAO is particularly important, lifting the degeneracy between w and Ω_k that




FIG. 8 (color online). Constraints on interesting parameter combinations in a variety of dark energy models: Λ CDM (upper left), $o\Lambda$ CDM (upper right), w CDM (middle left), ow CDM (middle right), w_0w_a CDM (bottom left), and ow_0w_a CDM (bottom right). Curves show 68%, 95%, and 99.7% confidence contours for the data combinations indicated in the legend. In the top panels the red contours are almost fully obscured by the green contours because the BAO + Planck combination is already as constraining as the BAO + SN + Planck combination, but for models with freedom in dark energy the SN and BAO constraints are complementary. The bottom panels, with evolving $w(z)$, display the value of w at $z = 0.266$, the “pivot” redshift where w is best constrained by BAO + SN + Planck in the w_0w_a CDM model. For our BAO + SN + Planck contours, the white zone interior to the dark green annulus marks the 68% confidence region, and the outer edge of the dark annulus is 95%.

arises for SN + CMB alone; we find $w = -0.98 \pm 0.06$ and $\Omega_k = -0.002 \pm 0.003$. Substituting WMAP9 for Planck again produces only slight shifts to central values and a minor increase of error bars.

Even with these powerful BAO, SN, and CMB data sets, constraining the evolution of w is difficult. The constraint on the evolution parameter from BAO + SN + Planck is $w_a = -0.2 \pm 0.4$ in w_0w_a CDM and weakens to $w_a = -0.6 \pm 0.6$ in ow_0w_a CDM. Both results are consistent

with constant w , but they allow order unity changes of w at $z < 1$. This data combination still provides a good constraint on the value of w at a “pivot” redshift $z_p = 0.266$ where it is uncorrelated with w_a (determined specifically for w_0w_a CDM for BAO + SN + Planck combinations): $w(0.266) = -0.97 \pm 0.05$ in w_0w_a CDM and -0.99 ± 0.06 in ow_0w_a CDM.

We note that the degradation of our ability to constrain the evolution of the equation of state is not accompanied

TABLE IV. Cosmological parameter constraints from Galaxy + Ly α BAO data combined with our compressed description of CMB constraints from Planck + WP or WMAP9 and the JLA SN data. Bold entries show our preferred constraints, from the BAO, SN, and Planck + WP data combination. Entries for which the parameter is fixed in the listed cosmological model are marked with a dash. For w_0w_a CDM and ow_0w_a CDM, column 7 lists the value of w at $z = 0.266$, which is the ‘‘pivot’’ redshift for w_0w_a CDM with the full data combination. For SlowRDE, this column lists $w = \delta w_0 - 1$.

Model	Data	Ω_m	$\Omega_b h^2$	h	Ω_k	w	w_a
Λ CDM	BAO + Planck	0.303 (8)	0.0223 (3)	0.682 (7)
Λ CDM	SN + Planck	0.295 (16)	0.0224 (3)	0.688 (13)
Λ CDM	BAO + SN + Planck	0.302 (8)	0.0223 (3)	0.682 (6)
Λ CDM	BAO + SN + WMAP	0.300 (8)	0.0224 (5)	0.681 (7)
$o\Lambda$ CDM	BAO + Planck	0.301 (8)	0.0225 (3)	0.679 (7)	-0.003 (3)
$o\Lambda$ CDM	SN + Planck	0.30 (4)	0.0224 (4)	0.68 (4)	-0.002 (10)
$o\Lambda$ CDM	BAO + SN + Planck	0.301 (8)	0.0225 (3)	0.679 (7)	-0.003 (3)
$o\Lambda$ CDM	BAO + SN + WMAP	0.295 (9)	0.0226 (5)	0.677 (8)	-0.004 (4)
w CDM	BAO + Planck	0.311 (13)	0.0225 (3)	0.669 (17)	...	-0.94 (8)	...
w CDM	SN + Planck	0.298 (18)	0.0225 (4)	0.685 (17)	...	-0.99 (6)	...
w CDM	BAO + SN + Planck	0.305 (10)	0.0224 (3)	0.676 (11)	...	-0.97 (5)	...
w CDM	BAO + SN + WMAP	0.303 (10)	0.0225 (5)	0.674 (12)	...	-0.96 (6)	...
ow CDM	BAO + Planck	0.308 (17)	0.0225 (4)	0.671 (19)	-0.001 (4)	-0.95 (11)	...
ow CDM	SN + Planck	0.28 (8)	0.0225 (4)	0.73 (11)	0.01 (3)	-0.97 (18)	...
ow CDM	BAO + SN + Planck	0.303 (10)	0.0225 (4)	0.676 (11)	-0.002 (3)	-0.98 (6)	...
ow CDM	BAO + SN + WMAP	0.299 (11)	0.0227 (5)	0.671 (12)	-0.004 (4)	-0.96(6)	...
w_0w_a CDM	BAO + Planck	0.34 (3)	0.0224 (3)	0.639 (25)	...	-0.77 (13)	-1.0 (6)
w_0w_a CDM	SN + Planck	0.292 (23)	0.0224 (4)	0.693 (24)	...	-1.00 (6)	-0.5 (8)
w_0w_a CDM	BAO + SN + Planck	0.307 (11)	0.0223 (3)	0.676 (11)	...	-0.97 (5)	-0.2 (4)
w_0w_a CDM	BAO + SN + WMAP	0.305 (11)	0.0224 (5)	0.674 (12)	...	-0.97 (6)	-0.2 (5)
ow_0w_a CDM	BAO + Planck	0.34 (3)	0.0225 (4)	0.640 (25)	-0.003 (4)	-0.81 (14)	-1.1 (6)
ow_0w_a CDM	SN + Planck	0.29 (8)	0.0225 (4)	0.72 (11)	0.01 (3)	-1.03 (21)	-0.3 (9)
ow_0w_a CDM	BAO + SN + Planck	0.307 (11)	0.0225 (4)	0.673 (11)	-0.005 (4)	-0.99 (6)	-0.6 (6)
ow_0w_a CDM	BAO + SN + WMAP	0.302 (11)	0.0227 (5)	0.670 (12)	-0.006 (5)	-0.99 (6)	-0.5 (5)
SlowRDE	BAO + SN + Planck	0.307 (10)	0.0224 (3)	0.676 (11)	...	-0.95 (7)	...

with significant degradation in our ability to measure the curvature of space: the constraint on curvature remains tight even when allowing an evolving equation of state, $\Omega_k = -0.005 \pm 0.004$.

By decoupling the time dependence of w from its present-day value, the $w_0 - w_a$ model allows flexible evolution of the dark energy density. Adopting a particular form for the dark energy potential reduces this freedom, and one can construct physically motivated models that have evolving dark energy but do not require an additional free parameter to describe it. The authors of [83] advocate an interesting example of this model class, in which dark energy is a slowly rolling scalar field with a $\frac{1}{2}m^2\phi^2$ potential, analogous to the inflaton of chaotic inflation models. The authors of [83] show that this model yields $\delta w(z) \equiv 1 + w(z) \approx \delta w_0 \times H_0^2/H^2(z)$ and therefore [84]

$$\frac{H^2(z)}{H_0^2} \approx \Omega_m(1+z)^3 + \Omega_{de} \left[\frac{(1+z)^3}{\Omega_m(1+z)^3 + \Omega_{de}} \right]^{\delta w_0/\Omega_{de}}, \quad (24)$$

where the approximation is first-order in $\delta w_0 = 1 + w_0$.

Figure 9 presents parameter constraints for the slow-roll dark energy scenario in a flat universe, a model that has the same number of parameters as w CDM. BAO and SN both contribute to the constraints of the joint fit, which yields $\delta w_0 = 0.05 \pm 0.07$, $h = 0.675 \pm 0.011$, and $\Omega_m = 0.306 \pm 0.010$. Results for this scenario are thus consistent with Λ CDM but allow small departures from $w_0 = -1$.




FIG. 9 (color online). Constraints on h and $\delta w_0 \equiv 1 + w_0$ in the slow-roll dark energy model [Eq. (24)], in the same format as Fig. 8.

A striking feature of Table IV is that the best-fit parameter values barely shift as additional freedom is added to the models. For the BAO + SN + Planck combination, the best-fit Ω_m values range from 0.301 to 0.307 and the best-fit h values from 0.673 to 0.682, while combinations with WMAP9 favor just slightly lower values of Ω_m and h . More importantly, models that allow dark energy evolution are all consistent with constant $w = -1.0$ at $\approx 1\sigma$, and Ω_k is consistent with zero at 1σ in all cases that allow curvature. The fact that models with additional freedom remain consistent with Λ CDM is a substantial argument in favor of this minimal model.

Figure 10 illustrates the goodness-of-fit for the models in Table IV, and for additional goodness models discussed below in Sec. VI. For the best-fit parameter values in each model, horizontal bars show the total χ^2 , with colors indicating the separate contributions from the JLA SN data, the various galaxy BAO data sets, and the LyaF autocorrelation and cross-correlation measurements. For visualization purposes, we have subtracted 30 from the SN χ^2 , which would otherwise dominate the total length of these bars because there are 31 SN data points and many fewer in other data sets. The constraints on ω_{cb} , ω_b , and $D_M(1090)/r_d$ from the CMB are sufficiently tight that parameter variations within the allowed range have minimal impact on other observables. Our minimization yields $\chi^2 \approx 0$ for the CMB data in essentially every case, since all the models have enough parameters to fit the three (compressed) CMB constraints perfectly. For this plot, we have




FIG. 10 (color online). χ^2 values for the best-fit versions of cosmological models considered in the paper. Each bar represents the minimum χ^2 for the model listed at the left axis, and colors show the χ^2 contributions of individual data sets. For better visualization, we subtract 30 from the SN χ^2 . CMB contributions are not included but (with our 3-parameter compression) are always close to zero. The total χ^2 and model degrees of freedom (d.o.f., 40 data points minus number of fit parameters, which includes the SNIa absolute magnitude normalization as well as cosmological quantities) are listed to the right of each bar. The bottom bar shows the number of d.o.f. associated with each data set. For the ΔN_{eff} model we use COSMOMC rather than our compressed CMB description, but we again omit CMB contributions to χ^2 .

chosen to omit the CMB constraints from both the χ^2 sum and the degrees-of-freedom (d.o.f.) computation, though these constraints are still used when determining model parameters.

The bottom bar in Fig. 10 indicates the number of d.o.f. associated with each data set: 31 for SNe, one each for the D_V measurements from LOWZ, MGS, and 6dFGS, two for the D_M and D_H measurements from CMASS, and two each (D_M and D_H) for LyaF auto and cross correlation, totaling 40. Numbers to the right of each model bar list the χ^2 of the model fit and the corresponding d.o.f. after subtracting the number of fit parameters. For Λ CDM, for example, we count as free parameters Ω_m , h , and the SNIa absolute magnitude normalization M_0 , yielding d.o.f. = 40 - 3 = 37. We omit ω_b because it is determined almost entirely by the CMB data, which we have excluded from the χ^2 sum. The total χ^2 for this model is 46.79, with a one-tailed p -value (probability of obtaining $\chi^2 \geq 46.79$) of 0.13 for 37 d.o.f. Thus, if we consider all of the data collectively, the fit of the Λ CDM model is acceptable, and for any of the more complex models considered so far the reduction in χ^2 is smaller than the number of additional free parameters in the model.

As already emphasized in our discussion, the Λ CDM model does not give a good fit to the LyaF BAO data. This tension is evident in Fig. 10 in the length of the yellow and green χ^2 bars relative to the corresponding d.o.f. Combining the Lyman- α auto- and cross-correlation measurements into a single likelihood because they measure the same quantities, the Λ CDM $\chi^2_{\text{Ly}\alpha} = 8.3$ for two d.o.f. has a p -value of 0.016, consistent with Fig. 7. It is unclear how much to make of this mild tension in the context of a fit that yields adequate-to-excellent agreement with multiple other data sets and an acceptable χ^2 overall. It is evident that none of the more complex models considered so far allows a significantly better fit to the LyaF BAO data. The partial exception is ow_0w_a CDM, which has the most freedom to adjust high-redshift behavior relative to low-redshift behavior, but even here the reduction in χ^2 relative to Λ CDM is only 1.33 (coming almost entirely from LyaF), for three additional model parameters. Omitting the LyaF data makes almost no difference to the best-fit parameter values or their error bars in any of these models, which are driven mainly by the high-precision CMB, CMASS, and SN constraints.

To conclude this section, we examine a model in which dark energy is characterized by specifying its energy density in four discrete bins of redshift: $z < 0.5$, $0.5 < z < 1.0$, $1.0 < z < 1.6$, and $z > 1.6$. This step-wise model is a useful complement to models that specify $\rho_{\text{de}}(z)$ through parametrized descriptions of $w(z)$. The bins are chosen to be the same as the ones considered in a similar analysis by [67] (their Table 8), who combined Union 2.1 SN data, WMAP7 CMB data [85], BAO from the combined analysis of SDSS DR7 and 2dFGRS [86], and the




FIG. 11 (color online). Constraints on $\rho_{DE}(z)$ assumed to be constant within redshift bins, in units of the present-day critical density ρ_c . Shaded areas represent 68% confidence levels. Yellow bands show constraints in the same bins from [67]. Our constraints in the $z = 1.0 - 1.6$ bin are omitted.

distance-ladder H_0 measurement of [73]. We fit simultaneously for $\rho_{de}(z)$ in each of these bins and for the values of Ω_m , Ω_b , and H_0 , assuming $\Omega_k = 0$ to match [67]. Within an individual bin, ρ_{de} is held constant and $H(z)$ evolves according to the Friedmann equation, but there are discontinuities in $H(z)$ at bin boundaries to accommodate the discontinuous changes in $\rho_{de}(z)$. The matter density evolves as $\rho_m(z) = \Omega_m \rho_{crit}(1+z)^3$, where Ω_m and ρ_{crit} denote $z = 0$ values as usual.

Constraints on this model from our BAO + Planck and BAO + SN + Planck data combinations appear in Fig. 11 and Table V. As in our other models that allow time-varying dark energy, BAO and SN data both contribute significantly to the parameter constraints. Our results show a clear detection of nonzero dark energy density in each of the first two redshift bins at $z < 1$, and they are consistent with a constant energy density across this redshift range. Compared to [67], we obtain a significantly tighter constraint in the $0.5 < z < 1.0$ bin, where the CMASS BAO measurement makes an important difference, but a slightly looser constraint in the $z < 0.5$ bin, where we do not incorporate a direct H_0 measurement. We obtain much poorer constraints in the $1 < z < 1.6$ bin because the JLA sample contains only 8 SNe with $z > 1$ compared to 29 for the Union 2.1 sample. At $z > 1.6$ our constraint is stronger thanks to the Ly α BAO measurement, but the uncertainty

is large nonetheless, and the low Ly α value of $H(z)$ leads to a preference for negative dark energy density in this bin, although consistent with zero at 1σ .

VI. ALTERNATIVE MODELS

We now turn to models with more unusual histories of the dark energy, matter, or radiation components. In part we want to know what constraints our combined data can place on interesting physical quantities, such as neutrino masses, extra relativistic species, dark energy that is dynamically significant at early times, or dark matter that decays into radiation over the history of the Universe. We also want to see whether any of these alternative models can resolve the tension with the Ly α measurements at $z = 2.34$, which persists in all of the models considered in Sec. V. We begin with the early dark energy model, because understanding the origin of the constraints on this model informs the discussion of subsequent models.

A. Early dark energy

In typical dark energy models, including all of those discussed in Sec. V, dark energy is dynamically negligible at high redshifts because its energy density grows with redshift much more slowly than $(1+z)^3$. However, some scalar field potentials yield a dark energy density that tracks the energy density of the dominant species during the radiation and matter dominated eras, then asymptotes towards a cosmological constant at late times [87,88]. These models ameliorate, to some degree, the ‘‘coincidence problem’’ of constant- w models because the ratio of dark energy density to total energy density varies over a much smaller range.

As a generic parametrized form of such early dark energy models, we adopt the formulation of Doran and Robbers [89], in which the density parameter of the dark energy component evolves with $a = (1+z)^{-1}$ as

$$\Omega_{de}(a) = \frac{\Omega_{de}^e - \Omega_{de}^e (1 - a^{-3w_0})}{\Omega_{de}^e + \Omega_m a^{3w_0}} + \Omega_{de}^e (1 - a^{-3w_0}), \quad (25)$$

where Ω_{de} and Ω_m denote $z = 0$ values as usual and Ω_{de}^e is the dark energy density parameter at early times. A flat universe is assumed, with $\Omega_{de} + \Omega_m = 1$. The quantity w_0 is the effective value of the equation-of-state parameter

TABLE V. Parameter constraints in the model in which $\rho_{de}(z)$ is held constant in four discrete bins of redshift. Uncertainties on each parameter are marginalized over all others, including $\Omega_b h^2$, which is not listed in the table. H_0 is in $\text{km s}^{-1} \text{Mpc}^{-1}$.

	Ω_m	H_0	ρ_{DE}/ρ_c			
			$z < 0.5$	$0.5 < z < 1.0$	$1.0 < z < 1.6$	$1.6 < z$
Suzuki <i>et al.</i> 2012 [67]	$0.731^{+0.015}_{-0.014}$	$0.880^{+0.240}_{-0.210}$	$0.330^{+1.900}_{-1.000}$	$0.700^{+2.400}_{-1.800}$
BAO + PLANCK	$0.317^{+0.009}_{-0.020}$	$66.4^{+1.5}_{-0.9}$	$0.667^{+0.026}_{-0.012}$	$0.844^{+0.194}_{-0.188}$	$8.581^{+5.237}_{-6.278}$	$-0.921^{+0.897}_{-0.611}$
BAO + SN + PLANCK	$0.307^{+0.012}_{-0.014}$	$67.3^{+1.5}_{-1.0}$	$0.685^{+0.022}_{-0.016}$	$0.765^{+0.146}_{-0.165}$	$5.154^{+4.761}_{-4.259}$	$-0.634^{+0.957}_{-0.601}$

today. At high redshift ($a \ll 1$), the denominator of the first term is $\gg 1$, and $\Omega_d(a)$ approaches the constant value Ω_{de}^e . This in turn requires a dark energy density that scales as a^{-3} in the matter-dominated era and as a^{-4} in the radiation-dominated era, though it is $\Omega_{\text{de}}(a)$ rather than $\rho_{\text{de}}(a)$ that is specified explicitly. For $w_0 = -1$, the model approaches Λ CDM as Ω_{de}^e goes to zero. There are other generic forms of models with early dark energy, as well as nonparametric descriptions (see discussion by [90]).

If dark energy is important in the pre-recombination era, then the boosted energy density in this era reduces the sound horizon by a factor $(1 - \Omega_{\text{de}}^e)^{1/2}$ relative to a conventional model with the same parameters [89,90]. Pre-recombination dark energy also influences the detailed shape of the CMB anisotropy spectrum by altering the early integrated Sachs-Wolfe contribution and the CMB damping tail [91]. Our analysis here incorporates the rescaling of the sound horizon, but we continue to use the compressed CMB description of Sec. II C and therefore ignore the more detailed changes to the power spectrum shape. Because of the exquisite precision of CMB measurements, the power spectrum shape may impose tighter constraints on early dark energy than the expansion history measurements employed here (see, e.g., [91]). However, those constraints are more dependent on the specifics of the models being examined, both the dark energy evolution and other parameters that describe the inflationary spectrum, tensor fluctuations, relativistic energy density, and reionization.

Figure 12 plots the evolution of $D_H(z)/r_d$ and $D_M(z)/r_d$ for models with $\Omega_{\text{de}}^e = 0, 0.02, 0.04$, and 0.08 . We always adopt $w_0 = -1$, but we constrain h , Ω_b , and $\Omega_m = 1 - \Omega_{\text{de}} - \Omega_{\nu+r}$ by fixing $\Omega_b h^2$, $\Omega_m h^2$, and $D_M(1090)/r_d$ to the values in the best-fit Planck + WP Λ CDM model, in effect forcing the errors in our compressed CMB description to zero. Solid curves incorporate the expected $(1 - \Omega_{\text{de}}^e)^{1/2}$ reduction of r_d . Dotted curves show the case in which we instead keep r_d fixed at its fiducial value of 147.49 Mpc. The latter case would be physically relevant in a model where dark energy is dynamically negligible in the pre-recombination era but approaches the evolution of Eq. (25) later in the matter dominated era. To highlight model differences, we scale all values to those of the fiducial Λ CDM model, which corresponds to $\Omega_{\text{de}}^e = 0$.

Remarkably, for the rescaled r_d case, the predicted values of $D_H(z)/r_d$ and $D_M(z)/r_d$ change by less than 0.5% at all redshifts, even for $\Omega_{\text{de}}^e = 0.08$. We can understand this insensitivity by considering the low and high-redshift limits for the simplified case of a flat cosmology with only matter and dark energy. The matter density at redshift z is

$$\begin{aligned} \rho_m(z) &= \rho_{\text{crit}} \times \Omega_m (1+z)^3 \\ &= \frac{3(100 \text{ km s}^{-1} \text{ Mpc}^{-1})^2}{8\pi G} \times (\Omega_m h^2)(1+z)^3, \end{aligned} \quad (26)$$




FIG. 12 (color online). Predicted BAO scales for early dark energy models with CMB observables $\Omega_b h^2$, $\Omega_m h^2$, and $D_M(1090)/r_d$ held fixed to the values of the best-fit Planck + WP Λ CDM model. We adopt Eq. (25) with $w_0 = -1$. Solid lines show the case in which r_d is rescaled by $(1 - \Omega_{\text{de}}^e)^{1/2}$ to represent the effect of early dark energy in the pre-recombination era, while dashed lines show the case in which r_d is held fixed at the fiducial model value of $r_d = 147.49$ Mpc. We show ratios of $D_H(z)/r_d$ (top) or $D_M(z)/r_d$ (bottom) relative to the fiducial ($\Omega_{\text{de}}^e = 0$) model. Points with error bars show the BAO measurements from CMASS galaxies at $z = 0.57$ (filled square) and from Ly α F autocorrelation (open circle) and cross correlation (filled circle) at $z = 2.34$. For visual clarity, the Ly α F cross-correlation points have been slightly shifted in redshift.

where ρ_{crit} denotes the $z = 0$ value as usual and we have used $H_0 = 100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ but relocated h to the second factor. Using $\rho_m(z) + \rho_{\text{de}}(z) = \rho_{\text{crit}}(z) = 3H^2(z)/8\pi G$ implies

$$\begin{aligned} H(z) &= (100 \text{ km s}^{-1} \text{ Mpc}^{-1}) \times [(\Omega_m h^2)(1+z)^3]^{1/2} \\ &\quad \times [1 + \rho_{\text{de}}(z)/\rho_m(z)]^{1/2}. \end{aligned} \quad (27)$$

For a cosmological constant, $\rho_{\text{de}}(z)/\rho_m(z) \propto (1+z)^{-3}$, so the ratio tends rapidly to zero at high redshift, but for the

early dark energy model this ratio asymptotes instead to $\Omega_{\text{de}}^e/\Omega_m(z) \approx \Omega_{\text{de}}^e/(1 - \Omega_{\text{de}}^e)$. Thus, at fixed $\Omega_m h^2$, $H(z)$ is higher in the early dark energy model by a factor $(1 - \Omega_{\text{de}}^e)^{-1/2}$, and $D_H(z)$ is smaller by the same factor. This reduction in $D_H(z)$ exactly compensates the $(1 - \Omega_{\text{de}}^e)^{1/2}$ rescaling of r_d , leaving $D_H(z)/r_d$ independent of Ω_{de}^e .

At low redshift, conversely,

$$D_M(z) = \frac{c}{H_0} \int_0^z \frac{H_0}{H(z')} dz' \quad (28)$$

depends mainly on H_0 , since the evolution of $H_0/H(z)$ is insensitive to moderate changes in Ω_m and Ω_{de}^e for $z \ll 1$. Therefore, to keep the value of $D_M(1090)/r_d$ fixed to the CMB constraint, one must increase H_0 by approximately $(1 - \Omega_{\text{de}}^e)^{-1/2}$ so that both the low and high-redshift contributions to $D_M(1090)$ shrink by the factor required to compensate the change in r_d . This change again forces $D_H(z)/r_d$ to nearly the same value as the fiducial model with $\Omega_{\text{de}}^e = 0$.

These scaling arguments are not perfect because they break down at intermediate redshifts and because a change in H_0 at fixed $\Omega_m h^2$ implies a change in Ω_m , which itself affects the low-redshift evolution of $H_0/H(z)$. Nonetheless, the full calculation in Fig. 12 demonstrates that for Ω_{de}^e as large as 0.08 there is minimal change in $D_H(z)$ at any redshift, and minimal change in $H(z)$ in turn implies minimal change in $D_M(z)$. However, the values of H_0 are larger, and Ω_m correspondingly smaller, for the successively higher Ω_{de}^e curves in Fig. 12. The combination $H_0 r_d$ is nearly constant, decreasing by just 0.14%, 0.24%, and 0.49% for $\Omega_{\text{de}}^e = 0.02, 0.04,$ and 0.08 , respectively.

Reversing these arguments explains why $D_H(z)$ and $D_M(z)$ change rapidly with Ω_{de}^e if r_d stays fixed instead of rescaling (dashed curves in Fig. 12). In this case, $D_M(1090)$ must stay fixed to keep the angular scale of the acoustic peaks unchanged, so the decrease of high-redshift contributions to $D_M(z)$ by $(1 - \Omega_{\text{de}}^e)^{1/2}$ requires a compensating increase of $D_M(z)$ at low redshift. This requires a large fractional reduction in H_0 , since most of the contribution to $D_M(1090)$ comes from high redshift (e.g., 75% from $z > 1$). This in turn leads to large deviations in $D_H(z)/r_d$ and $D_M(z)/r_d$ at low redshift. At high redshift, $D_H(z)$ is again smaller by $(1 - \Omega_{\text{de}}^e)^{1/2}$, but this now leads to a deviation in $D_H(z)/r_d$ because it is no longer compensated by a smaller r_d . Even for $\Omega_{\text{de}}^e = 0.02$, fixing CMB observables requires a 4.3% reduction in H_0 . Furthermore, adding early dark energy in this case moves model predictions further from the CMASS and Ly α F measurements of $D_H(z)/r_d$ and further from the Ly α F $D_M(z)/r_d$. We therefore expect tight constraints on Ω_{de}^e in the case of fixed r_d .

Figure 13 presents constraints on these early dark energy models from our MCMC analysis, with w_0 fixed to -1 .




FIG. 13 (color online). Constraints in the $\Omega_{\text{de}}^e - \Omega_m$ plane from the combination of our compressed CMB description with galaxy + Ly α F BAO data. Black contours (68%, 95%, and 99.7%) show the tight constraints on early dark energy for models with fixed $r_d = 147.49$ Mpc. Blue contours show the constraints for models with $r_d \propto (1 - \Omega_{\text{de}}^e)^{1/2}$ as expected if Ω_{de}^e is constant into the radiation-dominated era. The red solid line traces the parameter degeneracy $\Omega_m = \Omega_{m, \text{fid}}(1 - \Omega_{\text{de}}^e)$ predicted by the approximate scaling arguments described in the text.

We now account for uncertainties in the CMB constraints, using the compressed description of Sec. II C. Note that we assume that the CMB constraints on ω_m and ω_b are not altered by the introduction of early dark energy, which might not hold in a complete analysis that uses the full CMB spectrum. The nonrescaled case is tightly constrained as expected, with a 2σ upper limit $\Omega_{\text{de}}^e < 0.031$. SNe do not significantly improve these constraints, although they would play a larger role if we allowed w_0 as a free parameter. To summarize, adding early dark energy with fixed r_d worsens agreement with our BAO measurements, and a dynamically significant value of Ω_{de}^e is ruled out.

For rescaled r_d , which is the physically expected case if Ω_{de}^e remains constant back into the radiation-dominated era, we instead find a valley of near-perfect degeneracy between Ω_m and Ω_{de}^e . For $H_0 r_d = \text{const.}$ and fixed $\Omega_m h^2$, the expected degeneracy line is $\Omega_m \propto r_d^{-2} \propto (1 - \Omega_{\text{de}}^e)$, marked by the red solid line in Fig. 13. This prediction describes our numerical MCMC results extremely well. Along this line, there are models with $\Delta\chi^2 < 1$ relative to the best-fit Λ CDM ($\Omega_{\text{de}}^e = 0$) model, at least out to $\Omega_{\text{de}}^e = 0.32$. Including SNe again makes minimal difference to our constraints because the models along the degeneracy line predict nearly identical $D_M(z)$.

Although these models are degenerate with respect to our geometrical constraints, they predict different values of H_0 and different measures of structure growth. Intriguingly, nonzero Ω_{de}^e with rescaled r_d reduces tension with distance-ladder measurements of H_0 and with the level of matter clustering inferred from cluster masses, weak lensing, and redshift-space distortions. We discuss the impact on structure growth measures in Sec. VII.

As already emphasized, the detailed shape of the CMB power spectrum may impose much tighter constraints on early dark energy; e.g., for the specific case of the Doran-Robbers model, Ref. [91] infers $\Omega_{\text{de}}^e < 0.012$ at 95% confidence. However, the degeneracies identified here in the expansion history constraints are striking, and highlight the potential value of early dark energy studies that fully explore degeneracies with other parameters that affect the CMB power spectrum shape.

B. Decaying dark matter

If dark matter is a metastable particle that decays into undetected radiation on a time scale comparable to H_0^{-1} , where the undetected radiation could be neutrinos or some other low-mass particle that interacts weakly enough to avoid detection (note that decay of a significant fraction of the dark matter into photons would need to have a very small branching ratio to be consistent with upper limits on cosmic backgrounds), then the matter density will decrease faster at low redshift than simple $(1+z)^3$ dilution (for an early discussion, see [92]). While the radiation density is boosted by dark matter decay, it subsequently decreases as $(1+z)^4$, so the total energy density at low redshift is lower in a decaying dark matter (DDM) model than it would be for stable dark matter with the same high-redshift density. We initially considered this model as a potential explanation of the low $H(2.34)$ inferred from the LyaF BAO. The heights of the acoustic peaks constrain the value of $\Omega_m h^2$ at the recombination epoch, but the reduced matter density at low redshift implies a lower value of $H^2(z) = (8\pi G/3)\rho_{\text{crit}}(z)$. The sound horizon scale r_d is unchanged because the prerecombination densities are unchanged. However, the full impact of introducing DDM is complex, because the values of h and Ω_m must change to keep $D_M(1090)/r_d$ at its precisely measured value, and because these changes and the dark matter decay itself affect the galaxy BAO observables and the LyaF value of $D_M(2.34)$.

We assume exponential decay of the dark matter (i.e., a constant decay rate), so that the governing equations for the decaying matter density (marked with subscript x) and decay products' radiation density (marked with subscript g) are

$$\dot{\rho}_x = -3H\rho_x - \lambda H_0 \rho_x, \quad (29)$$

$$\dot{\rho}_g = -4H\rho_g + \lambda H_0 \rho_x. \quad (30)$$

The decay rate λ is dimensionless, and λ^{-1} is the decay time in units of H_0^{-1} . However, we allow for the possibility that there are two kinds of dark matter, only one of which is susceptible to decay, so we introduce an additional parameter $f_x = \lim_{z \rightarrow \infty} \Omega_x(z)/\Omega_{\text{dm}}(z)$ that is the ratio of this decaying component to the total dark matter density in the infinite past.⁴ The other components of the model remain

⁴The earlier arXiv version of this paper had f defined as a fraction of total matter density today.

the same as those used for Λ CDM. Initial conditions are chosen so that there is no energy density in the decay product radiation in the infinite past. We discuss some details of our solution technique in the Appendix.

An important subtlety in this analysis is that that the CMB peaks constrain the dark matter density at the time of recombination, and hence the ω_b and ω_c densities that we feed into the compressed CMB likelihood correspond to the densities the system would have had if the decay did not take place. Of course, the distance to the last scattering surface is still affected by the changes in the expansion history due to decaying dark matter.

Figure 14 shows two-dimensional posterior probability distributions in the $\lambda f_x - \Omega_m$ plane for several values of f_x . Although the data prefer no decaying dark matter, we see strong degeneracies that extend to surprisingly large values of λ . For $f_x = 1$, decay of nearly 50% of the primordial dark matter is allowed at 95% confidence. As expected, Ω_m is negatively correlated with λf_x : the CMB constrains ω_c in the early universe, and if more dark matter decays then Ω_m today is lower. There is also a weak correlation with the Hubble parameter (not shown), with h rising by ~ 0.01 for $\lambda f_x \sim 0.5$.

To gain some understanding of this degeneracy, one can calculate the effective w of the fluid composed of the combined decaying dark matter and the resulting radiation. At $f_x = 1$, $\Omega_m = 0.23$ and $\lambda = 0.4$ (the edge of our 68% contour), the effective w takes values 0.07 at $z = 0$, falling to 0.03 at $z = 2$ and 10^{-4} at $z = 100$. With no decay, this component (which starts at the same energy density, as fixed by the CMB) would evolve with $w = 0$. The surprisingly small corrections to the total w make it difficult




FIG. 14 (color online). Constraints on decaying dark matter from CMB, SN and galaxy + LyaF BAO data. For various choices of the fraction f_x of dark matter in the decaying component, we plot posterior probability distributions for the product λf_x , where λ is the decay constant, assuming a flat prior on λ vs value of Ω_m today (which, by definition, includes both the nondecaying component and the undecayed fraction of the decaying component). Removing SN data does not significantly relax these constraints.

to constrain the decaying dark matter from expansion history data alone.

Using combinations of CMB, SN, and large-scale structure data sets, the authors of [93] obtained a limit $\Gamma^{-1} > 100$ Gyr for the dark matter decay constant, using methodology similar to that described here but also including constraints from the full shape of the CMB power spectrum and the amplitude of matter clustering. A more recent analysis by [94], using the Planck + WP CMB power spectrum and BAO measurements from BOSS and WiggleZ, shows a somewhat stronger limit of $\Gamma^{-1} > 160$ Gyr. For a Hubble parameter $h = 0.68$, our limit for $f_x = 1$ corresponds to $\Gamma^{-1} > 28$ Gyr at 95% confidence level. A more detailed analysis by [95] calculates the velocity distributions of daughter particles for varying assumptions about the decay products. For a daughter relativistic fraction of 1% and higher, they find $\Gamma^{-1} > 10$ Gyr based on analysis of Union 2.1 SNIa data in the context of a CMB determined cosmological model. From these results we conclude that, somewhat surprisingly, the expansion history alone is not sufficient to significantly constrain the decay of dark matter into an unknown relativistic component.

We note that the authors of [96] state a limit $\Gamma^{-1} > 700$ Gyr based on only the CMB acoustic scale and SNIa data available in 2008. We do not understand how these more limited data could lead to a stronger bound on the decay time, which suggests that the analysis of [96] contains a hidden assumption.

C. Massive neutrinos

In addition to constraining dark energy and space curvature, measuring neutrino masses is a key objective of precision cosmology. Given CMB constraints that $\omega_m \approx 0.14$, the fractional contribution of neutrinos to the low-redshift matter density is $\omega_\nu/\omega_m \approx 0.07(\sum m_\nu/1 \text{ eV})$, so neutrino masses have a noticeable cosmological impact if they are a significant fraction of an eV. Atmospheric and laboratory measurements constrain the mass splittings among the three standard-model neutrino species to be $m_2^2 - m_1^2 = 7.54_{-0.22}^{+0.26} \times 10^{-5} \text{ eV}^2$ and $m_3^2 - (m_1^2 + m_2^2)/2 = \pm 2.43_{-0.10}^{+0.06} \times 10^{-3} \text{ eV}^2$ [97]. This sets the minimum neutrino mass in a normal hierarchy, where $m_1 < m_2 \ll m_3$, to $\sum m_\nu = 5.84_{-0.8}^{+1.2} \text{ meV}$, which motivates our assumption that $\sum m_\nu = 0.06 \text{ eV}$ in the standard Λ CDM model. In the case of an inverted hierarchy, where $m_1 \approx m_2 \gg m_3$, the sum of the neutrino masses must exceed 0.1 eV, and for the degenerate neutrino mass case where $m_1 \approx m_2 \approx m_3$, the minimum mass sum is approximately 0.15 eV. These masses are well within reach of the cosmological experiments in the coming decade.

Neutrinos affect the CMB and large-scale structure differently from cold dark matter because they are still relativistic at the epoch of matter-radiation equality,

because their linear clustering is suppressed on scales below $k_{\text{sup}} = 2\pi/\lambda_{\text{sup}} = 0.018\sqrt{m_\nu/1 \text{ eV}} h^{-1} \text{ Mpc}$ [98], and because their high thermal velocities prevent them from clustering in small potential wells even in the non-linear regime [99,100]. The relative suppression in the linear matter power spectrum is linear in the fraction density in neutrinos f_ν and is about $\Delta P/P \sim -8f_\nu \sim 0.063(\sum m_\nu/0.1 \text{ eV})$ (at best fit values of Ω_m and h).

Measurements of matter clustering can constrain $\sum m_\nu$ by detecting the suppression of small scale power. Expansion history measurements, which we focus on here, can constrain $\sum m_\nu$ because of their transition from a relativistic species whose energy density scales as $(1+z)^4$ to a nonrelativistic species whose energy density scales as $(1+z)^3$, effectively the converse of decaying dark matter. Specifically, the CMB acoustic peaks constrain ω_{cb} almost independently of ω_ν , but the matter density that affects late-time expansion rates and distances is $\omega_{cb} + \omega_\nu$. With other parameters held fixed, a 0.2 eV neutrino mass sum increases the late-time matter density by 1.4%, which is significant given the extremely precise CMB measurement of $D_M(1090)/r_d$ and precise distance scale measurements from BAO. In practice, changing $\sum m_\nu$ leads to adjustments in other parameters to seek a global best fit. Although the neutrino mass influences the sound horizon [Eq. (16)], this impact is small for the range of $\sum m_\nu$ allowed by our constraints (-0.26% for $\sum m_\nu = 0.5 \text{ eV}$).

The top row of Fig. 15 compares the CMASS and Ly α F BAO constraints to the predictions of CMB-constrained, flat Λ CDM models with $\sum m_\nu$ as a free parameter, in the same format as Fig. 7. The Planck CMB chain used here is `BASE_MNU_PLANCK_LOWL_LOWLIKE_HIGHL_ALENS`, where we have selected a chain that marginalizes over the lensing amplitude parameter A_L for reasons discussed below. The base Planck Λ CDM model adopts $\sum m_\nu = 0.06 \text{ eV}$, but this chain allows $\sum m_\nu$ down to zero. While $\sum m_\nu > 0.5 \text{ eV}$ is allowed by the CMB data alone, this mass significantly worsens agreement with the BAO data, both at $z = 0.57$ and at $z = 2.34$. Higher m_ν increases $\Omega_m h^2$ and thus decreases $D_H(2.34)$, moving further from the Ly α F measurement. Additionally, because of the tight CMB constraint on $D_M(1090)/r_d$, the reduction in $c/H(z)$ at high redshift must be compensated by changes in h and Ω_m that raise D_M at low redshift, so $D_M(2.34)$, $D_M(0.57)$, and $D_H(0.57)$ all increase, again moving away from the BAO measurements. Conversely, moving towards $\sum m_\nu = 0$ slightly improves agreement with the Ly α F BAO because $D_H(2.34)$ increases while $D_M(2.34)$, which has a large contribution from lower redshifts, decreases. However, the same change worsens agreement with the CMASS BAO, which are already well fit by the base model with $\sum m_\nu = 0.06 \text{ eV}$.

The red curve in the upper panel of Fig. 16 shows the purely geometric constraint that arises from combining just the compressed CMB description with galaxy and Ly α F




FIG. 15 (color online). BAO constraints in the $D_M - D_H$ planes at $z = 0.57$ (left) and $z = 2.34$ (middle) compared to predictions of CMB-constrained, flat Λ CDM models in which the neutrino mass $\sum m_\nu$ or the number of relativistic species N_{eff} is a free parameter. Black curves show 68%, 95%, and 99.7% likelihood contours from the CMASS and Ly α F BAO measurements, relative to the best-fit values (black dots). Colored points represent individual models from Planck + WP + ACT/SPT MCMC chains, which are color-coded by the value of $\sum m_\nu$ (top row) or N_{eff} (bottom row) as illustrated in the right panels. Green cross-hairs mark the predictions of the flat Λ CDM model with $\sum m_\nu = 0.06$ eV and $N_{\text{eff}} = 3.046$ that best fits the CMB data. White curves show 68% and 95% likelihood contours for the CMB data alone. CMB results in the top row are marginalized over the lensing parameter A_L .

BAO data. This constraint is surprisingly tight at $\sum m_\nu < 0.56$ eV (at 95% c.l.), and it is independent of mass constraints based on the suppression of structure growth by neutrino free-streaming. Adding SN data does not significantly improve this constraint. The geometrical constraint on neutrino mass weakens if we allow either curvature or $w \neq -1$, as shown by the blue and green curves in the lower panel of Fig. 16. Because neutrinos influence the observables only via the effect of ω_m on distances and expansion rates, adding another degree of freedom introduces degeneracy. In these cases, including SN data does improve the neutrino mass constraint, as shown by the thin curves. The improvement is not dramatic, indicating that our multiple BAO measurements can break the degeneracy themselves to a significant degree.

The 95% upper limits on $\sum m_\nu$ for the various models and data combinations we have considered are listed in Table VI. Direct measurements of matter clustering at low redshift can be a powerful diagnostic of neutrino masses

because they are sensitive to the distinctive effect of suppressing small scale power, which is not easily mimicked by other parameter variations. We return to this point in Sec. VII.

In the top panel of Fig. 16, the green curve shows the result of full fitting using the COSMOMC machinery. The constraint tightens significantly to $\sum m_\nu < 0.25$ eV (at 95% C.L.). This number is the same as the Planck Collaboration constraint for Planck + BAO [29], even though we use more and better constraining BAO data. This exercise demonstrates that the compression into ω_{cb} , ω_b , and $D_M(1090)/r_d$ is missing important information that the CMB provides on neutrino mass. The extra information is in the constraint on the amplitude of low-redshift matter clustering that comes from the smoothing of the acoustic peaks in the CMB power spectrum by weak gravitational lensing. The BAO information in this case fixes the matter density of the Universe, thus allowing inference on the amplitude of matter fluctuations and hence




FIG. 16 (color online). Neutrino mass constraints for several combinations of data and model freedom. In the top panel, the red curve shows the posterior pdf (with a flat prior) on $\sum m_\nu$ from the expansion history constraint on the neutrino mass, based on the combination of BAO with our compressed CMB description, which yields $\sum m_\nu < 0.56$ eV at 95% confidence. The green curve shows the result obtained by replacing our compressed CMB description with the full Planck + WP power spectrum using COSMOMC, which strengthens the upper limit to $\sum m_\nu < 0.25$ eV. The blue curve adopts the same data combination but additionally marginalizes over the parameter A_L , demonstrating that the difference between the green and red curves is driven mainly by the lensing amplitude information in the Planck data. In the lower panel, curves show the posterior pdf of $\sum m_\nu$ from our usual BAO + Planck (thick) or BAO + SN + Planck (thin) geometrical constraints, assuming ΛCDM , $w\text{CDM}$, or $\sigma\Lambda\text{CDM}$ (red, blue, green, respectively).

neutrino mass to be determined from the smoothness of the peaks (the CMB only constraint is $\sum m_\nu < 0.93$ eV [29]). The role of CMB lensing is further demonstrated by the blue curve in the top panel. Here we have run a COSMOMC chain that, in addition to having $\sum m_\nu$ as a free parameter, marginalizes over a parameter A_L that multiplies the predicted lensing signal, effectively removing the lensing information. (The base model fixes $A_L = 1$.) The result here is very similar to that found by using the compressed CMB description.

TABLE VI. 95% confidence limits for the sum of the mass of neutrino species. The first two cases use full Planck CMB chains, while other cases adopt our compressed CMB description.

Combination	Model	95% limit on $\sum m_\nu$
BAO + Planck(full)	ΛCDM	0.25 eV
BAO + Planck(full)	$\Lambda\text{CDM} + \text{free } A_L$	0.43 eV
BAO + Planck	ΛCDM	0.56 eV
BAO + Planck	$w\text{CDM}$	0.68 eV
BAO + Planck	$\sigma\Lambda\text{CDM}$	0.87 eV
BAO + SN + Planck	ΛCDM	0.56 eV
BAO + SN + Planck	$w\text{CDM}$	0.61 eV
BAO + SN + Planck	$\sigma\Lambda\text{CDM}$	0.84 eV

For ΛCDM , fitting the CMB temperature power spectrum from Planck (together with WMAP polarization and high- l data from ground-based experiments) yields $A_L = 1.23 \pm 0.11$ [29], showing that the lensing signal measured in the Planck power spectrum is significantly stronger than the predicted signal based on extrapolating the observed CMB fluctuations forward in time. A larger neutrino mass suppresses the low-redshift clustering, exacerbating this tension, which is why the $\sum m_\nu$ limit is considerably tighter when A_L is fixed to unity. However, the Planck measurement of lensing through the CMB 4-point function does agree with $A_L = 1$ [29]. These internal tensions on the lensing signal within the CMB data alone suggest that one should be cautious in using them to constrain $\sum m_\nu$. The red curve is thus a more conservative inference, using only geometric constraints plus the CMB constraints on ω_{cb} and ω_b . It is impressive that even a 3.5% contribution of neutrinos to ω_m ($\sum m_\nu = 0.5$ eV) is enough to be substantially disfavored by these expansion history measurements.

D. Extra relativistic species

If the Universe contains extra relativistic degrees of freedom beyond those in the standard model, these increase the expansion rate during the radiation-dominated era and shift the epoch of matter-radiation equality, thereby altering the sound horizon, the shape of the matter power spectrum, and the history of recombination. Extra radiation is usually parametrized by the quantity ΔN_{eff} , where $\Delta N_{\text{eff}} = 1$ corresponds to the amount of radiation that an extra massless thermalized neutrino species (i.e., a fermion that thermally decouples before electron-positron annihilation) would produce. In general, however, there is no requirement that ΔN_{eff} be an integer. The standard model has $N_{\text{eff}} = 3.046$ and $\Delta N_{\text{eff}} = 0$.

The bottom row of Fig. 15 shows the probability distribution for N_{eff} from CMB data alone (right panel, from the chain BASE_NNU_PLANCK_LOWL_LOWLIKE_HIGHL), which peaks at $N_{\text{eff}} \approx 3.4$, with a 95% confidence range $2.7 \leq N_{\text{eff}} \leq 4.04$. The middle panel shows that values of N_{eff} at the upper end of this range can noticeably improve consistency with the LyaF BAO measurement, pushing the

predicted values of $D_H(2.34)/r_d$ up and $D_M(2.34)/r_d$ down so that they lie within the 95% likelihood contour of the BAO data. However, high values of N_{eff} reduce the predicted values of D_H/r_d and D_M/r_d at $z = 0.57$, worsening agreement with the galaxy BAO measurements.

We can understand these trends by arguments similar to those given for early dark energy in Sec. VI A. An increase of $\Delta N_{\text{eff}} = 1$ reduces the sound horizon r_d by 3.2% [Eq. (17)] because of the higher expansion rate in the early universe. Maintaining the precisely measured value of $D_M(1090)/r_d$ requires changes in Ω_m and h to reduce $D_M(1090)$ by the same factor. Some of this reduction can be accomplished by raising $\Omega_m h^2$, and thus raising $H(z)$ at high redshift, but $\Omega_m h^2$ is already tightly constrained by the heights of the acoustic peaks. Therefore, the fractional change to $D_H(2.34)$ is much lower than the fractional change to r_d , and the value of $D_H(2.34)/r_d$ rises. To maintain $D_M(1090)/r_d$, the value of H_0 (which controls the low-redshift contribution to the D_M integral) must increase by *more* than the drop in r_d . Because $D_M(2.34)$ is an integral over all $z < 2.34$, the ratio $D_M(2.34)/r_d$ drops even as $D_H(2.34)/r_d$ rises. At $z = 0.57$, where the value of $H(z)$ retains sensitivity to H_0 , both D_H/r_d and D_M/r_d drop as N_{eff} increases.

A change in N_{eff} has multiple effects on the CMB, which renders our compression into a 3-variable matrix questionable. For this section of the paper, therefore, we have run COSMOMC chains using the full Planck + WP CMB information, while still treating the BAO data as measurements of D_M/r_d and D_H/r_d . We have checked that using our SIMPLEMC chains, which adopt the compressed CMB description, yields qualitatively similar but quantitatively different results. We assume a flat universe with a cosmological constant, but we treat the tensor-to-scalar ratio r as a free parameter and marginalize over it, since there is no theoretical reason to expect $r = 0$. While allowing free r would not alter the compressed CMB constraints used elsewhere in the paper, it has an impact here because r and N_{eff} have partially degenerate effects on the shape of the CMB power spectrum.

Red contours in Fig. 17 show confidence intervals in the $N_{\text{eff}} - h$ plane obtained by combining the CMB data with Ly α F BAO alone. The allowed range of N_{eff} is larger here than in Fig. 15 because we do not fix $r = 0$. As one would expect from Fig. 15, the addition of Ly α F BAO pulls the preferred value of N_{eff} upward, with a best-fit value of $N_{\text{eff}} \approx 4$ and $N_{\text{eff}} = 3$ significantly disfavored. However, the galaxy BAO measurements prefer $N_{\text{eff}} \approx 3$ and have higher precision, so when they are added (green contours) the allowed range shifts downward to $N_{\text{eff}} = 3.43 \pm 0.26$ (68%) or $N_{\text{eff}} = 3.43 \pm 0.53$ (95%). Higher N_{eff} correlates with higher h for the reasons discussed above, and with N_{eff} and r as an additional degree of freedom our BAO + CMB constraint is $h = 0.71 \pm 0.017$. Adding SN data does not significantly shift these contours once galaxy BAO are




FIG. 17 (color online). Constraints on the effective number of relativistic species N_{eff} , assuming $w = -1$ and $\Omega_k = 0$. These contours use full Planck + WP CMB constraints computed with COSMOMC, combined with Ly α F BAO only (red) or with our full set of BAO measurements (green). We marginalize over the tensor-to-scalar ratio r .

included. As shown in Fig. 10, introducing ΔN_{eff} as a parameter reduces χ^2 by 0.75 relative to Λ CDM, with nearly all of the change coming from a slightly better fit to the Ly α F BAO data.

One complication in constraining ΔN_{eff} models is that changing the radiation density can alter the broadband power spectrum shape enough to affect the BAO fitting procedure itself. The authors of [26] found that the compression of BAO data into the $\alpha_{\parallel} - \alpha_{\perp}$ plane (in other words, the fact that the inferred values of D_M/r_d and D_H/r_d are independent of the adopted fiducial model) breaks down in the presence of extra radiation at a $\sim 0.4\%$ level. This effect is negligible compared to the statistical errors in the Ly α F BAO data, but it is not completely negligible relative to the galaxy BAO errors. A more exact treatment of the N_{eff} constraints therefore requires refitting the BAO data themselves, but we would expect only small shifts relative to the constraints reported here. We plan to revisit this question when the final BOSS measurements are available.

E. A tuned oscillation

While the ΔN_{eff} model moderately reduces tension with the Ly α F BAO data, *none* of the models we have considered produces a truly good fit to all the measurements. To understand what is required to achieve a good fit, we have constructed an artificial model that maintains the mathematical link between $D_M(z)$ and $D_H(z)$ but has the freedom needed to fit all the BAO and CMB data at the $\approx 1\sigma$ level. We consider a flat universe whose angular diameter distance is given by

$$D_M^{(w)}(z) = D_M^{(\Lambda\text{CDM})}(z)[1 + A \times G(\log z; \log z_o, \sigma_o)], \quad (31)$$

where $G(\log z; \log z_m, \sigma_o)$ denotes a Gaussian in $\log z$ with mean z_o and variance σ_o and A is an amplitude parameter. It




FIG. 18 (color online). A “tuned oscillation” model in which a Gaussian perturbation of the Λ CDM $D_M(z)$ is introduced to allow a good simultaneous fit to the galaxy and Ly α BAO data. Solid lines show the same Λ CDM model plotted in Fig. 1, while the dashed line shows the perturbed model.

is clear that such model can, for a sufficiently localized Gaussian perturbation, fit the low-redshift and CMB data with sufficient goodness of fit. Given three parameters it has enough flexibility to also match our $z = 2.34$ points. Figure 18 plots the best-fitting model of this form. The Hubble parameter undergoes an oscillation between $z \approx 4$ and $z \approx 0.8$, which allows it to match the Ly α and CMASS values of D_H and to change D_M at $z = 2.34$ without altering the low and high redshift values.

This model reduces the overall χ^2 by 6.6 with 3 extra degrees of freedom, a considerable improvement over any other model we have investigated (see Fig. 10). Generically, any small perturbation to the Friedmann equation that is able to improve fits to our Ly α data runs afoul of CMB and/or galaxy BAO constraints. This model works because it is fine-tuned to change distances near $z = 2$ but not upset the distance to the last-scattering surface. However, the model is physically extreme, as demonstrated in Fig. 19, where we have converted $H(z)$ into an implied dark energy density via the Friedmann equation. Producing the desired oscillation in $H(z)$ requires a *negative* ρ_{de} between $z = 6$ and $z = 2$ [see Eq. (27)]. The authors of [101] argue that the BOSS Ly α measurements may be explained in a modified gravity model that alters the Friedmann equation itself in a physically motivated way, but more work is needed to determine whether any such model can provide a good fit to all of the BAO measurements while satisfying CMB constraints.

The difficulty in finding a well-motivated model that matches the BOSS Ly α measurements suggests that the tension with these measurements may be a statistical fluke, or a consequence of an unrecognized systematic that either biases the central values of $D_M(z)$ and $D_H(z)$ or causes their error bars to be underestimated. Analyses of the final BOSS data set will address both of the latter points, as they will allow more exhaustive investigation of analysis procedures and tests against larger suites of mock catalogs.




FIG. 19 (color online). Implied variation in the energy density of the dark energy component for the model shown in Fig. 18. The dotted line corresponds to the density becoming negative. These plots illustrate the difficulty of concurrently fitting the Ly α and galaxy BAO constraints on D_M and D_H while satisfying the CMB constraint on D_M .

Addressing the first point will require high-redshift BAO measurements from new data sets, such as the Ly α emission-line galaxy survey of HETDEX [102] or a much larger Ly α sample from DESI [103].

VII. COMPARISON TO STRUCTURE GROWTH CONSTRAINTS

The Planck cosmology papers highlighted a tension between the predictions of the CMB-normalized Λ CDM model and observational constraints on matter clustering at low redshifts, from cluster abundances, weak gravitational lensing, or redshift-space distortions. We now revisit this issue with our updated BAO and SN constraints and our broader set of models, to see whether these tensions persist and whether they are significantly reduced in some classes of models.

Low-redshift measurements of cluster abundances and weak lensing most tightly constrain the parameter combination $\sigma_8 \Omega_m^\alpha$ with $\alpha \approx 0.4 - 0.6$ (see discussions in [35] and references therein). As representative but not exhaustive examples of constraints at $z \approx 0$ we adopt $\sigma_8(\Omega_m/0.27)^{0.46} = 0.774^{+0.032}_{-0.041}$ from tomographic cosmic shear measurements in the CFHTLenS survey [104]; $\sigma_8(\Omega_m/0.27)^{0.5} = 0.86 \pm 0.035$ from cosmic shear measurements in the Deep Lens Survey⁵ [105]; $\sigma_8(\Omega_m/0.27)^{0.57} = 0.77 \pm 0.05$ from the combination of galaxy-galaxy lensing and galaxy clustering in the SDSS [106]; $\sigma_8(\Omega_m/0.25)^{0.47} = 0.813 \pm 0.013$ from the mass function of x-ray clusters observed with Chandra and ROSAT [107]; $\sigma_8(\Omega_m/0.25)^{0.41} = 0.832 \pm 0.033$ from stacked weak lensing of clusters in the SDSS [108]; and

⁵The authors do not quote their results in this form, so this constraint has been estimated from their Fig. 25.




FIG. 20 (color online). Predictions of matter clustering from our BAO + SN + CMB constrained models compared to observational estimates. The vertical location of the observational estimates (red points) is arbitrary. Labels for the model points (black) in all panels are indicated along the left vertical axis. Panel (a) shows the $z = 0$ parameter combination $\sigma_8(\Omega_m/0.30)^{0.4}$, which approximately describes the quantity best constrained by low-redshift measurements of the cluster mass function or weak lensing. Black points show the mean and 1σ range computed from our model chains, and red points show observational estimates discussed in the text. Panel (b) presents a similar comparison for $\sigma_8(z = 0.57)f(z = 0.57)$, constrained by redshift-space distortions in CMASS galaxy clustering. Panel (c) compares $\sigma_8(z = 2.5)$ to an estimate from the BOSS Ly α 1-d power spectrum. Observational sources are the cosmic shear measurements of Hey13 [104] and Jee13 [105], the galaxy-galaxy lensing measurement of Man13 [106], the cluster mass function measurements of Vik09 [107], Roz10 [108], Pla13 [109], and Man14 [110], the RSD measurements of Beu14 [111], Sam14 [112], and Rei14 [113], and the Ly α power spectrum measurement of Pal13 [114]. Dotted vertical lines are provided for visual reference.

$\sigma_8(\Omega_m/0.27)^{0.3} = 0.78 \pm 0.01$ from Sunyaev-Zeldovich clusters in Planck, where we have taken the value quoted for a 20% x-ray mass bias [109]. These estimates are shown as red points with 1σ error bars in Fig. 20(a), where we have scaled the amplitudes to $\Omega_m = 0.30$ using the formulas listed above and retained the original fractional errors. We compare to model predictions of $\sigma_8(\Omega_m/0.3)^{0.4}$, treating 0.4 as a representative slope for these constraints.

Recently the authors of [115] have completed an independent analysis of tomographic cosmic shear in the CFHTLenS data and confirmed the findings of [104]. Conversely, the authors of [110] have performed a cluster mass function analysis using extensive weak lensing calibration of x-ray cluster masses and found $\sigma_8(\Omega_m/0.3)^{0.17} = 0.81 \pm 0.03$, which corresponds to higher σ_8 for $\Omega_m \approx 0.3$ than the other cluster studies listed above. We also show this point in Fig. 20(a), with the caution that the scaling with Ω_m reported by [110] is quite different from that of the other analyses.

For redshift-space distortion (RSD), the point labeled Beu14 in Fig. 20(b) shows the recent BOSS CMASS measurement by [111], which yields $f(z)\sigma_8(z) = 0.422 \pm 0.027$ at $z = 0.57$, where $f(z) \approx [\Omega_m(z)]^{0.55}$ is the linear fluctuation growth rate. This analysis fits simultaneously for redshift-space distortion and the Alcock-Paczynski (AP) effect [116]. Here we have used the error for fixed value of the AP parameter $D_M(z)H(z)$ because the geometry is well constrained by our BAO + SN + CMB data, so that the fractional error in the AP parameter is much smaller

than the 6.4% error on $f(z)\sigma_8(z)$. The point labeled Sam14 shows the estimate $f(z)\sigma_8(z) = 0.447 \pm 0.028$ from the same data set using a power spectrum analysis instead of a correlation function analysis. Since the data are the same, the difference from Beu14 provides an indication of the uncertainties associated with modeling systematics. Other analyses of redshift-space clustering in BOSS [117,118] and the WiggleZ survey [119] yield compatible results. We also plot an estimate of $f(z)\sigma_8(z) = 0.450 \pm 0.011$ from an analysis of smaller scale redshift-space distortions in the CMASS sample by [113], which adopts more aggressive modeling assumptions and achieves a substantially smaller statistical error.

At higher redshift, the 1-dimensional flux power spectrum of the Lyman- α forest probes the underlying matter clustering, with the tightest constraints on comoving scales of a few Mpc [114,120,121]. Here we take the result from the BOSS analysis of [114], who find $\sigma_8 = 0.83 \pm 0.03$ when fitting a Λ CDM model to the 1-d $P(k)$ at redshifts $z = 2.2 - 4.0$. We translate this result to a constraint on $\sigma_8(z = 2.5) = 0.311 \pm 0.011$ by using the growth factor at $z = 3$ for their central value of $\Omega_m = 0.26$. This measurement is indicated by a point with 1σ error bar in Fig. 20(c).

Given the wide range of models that we wish to consider and the several-percent errors on the observational data, we have opted for an approximate method of computing clustering amplitude predictions that is accurate at the 0.5% level or better. Following the strategy of [122] and [35], we first use CAMB calculations to calibrate a Taylor

expansion for the value of $\sigma_8(z=9)$ about a fiducial Planck Λ CDM model, finding

$$\begin{aligned} \sigma_8(z=9) = & 0.1058 \times \left(\frac{A_s}{2.196 \times 10^{-9}} \right)^{1/2} \left(\frac{\Omega_m h^2}{0.1426} \right)^{0.520} \\ & \times \left(\frac{\Omega_b h^2}{0.02205} \right)^{-0.294} \left(\frac{h}{0.673} \right)^{0.683} \left(\frac{N_{\text{eff}}}{3.046} \right)^{-0.24} \\ & \times e^{0.3727(n_s-0.96)} (1 - \Omega_k)^{0.175}. \end{aligned} \quad (32)$$

Here A_s is the amplitude of primordial curvature perturbations at the scale $k_0 = 0.05 \text{ Mpc}^{-1}$. This formula updates Eqs. (46)–(47) of [35], which were expanded about a WMAP7 fiducial model. The fairly strong h -dependence arises because of the conversion from a power spectrum predicted in Mpc units based on cosmological parameter values to an amplitude defined on a scale of $8 h^{-1} \text{ Mpc}$. We have made numerous checks of this formula against full CAMB calculations for models in the parameter ranges allowed by Planck + WP data, finding accuracy of better than 0.1% for Λ CDM, for $\omega\Lambda$ CDM with $-0.2 < \Omega_k < 0.2$, and for w CDM with $-1.2 < w < -0.8$, and accuracy better than 0.5% for ΔN_{eff} models with $2.5 < N_{\text{eff}} < 4.5$. While Eq. (32) correctly reflects the response of $\sigma_8(z=9)$ to an isolated change in N_{eff} , in practice the CMB-preferred values of A_s , ω_m , ω_b , h , and n_s all increase when N_{eff} increases, with the result that higher N_{eff} models end up predicting higher clustering amplitudes. We have not attempted to incorporate the effects of nonzero neutrino mass in this formula because the suppression of clustering by neutrino free streaming is redshift and scale dependent (see [123] for useful representations).

Except in early dark energy models, the value of $\sigma_8(z=9)$ is essentially independent of dark energy parameters because dark energy is dynamically insignificant at $z > 9$ (e.g., $\rho_\Lambda/\rho_m < 0.003$ for a cosmological constant). To evolve σ_8 forward to $z=3$, 0.57, or 0, we use the approximate integral formulation of the growth factor from equation (16) of [35], which simply integrates the growth rate approximation of [124], $f(z) \approx [\Omega_m(z)]^\gamma$ with $\gamma = 0.55 + 0.05[1 + w(z=1)]$. Spot checks against exact calculations with COSMOMC indicate that this approach is accurate to 0.3% or better for models with $m_\nu = 0$ and other parameters in the range allowed by our CMB + BAO + SN data, although it becomes less accurate for more extreme parameter values (especially of Ω_k). For $\sum m_\nu = 0.06 \text{ eV}$, CAMB yields a ratio $\sigma_8(z=0)/\sigma_8(z=9)$ that is 0.5% lower than for $m_\nu = 0$, with little dependence on other parameters, so we also multiply all of our low-redshift σ_8 values by 0.995 to account for this effect.

We determine the mean values and error bars on the predicted growth observables for our models by computing the posterior-weighted mean and 1σ dispersion of $\sigma_8 \Omega_m^{0.4}$, $\sigma_8(z=0.57)[\Omega_m(z=0.57)]^{0.55}$, or $\sigma_8(z=2.5)$ for the parameter values in our MCMC chains, using the above

approximations for σ_8 . Because our chains do not actually use or include values of A_s , we compute σ_8 for the fiducial value in Eq. (32) and add a fractional error (based on the Planck + WP column in Table 5 of [29]) of 1.25% in quadrature to the MCMC error to account for the 2.5% error in A_s , which is proportional to σ_8^2 . Inspection of Planck chains indicates only weak correlations between A_s and other cosmological parameters, so the approximation of an independent error contribution added in quadrature should be adequate. We also add in quadrature a fractional error of 0.3% to represent potential errors of our approximate growth calculations, though our spot checks indicate higher accuracy than this.

Figure 20(a) shows a persistent offset between the predicted amplitude of matter clustering and the majority of observational estimates from weak lensing and cluster masses. For Λ CDM, where the model predictions are best constrained, the statistical significance of the tension with any given data set is usually only $\approx 2\sigma$ or smaller. However, the sign of discrepancy is usually the same, so the overall significance is high unless multiple analyses are affected by a common systematic. The important exceptions are the cosmic shear measurement from the Deep Lens Survey [105] and the recent cluster analysis of [110], which both agree well with the Λ CDM prediction. The authors of [110] emphasize that theirs was a “blind” analysis in which technical choices about data cuts and procedures were made without knowing their eventual impact on inferred cosmological parameter values. Our predicted value of $\sigma_8 \Omega_m^{0.4}$ is somewhat lower than the value inferred by [29] from CMB data alone (Planck + WP + highL), in part because the BAO data pull towards lower Ω_m , and in part because our compressed CMB description does not include the lensing information in the Planck power spectrum, which pulls towards higher σ_8 . For more flexible dark energy models, central values of $\sigma_8 \Omega_m^{0.4}$ remain within the 1σ range found for Λ CDM, and the error bars are moderately larger. The tension with the data is moderately reduced in these models, but not eliminated. A formal assessment of this reduced disagreement is difficult because the true level of systematic uncertainty in the low-redshift measurements is itself uncertain, but at a qualitative level this reduction appears too small to favor adopting one of these more complex models.

Figure 20(b) shows an offset of similar magnitude between the predictions of our standard dark energy models and the redshift-space distortion measurement of [111] at $z=0.57$. However, the statistical significance of this tension is low because of the statistical error on the measurement, and the power spectrum analysis of [112] yields a higher central value. The more precise determination from [113], which draws on simulation-based modeling of nonlinear scales, overlaps the Λ CDM prediction at $\approx 1\sigma$. A recent analysis that combines galaxy clustering and galaxy-galaxy lensing of CMASS galaxies

[125] yields constraints in the σ_8 - Ω_m plane that are consistent with the Λ CDM model predictions, but the current statistical errors are large enough to be compatible with the central values from any of the redshift-space distortion analyses shown here. In contrast to the lower redshift results, Fig. 20(c) shows that the predicted clustering amplitude at $z = 2.5$ in our standard models is *lower* than that inferred from the LyaF power spectrum, though consistent at $\approx 1\sigma$.

In the ΔN_{eff} model, where we assume a Λ CDM cosmology but allow extra relativistic species, the preferred value of N_{eff} is higher than the standard value of 3.046, as shown previously in Fig. 17. Because of the correlation of N_{eff} with other cosmological parameters in CMB fits, the central value of the clustering amplitude predictions shifts upwards, while the freedom in N_{eff} broadens the error bar relative to standard Λ CDM. These changes noticeably increase the tension with the clustering measurements at $z = 0$ and the RSD measurements at $z = 0.57$, though they improve agreement with the LyaF power spectrum at $z = 2.5$. Overall, current clustering measurements provide moderate evidence against extra relativistic species, though a firmer understanding of systematic uncertainties in these measurements will be needed to draw solid conclusions.

Massive neutrinos have a redshift- and scale-dependent impact on matter clustering, which changes between the linear and nonlinear regimes. We have not attempted a full examination of free- $\sum m_\nu$ models in this section because the summary of the observational results in terms of σ_8 - Ω_m constraints may not adequately capture the effect of massive neutrinos on the clustering observables. A value of $\sum m_\nu = 0.5$ eV, near the 95% upper bound inferred from our compressed CMB description and BAO constraints, would lower the predicted value of σ_8 in a Planck + WP-normalized Λ CDM model by about 12% relative to $\sum m_\nu = 0.06$ eV. A value $\sum m_\nu = 0.25$ eV, near the upper bound that we find when combining the full Planck likelihood with BAO data, would produce a 6% suppression of σ_8 . These numbers are somewhat different from what a naive expectation based on linear suppression would indicate because CMB degeneracies are important at these relatively large neutrino mass fractions. Even the lower value is enough to remove the tension seen in Fig. 20(a). However, the corresponding decrease in $\sigma_8(z = 2.5)$ produces a significant discrepancy with the LyaF measurement in Fig. 20(c), and a full analysis that models the LyaF power spectrum based on hydrodynamic simulations with a massive neutrino component leads to a stringent upper limit on neutrino mass [126].

As discussed in Sec. VI A, our geometric constraints are nearly degenerate with respect to the presence of an early dark energy component, provided this early dark energy is present in the radiation-dominated epoch as well as the matter-dominated epoch and therefore shrinks the scale of the sound horizon. Increasing the early dark energy fraction

reduces the value of Ω_m (see Fig. 13) and will also suppress growth of structure relative to Λ CDM. Predictions of structure for early dark energy are subtle because of the combined impacts of CMB normalization, the imprint of early dark energy fluctuations on the CMB itself, and the postrecombination growth rate. We therefore defer detailed investigation of early dark energy models to future work and make the qualitative observation that an early dark energy component will go in the direction of reducing tensions with low redshift clustering measurements.

VIII. CONCLUSIONS

In the decade since the first observational detection of baryon acoustic oscillations, BAO analysis has emerged as one of the sharpest tools of precision cosmology. Its power arises from the grounding of its absolute distance scale in straightforward underlying physics, from the distinctiveness of a feature that is localized in scale and thus not easily mimicked by observational systematics, and from its insensitivity to nonlinear gravitational evolution and galaxy formation physics (a consequence of the large scale of BAO). The principal challenge of the method is that one must map enormous cosmic volumes to obtain good statistical precision. Building on the legacy of 2dFGRS, SDSS-I/II, 6dFGS, and WiggleZ, BOSS has made major progress on this challenge, with distance scale measurements of 1%–2% precision at $z = 0.32, 0.57,$ and 2.34 . The combination of BAO measurements with Planck + WP CMB data and the JLA SNIa compilation leads to numerous significant constraints on dark energy, space curvature, and the cosmic matter and radiation density.

If we treat BAO as an *uncalibrated* standard ruler, assuming only that it is constant in time, then the combination of galaxy and LyaF BAO measurements yields a strong ($> 3\sigma$) detection of dark energy, independent of any other cosmological data. If we assume that the angular acoustic scale of the CMB represents the same standard ruler, then the resulting constraints in an ω Λ CDM model collapse around a flat universe dominated by dark energy, with $\Omega_m = 0.292 \pm 0.18$, $\Omega_k = -0.010 \pm 0.016$. Thus, high-precision measurements of a common standard ruler at $z < 0.7$, $z = 2.34$, and $z = 1090$ already lead to strong constraints on the cosmological model.

BAO become much more powerful when we incorporate the absolute calibration of the sound horizon r_d using CMB measurements of the matter, baryon, and radiation energy density [Eq. (16)]. With Planck + WP CMB data, residual uncertainties in ω_m and ω_b leave only 0.4% uncertainty in the acoustic scale $r_d = 147.49 \pm 0.59$ Mpc assuming a standard radiation background with three neutrino species. One particularly interesting application of this calibration is to combine galaxy BAO measurements with the high-precision measurements of *relative* distances from Type Ia SNe to infer H_0 . The addition of the SN data makes the inferred value of H_0 insensitive to uncertainties in the dark

energy model, which would otherwise affect the extrapolation of the distance scale from the moderate redshifts of the BAO measurements down to $z = 0$. With our standard BAO and SN data sets, this inverse distance ladder measurement yields $H_0 = 67.3 \pm 1.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$, where the 1.7% uncertainty includes the Planck + WP uncertainty in r_d . This value agrees perfectly with the value inferred from current CMB data under the much stronger assumption of a flat Λ CDM model, an important consistency test of the standard cosmology. It is lower than most recent estimates using a Cepheid-based distance ladder. Our measurement of H_0 does rely on the assumption of a standard cosmic radiation background, and the directly constrained parameter combination is $H_0 r_d$. A convincing discrepancy with conventional distance-ladder determinations of H_0 could not be resolved by appealing to the late-time behavior of dark energy. It would point instead to nonstandard physics in the pre-recombination universe, such as extra relativistic degrees of freedom or early dark energy, which can shrink r_d and thus raise the inferred value of H_0 .

The full combination of CMB, BAO, and SN data places strong constraints on dark energy and space curvature, as summarized in Fig. 8 and Table IV. In models that allow both $w \neq -1$ and nonzero curvature, the BAO and SN data are highly complementary. For the ow CDM model, we find $w = -0.98 \pm 0.06$ and $\Omega_k = -0.002 \pm 0.003$. For models with $w(a) = w_0 + w_a(1 - a)$, the constraint on evolution remains poor, with $w_a = -0.6 \pm 0.6$ in ow_0w_a CDM, but the value of w at the pivot redshift where it is best constrained remains close to -1 . A striking feature of Table IV is that as degrees of freedom are added to the cosmological model the best-fit values of parameters barely change, always remaining close to those of flat Λ CDM.

These models are fit to a total of 43 observables: 3 in our compressed description of the CMB, 31 for the compressed SN data, 5 for the galaxy BAO data (D_V from 6dFGS, MGS and BOSS LOWZ, D_M and D_H from BOSS CMASS), and 4 for the LyaF BAO (D_H and D_M from forest autocorrelation and from quasar-forest cross correlation). The Λ CDM model, with three free parameters (Ω_Λ , h , and the absolute magnitude normalization for SNIa), has $\chi^2 = 46.79$ for 40 d.o.f., which is statistically acceptable.⁶ The decrease in χ^2 for the alternative models is not enough to justify the addition of parameters; for example, the addition of three free parameters in ow_0w_a CDM reduces χ^2 by only 1.33. However, the best-fit models in all of these cases are in significant tension with the LyaF measurements on their own, typically at the 2–2.5 σ level. The LyaF data have little impact on the best-fit parameter values in any of these models, not because they agree well with the model predictions but because parameter changes that would

significantly improve agreement with the LyaF run afoul of the higher precision galaxy BAO measurements. Moreover, because the LyaF measurements have lower D_H but higher D_M than expected in the best-fit Λ CDM model, many parameter changes that would improve the fit to D_H worsen the fit to D_M , and vice versa.

We have examined several models with nonstandard dark energy or dark matter histories or nonstandard radiation backgrounds. Early dark energy that has constant Ω_{de}^e in the radiation and matter dominated eras (before evolving towards a cosmological constant at low redshift) alters the sound horizon r_d and evolution of $H(z)$ and $D_M(z)$. Remarkably, the cancellation of these effects leaves the BAO observables $D_H(z)/r_d$ and $D_M(z)/r_d$ nearly unchanged (including at $z = 1090$), even for Ω_{de}^e as large as 0.3, so that the observations incorporated in our fits still allow a substantial early dark energy component. Because of the smaller r_d , such model fits yield a higher H_0 and lower Ω_m , and the suppression of growth by early dark energy is likely to reduce the amplitude of low-redshift matter clustering. Full CMB power spectrum analyses yield stronger but more model-dependent constraints on early dark energy through its influence on the shape of the acoustic peaks and the structure of the damping tail [91]. Nonetheless, the ability of these models to match expansion history constraints while improving agreement with local H_0 and structure growth measurements suggests that they merit further investigation.

We also find surprisingly weak constraints on models in which cold dark matter decays into radiation at low redshifts. For a model with a single CDM species, our BAO+SN+CMB constraints allow nearly 50% of dark matter to decay by $z = 0$, corresponding to an inverse decay constant $\Gamma^{-1} > 28 \text{ Gyr}$ (at 95% confidence). Constraints on the decay products may allow much tighter limits for specific classes of dark matter models, but expansion history alone is not enough to show that dark matter is stable.

With respect to expansion history, massive neutrinos are in some sense the converse of decaying dark matter: they are relativistic at the epoch of recombination, but at low redshift they increase the matter density Ω_m relative to the value Ω_{cb} inferred from the CMB acoustic peaks. The purely geometric constraints that come from BAO and our compressed CMB description yield a 95% confidence upper limit of $\sum m_\nu < 0.56 \text{ eV}$ assuming Λ CDM, with moderately weaker limits for models that allow $w \neq -1$ or nonzero Ω_k (see Table VI). If we use full Planck CMB chains in place of our compressed description we obtain the significantly stronger limit $\sum m_\nu < 0.25 \text{ eV}$, a difference driven by the relatively high amplitude lensing signal detected in the Planck power spectrum. Measurements of low-redshift matter clustering can yield more sensitive limits on neutrino masses, and potentially a measurement of $\sum m_\nu$ through its impact on structure growth, but the

⁶Note that Fig. 10 omits the CMB data from the χ^2 accounting, so it has three fewer d.o.f.

expansion history constraints are robust and impressively stringent on their own.

Adding relativistic degrees of freedom can noticeably improve the agreement with the LyaF BAO, and if we combine only CMB and LyaF data the preferred N_{eff} is ≈ 4 . However, increasing N_{eff} worsens agreement with the galaxy BAO data, and when we consider our full data combination we find $N_{\text{eff}} = 3.43 \pm 0.26$. Increasing N_{eff} reduces the value of r_d and thereby leads to a higher inferred H_0 ; for a model with free N_{eff} and free tensor-to-scalar ratio r we find a marginalized constraint $H_0 = 70.1 \pm 1.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We caution that modifying the radiation background alters the shape of the acoustic peaks, an effect not accounted for in the BAO measurements used here; we expect this effect to be smaller than our statistical errors, but perhaps not negligible.

Among these alternative models, only the model with free N_{eff} can reduce the tension with the LyaF data, and even there the reduction is small once the galaxy BAO constraints are also imposed. We did construct a model with a tuned oscillation in $D_M(z)$ that reproduces both the D_M/r_d and D_H/r_d measurements from the LyaF BAO while continuing to satisfy all other constraints. However, this model requires nonmonotonic evolution of $H(z)$ and thus of $\rho_{\text{tot}}(z)$, which is difficult to achieve in any model with non-negative dark energy density. The artificiality and physical implausibility of this model, and the failure of our more physically motivated models, illustrate how difficult it is to obtain a good fit to the BOSS LyaF BAO measurements. This difficulty suggests that the tension of simpler models with the LyaF data is a statistical fluke, or perhaps reflects an unrecognized systematic in the BAO measurement, but it highlights the importance of further measurements of D_M and $H(z)$ at high redshifts.

The cosmological constraints considered here are essentially geometrical, tied to the expansion history of the homogeneous universe. As a further test, we have computed the predictions of our models for low redshift measurements of matter clustering. Confirming previous findings, but now with tighter cosmological parameter constraints, we find that a Λ CDM model normalized to the observed amplitude of CMB anisotropies predicts cluster masses, weak lensing signals, and redshift-space distortions that are higher than most observational estimates. The tension with individual data sets is only $\approx 2\sigma$, and the measurements themselves may be affected by systematics. However, the direction of the discrepancy is consistent across many analyses (though not all of them). The additional freedom in standard dark energy models does not reduce this tension because the parameter values allowed by our data are always close to those of Λ CDM. Massive neutrinos can reduce the tension by suppressing structure growth on small scales (lowering σ_8), an effect that is small but not negligible for neutrino masses in the range allowed by our fits. Conversely, increasing N_{eff}

above the standard value of 3.046 leads to higher predicted values of σ_8 because of correlation with other cosmological parameters, thus amplifying the tension. As previously noted, early dark energy may reduce the tension with the clustering data, both because it suppresses growth of structure during the matter dominated era and because the reduced r_d value leads to higher h and lower Ω_m when combined with CMB constraints. Decaying dark matter models might also reduce the tension through a lower value of Ω_m at low redshift. Our standard Λ CDM fits produce good agreement with the matter clustering amplitude inferred from the LyaF power spectrum at $z \approx 2.5$; this agreement is itself an important constraint on neutrino masses or other physical mechanisms that reduce small scale clustering [126].

The application of the BAO technique to large cosmological surveys has enabled the first percent-level measurements of absolute distances beyond the Milky Way. In combination with CMB and SN data, these measurements yield impressively tight constraints on the cosmic expansion history and correspondingly stringent tests of dark energy theories. Over the next year, the strength of these tests will advance significantly with the final results from BOSS and with CMB polarization and improved temperature maps from Planck. In the longer term, BAO measurements will gain in precision and redshift range through a multitude of ongoing or planned spectroscopic surveys, including SDSS-IV eBOSS, HETDEX, SuMIRE, DESI, WEAVE, Euclid, and WFIRST.⁷ These data sets also enable precise measurements of matter clustering through redshift-space distortion analyses, the shape of the 3-dimensional power spectrum, and other clustering statistics. In combination with the expansion history constraints, these measurements can test modified gravity explanations of cosmic acceleration and probe the physics of inflation, the masses of neutrinos, and the properties of dark matter. In parallel with these large spectroscopic surveys, supernova measurements of expansion history are gaining in precision, data quality, and redshift range, and weak lensing constraints on matter clustering are advancing to the percent and subpercent level as imaging surveys grow from millions of galaxy shape measurements to hundreds of millions, and eventually to billions. From the mid-1990s through the early 21st century, improving cosmological data sets transformed our picture of the Universe. The next decade—of time and of precision—could bring equally surprising changes to our understanding of the cosmos.

ACKNOWLEDGMENTS

We thank Eric Linder for useful discussions of early dark energy and structure growth. We also thank Savvas Koushiappas and Gordon Blackadder for alerting us to

⁷See [127] for a brief summary of these projects and references to more detailed descriptions.

an error in the decaying dark matter section of the preprint version of this paper and answering our questions as we corrected it. Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>. SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

APPENDIX: DECAYING DARK MATTER MODEL

We consider a model of dark matter decaying into radiation as

$$\dot{\rho}_x = -3H(t)\rho_x - \lambda H_0 \rho_x, \quad (\text{A1})$$

$$\dot{\rho}_g = -4H(t)\rho_g + \lambda H_0 \rho_x, \quad (\text{A2})$$

where ρ_x and ρ_g are the new decaying dark matter and radiation components and the decay time constant λ is made

dimensionless by expressing it in units of H_0 . The Hubble parameter is given by the usual expression for Λ CDM with two extra components

$$\left(\frac{H}{H_0}\right)^2 = \Omega_{cb}a^{-3} + \Omega_\Lambda + \rho_{\nu+r}(z)/\rho_{\text{crit}} + \frac{\rho_x(a) + \rho_g(a)}{\rho_{\text{crit}}}. \quad (\text{A3})$$

Writing $\rho_x = \rho_{\text{crit}} r_x a^{-3}$ and $\rho_g = \rho_{\text{crit}} r_r a^{-4}$, the system of equations can be rewritten as

$$\frac{dr_x}{d \ln a} = -\lambda r_x \left(\frac{H}{H_0}\right)^{-1}, \quad (\text{A4})$$

$$\frac{dr_r}{d \ln a} = +\lambda r_x \left(\frac{H}{H_0}\right)^{-1}, \quad (\text{A5})$$

$$\left(\frac{H}{H_0}\right)^2 = \Omega_{cb}a^{-3} + \Omega_\Lambda + \rho_{\nu+r}(z)/\rho_{\text{crit}} + r_x(a)a^{-3} + r_r(a)a^{-4}, \quad (\text{A6})$$

with initial conditions $r_x(a=1) = \Omega_x$ and $r_r(a=1) = \Omega_r$. We can solve this system of differential equations starting at $a=1$ and going backwards in time for a given choice of Ω_x , Ω_r and λ .

However, in our parametrization, boundary conditions are specified in the infinite past. We therefore use a minimizing routine that determines the values of Ω_x and Ω_r today that are required to obtain the right fraction of decaying dark matter fraction and zero initial density in the decay product in the infinite past (assumed to be $a \sim 10^{-4}$ in the code). At each step in minimization, the evolution equations are solved numerically and a suitable penalty function is evaluated.

-
- [1] A. D. Sakharov, *Soviet Physics JETP*. **22**, 241 (1966).
 [2] P. J. E. Peebles and J. T. Yu, *Astrophys. J.* **162**, 815 (1970).
 [3] R. A. Sunyaev and Y. B. Zeldovich, *Astrophys. Space Sci.* **7**, 3 (1970).
 [4] C. Blake and K. Glazebrook, *Astrophys. J.* **594**, 665 (2003).
 [5] H.-J. Seo and D. J. Eisenstein, *Astrophys. J.* **598**, 720 (2003).
 [6] M. Hamuy, M. M. Phillips, N. B. Suntzeff, R. A. Schommer, J. Maza, A. R. Antezan, M. Wischnjewsky, G. Valladares, C. Muena, L. E. Gonzales *et al.*, *Astron. J.* **112**, 2408 (1996).
 [7] A. G. Riess, A. V. Filippenko, P. Challis, A. Clocchiatti, A. Diercks, P. M. Garnavich *et al.*, *Astron. J.* **116**, 1009 (1998).
 [8] S. Perlmutter, G. Aldering, G. Goldhaber, R. A. Knop, P. Nugent, P. G. Castro *et al.*, *Astrophys. J.* **517**, 565 (1999).
 [9] D. W. Hogg, [arXiv:astro-ph/9905116](https://arxiv.org/abs/astro-ph/9905116).
 [10] S. Cole, W. J. Percival, J. A. Peacock, P. Norberg, C. M. Baugh, C. S. Frenk, I. Baldry, J. Bland-Hawthorn, T. Bridges, R. Cannon *et al.*, *Mon. Not. R. Astron. Soc.* **362**, 505 (2005).
 [11] D. J. Eisenstein, I. Zehavi, D. W. Hogg, R. Scoccimarro, M. R. Blanton, R. C. Nichol, R. Scranton, H.-J. Seo, M. Tegmark, Z. Zheng *et al.*, *Astrophys. J.* **633**, 560 (2005).

- [12] M. Colless, G. Dalton, S. Maddox, W. Sutherland, P. Norberg, S. Cole *et al.*, *Mon. Not. R. Astron. Soc.* **328**, 1039 (2001).
- [13] D. J. Eisenstein, J. Annis, J. E. Gunn, A. S. Szalay, A. J. Connolly, R. C. Nichol *et al.*, *Astron. J.* **122**, 2267 (2001).
- [14] D. G. York, J. Adelman, J. E. Anderson, Jr., S. F. Anderson, J. Annis, N. A. Bahcall, J. A. Bakken, R. Barkhouser, S. Bastian, E. Berman *et al.*, *Astron. J.* **120**, 1579 (2000).
- [15] W. J. Percival, B. A. Reid, D. J. Eisenstein, N. A. Bahcall, T. Budavari, J. A. Frieman *et al.*, *Mon. Not. R. Astron. Soc.* **401**, 2148 (2010).
- [16] N. Padmanabhan, X. Xu, D. J. Eisenstein, R. Scalzo, A. J. Cuesta, K. T. Mehta, and E. Kazin, *Mon. Not. R. Astron. Soc.* **427**, 2132 (2012).
- [17] D. J. Eisenstein, H.-J. Seo, E. Sirko, and D. N. Spergel, *Astrophys. J.* **664**, 675 (2007).
- [18] M. J. Drinkwater, R. J. Jurek, C. Blake, D. Woods, K. A. Pimblett, K. Glazebrook, R. Sharp, M. B. Pracy, S. Brough, M. Colless *et al.*, *Mon. Not. R. Astron. Soc.* **401**, 1429 (2010).
- [19] C. Blake, E. A. Kazin, F. Beutler, T. M. Davis, D. Parkinson, S. Brough *et al.*, *Mon. Not. R. Astron. Soc.* **418**, 1707 (2011).
- [20] D. H. Jones, M. A. Read, W. Saunders, M. Colless, T. Jarrett, Q. A. Parker, A. P. Fairall, T. Mauch, E. M. Sadler, F. G. Watson *et al.*, *Mon. Not. R. Astron. Soc.* **399**, 683 (2009).
- [21] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders, and F. Watson, *Mon. Not. R. Astron. Soc.* **416**, 3017 (2011).
- [22] M. A. Strauss, D. H. Weinberg, R. H. Lupton, V. K. Narayanan, J. Annis, M. Bernardi, M. Blanton, S. Burles, A. J. Connolly, J. Dalcanton *et al.*, *Astron. J.* **124**, 1810 (2002).
- [23] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden, and M. Manera, *Mon. Not. R. Astron. Soc.* **449**, 835 (2015).
- [24] K. S. Dawson, D. J. Schlegel, C. P. Ahn, S. F. Anderson, É. Aubourg, S. Bailey, R. H. Barkhouser, J. E. Bautista, A. Beifiori, A. A. Berlind *et al.*, *Astron. J.* **145**, 10 (2013).
- [25] D. J. Eisenstein, D. H. Weinberg, E. Agol, H. Aihara, C. Allende Prieto, S. F. Anderson *et al.*, *Astron. J.* **142**, 72 (2011).
- [26] L. Anderson, É. Aubourg, S. Bailey, F. Beutler, V. Bhardwaj, M. Blanton, A. S. Bolton, J. Brinkmann, J. R. Brownstein, A. Burden *et al.*, *Mon. Not. R. Astron. Soc.* **441**, 24 (2014).
- [27] T. Delubac, J. E. Bautista, N. G. Busca, J. Rich, D. Kirkby, S. Bailey, A. Font-Ribera, A. Slosar, K.-G. Lee, M. M. Pieri *et al.*, *Astron. Astrophys.* **574**, A59 (2015).
- [28] A. Font-Ribera, D. Kirkby, N. Busca, J. Miralda-Escudé, N. P. Ross, A. Slosar, J. Rich, É. Aubourg, S. Bailey, V. Bhardwaj *et al.*, *J. Cosmol. Astropart. Phys.* **05** (2014) 027.
- [29] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday *et al.*, *Astron. Astrophys.* **571**, A16 (2014).
- [30] M. Betoule, R. Kessler, J. Guy, J. Mosher, D. Hardin, R. Biswas, P. Astier, P. El-Hage, M. Konig, S. Kuhlmann *et al.*, *Astron. Astrophys.* **568**, A22 (2014).
- [31] A. Conley, J. Guy, M. Sullivan, N. Regnault, P. Astier, C. Balland, S. Basa, R. G. Carlberg, D. Fouchez, D. Hardin *et al.*, *Astrophys. J. Suppl. Ser.* **192**, 1 (2011).
- [32] J. A. Frieman, B. Bassett, A. Becker, C. Choi, D. Cinabro, F. DeJongh, D. L. Depoy, B. Dilday, M. Doi, P. M. Garnavich *et al.*, *Astron. J.* **135**, 338 (2008).
- [33] M. Sako, B. Bassett, A. C. Becker, P. J. Brown, H. Campbell, R. Cane, D. Cinabro, C. B. D'Andrea, K. S. Dawson, F. DeJongh *et al.*, arXiv:1401.3317.
- [34] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett *et al.*, arXiv:1502.01589.
- [35] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, *Phys. Rep.* **530**, 87 (2013).
- [36] A. Friedmann, *Z. Phys.* **10**, 377 (1922).
- [37] See <https://github.com/slosar/april>.
- [38] A. Lewis, A. Challinor, and A. Lasenby, *Astrophys. J.* **538**, 473 (2000).
- [39] J. E. Gunn, W. A. Siegmund, E. J. Mannery, R. E. Owen, C. L. Hull, R. F. Leger, L. N. Carey, G. R. Knapp, D. G. York, W. N. Boroski *et al.*, *Astron. J.* **131**, 2332 (2006).
- [40] S. A. Smee, J. E. Gunn, A. Uomoto, N. Roe, D. Schlegel, C. M. Rockosi, M. A. Carr, F. Leger, K. S. Dawson, M. D. Olmstead *et al.*, *Astron. J.* **146**, 32 (2013).
- [41] A. S. Bolton, D. J. Schlegel, É. Aubourg, S. Bailey, V. Bhardwaj, J. R. Brownstein, S. Burles, Y.-M. Chen, K. Dawson, D. J. Eisenstein *et al.*, *Astron. J.* **144**, 144 (2012).
- [42] R. Tojeiro, A. J. Ross, A. Burden, L. Samushia, M. Manera, W. J. Percival, F. Beutler, J. Brinkmann, J. R. Brownstein, A. J. Cuesta *et al.*, *Mon. Not. R. Astron. Soc.* **440**, 2222 (2014).
- [43] N. G. Busca, T. Delubac, J. Rich, S. Bailey, A. Font-Ribera, D. Kirkby, J.-M. Le Goff, M. M. Pieri, A. Slosar, É. Aubourg *et al.*, *Astron. Astrophys.* **552**, A96 (2013).
- [44] A. Slosar, V. Iršič, D. Kirkby, S. Bailey, N. G. Busca, T. Delubac, J. Rich, É. Aubourg, J. E. Bautista, V. Bhardwaj *et al.*, *J. Cosmol. Astropart. Phys.* **04** (2013) 026.
- [45] D. Kirkby, D. Margala, A. Slosar, S. Bailey, N. G. Busca, T. Delubac, J. Rich, J. E. Bautista, M. Blomqvist, J. R. Brownstein *et al.*, *J. Cosmol. Astropart. Phys.* **03** (2013) 024.
- [46] A. Slosar, A. Font-Ribera, M. M. Pieri, J. Rich, J.-M. Le Goff, É. Aubourg, J. Brinkmann, N. Busca, B. Carithers, R. Charlassier *et al.*, *J. Cosmol. Astropart. Phys.* **09** (2011) 001.
- [47] I. Pâris, P. Petitjean, É. Aubourg, N. P. Ross, A. D. Myers, A. Streblyanska, S. Bailey, P. B. Hall, M. A. Strauss, S. F. Anderson *et al.*, *Astron. Astrophys.* **563**, A54 (2014).
- [48] N. P. Ross, A. D. Myers, E. S. Sheldon, C. Yèche, M. A. Strauss, J. Bovy, J. A. Kirkpatrick, G. T. Richards, É. Aubourg, M. R. Blanton *et al.*, *Astrophys. J. Suppl. Ser.* **199**, 3 (2012).
- [49] J. Bovy, A. D. Myers, J. F. Hennawi, D. W. Hogg, R. G. McMahon, D. Schiminovich, E. S. Sheldon, J. Brinkmann,

- D. P. Schneider, and B. A. Weaver, *Astrophys. J.* **749**, 41 (2012).
- [50] J. A. Kirkpatrick, D. J. Schlegel, N. P. Ross, A. D. Myers, J. F. Hennawi, E. S. Sheldon, D. P. Schneider, and B. A. Weaver, *Astrophys. J.* **743**, 125 (2011).
- [51] G. T. Richards, A. D. Myers, A. G. Gray, R. N. Riegel, R. C. Nichol, R. J. Brunner, A. S. Szalay, D. P. Schneider, and S. F. Anderson, *Astrophys. J. Suppl. Ser.* **180**, 67 (2009).
- [52] C. Yèche, P. Petitjean, J. Rich, E. Aubourg, N. Busca, J.-C. Hamilton, J.-M. Le Goff, I. Paris, S. Peirani, C. Pichon *et al.*, *Astron. Astrophys.* **523**, A14 (2010).
- [53] See <http://darkmatter.ps.uci.edu/baofit/>.
- [54] A. Pontzen, *Phys. Rev. D* **89**, 083010 (2014).
- [55] S. Gontcho A Gontcho, J. Miralda-Escudé, and N. G. Busca, *Mon. Not. R. Astron. Soc.* **442**, 187 (2014).
- [56] A. Lewis and S. Bridle, *Phys. Rev. D* **66**, 103511 (2002).
- [57] See http://wiki.cosmos.esa.int/planckpla/index.php/Cosmological_Parameters.
- [58] G. Efstathiou and J. R. Bond, *Mon. Not. R. Astron. Soc.* **304**, 75 (1999).
- [59] A. Kosowsky, M. Milosavljevic, and R. Jimenez, *Phys. Rev. D* **66**, 063007 (2002).
- [60] Y. Wang and S. Wang, *Phys. Rev. D* **88**, 043522 (2013); **88**, 069903 (2013).
- [61] C. Contreras, M. Hamuy, M. M. Phillips, G. Folatelli, N. B. Suntzeff, S. E. Persson, M. Stritzinger, L. Boldt, S. González, W. Krzeminski *et al.*, *Astron. J.* **139**, 519 (2010).
- [62] M. Hicken, P. Challis, S. Jha, R. P. Kirshner, T. Matheson, M. Modjaz, A. Rest, W. M. Wood-Vasey, G. Bakos, E. J. Barton *et al.*, *Astrophys. J.* **700**, 331 (2009).
- [63] S. Jha, R. P. Kirshner, P. Challis, P. M. Garnavich, T. Matheson, A. M. Soderberg, G. J. M. Graves, M. Hicken, J. F. Alves, H. G. Arce *et al.*, *Astron. J.* **131**, 527 (2006).
- [64] G. Altavilla, G. Fiorentino, M. Marconi, I. Musella, E. Cappellaro, R. Barbon, S. Benetti, A. Pastorello, M. Rielo, M. Turatto *et al.*, *Mon. Not. R. Astron. Soc.* **349**, 1344 (2004).
- [65] A. G. Riess, R. P. Kirshner, B. P. Schmidt, S. Jha, P. Challis, P. M. Garnavich, A. A. Esin, C. Carpenter, R. Grashius, R. E. Schild *et al.*, *Astron. J.* **117**, 707 (1999).
- [66] A. G. Riess, L.-G. Strolger, S. Casertano, H. C. Ferguson, B. Mobasher, B. Gold, P. J. Challis, A. V. Filippenko, S. Jha, W. Li *et al.*, *Astrophys. J.* **659**, 98 (2007).
- [67] N. Suzuki, D. Rubin, C. Lidman, G. Aldering, R. Amanullah, K. Barbary, L. F. Barrientos, J. Botyanszki, M. Brodwin, N. Connolly *et al.*, *Astrophys. J.* **746**, 85 (2012).
- [68] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith, R. S. Hill, B. Gold, M. Halpern *et al.*, *Astrophys. J. Suppl. Ser.* **208**, 20 (2013).
- [69] F. Melia and A. S. H. Shevchuk, *Mon. Not. R. Astron. Soc.* **419**, 2579 (2012).
- [70] G. E. Addison, G. Hinshaw, and M. Halpern, *Mon. Not. R. Astron. Soc.* **436**, 1674 (2013).
- [71] R. J. Cooke, M. Pettini, R. A. Jorgenson, M. T. Murphy, and C. C. Steidel, *Astrophys. J.* **781**, 31(2014).
- [72] W. L. Freedman and B. F. Madore, *Annu. Rev. Astron. Astrophys.* **48**, 673 (2010).
- [73] A. G. Riess, L. Macri, S. Casertano, H. Lampeitl, H. C. Ferguson, A. V. Filippenko, S. W. Jha, W. Li, and R. Chornock, *Astrophys. J.* **730**, 119 (2011); **732**, 129 (2011).
- [74] W. L. Freedman, B. F. Madore, V. Scowcroft, C. Burns, A. Monson, S. E. Persson, M. Seibert, and J. Rigby, *Astrophys. J.* **758**, 24 (2012).
- [75] E. M. L. Humphreys, M. J. Reid, J. M. Moran, L. J. Greenhill, and A. L. Argon, *Astrophys. J.* **775**, 13 (2013).
- [76] G. Efstathiou, *Mon. Not. R. Astron. Soc.* **440**, 1138 (2014).
- [77] C. Cheng and Q.-G. Huang, *Science China: Physics, Mechanics and Astronomy* **58**, 9 (2015).
- [78] A. Cuesta, L. Verde, A. Riess, and R. Jimenez, *Mon. Not. R. Astron. Soc.* **448**, 3463 (2015).
- [79] A. Heavens, R. Jimenez, and L. Verde, *Phys. Rev. Lett.* **113**, 241302 (2014).
- [80] J. R. Gott, III, M. S. Vogeley, S. Podariu, and B. Ratra, *Astrophys. J.* **549**, 1 (2001).
- [81] G. Chen and B. Ratra, *Publ. Astron. Soc. Pac.* **123**, 1127 (2011).
- [82] H. Lampeitl, R. C. Nichol, H.-J. Seo, T. Giannantonio, C. Shapiro, B. Bassett, W. J. Percival, T. M. Davis, B. Dilday, J. Frieman *et al.*, *Mon. Not. R. Astron. Soc.* **401**, 2331 (2010).
- [83] J. R. Gott and Z. Slepian, *Mon. Not. R. Astron. Soc.* **416**, 907 (2011).
- [84] Z. Slepian, J. R. Gott, and J. Zinn, *Mon. Not. R. Astron. Soc.* **438**, 1948 (2014).
- [85] E. Komatsu, K. M. Smith, J. Dunkley, C. L. Bennett, B. Gold, G. Hinshaw, N. Jarosik, D. Larson, M. R. Nolta, L. Page *et al.*, *Astrophys. J. Suppl. Ser.* **192**, 18 (2011).
- [86] W. J. Percival, W. Sutherland, J. A. Peacock, C. M. Baugh, J. Bland-Hawthorn, T. Bridges, R. Cannon, S. Cole, M. Colless, C. Collins *et al.*, *Mon. Not. R. Astron. Soc.* **337**, 1068 (2002).
- [87] A. Albrecht and C. Skordis, *Phys. Rev. Lett.* **84**, 2076 (2000).
- [88] A. Hebecker and C. Wetterich, *Phys. Lett. B* **497**, 281 (2001).
- [89] M. Doran and G. Robbers, *J. Cosmol. Astropart. Phys.* **06** (2006) 026.
- [90] J. Samsing, E. V. Linder, and T. L. Smith, *Phys. Rev. D* **86**, 123504 (2012).
- [91] A. Hojjati, E. V. Linder, and J. Samsing, *Phys. Rev. Lett.* **111**, 041301 (2013).
- [92] M. S. Turner, *Phys. Rev. D* **31**, 1212 (1985).
- [93] S. DeLope Amigo, W. Man-Yin Cheung, Z. Huang, and S.-P. Ng, *J. Cosmol. Astropart. Phys.* **06** (2009) 005.
- [94] B. Audren, J. Lesgourgues, G. Mangano, P. Dario Serpico, and T. Tram, *J. Cosmol. Astropart. Phys.* **12** (2014) 028.
- [95] G. Blackadder and S. M. Koushiappas, *Phys. Rev. D* **90**, 103527 (2014).
- [96] Y. Gong and X. Chen, *Phys. Rev. D* **77**, 103511 (2008).
- [97] G. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, and A. M. Rotunno, and A. M. Rotunno, *Phys. Rev. D* **86**, 013012 (2012).
- [98] J. Lesgourgues and S. Pastor, *Phys. Rep.* **429**, 307 (2006).
- [99] M. LoVerde and M. Zaldarriaga, *Phys. Rev. D* **89**, 063502 (2014).
- [100] M. LoVerde, *Phys. Rev. D* **90**, 083518 (2014).

- [101] V. Sahni, A. Shafieloo, and A. A. Starobinsky, *Astrophys. J.* **793**, L40 (2014).
- [102] G. J. Hill, K. Gebhardt, E. Komatsu, and P. J. MacQueen, in *The New Cosmology: Conference on Strings and Cosmology*, edited by R. E. Allen, D. V. Nanopoulos, and C. N. Pope, American Institute of Physics Conference Series (American Institute of Physics (AIP), College Park, USA, 2004), Vol. 743, pp. 224–233.
- [103] M. Levi, C. Bebek, T. Beers, R. Blum, R. Cahn, D. Eisenstein, B. Flaugher, K. Honscheid, R. Kron, O. Lahav *et al.*, [arXiv:1308.0847](https://arxiv.org/abs/1308.0847).
- [104] C. Heymans, E. Grocutt, A. Heavens, M. Kilbinger, T. D. Kitching, F. Simpson, J. Benjamin, T. Erben, H. Hildebrandt, H. Hoekstra *et al.*, *Mon. Not. R. Astron. Soc.* **432**, 2433 (2013).
- [105] M. J. Jee, J. A. Tyson, M. D. Schneider, D. Wittman, S. Schmidt, and S. Hilbert, *Astrophys. J.* **765**, 74 (2013).
- [106] R. Mandelbaum, A. Slosar, T. Baldauf, U. Seljak, C. M. Hirata, R. Nakajima, R. Reyes, and R. E. Smith, *Mon. Not. R. Astron. Soc.* **432**, 1544 (2013).
- [107] A. Vikhlinin, A. V. Kravtsov, R. A. Burenin, H. Ebeling, W. R. Forman, A. Hornstrup, C. Jones, S. S. Murray, D. Nagai, H. Quintana *et al.*, *Astrophys. J.* **692**, 1060 (2009).
- [108] E. Rozo, R. H. Wechsler, E. S. Rykoff, J. T. Annis, M. R. Becker, A. E. Evrard, J. A. Frieman, S. M. Hansen, J. Hao, D. E. Johnston *et al.*, *Astrophys. J.* **708**, 645 (2010).
- [109] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday *et al.*, *Astron. Astrophys.* **571**, A20 (2014).
- [110] A. B. Mantz, A. von der Linden, S. W. Allen, D. E. Applegate, P. L. Kelly, R. G. Morris, D. A. Rapetti, R. W. Schmidt, S. Adhikari, M. T. Allen *et al.*, *Mon. Not. R. Astron. Soc.* **446**, 2205 (2015).
- [111] F. Beutler, S. Saito, H.-J. Seo, J. Brinkmann, K. S. Dawson, D. J. Eisenstein, A. Font-Ribera, S. Ho, C. K. McBride, F. Montesano *et al.*, *Mon. Not. R. Astron. Soc.* **443**, 1065 (2014).
- [112] L. Samushia, B. A. Reid, M. White, W. J. Percival, A. J. Cuesta, G.-B. Zhao, A. J. Ross, M. Manera, É. Aubourg, F. Beutler *et al.*, *Mon. Not. R. Astron. Soc.* **439**, 3504 (2014).
- [113] B. A. Reid, H.-J. Seo, A. Leauthaud, J. L. Tinker, and M. White, *Mon. Not. R. Astron. Soc.* **444**, 476 (2014).
- [114] N. Palanque-Delabrouille, C. Yèche, A. Borde, J.-M. Le Goff, G. Rossi, M. Viel, É. Aubourg, S. Bailey, J. Bautista, M. Blomqvist *et al.*, *Astron. Astrophys.* **559**, A85 (2013).
- [115] N. MacCrann, J. Zuntz, S. Bridle, B. Jain, and M. R. Becker, *Mon. Not. R. Astron. Soc.* **451**, 2877 (2015).
- [116] C. Alcock and B. Paczynski, *Nature (London)* **281**, 358 (1979).
- [117] C.-H. Chuang, F. Prada, A. J. Cuesta, D. J. Eisenstein, E. Kazin, N. Padmanabhan, A. G. Sánchez, X. Xu, F. Beutler, M. Manera *et al.*, *Mon. Not. R. Astron. Soc.* **433**, 3559 (2013).
- [118] A. G. Sánchez, E. A. Kazin, F. Beutler, C.-H. Chuang, A. J. Cuesta, D. J. Eisenstein, M. Manera, F. Montesano, R. C. Nichol, N. Padmanabhan *et al.*, *Mon. Not. R. Astron. Soc.* **433**, 1202 (2013).
- [119] C. Blake, S. Brough, M. Colless, C. Contreras, W. Couch, S. Croom, T. Davis, M. J. Drinkwater, K. Forster, D. Gilbank *et al.*, *Mon. Not. R. Astron. Soc.* **415**, 2876 (2011).
- [120] R. A. C. Croft, D. H. Weinberg, M. Bolte, S. Burles, L. Hernquist, N. Katz, D. Kirkman, and D. Tytler, *Astrophys. J.* **581**, 20 (2002).
- [121] P. McDonald, U. Seljak, R. Cen, D. Shih, D. H. Weinberg, S. Burles, D. P. Schneider, D. J. Schlegel, N. A. Bahcall, J. W. Briggs *et al.*, *Astrophys. J.* **635**, 761 (2005).
- [122] W. Hu, [arXiv:astro-ph/0402060](https://arxiv.org/abs/astro-ph/0402060).
- [123] G. Rossi, N. Palanque-Delabrouille, A. Borde, M. Viel, C. Yèche, J. S. Bolton, J. Rich, and J.-M. Le Goff, *Astron. Astrophys.* **567**, A79 (2014).
- [124] E. V. Linder, *Phys. Rev. D* **72**, 043529 (2005).
- [125] S. More, H. Miyatake, R. Mandelbaum, M. Takada, D. Spergel, J. Brownstein, and D. P. Schneider, *Astrophys. J.* **806**, 2 (2015).
- [126] N. Palanque-Delabrouille, C. Yèche, J. Lesgourgues, G. Rossi, A. Borde, M. Viel, E. Aubourg, D. Kirkby, J.-M. LeGoff, J. Rich *et al.*, *J. Cosmol. Astropart. Phys.* **02** (2015) 045.
- [127] D. Weinberg, D. Bard, K. Dawson, O. Dore, J. Frieman, K. Gebhardt, M. Levi, and J. Rhodes, [arXiv:1309.5380](https://arxiv.org/abs/1309.5380).