# Computing finite semigroups

# J. East, A. Egri-Nagy, J. D. Mitchell, and Y. Péresse August 24, 2018

#### Abstract

Using a variant of Schreier's Theorem, and the theory of Green's relations, we show how to reduce the computation of an arbitrary subsemigroup of a finite regular semigroup to that of certain associated subgroups. Examples of semigroups to which these results apply include many important classes: transformation semigroups, partial permutation semigroups and inverse semigroups, partition monoids, matrix semigroups, and subsemigroups of finite regular Rees matrix and 0-matrix semigroups over groups. For any subsemigroup of such a semigroup, it is possible to, among other things, efficiently compute its size and Green's relations, test membership, factorize elements over the generators, find the semigroup generated by the given subsemigroup and any collection of additional elements, calculate the partial order of the  $\mathscr{D}$ -classes, test regularity, and determine the idempotents. This is achieved by representing the given subsemigroup without exhaustively enumerating its elements. It is also possible to compute the Green's classes of an element of such a subsemigroup without determining the global structure of the semigroup.

## Contents

1	Introduction	2								
2	Mathematical prerequisites	4								
3	From transformation semigroups to arbitrary regular semigroups									
	3.1 Actions on Green's classes	6								
	3.2 Faithful representations of stabilisers	7								
	3.3 A decomposition for Green's classes	8								
	3.4 Membership testing	11								
	3.5 Classes within classes	13								
4	Specific classes of semigroups	<b>1</b> 4								
	4.1 Transformation semigroups	15								
	4.2 Partial permutation semigroups and inverse semigroups	16								
	4.3 Matrix semigroups	16								
	4.4 Subsemigroups of a Rees 0-matrix semigroup	17								
	4.5 Partition monoids	19								
5	Algorithms	21								
	5.1 Assumptions	22								
	5.2 Computational prerequisites	23								
	5.3 Components of the action	24								
	5.4 Individual Green's classes	26								
	5.5 The global structure of a semigroup	30								
	5.6 Optimizations for regular and inverse semigroups	34								
6	Examples	36								
7	Experimental results 42									
Λ.	knowlodgoments	45								

# List of Algorithms

1	Compute a component of an action	25				
2	Trace a Schreier tree	25				
3	Compute Schreier generators for a stabiliser	25				
4	Compute Schreier generators for $S_x$	26				
5	Elements of an $\mathscr{R}$ -class	27				
6	$\mathscr{R}$ -classes in a $\mathscr{D}$ -class	28				
7	Test membership in an $\mathscr{R}$ -class	28				
8	Test membership in a $\mathscr{D}$ -class	28				
9	Regularity of an $\mathscr{R}$ -class	29				
10	Idempotents in an $\mathscr{R}$ -class	29				
11	Enumerate the $\mathscr{R}$ -classes of a semigroup	31				
12	Test membership in a semigroup	2				
13	Factorise an element over the generators	3				
14	The partial order of the $\mathscr{D}$ -classes of a semigroup	34				
15	The closure of a semigroup and some elements	5				
List	of Figures					
1	Graphical representations of the partitions $x, y \in P_6$					
2		20				
3	The orbit graph of $(S)\lambda$ with loops and (all but one of the) edges to $\alpha_9$ omitted, and the					
	v e	37				
4		8				
5	Schreier trees for the strongly connected component of $(x_1x_3x_4)\lambda$ in $(S)\lambda$ , and for $(x_1x_3x_4)\rho$					
	\ /1	<b>8</b> 9				
6	The orbit graph of $(T)\rho$ and $\mathfrak{R}$ with loops omitted, and the Schreier tree indicated by solid					
_		ŀ0				
7	The partial order of the $\mathscr{D}$ -classes of $S$ (left) and $T$ (right), group $\mathscr{H}$ -classes indicated by					
0	shaded boxes					
8	1	13				
9	Time in milliseconds to find the size of the full transformation monoid $T_n$					
10	v	14				
11	,, 1					
		4				
	on $\{1,\ldots,n\}$					
12	on $\{1, \ldots, n\}$					

## 1 Introduction

A semigroup is a set with an associative binary operation. There are many articles in the literature concerned with the idea of investigating semigroups using a computer; early examples are [5,6,12,18,34]. There are also several examples of software packages specifically for computing with semigroups, such as AUTOMATE [7], Monoid [22], SgpWin [27], Semigroupe [35], and more general computational algebra systems, such as Magma [4], GAP [15], and Sage [43], with some functionality relating to semigroups; see also [8].

Semigroups are commonly represented either by presentations (abstract generators and defining relations) or by a generating set consisting of a specific type of element, such as transformations, matrices, or binary relations. In this paper, we are solely concerned with semigroups represented by a generating set.

Computing with semigroups defined by generators or with finitely presented semigroups is **hard**; it is shown in [3] that testing membership in a finite commutative transformation semigroup is NP-complete, and it is well-known that determining any 'sensible' property of a finitely presented semigroup is undecidable by the famous results of Post [36] and Markov [26]. However, in spite of the fact that the general case is hard, it is still possible to compute with semigroups efficiently in many particular instances. Perhaps

more importantly, it is possible to perform calculations using a computer that it would be impossible (several times over) to do by hand.

Algorithms, and their implementations, for computing semigroups defined by a generating set fall into two classes: those that exhaustively enumerate the elements, and those that do not. Examples of the first type are the algorithms described in [13] and implemented in Semigroupe [35], and those in SgpWin [27]. Exhaustively enumerating and storing the elements of a semigroup quickly becomes impractical. To illustrate, a transformation is a function from the set  $\{1, \ldots, n\}$  to itself for some  $n \in \mathbb{N}$ . A semigroup whose elements are transformations and whose operation is composition of functions is called a transformation semigroup. For example, if each of the  $9^9 = 387420489$  transformations on a 9-element set is stored as a tuple of 9 integers from 1 to 9 and  $\alpha$  is the number of bits required to store such an integer, then

$$9^{9+1} \cdot \alpha = 3486784401 \cdot \alpha$$
 bits

are required to store these transformations. In GAP, for example, such an integer requires 16-bits, and so approximately 6 gigabytes of memory would be required in this case. Therefore if we want to exhaustively enumerate a semigroup, then we must be happy to do so with relatively small semigroups. Exhaustive algorithms have the advantage that they are relatively straightforward to apply; if the multiplication of a class of semigroups can be defined to a computer, then these algorithms can be applied. For example, in Semigroupe [35] it is possible to compute with semigroups of transformations, partial transformations, and several types of matrix semigroups including boolean matrices.

Non-exhaustive algorithms are described in [19, 20, 23, 24], and the latter were implemented in the Monoid package [22] for GAP 3 and its later incarnations in GAP 4. In examples where it is not possible to store the elements, these methods can be used to determine structural information about a semigroup, such as its size and Green's relations (see Section 2 for the relevant definitions). In many examples, the non-exhaustive algorithms have better performance than their exhaustive analogues. However, on the down side, the non-exhaustive algorithms described in [20,23,24] only apply to transformation semigroups. The methods in [19] are analogues of the methods in [20] in the context of semigroups of binary relations; but an implementation does not appear to be readily available.

To one degree or another, the articles [3, 19, 20, 23, 24] use variants of Schreier's Theorem [46, Theorem 2.57 and the theory of Green's relations to reduce the computation of the semigroup to that of its Schützenberger groups. It is then possible to use the well-developed and efficient algorithms from Computational Group Theory [40, 42, 46], stemming from the Schreier-Sims Algorithm, to compute with these subgroups. In this paper, we go one step further by giving a computational paradigm for arbitrary subsemigroups of finite regular semigroups. Semigroups to which the paradigm can be efficiently applied include many of the most important classes: transformation semigroups, partial permutation semigroups and inverse semigroups, partition monoids, matrix semigroups, and subsemigroups of finite regular Rees matrix and 0-matrix semigroups. We generalise and improve the central notions in [20, 23, 24] from transformation semigroups to arbitrary subsemigroups of an arbitrary finite regular semigroup. For such a subsemigroup, it is possible to efficiently compute its size and Green's classes, test membership, factorize elements over the generators, find the semigroup generated by the given subsemigroup and any collection of additional elements, calculate the partial order of the  $\mathcal{D}$ -classes, test regularity, and determine the idempotents. This is achieved by representing the given subsemigroup without exhaustively enumerating its elements. In particular, our methods can be used to determine properties of semigroups, where it is not possible to store every element of that semigroup. It is also possible to compute the Green's classes of an element of such a subsemigroup without determining the global structure of the semigroup.

Although not described here, it is also possible to use the data structures provided to find the group of units (if it exists), minimal ideal, find a small generating set, and test if a semigroup satisfies various properties such as being simple, completely regular, Clifford and so on. The algorithms described in this paper are implemented in their full generality in the GAP [15] package SEMIGROUPS [29], which is open source software.

The analogue of Cayley's Theorem for semigroups states that every finite semigroup is isomorphic to a transformation semigroup. Consequently, it could be argued that it is sufficient to have computational tools available for transformation semigroups only. An analogous argument could be made for arbitrary groups with respect to permutation groups, but developments in computational group theory suggest otherwise. For example, the Matrix Group Recognition Project has produced efficient algorithms for computing with groups of matrices over finite fields; [1] and [21]. Similarly, for some classes of semigroups, such as subsemigroups of the partition monoid or a Rees matrix semigroup, the only known faithful transformation

representations are those that act on the elements of the semigroup itself. Hence, it is necessary in such examples to exhaustively compute the elements of the semigroup before a transformation representation is available. At this point any transformation representation, and the non-exhaustive methods that could be applied to it, are redundant. Therefore, to compute with such semigroups without exhaustively enumerating them, it is necessary to have non-exhaustive algorithms that apply directly to the given semigroups and, in particular, do not require a transformation representation. It is such algorithms that we present in this paper.

When considering matrix semigroups, it is straightforward to determine a transformation representation. Even in these cases, it is sometimes preferable to compute in the native matrix representation: in particular, where there are methods for matrix groups that are more efficient than computing a permutation representation.

The algorithms in this paper only apply to subsemigroups of a regular semigroup. However, it would be possible to modify several of these algorithms, including the main one (Algorithm 11), so that they apply to subsemigroups S of a non-regular semigroup U. In particular, if it were possible to determine whether a given element was regular or not in U, then we could use the data structure for  $\mathcal{R}$ -classes described in Section 5 for the regular elements, and perform an exhaustive enumeration of  $\mathcal{R}$ -classes of non-regular elements in U. Or alternatively, it might be possible to use a combination of the approaches described in this paper and those in [19,20]. Such an approach would be possible with, say, semigroups of binary relations. It is possible to check that a binary relation is regular as an element of the semigroup of all binary relations in polynomial time; see [10,38]. However, we did not yet follow this approach either in the paper or in the SEMIGROUPS package since it is relatively easy to find a transformation representation of a semigroup of binary relations.

This paper is organised as follows. In Section 2, we recall some well-known mathematical notions, and establish some notation that is required in subsequent sections. In Section 3, we provide the mathematical basis that proves the validity of the algorithms presented in Sections 5. In Section 4, we show that transformation semigroups, partial permutation semigroups and inverse semigroups, partition monoids, semigroups of matrices over a finite field, and subsemigroups of finite regular Rees matrix or 0-matrix semigroups satisfy the conditions from Section 3 and, hence, belong to the class of semigroups with which we can compute efficiently. Detailed algorithms are presented as pseudocode in Section 5, including some remarks about how the main algorithms presented can be simplified in the case of regular and inverse semigroups. In Section 6 we give several detailed examples. The final section, Section 7 is devoted to a discussion of the performance of the algorithms presented herein, and a comparison with the algorithm described in [13].

# 2 Mathematical prerequisites

A semigroup S is regular if for every  $x \in S$  there exists  $x' \in S$  such that xx'x = x. A semigroup S is a monoid if it has an identity element, i.e. an element  $e \in S$  such that es = se = s for all  $s \in S$ . If S is a semigroup, we write  $S^1$  for the monoid obtained by adjoining an identity  $1_S \notin S$  to S. Note that with this definition  $S^1$  always contains one more element  $(1_S)$  than S regardless of whether or not S has an existing identity element.

For any set  $\Omega$ , the set  $\Omega^{\Omega}$  of transformations of  $\Omega$  is a semigroup under composition of functions, known as the *full transformation monoid on*  $\Omega$ . The identity element of  $\Omega^{\Omega}$  is the identity function on  $\Omega$ , which will be denoted  $\mathrm{id}_{\Omega}$ . We denote the full transformation monoid on the finite set  $\{1,\ldots,n\}$  by  $T_n$ . Throughout this article, we will write functions to the right of their arguments and compose functions from left to right.

A subsemigroup T of a semigroup S is just a subset which is a semigroup under the same operation as S; we denote this by  $T \leq S$ . If X is a subset of a semigroup S, then the least subsemigroup of S containing X is denoted by  $\langle X \rangle$ ; this is also referred to as the subsemigroup generated by X. We denote the cardinality of a set X by |X|.

Let S be a semigroup and let  $x, y \in S$  be arbitrary. We say that x and y are  $\mathcal{L}$ -related if the principal left ideals generated by x and y in S are equal; in other words,  $S^1x = S^1y$ . We write  $x\mathcal{L}y$  to denote that x and y are  $\mathcal{L}$ -related. In Section 3, we often want to distinguish between the cases when elements are  $\mathcal{L}$ -related in a semigroup U or in a subsemigroup S of U. We write  $x\mathcal{L}^Sy$  or  $x\mathcal{L}^Uy$  to differentiate these cases.

Green's  $\mathcal{R}$ -relation is defined dually to Green's  $\mathcal{L}$ -relation; Green's  $\mathcal{H}$ -relation is the meet, in the

lattice of equivalence relations on S, of  $\mathscr{L}$  and  $\mathscr{R}$ ; and  $\mathscr{D}$  is the join. We will refer to the equivalence classes as  $\mathscr{K}$ -classes where  $\mathscr{K}$  is any of  $\mathscr{R}$ ,  $\mathscr{L}$ ,  $\mathscr{H}$ , or  $\mathscr{D}$ , and the  $\mathscr{K}$ -class of  $x \in S$  will be denoted by  $K_x$ , or  $K_x^S$  if it is necessary to explicitly refer to the semigroup where the relation is defined. We denote the set of  $\mathscr{K}$ -classes of a semigroup S by  $S/\mathscr{K}$ .

If S is a semigroup and  $x,y\in S$ , then x and y are  $\mathscr{J}$ -related if they generate the same principal two-sided ideal of S, i.e.  $S^1xS^1=S^1yS^1$ . In a finite semigroup,  $x\mathscr{J}y$  if and only if  $x\mathscr{D}y$ . Containment of principal ideals induces a partial order on the  $\mathscr{J}$ -classes of S (and hence  $\mathscr{D}$ -classes if S is finite), sometimes denoted  $\leq_{\mathscr{J}}$ ; that is,  $J_x\leq_{\mathscr{J}}J_y$  if and only if  $S^1xS^1\subseteq S^1yS^1$ .

**Proposition 2.1** (cf. Proposition A.1.16 in [37]). Let U be a semigroup and let S be a subsemigroup of U. Suppose that x and y are regular elements of S. Then  $x\mathcal{K}^U y$  if and only if  $x\mathcal{K}^S y$ , where  $\mathcal{K}$  is any of  $\mathcal{R}$ ,  $\mathcal{L}$  or  $\mathcal{H}$ .

Note that the previous result does not necessarily hold for  $\mathcal{K} = \mathcal{D}$ .

Let S be a semigroup and let  $\Omega$  be a set. A function  $\Psi: \Omega \times S^1 \longrightarrow \Omega$  is a right action of S on  $\Omega$  if

- $((\alpha, s)\Psi, t)\Psi = (\alpha, st)\Psi;$
- $(\alpha, 1_S)\Psi = \alpha$ .

For the sake of brevity, we will write  $\alpha \cdot s$  instead of  $(\alpha, s)\Psi$ , and we will say that S acts on  $\Omega$  on the right. Left actions are defined analogously, and we write  $s \cdot \alpha$  in this case, and say S acts on  $\Omega$  on the left. The kernel of a function  $f: X \longrightarrow Y$ , where X and Y are any sets, is the equivalence relation  $\{(x,y) \in X \times X : (x)f = (y)f\}$ . A right action of a semigroup S on a set  $\Omega$  induces a homomorphism from S to the full transformation monoid on  $\Omega$  defined by mapping  $s \in S$  to the transformation defined by

$$\alpha \mapsto \alpha \cdot s$$
 for all  $\alpha \in \Omega$ .

An action is called *faithful* if the induced homomorphism is injective. The *kernel of a right action* of a semigroup S on a set  $\Omega$  is just the kernel of the induced homomorphism, i.e. the equivalence relation  $\{(s,t) \in S \times S : \alpha \cdot s = \alpha \cdot t \ (\forall \alpha \in \Omega)\}$ . The kernel of a left action is defined analogously.

If S acts on the sets  $\Omega$  and  $\Omega'$  on the right, then we say that  $\lambda:\Omega\longrightarrow\Omega'$  is a homomorphism of right actions if  $(\alpha \cdot s)\lambda=(\alpha)\lambda \cdot s$  for all  $\alpha\in\Omega$  and  $s\in S^1$ . Homomorphisms of left actions are defined analogously. An isomorphism of (left or right) actions is a bijective homomorphism of (left or right) actions.

If  $\Omega$  is a set, then we denote the set of subsets of  $\Omega$  by  $\mathcal{P}(\Omega)$ . If S is a semigroup acting on the right on  $\Omega$ , then the action of S on  $\Omega$  induces a natural action of S on  $\mathcal{P}(\Omega)$ , which we write as:

$$\Sigma \cdot s = \{\alpha \cdot s : \alpha \in \Sigma\} \qquad \text{for each } \Sigma \subseteq \Omega. \tag{2.2}$$

We will denote the function from  $\Sigma$  to  $\Sigma \cdot s$  defined by  $\alpha \mapsto \alpha \cdot s$  by  $s|_{\Sigma}$ . We define the *stabiliser* of  $\Sigma$  under S to be

$$\operatorname{Stab}_{S}(\Sigma) = \{ s \in S^{1} : \Sigma \cdot s = \Sigma \}.$$

The quotient of the stabiliser by the kernel of its action on  $\Sigma$ , i.e. the congruence

$$\{(s,t): s,t \in \operatorname{Stab}_S(\Sigma), \ s|_{\Sigma} = t|_{\Sigma}\},\$$

is isomorphic to

$$S_{\Sigma} = \{s|_{\Sigma} : s \in \operatorname{Stab}_{S}(\Sigma)\}\$$

which in the case that  $\Sigma$  is finite, is a subgroup of the symmetric group  $\operatorname{Sym}(\Sigma)$  on  $\Sigma$ . The stabiliser  $S_{\Sigma}$  can also be seen as a subgroup of  $\operatorname{Sym}(\Omega)$  by extending the action of its elements so that they fix  $\Omega \setminus \Sigma$  pointwise. It is immediate that  $s|_{\Sigma} \cdot t|_{\Sigma} = (st)|_{\Sigma}$  for all  $s, t \in \operatorname{Stab}_{S}(\Sigma)$ .

If S acts on  $\Omega$ , the strongly connected component (usually abbreviated to s.c.c.) of an element  $\alpha \in \Omega$  is the set of all  $\beta \in \Omega$  such that  $\beta = \alpha \cdot s$  and  $\alpha = \beta \cdot t$  for some  $s, t \in S^1$ . We write  $\alpha \sim \beta$  if  $\alpha$  and  $\beta$  belong to the same s.c.c. and the action is clear from the context.

If  $\alpha \in \Omega$ , then

$$\alpha \cdot S^1 = \{\alpha \cdot s : s \in S^1\},\$$

is a disjoint union of strongly connected components of the action of S. Note that  $\alpha \cdot S^1$  might consist of more than one strongly connected component. If S is a group, then  $\alpha \cdot S^1$  has only one strongly connected component, which is usually called the *orbit* of  $\alpha$  under S.

**Proposition 2.3.** Let  $S = \langle X \rangle$  be a semigroup that acts on a finite set  $\Omega$  on the right and let  $\Sigma_1, \ldots, \Sigma_n \subseteq \Omega$  be the elements of a strongly connected component of the action of S on  $\mathcal{P}(\Omega)$ . Then the following hold:

- (a) if  $\Sigma_1 \cdot u_i = \Sigma_i$  for some  $u_i \in S^1$ , then there exists  $\overline{u_i} \in S^1$  such that  $\Sigma_i \cdot \overline{u_i} = \Sigma_1$ ,  $(u_i \overline{u_i})|_{\Sigma_1} = \mathrm{id}_{\Sigma_1}$ , and  $(\overline{u_i}u_i)|_{\Sigma_i} = \mathrm{id}_{\Sigma_i}$ ;
- (b)  $S_{\Sigma_i}$  and  $S_{\Sigma_j}$  are conjugate subgroups of  $Sym(\Omega)$  for all  $i, j \in \{1, ..., n\}$ ;
- (c) if  $u_1 = \overline{u_1} = 1_S$  and  $u_i, \overline{u_i} \in S$  are as in part (a) for i > 1, then  $S_{\Sigma_1}$  is generated by

$$\{(u_i x \overline{u_j})|_{\Sigma_1} : 1 \le i, j \le n, \ x \in X, \ \Sigma_i \cdot x = \Sigma_j\}.$$

*Proof.* Let  $\theta: S \longrightarrow \Omega^{\Omega}$  be the homomorphism induced by the action of S on  $\Omega$ . Then  $(S)\theta$  is a transformation semigroup and the actions of  $(S)\theta$  and S on  $\Omega$  are equal. Hence (a), (b), and (c) are just Lemma 2.2 and Theorems 2.1 and 2.3 in [23], respectively.

We will refer to the generators of  $S_{\Sigma_1}$  in Proposition 2.3(c) as Schreier generators of  $S_{\Sigma_1}$ , due to the similarity of this proposition and Schreier's Theorem [46, Theorem 2.57]. If S is a semigroup acting on a set  $\Omega$ , if  $\Sigma$ ,  $\Gamma \subseteq \Omega$  are such that  $\Sigma \sim \Gamma$  and if  $u \in S$  is any element such that  $\Sigma \cdot u = \Gamma$ , then we write  $\overline{u}$  to denote an element of S with the properties in Proposition 2.3(a).

The analogous definitions can be made, and an analogous version of Proposition 2.3 holds for left actions. In the next section there are several propositions involving left and right actions in the same statement, and so we define the following notation for left actions. If S is a semigroup acting on the left on a set  $\Omega$  and  $\Sigma \subseteq \Omega$ , then the induced left action of S on  $\mathcal{P}(\Omega)$  is defined analogously to (2.2); the function  $\alpha \mapsto s \cdot \alpha$  is denoted by  $\Sigma | s$ ; and we define

$$_{\Sigma}S = \{_{\Sigma}|s: s \cdot \Sigma = \Sigma\}$$

and in the case that  $\Omega$  is finite,  $\Sigma S \leq \operatorname{Sym}(\Omega)$ .

# 3 From transformation semigroups to arbitrary regular semigroups

In this section, we will generalise the results of [23] from transformation semigroups to subsemigroups of an arbitrary finite regular semigroup.

Suppose that U is a regular semigroup whose Green's structure is known in the sense that it is possible to efficiently verify if two elements are  $\mathcal{L}^U$ -,  $\mathcal{R}^U$ -,  $\mathcal{R}^U$ -, or  $\mathcal{D}^U$ -related. For instance, U might be the full transformation monoid or the symmetric inverse monoid on a finite set; the Green's relations of which are described in Propositions 4.1 and 4.3, respectively. In this paper, the central notion is to use our knowledge of the Green's structure of U to produce algorithms to compute any subsemigroup S of U specified by a generating set. Roughly speaking, the subsemigroup S can be decomposed into its  $\mathscr{R}$ -classes, and an  $\mathscr{R}$ -class can be decomposed into the stabiliser  $S_L$  (under right multiplication on  $\mathcal{P}(U)$ ) of an  $\mathscr{L}^U$ -class L and the s.c.c. of L under the action of S on the  $U/\mathscr{L}^U$ . Decomposing S in this way will permit us to efficiently compute many aspects of the structure of S without enumerating its elements exhaustively. For instance, using this decomposition, we can test membership in S, compute the size, Green's structure, idempotents, elements, and maximal subgroups of S.

Throughout the remainder of this section we suppose that U is an arbitrary finite semigroup, and S is a subsemigroup of U.

### 3.1 Actions on Green's classes

We require the following actions of S: the action on  $\mathcal{P}(U)$  induced by multiplying elements of U on the right by  $s \in S^1$ , i.e.:

$$As = \{as : a \in A\} \tag{3.1}$$

where  $s \in S^1$  and  $A \subseteq U$ ; and the action of S on  $U/\mathcal{L}^U$  defined by

$$L_x \cdot s = L_{xs} \tag{3.2}$$

for all  $x \in U$  and  $s \in S^1$ . The latter defines an action since  $\mathcal{L}^U$  is a right congruence.

In general, the actions defined in (3.1) and (3.2) are not equal when restricted to  $U/\mathcal{L}^U$ . For example, it can be the case that  $L_x s \subseteq L_{xs}$  and, in particular,  $L_x s \notin U/\mathcal{L}^U$ . However, the actions do coincide in one case, as described in the next lemma, which is particularly important here.

**Lemma 3.3.** Let  $x, y \in U$  be arbitrary. Then  $L_x, L_y \in U/\mathcal{L}^U$  belong to the same s.c.c. of the action of S defined by (3.2) if and only if  $L_x$  and  $L_y$  belong to the same s.c.c. of the action of S defined by (3.1).

*Proof.* The converse implication is trivial. If  $L_x, L_y \in U/\mathscr{L}^U$  belong to the same strongly connected component under the action (3.2), then, there exists  $s \in S^1$  such that  $L_{xs} = L_y$ . Hence, by Green's Lemma [17, Lemma 2.2.1], the function from  $L_x$  to  $L_{xs} = L_y$  defined by  $z \mapsto zs$  for all  $z \in L_x$  is a bijection and so  $L_x \cdot s = L_{xs} = \{ys : y \in L_x\}$ .

Although S does not, in general, act on the  $\mathscr{L}$ -classes of U by right multiplication, by Lemma 3.3, it does act by right multiplication on the set of  $\mathscr{L}^U$ -classes within a strongly connected component of its action. We will largely be concerned with the strongly connected components of the restriction of the action on  $\mathcal{P}(U)$  in (3.1) to  $U/\mathscr{L}^U$ . In this context, by Lemma 3.3, we may, without ambiguity, use the actions defined in (3.1) and (3.2) interchangeably.

actions defined in (3.1) and (3.2) interchangeably.

If  $x, y \in S$ , then we write  $L_x^U \sim L_y^U$  and  $R_x^U \sim R_y^U$  to denote that these  $\mathcal{L}^U$ - and  $\mathcal{R}^U$ -classes belong to the same strongly connected component of an action defined in (3.1) or (3.2). We will make repeated use of the following lemma later in this section.

**Lemma 3.4.** Let  $x \in S$  and  $s, t \in S^1$  be arbitrary. Then:

- (a)  $L_x^U \sim L_{xs}^U$  if and only if  $x\mathscr{R}^S xs$ ;
- (b)  $R_x^U \sim R_{tx}^U$  if and only if  $x \mathcal{L}^S tx$ ;
- (c)  $L_x^U \sim L_{xs}^U$  and  $R_x^U \sim R_{tx}^U$  together imply that  $x\mathscr{D}^S txs$ .

*Proof.* We prove only parts (a) and (c), since part (b) is the dual of (a).

- (a). ( $\Rightarrow$ ) By assumption,  $L_{xs}^U$  and  $L_x^U$  belong to the same s.c.c. of the action of S on  $U/\mathscr{L}^U$  by right multiplication. Hence, by Proposition 2.3(a), there exists  $\overline{s} \in S^1$  such that  $L_{xs}^U \overline{s} = L_x^U$  and  $s\overline{s}$  acts on  $L_x^U$  as the identity. Hence, in particular,  $xs\overline{s} = x$  and so  $xs\mathscr{R}^S x$ .
- $(\Leftarrow)$  Suppose  $x\mathscr{R}^S xs$ . Then there exists  $t \in S^1$  such that xst = x. It follows that  $L_x^U \cdot s = L_{xs}^U$  and  $L_{xs}^U \cdot t = L_x^U$ . Hence  $L_x^U \sim L_{xs}^U$ .
- (c). Suppose that  $L_x^U \sim L_{xs}^U$  and  $R_x^U \sim R_{tx}^U$ . Then, by parts (a) and (b),  $x\mathscr{R}^S xs$  and  $x\mathscr{L}^S tx$ . The latter implies that  $xs\mathscr{L}^S txs$ , and so  $x\mathscr{D}^S txs$ .

## 3.2 Faithful representations of stabilisers

Let U be an arbitrary finite semigroup, and let S be any subsemigroup of U.

The algorithms presented in this paper involve computing with the groups  $S_L$  arising from the stabiliser  $\operatorname{Stab}_S(L)$  of an  $\mathscr{L}^U$ -class L. It is impractical to compute directly with  $S_L$  or  $U_L$  with their natural actions on L. For example, the  $\mathscr{L}$ -class L of any transformation  $x \in T_{10}$  with 5 points in its image has 5, 103, 000 elements but, in this case,  $U_L$  has a faithful permutation representation on only 5 points.

With the preceding comments in mind, throughout this section we will make statements about arbitrary faithful representations of  $U_L$  or  $S_L$ ,  $L \in U/\mathscr{L}^U$ , rather than about the natural action of  $U_L$  (or  $S_L$ ) on L. More specifically, if  $x \in U$  and  $\zeta$  is any faithful representation of  $U_{L^U}$ , then we define

$$\mu_x : \operatorname{Stab}_U(L_x^U) \longrightarrow (U_{L_x^U})\zeta \quad \text{by} \quad (u)\mu_x = (u|_{L_x^U})\zeta \quad \text{for all } u \in U.$$
 (3.5)

It is clear that  $\mu_x$  is a homomorphism. Since S is a subsemigroup of U, it follows that  $\operatorname{Stab}_S(L_x^U)$  is a subsemigroup of  $\operatorname{Stab}_U(L_x^U)$  and  $S_{L_x^U}$  is a subgroup of  $U_{L_x^U}$ . Hence  $\mu_x : \operatorname{Stab}_U(L_x^U) \longrightarrow (U_{L_x^U})\zeta$  restricted to  $\operatorname{Stab}_S(L_x^U)$  is a homomorphism from  $\operatorname{Stab}_S(L_x^U)$  to  $(S_{L_x^U})\zeta$ .

It is possible that, since  $\zeta$  depends on the  $\mathscr{L}^U$ -class of x in U, we should use  $\zeta_{L_x^U}$  to denote the faithful representation given above. However, to simplify our notation we will not do this. Note that, with this definition,  $(s,t) \in \ker(\mu_x)$  if and only if  $s|_{L_x^U} = t|_{L_x^U}$ . To further simplify our notation, we will write  $S_x$  and  $U_x$  to denote  $(\operatorname{Stab}_S(L_x^U))\mu_x = (S_{L_x^U})\zeta$  and  $(\operatorname{Stab}_U(L_x^U))\mu_x = (U_{L_x^U})\zeta$ , respectively. Analogously, for

every  $x \in U$  we suppose that we have a homomorphism  $\nu_x$  from  $\operatorname{Stab}_U(R_x^U)$  into a group where the image of  $\nu_x$  is isomorphic to  $\nu_x^U$ . We write  $\nu_x^U$  and  $\nu_x^U$  for  $(\operatorname{Stab}_S(R_x^U))\nu_x$  and  $(\operatorname{Stab}_U(R_x^U))\nu_x$ , respectively.

In the case that S is a transformation, partial permutation, matrix, or partition semigroup, or a subsemigroup of a Rees 0-matrix semigroup, we will show in Section 4 how to obtain faithful representations of the stabilisers of  $\mathcal{L}^U$ - and  $\mathcal{R}^U$ -classes as permutation or matrix groups. It is then possible to use algorithms, such as the Schreier-Sims algorithm, from Computational Group Theory to compute with these groups.

We will make repeated use of the following straightforward lemma.

**Lemma 3.6.** Let  $x \in U$  and  $s, t \in Stab_U(L_x^U)$  be arbitrary. Then the following are equivalent:

- (a)  $(s)\mu_x = (t)\mu_x$ ;
- (b)  $ys = yt \text{ for all } y \in L_x^U;$
- (c) there exists  $y \in L_x^U$  such that ys = yt.

*Proof.* (a)  $\Rightarrow$  (b) If  $(s)\mu_x = (t)\mu_x$ , then  $(s|_{L_x^U})\zeta = (t|_{L_x^U})\zeta$  and so  $s|_{L_x^U} = t|_{L_x^U}$ . It follows that ys = yt for all  $y \in L_x^U$ .

- (b)  $\Rightarrow$  (c) is trivial.
- (c)  $\Rightarrow$  (a) Suppose that  $y \in L_x^U$  is such that ys = yt. If  $z \in L_x^U$  is arbitrary, then there exists  $u \in U^1$  such that z = uy. Hence zs = uys = uyt = zt and so  $s|_{L_x^U} = t|_{L_x^U}$ . It follows, by the definition of  $\mu_x$ , that  $(s,t) \in \ker(\mu_x)$  and so  $(s)\mu_x = (t)\mu_x$ .

The analogue of Lemma 3.6 also holds for  $\operatorname{Stab}_U(R_x^U)$  and  $\nu_x$ ; the details are omitted.

## 3.3 A decomposition for Green's classes

In this section, we show how to decompose an  $\mathcal{R}$ - or  $\mathcal{L}$ -class of our subsemigroup S as briefly discussed above.

**Proposition 3.7** (cf. Theorems 3.3 and 4.3 in [23]). If  $x \in S$  is arbitrary, then:

- (a)  $\{L_y^U:y\mathscr{R}^Sx\}$  is a strongly connected component of the right action of S on  $\{L_x^U:x\in S\}$ ;
- (b)  $\{R_y^U: y\mathscr{L}^S x\}$  is a strongly connected component of the left action of S on  $\{R_x^U: x \in S\}$ .

*Proof.* We prove only the first statement, since the second is dual. Suppose  $y \in S$  and  $x \neq y$ . Then  $y\mathscr{R}^S x$  if and only if there exists  $s \in S$  such that xs = y and  $xs\mathscr{R}^S x$ . By Lemma 3.4(a),  $xs\mathscr{R}^S x$  if and only if  $L_x^U \sim L_{xs}^U$ .

**Corollary 3.8.** Let  $x, y, s, t \in S$ . Then the following hold:

- (a) if  $x\mathcal{R}^S y$  and  $xs\mathcal{L}^U y$ , then  $xs\mathcal{R}^S y$ :
- (b) if  $x\mathcal{L}^S y$  and  $tx\mathcal{R}^U y$ , then  $tx\mathcal{L}^S y$ .

*Proof.* Again, we only prove part (a). Since  $xs\mathscr{L}^Uy$ , it follows that  $L^U_{xs} = L^U_y$  and, since  $x\mathscr{R}^Sy$ , by Proposition 3.7(a),  $L^U_x \sim L^U_y = L^U_{xs}$ . Hence  $xs\mathscr{R}^Sx$  by Lemma 3.4(b), and, since  $x\mathscr{R}^Sy$  by assumption, the proof is complete.

**Proposition 3.9** (cf. Theorems 3.7 and 4.6 in [23]). Suppose that  $x \in S$  and there exists  $x' \in U$  where xx'x = x (i.e. x is regular in U). Then the following hold:

- (a)  $L_x^U \cap R_x^S$  is a group under the multiplication \* defined by s \* t = sx't for all  $s, t \in L_x^U \cap R_x^S$  and its identity is x;
- (b)  $\phi: S_x \longrightarrow L_x^U \cap R_x^S$  defined by  $((s)\mu_x)\phi = xs$ , for all  $s \in \operatorname{Stab}_S(L_x^U)$ , is an isomorphism;
- (c)  $\phi^{-1}: L_x^U \cap R_x^S \longrightarrow S_x$  is defined by  $(s)\phi^{-1} = (x's)\mu_x$  for all  $s \in L_x^U \cap R_x^S$ .

*Proof.* We begin by showing that x is an identity under the multiplication \* of  $L_x^U \cap R_x^S$ . Since  $x'x \in L_x^U$  and  $xx' \in R_x^U$  are idempotents, it follows that x'x is a right identity for  $L_x^U$  and xx' is a left identity for  $R_x^S \subseteq R_x^U$ . So, if  $s \in L_x^U \cap R_x^S$  is arbitrary, then

$$x * s = xx's = s = sx'x = s * x,$$

as required.

We will prove that part (b) holds, which implies part (a).

 $\phi$  is well-defined. If  $s \in \operatorname{Stab}_S(L_x^U)$ , then  $xs\mathscr{L}^Ux$ . Hence, by Corollary 3.8(a),  $xs\mathscr{R}^Sx$  and so  $((s)\mu_x)\phi = xs \in L_x^U \cap R_x^S$ . If  $t \in \operatorname{Stab}_S(L_x^U)$  is such that  $(t)\mu_x = (s)\mu_x$ , then, by Lemma 3.6, xt = xs.

 $\phi$  is surjective. Let  $s \in L_x^U \cap R_x^S$  be arbitrary. Then xx's = x \* s = s since x is the identity of  $L_x^U \cap R_x^S$ . It follows that

$$L_x^U \cdot x's = L_s^U = L_x^U$$

and so  $x's \in \operatorname{Stab}_U(L_x^U)$ . Since  $x\mathscr{R}^S s$ , there exists  $u \in S^1$  such that xu = s = xx's. It follows that  $u \in \operatorname{Stab}_S(L_x^U)$  and, by Lemma 3.6,  $(u)\mu_x = (x's)\mu_x$ . Thus  $((u)\mu_x)\phi = xu = s$  and  $\phi$  is surjective.

 $\phi$  is a homomorphism. Let  $s, t \in \operatorname{Stab}_S(L_x^U)$ . Then, since  $xs \in L_x^U$  and x'x is a right identity for  $L_x^U$ ,

$$((s)\mu_x)\phi * ((t)\mu_x)\phi = xs * xt = xsx'xt = xst = ((st)\mu_x)\phi = ((s)\mu_x \cdot (t)\mu_x)\phi,$$

as required.

 $\phi$  is injective. Let  $\theta: L_x^U \cap R_x^S \longrightarrow S_x$  be defined by  $(y)\theta = (x'y)\mu_x$  for all  $y \in L_x^U \cap R_x^S$ . We will show that  $\phi\theta$  is the identity mapping on  $S_x$ , which implies that  $\phi$  is injective, that  $(y)\theta \in S_x$  for all  $y \in L_x^U \cap R_x^S$  (since  $\phi$  is surjective), and also proves part (c) of the proposition. If  $s \in \operatorname{Stab}_S(L_x^U)$ , then  $((s)\mu_x)\phi\theta = (xs)\theta = (x'xs)\mu_x$ . But xx'xs = xs and so  $(x'xs)\mu_x = (s)\mu_x$  by Lemma 3.6. Therefore,  $((s)\mu_x)\phi\theta = (s)\mu_x$ , as required.

We state the analogue of Proposition 3.9 for the action of S on  $U/\mathcal{R}^U$  by left multiplication.

**Proposition 3.10.** Suppose that  $x \in S$  and there exists  $x' \in U$  where xx'x = x. Then the following hold:

- (a)  $L_x^S \cap R_x^U$  is a group under the multiplication \* defined by s \* t = sx't for all  $s, t \in L_x^S \cap R_x^U$  and its identity is x;
- (b)  $\phi: {}_xS \longrightarrow L_x^S \cap R_x^U$  defined by  $((s)\nu_x)\phi = sx$ , for all  $s \in \operatorname{Stab}_S(R_x^U)$ , is an isomorphism;
- (c)  $\phi^{-1}: L_x^S \cap R_x^U \longrightarrow {}_xS$  is defined by  $(s)\phi^{-1} = (sx')\nu_x$  for all  $s \in L_x^S \cap R_x^U$ .

We can also characterise an  $\mathscr{H}^S$ -class H in our subsemigroup S in terms of the stabilisers of the  $\mathscr{L}^U$ - and  $\mathscr{R}^U$ -class containing H. Note that in the special case that S=U, it follows immediately from Proposition 3.9 that  $U_x$  is isomorphic to  $H_x^U = L_x^U \cap R_x^U$  under the operation \* defined in the proposition.

**Proposition 3.11** (cf. Theorem 5.1 in [23]). Suppose that  $x \in S$  and there exists  $x' \in U$  where xx'x = x. Then the following hold:

- (a)  $\Psi: {}_xS \longrightarrow U_x$  defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$ ,  $s \in \operatorname{Stab}_S(R_x^U)$ , is an embedding;
- (b)  $H_x^S$  is a group under the multiplication s\*t = sx't, with identity x, and it is isomorphic to  $S_x \cap (_xS)\Psi$ ;
- (c)  $H_r^S$  under \* is isomorphic to the Schützenberger group of  $H_r^S$ .

*Proof.* (a). Let  $\phi: {}_xS \longrightarrow L^S_x \cap R^U_x$  be the isomorphism defined in Proposition 3.10(b), and let  $\theta: L^U_x \cap R^U_x \longrightarrow U_x$  be the isomorphism defined in Proposition 3.9(c) (applied to U as a subsemigroup of itself). Then, since  $L^S_x \cap R^U_x \subseteq L^U_x \cap R^U_x$ ,  $\phi \theta: {}_xS \longrightarrow U_x$  is a embedding (being the composition of injective homomorphisms). By definition,  $((s)\nu_x)\phi\theta = (x'sx)\mu_x = ((s)\nu_x)\Psi$ , for all  $s \in \operatorname{Stab}_S(R^U_x)$ .

(b). Note that  $H_x^S = L_x^S \cap R_x^S = (L_x^U \cap R_x^S) \cap (R_x^U \cap L_x^S)$ . From Proposition 3.9(c),  $\phi_1^{-1} : R_x^S \cap L_x^U \longrightarrow S_x \leq U_x$  defined by

$$(s)\phi_1^{-1} = (x's)\mu_x$$

is an isomorphism (where  $R_x^S \cap L_x^U$  has multiplication \* defined in Proposition 3.9(a) as s\*t = sx't for all  $s,t \in R_x^S \cap L_x^U$ ). Similarly, by Proposition 3.10(c),  $\phi_2^{-1}: R_x^U \cap L_x^S \longrightarrow {}_xS$  defined by

$$(s)\phi_2^{-1} = (sx')\nu_x$$

is an isomorphism. Hence if  $s \in H_x^S$ , then, by Proposition 3.9(a),

$$(s)\phi_2^{-1}\Psi = (x'sx'x)\mu_x = (x's)\mu_x = (s)\phi_1^{-1}$$

and so  $\phi_1^{-1}$ , restricted to  $H_x^S$ , is an injective homomorphism from  $H_x^S$  under \* into  $S_x \cap (_xS)\Psi$ .

If  $g \in S_x \cap (_xS)\Psi$ , then there exists  $a \in \operatorname{Stab}_S(R_x^U)$  such that  $(x'ax)\mu_x = g$ . Since  $ax\mathcal{R}^Ux$  and xx' is a left identity in  $R_x^U$ , it follows that xx'ax = ax and so  $ax = xx'ax = ((x'ax)\mu_x)\phi_1 \in R_x^S \cap L_x^U \subseteq R_x^S$  where  $\phi_1$  is given in Proposition 3.9(b). Similarly,  $(a)\nu_x \in _xS$  implies that  $ax = ((a)\nu_x)\phi_2 \in R_x^U \cap L_x^S \subseteq L_x^S$ . Therefore  $ax \in H_x^S$  and  $(ax)\phi_1^{-1} = (x'ax)\mu_x = g$ , and so  $\phi_1^{-1}$  is surjective and thus an isomorphism from  $H_x^S$  to  $S_x \cap (_xS)\Psi$ , as required.

(c). The Schützenberger group of  $H = H_x^S$  is defined to be the quotient of  $\operatorname{Stab}_S(H)$  by the kernel of its action on H (by right multiplication on the elements of H). In our notation the Schützenberger group of H is denoted  $S_H$ , however in the literature it is usually denoted by  $\Gamma_R(H)$ . It is well-known that  $S_H$  acts transitively and freely on H, see for example [37, Section A.3.1]. It follows that  $\phi: S_H \longrightarrow H$  defined by

$$(s|_H)\phi = xs$$

is a bijection. If  $s|_H, t|_H \in S_H$ , then  $xs \in H$  and so xsx'x = xs \* x = xs, since x is the identity of H by Proposition 3.9(a). Thus

$$(s|_H \cdot t|_H)\phi = xst = xsx'xt = (s|_H)\phi * (t|_H)\phi$$

and  $\phi$  is an isomorphism, as required.

The statement of Proposition 3.11 can be simplified somewhat in the case that the element  $x \in S$  is regular in S and not only in U.

Corollary 3.12. Suppose that  $x \in S$  and there exists  $x' \in S$  where xx'x = x. Then the following three groups are isomorphic:  $H_x^S$  under the multiplication s \* t = sx't,  $S_x$ , and  $_xS$ . Furthermore,  $\Psi : _xS \longrightarrow S_x$  defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$  is an isomorphism.

*Proof.* Since x is regular in S, it follows that  $H_x^S = L_x^S \cap R_x^S = L_x^U \cap R_x^S$  and, similarly,  $H_x^S = R_x^U \cap L_x^S$ . So, the first part of the statement follows by Propositions 3.9(b) and 3.10(b).

By Propositions 3.10(b) and 3.9(c), respectively, there exist isomorphisms  $\phi: {}_xS \longrightarrow L_x^S \cap R_x^U$  and  $\theta: L_x^U \cap R_x^S \longrightarrow S_x$ . Therefore since  $H_x^S = L_x^U \cap R_x^S = R_x^U \cap L_x^S$ , it follows that  $\Psi = \phi\theta: {}_xS \longrightarrow S_x$  is an isomorphism.

We collect some corollaries of what we have proved so far.

**Corollary 3.13.** If  $x, y \in S$  are regular elements of U, then the following hold:

- (a) If  $x\mathscr{R}^S y$ , then  $S_{L^U_x}$  and  $S_{L^U_y}$  are conjugate subgroups of  $\operatorname{Sym}(U)$ . In particular,  $S_x$  and  $S_y$  are isomorphic;
- (b)  $|R_x^S|$  equals the size of the group  $S_x$  multiplied by the size of the s.c.c. of  $L_x^U$  under the action of S on  $\{L_x^U : x \in S\}$ ;
- (c) If  $L_x^U \sim L_y^U$ , then  $|R_x^S| = |R_y^S|$ ;
- (d) If  $x\mathscr{R}^S y$  and  $u \in S^1$  is such that  $L_x^U \cdot u = L_y^U$ , then the function from  $L_x^U \cap R_x^S$  to  $L_y^U \cap R_x^S$  defined by  $s \mapsto su$  is a bijection.
- *Proof.* (a). Since  $x\mathscr{R}^S y$ , it follows by Proposition 3.7(a) that  $L^U_x$  and  $L^U_y$  are in the same s.c.c. of the action of S on the  $\mathscr{L}$ -classes of U. Thus, by Proposition 2.3(b), it follows that  $S_{L^U_x}$  and  $S_{L^U_y}$  are conjugate subgroups of the symmetric group on U, and so, in particular, are isomorphic.
- (b). The set  $R_x^S$  is partitioned by the sets  $R_x^S \cap L_y^U = R_y^S \cap L_y^U$  for all  $y \in R_x^S$ . By Proposition 3.9(b),  $|R_y^S \cap L_y^U| = |S_y|$  and by part (a),  $|S_y| = |S_x|$  for all  $y \in R_x^S$ . Thus  $|R_x^S|$  equals  $|S_x|$  multiplied by the number of distinct  $\mathcal{L}^U$ -classes of elements of  $R_x^S$ . Proposition 3.7(a) says that  $\{L_y^U : y \in R_x^S\}$  is a s.c.c. of the right action of S on  $\{L_x^U : x \in S\}$ .
  - (c). This follows immediately from parts (a) and (b).
- (d). Let  $s \in L_x^U \cap R_x^S$  be arbitrary. Then  $s\mathscr{R}^S x\mathscr{R}^S y$  and  $su\mathscr{L}^U xu\mathscr{L}^U y$ , and so, by Corollary 3.8(a),  $su\mathscr{R}^S y$ , i.e.  $su \in L_y^U \cap R_x^S$ . By Proposition 2.3(a), there exists  $\overline{u} \in S^1$  such that  $su\overline{u} = s$  for all  $s \in L_x^U \cap R_x^S$ . Therefore  $s \mapsto su$  and  $t \mapsto t\overline{u}$  are mutually inverse bijections from  $L_x^U \cap R_x^S$  to  $L_y^U \cap R_x^S$  and back.  $\square$

For the sake of completeness, we state the analgoue of Corollary 3.13 for  $\mathcal{L}$ -classes.

**Corollary 3.14.** If  $x, y \in S$  are regular elements of U, then the following hold:

- (a) If  $x\mathscr{L}^S y$ , then  $_{R_x^U} S$  and  $_{R_y^U} S$  are conjugate subgroups of  $\operatorname{Sym}(U)$ . In particular,  $_x S$  and  $_y S$  are isomorphic;
- (b)  $|L_x^S|$  equals the size of the group  $_xS$  multiplied by the size of the s.c.c. of  $R_x^U$  under the action of S on  $\{R_x^U: x \in S\}$ ;
- (c) If  $R_x^U \sim R_y^U$ , then  $|L_x^S| = |L_y^S|$ ;
- (d) If  $x\mathscr{L}^S y$  and  $u \in S^1$  is such that  $u \cdot R_x^U = R_y^U$ , then the function from  $R_x^U \cap L_x^S$  to  $R_y^U \cap L_x^S$  defined by  $s \mapsto us$  is a bijection.

## 3.4 Membership testing

Let U be a semigroup and let S be a subsemigroup of U. The next proposition shows that testing membership in an  $\mathcal{R}$ -class of S is equivalent to testing membership in a stabiliser of an  $\mathcal{L}$ -class of U. Since the latter is a group, this reduces the problem of membership testing in  $\mathcal{R}^S$ -classes to that of membership testing in a group, so we can then take advantage of efficient algorithms from computational group theory; such as the Schreier-Sims algorithm [46, Section 4.4].

**Proposition 3.15.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x. If  $y \in U$  is arbitrary, then  $y\mathscr{R}^S x$  if and only if  $y\mathscr{R}^U x$ ,  $L_y^U \sim L_x^U$ , and  $(x'yv)\mu_x \in S_x$  where  $v \in S^1$  is any element such that  $L_y^U \cdot v = L_x^U$ .

Proof. ( $\Rightarrow$ ) Since  $R_x^S \subseteq R_x^U$ ,  $y\mathscr{R}^U x$  and from Proposition 3.7(a),  $L_y^U \sim L_x^U$ . Suppose that  $v \in S^1$  is such that  $L_y^U \cdot v = L_x^U$ . Then, by Corollary 3.13(d),  $yv \in L_x^U \cap R_x^S$  and so, by Proposition 3.9(c),  $(x'yv)\mu_x \in S_x$ .

 $(\Leftarrow)$  Since  $y \in R^U_x$  and xx' is a left identity in its  $\mathscr{R}^U$ -class, it follows that xx'y = y. Suppose that  $v \in S^1$  is any element such that  $L^U_y \cdot v = L^U_x$  (such an element exists by assumption). Then, by assumption,  $(x'yv)\mu_x \in S_x$  and so by Proposition 3.9(b),  $yv = x \cdot x'yv \in L^U_x \cap R^S_x$ . But  $L^U_y \sim L^U_{yv}$ , and so, by Lemma 3.4(a),  $y\mathscr{R}^S yv$ , and so  $x\mathscr{R}^S yv\mathscr{R}^S y$ , as required.

The following corollary of Proposition 3.15 will be important in Section 5.

Corollary 3.16. Let  $x \in S$  be such that there is  $x' \in U$  with xx'x = x, and let  $y \in U$  be such that there exist  $u \in S^1$  and  $v \in U^1$  with  $L_x^U \cdot u = L_y^U$  and xuv = x. Then  $y\mathscr{R}^S x$  if and only if  $y\mathscr{R}^U x$ ,  $L_y^U \sim L_x^U$ , and  $(x'yv)\mu_x \in S_x$ .

Proof. ( $\Rightarrow$ ) That  $y\mathscr{R}^Ux$  and  $L^U_y\sim L^U_x$  follows from Proposition 3.15. Suppose  $u\in S$  and  $v\in U$  are such that  $L^U_x\cdot u=L^U_y$  and xuv=x. By Proposition 2.3(a) there exists  $\overline{u}\in S^1$  such that  $xu\overline{u}=x=xuv$ . It follows that  $z\overline{u}=zv$  for all  $z\in L^S_{xu}$ , and, in particular,  $y\overline{u}=yv$ . Hence, by Proposition 3.15,  $(x'yv)\mu_x=(x'y\overline{u})\mu_x\in S_x$ .

 $(x,y)\mu_x = (x,y)\mu_x \in S_x$ .  $(\Leftarrow)$  It suffices by Proposition 3.15 to show that there exists  $w \in S^1$  such that  $L_y^U \cdot w = L_x^U$  and  $(x'yw)\mu_x \in S_x$ . Since  $L_x^U \sim L_y^U$  and  $L_x^U \cdot u = L_y^U$ , by Proposition 2.3(a) and Lemma 3.6, there exists  $\overline{u} \in S^1$  such that  $xu\overline{u} = x = xuv$ . Hence, as above,  $y\overline{u} = yv$  and so  $(x'y\overline{u})\mu_x = (x'yv)\mu_x \in S_x$ , as required.

We state an analogue of Proposition 3.15 for  $\mathcal{L}$ -classes, with a slight difference.

**Proposition 3.17.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x. If  $y \in U$  is arbitrary, then  $y \mathcal{L}^S x$  if and only if  $y \mathcal{L}^U x$ ,  $R_y^U \sim R_x^U$ , and  $(x'vy)\mu_x \in (_xS)\Psi$  where  $v \in S^1$  is any element such that  $v \cdot R_y^U = R_x^U$  and  $\Psi : _xS \longrightarrow U_x$  defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$  is the embedding from Proposition 3.11(a).

*Proof.* The direct analogue of Proposition 3.15 states that  $y\mathscr{L}^S x$  if and only if  $y\mathscr{L}^U x$ ,  $R_y^U \sim R_x^U$ , and  $(vyx')\nu_x \in {}_xS$ . The last part is equivalent to  $((vyx')\nu_x)\Psi = (x'vyx'x)\mu_x = (x'vy)\mu_x \in ({}_xS)\Psi$ , as required.

Propositions 3.15 and 3.17 allow us to express the elements of an  $\mathcal{R}^{S}$ - and  $\mathcal{L}^{S}$ -class in a particular form, which will be of use in the algorithms later in the paper.

**Corollary 3.18.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x. Then

(a) if  $\mathcal{U}$  is any subset of  $S^1$  such that  $\{L_{xu}^U : u \in \mathcal{U}\} = \{L_y^U : L_y^U \sim L_x^U\}$ , then

$$R_x^S = \{xsu : s \in \operatorname{Stab}_S(L_x^U), u \in \mathcal{U}\};$$

(b) if  $\mathcal V$  is any subset of  $S^1$  such that  $\{R^U_{vx}:v\in\mathcal V\}=\{R^U_y:R^U_y\sim R^U_x\}$ , then

$$L_x^S = \{vtx : t \in \operatorname{Stab}_S(R_x^U), \ v \in \mathcal{V}\}.$$

*Proof.* We prove only part (a), since part (b) is dual.

Let  $s \in \operatorname{Stab}_S(L_x^U)$  and let  $u \in \mathcal{U}$  be arbitrary. Since  $L_{xsu}^U = L_{xu}^U \sim L_x^U$ , it follows, by Lemma 3.4(a), that  $xsu\mathcal{R}^S x$ .

If  $y\mathscr{R}^S x$ , then, by Proposition 3.7,  $L_y^U \sim L_x^U$ , and so there exists  $u \in \mathcal{U}$  such that  $L_x^U \cdot u = L_y^U$ . Since xx' is a left identity for  $R_x^U$ , xx'y=y. By Proposition 2.3(a), there is  $\overline{u}\in S^1$  such that  $L_y^U\cdot \overline{u}=L_x^U$  and  $y\overline{u}u=y$ . Hence, by Proposition 3.15,  $x'y\overline{u}\in\operatorname{Stab}_S(L_x^U)$  and  $y=x\cdot x'y\overline{u}\cdot u$ .

We will prove the analogue of Proposition 3.15 for  $\mathcal{D}$ -classes, for which we require following proposition.

**Proposition 3.19** (cf. Theorem 6.2 in [22]). If  $x \in S$  is such that there is  $x' \in U$  with xx'x = x, then  $D_x^S \cap H_x^U = \{sxt : s \in \operatorname{Stab}_S(R_x^U), \ t \in \operatorname{Stab}_S(L_x^U)\}.$ 

Proof. Let  $s \in \operatorname{Stab}_S(R_x^U)$  and  $t \in \operatorname{Stab}_S(L_x^U)$  be arbitrary. It follows that  $L_{xt}^U = L_x^U$  and  $R_{sx}^U = R_x^U$ , and so, by Lemma 3.4,  $xt\mathscr{R}^S x$ ,  $sx\mathscr{L}^S x$ , and  $x\mathscr{D}^S sxt$ . Also  $xt\mathscr{R}^S x$  implies that  $sxt\mathscr{R}^S sx\mathscr{R}^U x$  and  $sx\mathscr{L}^S x$ 

implies  $sxt\mathcal{L}^Sxt\mathcal{L}^Ux$ , and so  $sxt\in H^U_x$ , as required.

For the other inclusion, let  $y\in D^S_x\cap H^U_x$  be arbitrary. Then  $x\mathcal{D}^Sy$  and so there is  $s\in S^1$  such that  $sx\mathcal{L}^Sx$  and  $sx\mathcal{R}^Sy$ . Hence  $s\cdot R^U_x=R^U_x=R^U_y=R^U_x$  and so  $s\in \mathrm{Stab}_S(R^U_x)$ . Since  $sx\mathcal{R}^Sy$ , there exists  $t\in S^1$  such that sxt=y and so  $L^U_x\cdot t=L^U_{sx}\cdot t=L^U_y=L^U_x$ , which implies that  $t\in \mathrm{Stab}_S(L^U_x)$ , as required.

**Proposition 3.20.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x. If  $y \in U$  is arbitrary, then  $y\mathscr{D}^S x$  if and only if  $L_y^U \sim L_x^U$ ,  $R_y^U \sim R_x^U$ , and for any  $u, v \in S^1$  such that  $L_y^U \cdot u = L_x^U$  and  $v \cdot R_y^U = R_x^U$ there exists  $t \in \operatorname{Stab}_S(L_x^U)$  such that

$$(x'vyu)\mu_x \cdot ((t)\mu_x)^{-1} \in ({}_xS)\Psi,$$

where  $\Psi: {}_xS \longrightarrow U_x$ , defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$ , is the embedding from Proposition 3.11(a).

Proof. ( $\Rightarrow$ ) Let  $y \in D_x^S$ . Then there exists  $w \in S$  such that  $y\mathscr{R}^S w\mathscr{L}^S x$ . By Proposition 3.7(a) and (b), respectively, it follows that  $L_y^U \sim L_w^U = L_x^U$ , and  $R_x^U \sim R_w^U = R_y^U$ . Suppose that  $u, v \in S^1$  are any elements such that  $L_y^U \cdot u = L_x^U$  and  $v \cdot R_y^U = R_x^U$ . Then, by Lemma 3.4,  $vy\mathscr{L}^S y$  and  $yu\mathscr{R}^S y$ , and so  $vyu\mathscr{L}^S yu\mathscr{L}^U x$  and  $vyu\mathscr{R}^S vy\mathscr{R}^U x$ . Thus  $vyu\mathscr{H}^U x$  and, since  $y\mathscr{R}^S yu\mathscr{L}^S vyu$ , it follows that  $vyu\mathcal{D}^S x$ .

By Proposition 3.19, there exist  $s \in \operatorname{Stab}_S(R_x^U)$  and  $t \in \operatorname{Stab}_S(L_x^U)$  such that vyu = sxt. Since  $s \in \operatorname{Stab}_S(R_x^U)$ , it follows that  $((s)\nu_x)\Psi = (x'sx)\mu_x \in (x'sx)\Psi$  (by Proposition 3.11(a)). In particular,  $x'sx \in \operatorname{Stab}_U(L_x^U)$  and so  $(x'vyu)\mu_x = (x'sxt)\mu_x = (x'sx)\mu_x \cdot (t)\mu_x$  and so

$$(x'vyu)\mu_x \cdot ((t)\mu_x)^{-1} = (x'sx)\mu_x \in (_xS)\Psi,$$

as required.

( $\Leftarrow$ ) Suppose that  $u,v\in S^1$  are any elements such that  $L^U_y\cdot u=L^U_x$  and  $v\cdot R^U_y=R^U_x$ . Since  $L^U_y\sim L^U_x=L^U_{yu}$  and  $R^U_y\sim R^U_x=R^U_{vy}$ , it follows from Lemma 3.4(c), that  $y\mathscr{D}^Svyu$ . By assumption, there exist  $s \in \operatorname{Stab}_{S}(R_{x}^{U}), t \in \operatorname{Stab}_{S}(L_{x}^{U})$  such that

$$(x'vyu)\mu_x = ((s)\nu_x)\Psi \cdot (t)\mu_x = (x'sxt)\mu_x.$$

In particular, by Lemma 3.6, xx'vyu = xx'sxt. Since xx' is a left identity for  $R_x^U = R_{vy}^U$ , we deduce that xx'vy = vy. Also, by Proposition 3.19,  $sxt \in D_x^S \cap H_x^U$  implies that  $R_{sxt}^U = R_x^U$  and so xx'sxt = sxt. Thus  $y\mathscr{D}^S vyu = xx'vyu = xx'sxt = sxt\mathscr{D}^S x.$ 

#### 3.5 Classes within classes

The next two propositions allow us to determine the  $\mathscr{R}^S$ -,  $\mathscr{L}^S$ - and  $\mathscr{H}^S$ -classes within a given  $\mathscr{D}^S$ -class in terms of the groups  $S_x$ ,  $_xS$ , and the action of S on  $\{L_x^U: x \in S\}$  and  $\{R_x^U: x \in S\}$ .

If G is a group and H is a subgroup of G, then a *left transversal* of H in G is a set of left coset representatives of H in G. Right transversals are defined analogously.

**Proposition 3.21.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x and that:

- (a) C is a minimal subset of  $\operatorname{Stab}_U(L_x^U)$  such that  $\{(c)\mu_x : c \in C\}$  is a left transversal of  $S_x \cap (xS)\Psi$  in  $(xS)\Psi$  where  $\Psi : xS \longrightarrow U_x$ , defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$ , is the embedding from Proposition 3.11(a);
- (b)  $\{u_1, \ldots, u_m\}$  is a minimal subset of  $S^1$  such that  $\{u_i \cdot R_x^U : 1 \le i \le m\}$  equals the s.c.c. of  $R_x^U$  under the left action of S on  $\{R_x^U : x \in S\}$ .

Then  $\{u_ixc:c\in\mathcal{C},\ 1\leq i\leq m\}$  is a minimal set of  $\mathcal{H}^S$ -class representatives of  $L_x^S$ , and hence a minimal set of  $\mathcal{H}^S$ -class representatives for  $D_x^S$ .

Proof. We start by proving that for all  $c \in \mathcal{C}$ , there exists  $c^* \in \operatorname{Stab}_S(R_x^U)$  such that  $c^*x = xc$  and that  $xc\mathscr{L}^Sx$ . Suppose that  $c \in \mathcal{C}$  is arbitrary. Then  $(c)\mu_x \in (_xS)\Psi$  and so there exists  $c^* \in \operatorname{Stab}_S(R_x^U)$  such that  $((c^*)\nu_x)\Psi = (x'c^*x)\mu_x = (c)\mu_x$ . Hence, by Lemma 3.6,  $xx'c^*x = xc$ . Since  $c^* \in \operatorname{Stab}_S(R_x^U)$ , Proposition 3.10(b) implies that  $c^*x\mathscr{R}^Ux$  and  $c^*x\mathscr{L}^Sx$ , and so  $c^*x = xx'c^*x = xc$ . It follows that  $xc = c^*x\mathscr{L}^Sx$ .

By the assumption in part (b) and by Lemma 3.4(b),  $u_i x \mathcal{L}^S x$  for all i, and so  $u_i x c \mathcal{L}^S x c \mathcal{L}^S x$  for all i. Hence it suffices to show that  $\{u_i x c : c \in \mathcal{C}, 1 \leq i \leq m\}$  is a minimal set of  $\mathcal{R}^S$ -class representatives for  $D_x^S$ . In other words, if  $y \mathcal{D}^S x$ , then we must show that  $y \mathcal{R}^S u_i x c$  for some  $i \in \{1, \ldots, m\}$  and  $c \in \mathcal{C}$ , and that  $(u_i x c, u_j x d) \notin \mathcal{R}^S$  if  $i \neq j$  or  $c \neq d$ .

 $sxg\mathscr{R}^Ssx$ . But  $sxg\mathscr{R}^Ssx$  and so  $y=sxt\mathscr{R}^Ssx\mathscr{R}^Ssxg=sxc$ . If  $y\mathscr{D}^Sx$  is arbitrary, then  $R_y^U\sim R_x^U$ , by Proposition 3.20, and so there exists  $i\in\{1,\ldots,m\}$  such that  $R_y^U=u_i\cdot R_x^U$ . By Proposition 2.3(a), there exists  $\overline{u}_i\in S^1$  such that  $u_i\overline{u}_iy=y$  and  $\overline{u}_i\cdot R_y^U=R_x^U$ . Again by Proposition 3.20,  $L_x^U\sim L_y^U$  and so there exists  $v\in S^1$  such that  $L_y^U\cdot v=L_x^U$ . It follows that  $y\mathscr{D}^S\overline{u}_iyv$  by Lemma 3.4(c). From Lemma 3.4(a), since  $L_y^U\sim L_x^U=L_y^U$ , it follows that  $y\mathscr{R}^Syv$  and so  $R_y^U=R_y^U$ . Thus  $R_{\overline{u}_iyv}^U=\overline{u}_i\cdot R_y^U=\overline{u}_i\cdot R_y^U=R_x^U$  and so  $\overline{u}_iyv\mathscr{R}^Ux$ . Dually, from Lemma 3.4(b),  $\overline{u}_iyv\mathscr{L}^Ux$  and so  $\overline{u}_iyv\in D_x^S\cap H_x^U$ . Hence there exists  $c\in \mathcal{C}$  such that  $\overline{u}_iyv\mathscr{R}^Sxc$  and so  $y\mathscr{R}^Syv=u_i\overline{u}_iyv\mathscr{R}^Su_ixc$ , as required.

Suppose there exist  $i, j \in \{1, 2, ..., m\}$  and  $c, d \in \mathcal{C}$  such that  $u_i x c \mathscr{R}^S u_j x d$ . Then, since  $x c \mathscr{R}^U x \mathscr{R}^U x d$  (from the first paragraph), it follows that  $R_{xc}^U = R_x^U = R_{xd}^U$ . Thus

$$u_i \cdot R_x^U = u_i \cdot R_{xc}^U = R_{u_ixc}^U = R_{u_ixd}^U = u_j \cdot R_{xd}^U = u_j \cdot R_x^U$$

and, by the minimality of  $\{u_1, \ldots, u_m\}$ , it follows that  $u_i = u_j$  and i = j. By the analogue of Proposition 2.3(a), there exists  $\overline{u_i}$  such that  $\overline{u_i}u_ixc = xc$  and  $\overline{u_i}u_ixd = xd$ . Hence since  $u_ixc\mathscr{R}^Su_jxd$  and  $\mathscr{R}^S$  is a left congruence, it follows that  $xc\mathscr{R}^Sxd$ . If xc = xd, then, by Lemma 3.6,  $(c)\mu_x = (d)\mu_x$ , and by the minimality of  $\mathcal{C}$ , c = d. Suppose that  $xc \neq xd$ . Then there exists  $y \in S$  such that xcy = xd. We showed above that  $xc\mathscr{L}^Sx\mathscr{L}^Sxd$ , and so  $x\mathscr{L}^Sxd = xcy\mathscr{L}^Sxy$ , and, in particular,  $y \in \operatorname{Stab}_S(L_x^U)$ . From Lemma 3.6 applied to xcy = xd, we deduce that  $(c)\mu_x(y)\mu_x = (cy)\mu_x = (d)\mu_x$ . This implies that  $((c)\mu_x)^{-1}(d)\mu_x = (y)\mu_x \in S_x$ . Therefore  $(c)\mu_x$  and  $(d)\mu_x$  are representatives of the same left coset of  $S_x \cap (_xS)\Psi$  in  $(_xS)\Psi$ , and again by the minimality of  $\mathcal{C}$ , c = d.

Next, we give an analogue of Proposition 3.21 for  $\mathcal{L}^S$ - and  $\mathcal{H}^S$ -class representatives.

**Proposition 3.22.** Suppose that  $x \in S$  and there is  $x' \in U$  with xx'x = x and that:

- (a) C is a minimal subset of  $\operatorname{Stab}_S(L_x^U)$  such that  $\{(c)\mu_x : c \in C\}$  is a right transversal of  $S_x \cap (_xS)\Psi$  in  $S_x$  where  $\Psi : _xS \longrightarrow U_x$ , defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$ , is the embedding from Proposition 3.11(a);
- (b)  $\{v_1,\ldots,v_m\}$  is a minimal subset of  $S^1$  such that  $\{R_x^U\cdot v_i:1\leq i\leq m\}$  equals the s.c.c. of  $L_x^U$  under the right action of S on  $\{L_x^U:x\in S\}$ .

Then  $\{xcv_i : c \in \mathcal{C}, \ 1 \leq i \leq m\}$  is a minimal set of  $\mathcal{H}^S$ -class representatives of  $R_x^S$ , and hence a minimal set of  $\mathcal{L}^S$ -class representatives for  $D_x^S$ .

*Proof.* It follows from part (b) and Lemma 3.4(a) that  $xcv_i\mathscr{R}^Sxc\mathscr{R}^Sx$ , and so it suffices to show that  $\{xcv_i:c\in\mathcal{C},\ 1\leq i\leq m\}$  is a set of  $\mathscr{L}^S$ -class representatives for  $D^S_x$ . The proof is somewhat similar to that of Proposition 3.21, and so we will omit some details.

If  $y \in D_x^S \cap H_x^U$ , then we will show that there is  $c \in \mathcal{C}$  such that  $y \mathscr{L}^S xc$ . By Proposition 3.19, there exist  $s \in \operatorname{Stab}_S(R_x^U)$  and  $t \in \operatorname{Stab}_S(L_x^U)$  such that y = sxt. As in the proof of Proposition 3.21, it follows that  $sx\mathscr{L}^S x$  and  $sx\mathscr{L}^S x$  and  $sx \in \operatorname{Stab}_S(L_x^U)$  such that  $sx \in \operatorname{L}_x^U \cap R_x^S$  and so  $(s'xt)\mu_x \in S_x$ . If  $s \in \mathcal{C}$  is such that  $(s)\mu_x$  is the representative of the right coset containing  $(s'xt)\mu_x$ , then there exists  $s \in \operatorname{Stab}_S(R_x^U)$  such that  $(s'gx)\mu_x \in S_x \cap (s)\Psi$  and  $(s'gxt)\mu_x = (s'gxx'xt)\mu_x = (s)\mu_x$ . Hence, by Lemma 3.6, sx'gxt = sxt. But  $sx^S x$  and  $sx^S x$  and

The proof that an arbitrary  $y \in D_x^S$  is  $\mathscr{L}^S$ -related to  $xcv_i$  for some i and that  $(xcv_i, xdv_j) \notin \mathscr{L}^S$  if  $i \neq j$  or  $c \neq d$ , is directly analogous to the final part of the proof of Proposition 3.21, and so we omit it

# 4 Specific classes of semigroups

In this section, we show how the results in Section 3 can be efficiently applied to transformation, partial permutation, matrix, and partition semigroups; and also to subsemigroups of finite regular Rees 0-matrix semigroups. More precisely, suppose that U is any of the full transformation monoid, the symmetric inverse monoid, the general linear monoid over any finite field, the partition monoid, or a finite regular Rees 0-matrix semigroup (the definitions of these semigroups can be found below) and that S is any subsemigroup of U.

In general, computing with the action of S on  $U/\mathscr{L}^U$  defined in (3.2) directly has the disadvantage that the points acted on are  $\mathscr{L}^U$ -classes, which are relatively complex to represent on a computer, and for the purposes of checking equality, and hence for determining the value of a point under the action. Instead, we represent each  $\mathscr{L}^U$ -class by a point in some convenient set  $\Omega$  and define an isomorphic action on  $\Omega$ .

Let  $\Omega$  be any set such that  $|\Omega| = |U/\mathcal{L}^U|$  and let  $\lambda: U \longrightarrow \Omega$  be a surjective function such that  $\ker(\lambda) = \mathcal{L}^U$ . We define a right action of S on  $\Omega$  by

$$(x)\lambda \cdot s = (xs)\lambda$$

for all  $x \in U$  and  $s \in S^1$ . This action is well-defined since  $\mathcal{L}^U$  is a right congruence on S. It follows immediately from the definition that  $\lambda$  induces an isomorphism of the action of S on  $U/\mathcal{L}^U$  (defined in (3.2)) and the action of S on  $\Omega$ .

Clearly, any statement about the action of S defined in (3.2) can be replaced with an equivalent statement about the action of S on  $\Omega$ . Furthermore, any statement about the action of S defined in (3.1) within a strongly connected component can be replaced with an equivalent statement about the action of S within a strongly connected component of  $\Omega$ .

As an aide-mémoire, we will write  $(U)\lambda$  or  $(S)\lambda$  instead of  $\Omega$ . We also write  $\rho$  to denote the analogue of  $\lambda$  for left actions. More precisely, we suppose that S has a left action on a set  $(U)\rho$ , the kernel of this action is  $\mathscr{R}^U$ , and there is a homomorphism  $\rho: U \longrightarrow (U)\rho$  of the actions of S on U by left multiplication and of S on  $(U)\rho$ .

In this section, we define the functions  $\lambda:U\longrightarrow (U)\lambda$  and  $\rho:U\longrightarrow (U)\rho$  where U is any of the monoids mentioned above. The functions  $\lambda$  and  $\rho$  are defined so that if S is any subsemigroup of U, then we can compute with the action of S on  $(S)\lambda$  and  $(S)\rho$  efficiently. For such subsemigroups S, we also show how to obtain faithful representations of relatively small degrees of the stabilisers of  $\mathcal{L}^U$ - and  $\mathcal{R}^U$ -classes under the action of S.

## 4.1 Transformation semigroups

Let  $n \in \mathbb{N}$  and write  $\mathbf{n} = \{1, ..., n\}$ . As already stated, a transformation of  $\mathbf{n}$  is a function from  $\mathbf{n}$  to itself, and the full transformation monoid of degree n, denoted  $T_n$ , is the monoid of all transformations on  $\mathbf{n}$  under composition. We refer to subsemigroups of  $T_n$  as transformation semigroups of degree n. It is well-known that the full transformation monoid is regular; see [17, Exercise 2.6.15]. Hence, it is possible to apply the results from Section 3 to any transformation semigroup S.

Let  $f \in T_n$  be arbitrary. Then the *image* of f is defined to be

$$im(f) = \{(i)f : i \in \mathbf{n}\} \subseteq \mathbf{n}$$

and the kernel of f is defined by

$$\ker(f) = \{(i,j) : (i)f = (j)f\} \subseteq \mathbf{n} \times \mathbf{n}.$$

The kernel of a transformation is an equivalence relation, and every equivalence relation on  $\mathbf{n}$  is the kernel of some transformation on  $\mathbf{n}$ . We will denote by  $\mathcal{K}$  the set of all equivalence relations on  $\mathbf{n}$ . The kernel classes of a transformation  $f \in T_n$ , are just the equivalence classes of the equivalence relation  $\ker(f)$ .

The following well-known result characterises the Green's relations on the full transformation monoid.

**Proposition 4.1** (Exercise 2.6.16 in [17].). Let  $n \in \mathbb{N}$  and let  $f, g \in T_n$ . Then the following hold:

- (a)  $f \mathcal{L}^{T_n} g$  if and only if  $\operatorname{im}(f) = \operatorname{im}(g)$ ;
- (b)  $f \mathcal{R}^{T_n} g$  if and only if  $\ker(f) = \ker(g)$ ;
- (c)  $f \mathcal{D}^{T_n} g$  if and only if  $|\operatorname{im}(f)| = |\operatorname{im}(g)|$ .

**Proposition 4.2.** Let S be an arbitrary transformation semigroup of degree  $n \in \mathbb{N}$ . Then:

- (a)  $\lambda: T_n \longrightarrow \mathcal{P}(\mathbf{n})$  defined by  $(x)\lambda = \operatorname{im}(x)$  is a homomorphism of the actions of S on  $T_n$  by right multiplication and the natural action on  $\mathcal{P}(\mathbf{n})$ , and  $\ker(\lambda) = \mathcal{L}^{T_n}$ ;
- (b) if L is any  $\mathcal{L}$ -class of  $T_n$ , then  $S_L$  acts faithfully on  $\operatorname{im}(x)$  for each  $x \in L$ ;
- (c)  $\rho: T_n \longrightarrow \mathcal{K}$  defined by  $(x)\rho = \ker(x)$  is a homomorphism of the actions of S on  $T_n$  by left multiplication and the left action of S on  $\mathcal{K}$  defined by

$$x \cdot K = \ker(xy)$$
 where  $y \in T_n$ ,  $\ker(y) = K$ ,

and  $\ker(\rho) = \mathscr{R}^U$ ;

(d) if R is any  $\mathscr{R}$ -class of  $T_n$ , then  $T_n$  acts faithfully on the set of kernel classes of  $T_n$  for each  $x \in R$ .

*Proof.* We will prove only parts (a) and (b); the proofs of parts (c) and (d) are analogous.

- (a). It follows from Proposition 4.1 that  $\ker(\lambda) = \mathscr{L}^{T_n}$ . If  $x \in T_n$  and  $s \in S$  are arbitrary, then  $(xs)\lambda = \operatorname{im}(xs) = \operatorname{im}(x) \cdot s = (x)\lambda \cdot s$  and so  $\lambda$  is a homomorphism of the actions in part (a).
- (b). Let  $x \in L$  and let  $\zeta : S_L \longrightarrow S_{\operatorname{im}(x)}$  be defined by  $(s|_L)\zeta = s|_{\operatorname{im}(x)}$  where the action of  $s|_{\operatorname{im}(x)}$  on  $\operatorname{im}(x)$  (on the right) is defined by:  $i \cdot (s|_{\operatorname{im}(x)}) = (i)s$ , for all  $i \in \operatorname{im}(x)$ . Let  $s \in \operatorname{Stab}_S(L)$  be arbitrary. Then  $xs \in L$  and so  $\operatorname{im}(xs) = \operatorname{im}(x)$ , and, in particular, s acts on  $\operatorname{im}(x)$ . Thus  $\zeta$  is well-defined. It is routine to verify that  $\zeta$  is a homomorphism. From the definition of  $\zeta$ ,  $s, t \in S$  have the same action on  $\operatorname{im}(x)$  if and only if xs = xt. But, by Lemma 3.6, xs = xt if and only if  $s|_L = t|_L$  and the action of  $s|_L$  on  $\operatorname{im}(x)$  is faithful.

## 4.2 Partial permutation semigroups and inverse semigroups

A partial permutation on  $\mathbf{n} = \{1, \dots, n\}$  is an injective function from a subset of  $\mathbf{n}$  to another subset of equal cardinality. The symmetric inverse monoid of degree n, denoted  $I_n$ , is the monoid of all partial permutations on  $\mathbf{n}$  under composition (as binary relations). We refer to subsemigroups of  $I_n$  as partial permutation semigroups of degree n. A semigroup U is called inverse if for every  $x \in U$  there exists a unique  $y \in U$  such that xyx = x and yxy = y. Every inverse semigroup is isomorphic to an inverse subsemigroup of a symmetric inverse monoid by the Vagner-Preston Theorem; see [17, Theorem 5.1.7]. Since every inverse semigroup is regular, we may apply the results from Section 3 to arbitrary subsemigroups of the symmetric inverse monoid. We will give an analogue of Proposition 4.2 for subsemigroups of any symmetric inverse monoid over a finite set, for which we require a description of the Green's relations in  $I_n$ .

Let  $f \in I_n$  be arbitrary. Then the *domain* of f is defined to be

$$dom(f) = \{i \in \mathbf{n} : (i)f \text{ is defined}\} \subseteq \mathbf{n}$$

and the image of f is

$$im(f) = \{(i)f : i \in dom(f)\} \subseteq \mathbf{n}.$$

The *inverse* of f is the unique partial permutation  $f^{-1}$  with the property that  $ff^{-1}f = f$  and  $f^{-1}ff^{-1} = f^{-1}$ ; note that  $f^{-1}$  coincides with the usual inverse mapping  $\operatorname{im}(f) \longrightarrow \operatorname{dom}(f)$ .

**Proposition 4.3** (Exercise 5.11.2 in [17].). Let  $n \in \mathbb{N}$  and let  $f, g \in I_n$  be arbitrary. Then the following hold:

- (a)  $f \mathcal{L}^{I_n} g$  if and only if  $\operatorname{im}(f) = \operatorname{im}(g)$ ;
- (b)  $f\mathcal{R}^{I_n}g$  if and only if dom(f) = dom(g);
- (c)  $f \mathcal{D}^{I_n} g$  if and only if  $|\operatorname{im}(f)| = |\operatorname{im}(g)|$ .

**Proposition 4.4.** Let S be an arbitrary partial permutation semigroup of degree  $n \in \mathbb{N}$ . Then:

(a)  $\lambda: I_n \longrightarrow \mathcal{P}(\mathbf{n})$  defined by  $(x)\lambda = \operatorname{im}(x)$  is a homomorphism of the actions of S on  $I_n$  by right multiplication, and the right action on  $\mathcal{P}(\mathbf{n})$  defined by

$$A \cdot x = \{(a)x : a \in A \cap dom(x)\}$$
 for  $A \in \mathcal{P}(\mathbf{n})$  and  $x \in I_n$ 

and  $\ker(\lambda) = \mathcal{L}^{I_n}$ :

- (b) if L is any  $\mathcal{L}$ -class of  $I_n$ , then  $(I_n)_L$  acts faithfully on the right of  $\operatorname{im}(x)$  for each  $x \in L$ ;
- (c)  $\rho: I_n \longrightarrow \mathcal{P}(\mathbf{n})$  defined by  $(x)\rho = \operatorname{dom}(x)$  is a homomorphism of the actions of S on  $I_n$  by left multiplication, and the left action on  $\mathcal{P}(\mathbf{n})$  defined by

$$x \cdot A = \{(a)x^{-1} : a \in A \cap \operatorname{im}(x)\}$$
 for  $A \in \mathcal{P}(\mathbf{n})$  and  $x \in I_n$ ;

and  $\ker(\rho) = \mathcal{R}^{I_n}$ ;

(d) if R is any  $\mathscr{R}$ -class of  $I_n$ , then  ${}_RS$  acts faithfully on the left of dom(x) for each  $x \in R$ .

*Proof.* The proof of this proposition is very similar to that of Proposition 4.2 and is omitted.  $\Box$ 

## 4.3 Matrix semigroups

Let R be a finite field, let  $n \in \mathbb{N}$ , and let  $M_n(R)$  denote the monoid of  $n \times n$  matrices with entries in R (under the usual matrix multiplication). The monoid  $M_n(R)$  is called a *general linear monoid*. In this paper, a *matrix semigroup* is a subsemigroup of some general linear monoid. It is well-known that  $M_n(R)$  is a regular semigroup [33, Lemma 2.1].

If  $\alpha \in M_n(R)$  is arbitrary, then denote by  $r(\alpha)$  the row space of  $\alpha$  (i.e. the subspace of the n-dimensional vector space over R spanned by the rows of  $\alpha$ ). We denote the dimension of  $r(\alpha)$  by  $\dim(r(\alpha))$ . The notion of a column space and its dimension are defined dually. We denote the column space of  $\alpha \in M_n(R)$  by  $c(\alpha)$ .

**Proposition 4.5** (Lemma 2.1 in [33]). Let R be a finite field, let  $n \in \mathbb{N}$ , and let  $\alpha, \beta \in M_n(R)$  be arbitrary. Then the following hold:

- (a)  $\alpha \mathcal{L}^{M_n(R)} \beta$  if and only if  $r(\alpha) = r(\beta)$ ;
- (b)  $\alpha \mathcal{R}^{M_n(R)} \beta$  if and only if  $c(\alpha) = c(\beta)$ :
- (c)  $\alpha \mathcal{D}^{M_n(R)} \beta$  if and only if  $\dim(r(\alpha)) = \dim(r(\beta))$ .

**Proposition 4.6.** Let R be a finite field, let  $n \in \mathbb{N}$ , and let S be an arbitrary subsemigroup of the general linear monoid  $M_n(R)$ . Then the following hold:

- (a) if  $\Omega$  denotes the collection of subspaces of  $R^n$  as row vectors, then  $\lambda : M_n(R) \longrightarrow \Omega$  defined by  $(\alpha)\lambda = r(\alpha)$  is a homomorphism of the actions of S on  $M_n(R)$  by right multiplication, and the action on  $\Omega$  by right multiplication, and  $\ker(\lambda) = \mathcal{L}^{M_n(R)}$ ;
- (b) if L is any  $\mathscr{L}$ -class of  $M_n(R)$ , then  $S_L$  acts faithfully on  $r(\alpha)$  for each  $\alpha \in L$ ;
- (c) if  $\Omega$  denotes the collection of subspaces of  $R^n$  as column vectors, then  $\rho: M_n(R) \longrightarrow \Omega$  defined by  $(\alpha)\rho = c(\alpha)$  is a homomorphism of the actions of S on  $M_n(R)$  by left multiplication, and the action on  $\Omega$  by left multiplication, and  $\ker(\rho) = \mathscr{R}^{M_n(R)}$ ;
- (d) if R is any  $\mathscr{R}$ -class of  $M_n(R)$ , then  ${}_RS$  acts faithfully on  $c(\alpha)$  for each  $\alpha \in R$ .

*Proof.* We will prove only parts (a) and (b); the proofs of parts (c) and (d) follow by analogous arguments. We will write  $L_{\alpha}$  to mean the  $\mathscr{L}$ -class of  $\alpha \in M_n(R)$  in  $M_n(R)$  throughout this proof.

(a). It follows from Proposition 4.5(a) that  $(\alpha)\lambda = (\beta)\lambda$  if and only if  $\alpha \mathcal{L}^{M_n(R)}\beta$ , and so  $\ker(\lambda) = \mathcal{L}^{M_n(R)}$ . We also have

$$(\alpha\beta)\lambda = r(\alpha\beta) = r(\alpha) \cdot \beta = (\alpha)\lambda \cdot \beta$$

for all  $\alpha \in M_n(R)$  and  $\beta \in S$ , and so  $\lambda$  is a homomorphism of actions.

(b). Let L be any  $\mathscr{L}$ -class in  $M_n(R)$ , let  $\alpha \in L$  and let  $\beta, \gamma \in \operatorname{Stab}_S(L)$ . Then  $\alpha\beta \in L$  and so  $\beta$  acts on  $r(\alpha)$  by right multiplication. By Lemma 3.6, it follows that  $\beta$  and  $\gamma$  have equal action on  $r(\alpha)$  if and only if  $\alpha\beta = \alpha\gamma$  if and only if  $\alpha|_L = \beta|_L$ .

## 4.4 Subsemigroups of a Rees 0-matrix semigroup

In this section, we describe how the results from Section 3 can be applied to subsemigroups of a Rees 0-matrix semigroup. We start by recalling the relevant definitions.

Let T be a semigroup, let 0 be an element not in T, let I and J be sets, and let  $P = (p_{j,i})_{j \in J, i \in I}$  be a  $|J| \times |I|$  matrix with entries from  $T \cup \{0\}$ . Then the Rees 0-matrix semigroup  $\mathcal{M}^0[T; I, J; P]$  is the set  $(I \times T \times J) \cup \{0\}$  with multiplication defined by

$$0x = x0 = 0 \text{ for all } x \in \mathcal{M}^0[T; I, J; P] \qquad \text{and} \qquad (i, g, j)(k, h, l) = \begin{cases} (i, gp_{j,k}h, l) & \text{if } p_{j,k} \neq 0 \\ 0 & \text{if } p_{j,k} = 0. \end{cases}$$

A semigroup U with a zero element 0 is 0-simple if U and  $\{0\}$  are its only ideals.

**Theorem 4.7** (Theorem 3.2.3 in [17] or Theorem A.4.15 in [37]). A finite semigroup U is 0-simple if and only if it is isomorphic to a Rees 0-matrix semigroup  $\mathcal{M}^0[G;I,J;P]$ , where G is a group, and P is regular, in the sense that every row and every column contains at least one non-zero entry.

Green's relations of a regular Rees 0-matrix semigroup are described in the following proposition.

**Proposition 4.8.** Let  $U = \mathcal{M}^0[G; I, J; P]$  be a finite Rees 0-matrix semigroup where G is a group and P is regular. Then the following hold for all  $x, y \in U$ :

- (a)  $x\mathcal{L}^U y$  if and only if  $x, y \in I \times G \times \{j\}$  for some  $j \in J$  or x = y = 0.
- (b)  $x\mathscr{R}^U y$  if and only if  $x, y \in \{i\} \times G \times J$  for some  $i \in I$  or x = y = 0;

Obviously, we do not require any theory beyond that given above to compute with Rees 0-matrix semigroups, since their size, elements, and in the case that they are regular, their Green's structure and maximal subgroups too, are part of their definition. However, it might be that we would like to compute with a proper subsemigroup of a Rees 0-matrix semigroup. Several computational problems for arbitrary finite semigroups can be reduced, in part, to problems for associated Rees 0-matrix semigroups (the principal factors of certain  $\mathscr{D}$ -classes). For example, this is the case for finding the automorphism group [2], minimal (idempotent) generating sets [9,16], or the maximal subsemigroups of a finite semigroup. In the latter example, we may wish to determine the structure of the maximal subsemigroups, which are not necessarily Rees 0-matrix semigroups themselves. In the absence of a method to find a convenient representation of a subsemigroup of a Rees 0-matrix semigroup, as, for example, a transformation semigroup, we would have to compute directly with the subsemigroup.

**Proposition 4.9.** Let S be an arbitrary subsemigroup of a finite regular Rees 0-matrix semigroup  $U = \mathcal{M}^0[G; I, J; P]$  over a permutation group G acting faithfully on  $\mathbf{n}$  for some  $n \in \mathbb{N}$ . Then:

(a)  $\lambda: U \longrightarrow J \cup \{0\}$  defined by  $(i, g, j)\lambda = j$  and  $(0)\lambda = 0$  is a homomorphism of the actions of S on U by right multiplication, and the right action of S on  $J \cup \{0\}$  defined by

$$0 \cdot (i, g, j) = 0 \cdot 0 = 0 = k \cdot 0 \quad and \quad k \cdot (i, g, j) = \begin{cases} j & \text{if } p_{k,i} \neq 0 \\ 0 & \text{if } p_{k,i} = 0 \end{cases}$$

for all  $k \in J$ , and  $\ker(\lambda) = \mathcal{L}^U$ ;

(b) if L is any non-zero  $\mathscr{L}$ -class of U, then the action of  $S_L$  on  $\mathbf n$  defined by

$$m \cdot (i, g, j)|_{L} = (m)p_{j,i}g$$
 for all  $m \in \mathbf{n}$ 

 $is\ faithful;$ 

(c)  $\rho: U \longrightarrow I \cup \{0\}$  defined by  $(i, g, j)\rho = i$  and  $(0)\rho = 0$  is a homomorphism of the actions of S on U by left multiplication, and the left action of S on  $I \cup \{0\}$  defined by

$$(i, g, j) \cdot 0 = 0 \cdot 0 = 0 = 0 \cdot k$$
 and  $(i, g, j) \cdot k = \begin{cases} i & \text{if } p_{j,k} \neq 0 \\ 0 & \text{if } p_{j,k} = 0 \end{cases}$ 

for all  $k \in I$ , and  $\ker(\rho) = \mathscr{R}^U$ ;

(d) if R is any non-zero  $\mathcal{R}$ -class of U, then the action of  $_RS$  on  $\mathbf n$  defined by

$$_{R}|(i,g,j)\cdot m=(m)g^{-1}p_{j,i}^{-1}\quad \textit{ for all }\quad m\in\mathbf{n}$$

is faithful.

Proof. We prove only parts (a) and (b); parts (c) and (d) follow by analogous arguments.

(a). It follows by Proposition 4.8(a) that  $(x)\lambda = (y)\lambda$  if and only if  $x\mathcal{L}^U y$ , for each  $x, y \in U$ , and so the kernel of  $\lambda$  is  $\mathcal{L}^U$ . We will show that  $\lambda$  is a homomorphism of actions.

Let  $x \in U$  and  $s \in S$  be arbitrary. We must show that  $(xs)\lambda = (x)\lambda \cdot s$ . If x = 0 or s = 0, then  $(xs)\lambda = (0)\lambda = 0 = (x)\lambda \cdot s$ . Suppose that  $x = (i, g, j) \in U \setminus \{0\}$  and  $s = (k, h, l) \in S \setminus \{0\}$ . If  $p_{j,k} = 0$ , then xs = 0 and so

$$(xs)\lambda = (0)\lambda = 0 = j \cdot (k, h, l) = (x)\lambda \cdot s.$$

If  $p_{j,k} \neq 0$ , then

$$(xs)\lambda = l = j \cdot (k, h, l) = (x)\lambda \cdot s.$$

(b). Let  $x = (i, g, j) \in U \setminus \{0\}$  and let  $L = L_x^U = \{(i', g', j) : i' \in I, g' \in G\}$ . If  $(k, h, l) \in \operatorname{Stab}_S(L)$  is arbitrary, then, since  $L \cdot (k, h, l) = L$ , it follows that  $p_{j,k} \neq 0$  and l = j. It follows that we may define a mapping  $\zeta : S_L \longrightarrow G$  by  $((k, h, l)|_L)\zeta = p_{j,k}h$ .

If  $(k_1, h_1, j), (k_2, h_2, j) \in \operatorname{Stab}_S(L)$ , then, by Lemma 3.6, it follows that

$$\begin{split} (k_1,h_1,j)|_L &= (k_2,h_2,j)|_L & \quad \text{if and only if} & \quad (i,g,j)(k_1,h_1,j) = (i,g,j)(k_2,h_2,j) \\ & \quad \text{if and only if} & \quad (i,gp_{j,k_1}h_1,j) = (i,gp_{j,k_2}h_2,j) \\ & \quad \text{if and only if} & \quad p_{j,k_1}h_1 = p_{j,k_2}h_2. \end{split}$$

Hence,  $\zeta$  is well-defined and injective.

To show that  $\zeta$  is a homomorphism, suppose that  $(k_1, h_1, j), (k_2, h_2, j) \in \operatorname{Stab}_S(L)$ . Then

$$\begin{aligned} ((k_1,h_1,j)|_L(k_2,h_2,j)|_L)\zeta &=& (((k_1,h_1,j)(k_2,h_2,j))|_L)\zeta \\ &=& ((k_1,h_1p_{j,k_2}h_2,j)|_L)\zeta \\ &=& p_{j,k_1}h_1p_{j,k_2}h_2 \\ &=& ((k_1,h_1,j)|_L)\zeta \cdot ((k_2,h_2,j)|_L)\zeta, \end{aligned}$$

as required.

#### 4.5 Partition monoids

Let  $n \in \mathbb{N}$ , let  $\mathbf{n} = \{1, \dots, n\}$ , and let  $-\mathbf{n} = \{-1, \dots, -n\}$ . A partition of  $\mathbf{n} \cup -\mathbf{n}$  is a set of pairwise disjoint non-empty subsets of  $\mathbf{n} \cup -\mathbf{n}$  (called *blocks*) whose union is  $\mathbf{n} \cup -\mathbf{n}$ . If  $i, j \in \mathbf{n} \cup -\mathbf{n}$  belong to the same block of a partition x, then we write  $(i, j) \in x$ .

If x and y are partitions of  $\mathbf{n} \cup -\mathbf{n}$ , then we define the product xy of x and y to be the partition where for  $i, j \in \mathbf{n}$ 

(i)  $(i,j) \in xy$  if and only if  $(i,j) \in x$  or there exist  $a_1, \ldots, a_{2r} \in \mathbf{n}$ , for some  $r \geq 1$ , such that

$$(i, -a_1) \in x$$
,  $(a_1, a_2) \in y$ ,  $(-a_2, -a_3) \in x$ , ...,  $(a_{2r-1}, a_{2r}) \in y$ ,  $(-a_{2r}, j) \in x$ 

(ii)  $(i, -j) \in xy$  if and only if there exist  $a_1, \ldots, a_{2r-1} \in \mathbf{n}$ , for some  $r \geq 1$ , such that

$$(i, -a_1) \in x$$
,  $(a_1, a_2) \in y$ ,  $(-a_2, -a_3) \in x$ , ...,  $(-a_{2r-2}, -a_{2r-1}) \in x$ ,  $(a_{2r-1}, -j) \in y$ 

(iii)  $(-i, -j) \in xy$  if and only if  $(-i, -j) \in y$  or there exist  $a_1, \ldots, a_{2r} \in \mathbf{n}$ , for some  $r \ge 1$ , such that

$$(-i, a_1) \in y$$
,  $(-a_1, -a_2) \in x$ ,  $(a_2, a_3) \in y$ , ...,  $(-a_{2r-1}, -a_{2r}) \in x$ ,  $(a_{2r}, -j) \in y$ 

for  $i, j \in \mathbf{n}$ .

This product can be shown to be associative, and so the collection of partitions of  $\mathbf{n} \cup -\mathbf{n}$  is a monoid; the identity element is the partition  $\{\{i, -i\} : i \in \mathbf{n}\}$ . This monoid is called the *partition monoid* and is denoted  $P_n$ .

It can be useful to represent a partition as a graph with vertices  $\mathbf{n} \cup -\mathbf{n}$  and the minimum number of edges so that the connected components of the graph correspond to the blocks of the partition. Of course, such a representation is not unique in general. An example is given in Figure 1 for the partitions:

$$x = \{\{1, -1\}, \{2\}, \{3\}, \{4, -3\}, \{5, 6, -5, -6\}, \{-2, -4\}\}\}$$

$$y = \{\{1, 4, -1, -2, -6\}, \{2, 3, 5, -4\}, \{6, -3\}, \{-5\}\}$$

and the product

$$xy = \big\{\{1,4,5,6,-1,-2,-3,-4,-6\},\{2\},\{3\},\{-5\}\big\}$$

is shown in Figure 2.

A block of a partition containing elements of both  $\mathbf{n}$  and  $-\mathbf{n}$  is called a transverse block. If  $x \in P_n$ , then we define  $x^*$  to be the partition obtained from x by replacing i by -i and -i by i in every block of x for all  $i \in \mathbf{n}$ . It is routine to verify that if  $x, y \in P_n$ , then

$$(x^*)^* = x$$
,  $xx^*x = x$ ,  $x^*xx^* = x^*$ ,  $(xy)^* = y^*x^*$ .

In this way, the partition monoid is a regular \*-semigroup in the sense of [32].




Figure 1: Graphical representations of the partitions  $x, y \in P_6$ .




Figure 2: A graphical representation of the product  $xy \in P_6$ .

If  $x \in P_n$  is arbitrary, then  $xx^*$  and  $x^*x$  are idempotents; called the *projections* of x. We will write

$$Proj(P_n) = \{xx^* : x \in P_n\} = \{x^*x : x \in P_n\}.$$

If B is a transverse block of  $xx^*$  (or  $x^*x$ ), then  $i \in B$  if and only if  $-i \in B$ . If B is a non-transverse block of  $xx^*$ , then  $-B = \{-b : b \in B\}$  is also a block of  $xx^*$ .

**Proposition 4.10** (cf. [11,45]). Let  $n \in \mathbb{N}$  and let  $x, y \in P_n$ . Then the following hold:

- (a)  $x\mathcal{L}^{P_n}y$  if and only if  $x^*x = y^*y$ ;
- (b)  $x\mathscr{R}^{P_n}y$  if and only if  $xx^* = yy^*$ ;
- (c)  $x\mathcal{D}^{P_n}y$  if and only if x and y have the same number of transverse blocks.

The characterisation in Proposition 4.10 can be used to define representations of the actions mentioned above.

**Proposition 4.11.** Let S be an arbitrary subsemigroup of  $P_n$ . Then:

(a)  $\lambda: P_n \longrightarrow \operatorname{Proj}(P_n)$  defined by  $(x)\lambda = x^*x$  is a homomorphism between the action of S on  $P_n$  by right multiplication and the right action of S on  $\operatorname{Proj}(P_n)$  defined by

$$x^*x \cdot y = (xy)^*xy = y^*x^*xy$$

and the kernel of  $\lambda$  is  $\mathcal{L}^{P_n}$ ;

- (b) if L is any  $\mathscr{L}$ -class of  $P_n$ , then  $S_L$  acts faithfully on the transverse blocks of  $x^*x$  for each  $x \in L$ ;
- (c)  $\rho: P_n \longrightarrow \operatorname{Proj}(P_n)$  defined by  $(x)\rho = xx^*$  is a homomorphism between the action of S on  $P_n$  by left multiplication and the left action of S on  $\operatorname{Proj}(P_n)$  defined by

$$y \cdot xx^* = yx(yx)^* = yxx^*y^*$$

and the kernel of  $\rho$  is  $\mathscr{R}^{P_n}$ ;

(d) if R is any  $\mathscr{R}$ -class of  $P_n$ , then RS acts faithfully on the transverse blocks of  $xx^*$  for each  $x \in R$ .

*Proof.* We prove only parts (a) and (b), since the other parts are dual.

(a). Let  $x \in P_n$  and  $s \in S$  be arbitrary. Then

$$(xs)\lambda = (xs)^*xs = s^*(x^*x)s = (x)\lambda \cdot s.$$

Together with Proposition 4.10, this completes the proof of (a).

(b). Let  $x \in P_n$  be arbitrary and suppose that  $y \in \operatorname{Stab}_S(L_x^{P_n})$ . It follows that  $x^*x = (x)\lambda = (xy)^*xy = y^*x^*xy$ . We denote the intersection of the transverse blocks of  $x^*x$  with  $\mathbf{n}$  by  $B_1, \ldots, B_r$  and we define the binary relation

$$p_y = \{(i,j) \in \mathbf{r} \times \mathbf{r} : \exists k \in B_i, \exists l \in B_j, (k,-l) \in x^*xy\}.$$

We will show that  $p_y$  is a permutation, and that  $\zeta: S_{L_x^{P_n}} \longrightarrow \operatorname{Sym}(\mathbf{r})$  defined by  $(y|_{L_x^{P_n}})\zeta = p_y$  is a monomorphism.

Seeking a contradiction, assume that there exist  $i, j, j' \in \mathbf{r}$  such that  $j \neq j'$  and  $(i, j), (i, j') \in p_y$ . Then there exist  $k, k' \in B_i$ ,  $l \in B_j$ , and  $l' \in B_{j'}$  such that  $(k, -l), (k', -l') \in x^*xy$ . Since  $k, k' \in B_i$ , it follows that  $(k, k') \in x^*x$  and so  $(k, k') \in x^*xy$ . Since  $x^*xy$  is an equivalence relation, it follows that  $(-l, -l') \in x^*xy$ , which implies that  $(-l, -l') \in y^*x^*xy = x^*x$ , and so  $(l, l') \in x^*x$ , a contradiction. Hence  $p_y$  is a function.

Since **n** is finite, to show that  $p_y$  is a permutation it suffices to show that it is surjective. Suppose that  $i \in \mathbf{r}$  and  $l \in B_i$  are arbitrary. Then  $(l, -l) \in x^*x = y^*x^*xy$ , and so by part (ii) of the definition of the multiplication of the partitions  $y^*$  and  $x^*xy$ , there exists  $k \in \mathbf{n}$  such that  $(k, -l) \in x^*xy$ . In other words, k belongs to a transverse block of  $x^*xy$ , and hence to a transverse block of  $x^*x$ . Thus there exists  $j \in \mathbf{r}$  such that  $k \in B_j$  and so  $(j)p_y = i$ , as required.

Note that, since  $x^*xy$  is an equivalence relation and  $p_y$  is a permutation, the transverse blocks of  $x^*xy$  are of the form  $B_i \times -B_{(i)p_y}$ .

Next, we show that  $\zeta$  :  $S_{L_x^{P_n}} \longrightarrow \operatorname{Sym}(\mathbf{r})$  defined by  $(y|_{L_x^{P_n}})\zeta = p_y$  is a homomorphism. Let  $y,z \in \operatorname{Stab}_S(L_x^{P_n})$ . It suffices to prove that  $p_yp_z = p_{yz}$ . If  $i \in \mathbf{r}$ , then there exist  $k \in B_i$ ,  $l,l' \in B_{(i)p_y}$ , and  $m \in B_{((i)p_y)p_z}$  such that  $(k,-l) \in x^*xy$  and  $(l',-m) \in x^*xz$ . Since  $l,l' \in B_{(i)p_y}$  (a transverse block of  $x^*x$ ),  $(l,-l') \in x^*x$ , and so  $(k,-m) \in x^*xy \cdot x^*x \cdot x^*xz = x^*xyx^*xz$ . Since  $xy\mathscr{L}^{P_n}x$  and  $x^*x$  is a right identity in its  $\mathscr{L}^{P_n}$ -class, it follows that  $xyx^*x = xy$  and so  $x^*xyx^*xz = x^*xyz$ . In particular,  $(k,-m) \in x^*xyz$ , and so  $(i)p_{yz} = ((i)p_y)p_z$ , as required.

It remains to prove that  $\zeta$  is injective. Suppose that  $y, z \in \operatorname{Stab}_S(L_x^{P_n})$  are such that  $p_y = p_z$ . It suffices, by Lemma 3.6, to show that xy = xz. We will prove that  $x^*xy = x^*xz$  so that  $xy = xx^*xy = xx^*xz = xz$ . Since the transverse blocks of  $x^*xy$  are  $B_i \times -B_{(i)p_y} = B_i \times -B_{(i)p_z}$  where  $i \in \mathbf{r}$ , it follows that the transverse blocks of  $x^*xy$  and  $x^*xz$  coincide. Suppose that  $(k,l) \in x^*xy$  where neither k nor l belongs to a transverse block of  $x^*xy$ . It follows from the form of the transverse blocks of  $x^*xy$  that neither k nor l belongs to a transverse block of  $x^*x$ . There are two cases to consider: k,l>0 and k,l<0. In the first case, by part (i) of the definition of the mulitplication of  $x^*x$  and y, either  $(k,l) \in x^*x$  or k and l belong to transverse blocks of  $x^*x$ . Since the latter is not the case,  $(k,l) \in x^*x$  and so  $(k,l) \in x^*xz$ . In the second case, when k,l<0, it follows from part (iii) of the definition of the multiplication of  $x^*x$  and  $x^*xz$  and so  $x^*xy$  and so  $x^*xy$  and so  $x^*xy$  or both  $x^*xy$  and  $x^*xz$  or both  $x^*xy$  or both  $x^*xy$  and  $x^*xz$ . Since the transverse blocks of  $x^*xy$  and  $x^*xz$  coincide and contain neither  $x^*xy$  are  $x^*xy$ . Therefore  $x^*y$  are  $x^*y$  are  $x^*y$  and  $x^*xy$  and  $x^*xz$  coincide and contain neither  $x^*y$  are  $x^*y$  are  $x^*y$  and  $x^*yz$  coincide and contain neither  $x^*y$  are  $x^*y$ . Therefore  $x^*y$  are  $x^*y$  are  $x^*y$  and  $x^*yz$  coincide and contain neither  $x^*y$  are  $x^*y$  are  $x^*yz$  and  $x^*yz$  coincide and contain neither  $x^*yz$  are required.

# 5 Algorithms

In this section, we outline some algorithms for computing with semigroups that utilise the results in Section 3. The algorithms described in this section are implemented in the GAP [15] package Semigroups [29] in their full generality, and can currently be applied to semigroups of transformations, partial permutations, partitions, and to subsemigroups of regular Rees 0-matrix semigroups over groups.

Throughout this section, we will suppose that U is a finite regular semigroup, and that S is subsemigroup of U generated by  $X = \{x_1, \ldots, x_m\} \subseteq U$ . The purpose of the algorithms described in this section is to answer various questions about the structure of the semigroup S.

Recall that  $U^1$  has an identity  $1_U \notin U$  adjoined regardless of whether or not U is already a monoid; the analogous statement holds for S where the adjoined identity is denoted  $1_S$ . We note that it is not necessary for  $1_S$  and  $1_U$  to be equal.

Apart from this introduction, this section has 6 subsections. In Subsection 5.1, we outline some basic operations that we must be able to compute in order to apply the algorithms later in this section. We then show how to perform these basic operations in the examples of semigroups of transformations, partial

permutations, matrices, partitions, and in subsemigroups of a Rees 0-matrix semigroup in Subsection 5.2. In Subsection 5.3, we describe how to calculate the components of the action of a semigroup on a set, and how to use this to obtain the Schreier generators from Proposition 2.3(c). In Subsection 5.4, we describe a data structure for individual Green's classes of S and give algorithms showing how this data structure can be used to compute various properties of these classes. In Subsection 5.5, we describe algorithms that can be used to find global properties of S, such as its size,  $\mathcal{R}$ -classes, and so on. In Subsection 5.6, we give details of how some of the algorithms in Subsection 5.5 can be optimised when it is known a priori that S is a regular or inverse semigroup.

At several points in this section it is necessary to be able to determine the strongly connected components (s.c.c.) of a directed graph. This can be achieved using Tarjan's [44] or Gabow's [14] algorithms, for example; see also Sedgwick [39].

In the algorithms in this section, ":=" indicates that we are assigning a value (the right hand side of the expression) to a variable (the left hand side), while "=" denotes a comparison of variables. The symbol "\(—\)" is the replacement operator, used to indicated that the value of the variable on the left hand side is replaced by the value on the right hand side.

## 5.1 Assumptions

Suppose that U is a finite regular semigroup and that S is a subsemigroup of U. As discussed at the start of Section 4, we suppose that we have a right action of S on a set  $(U)\lambda$  and a homomorphism  $\lambda:U\longrightarrow (U)\lambda$  of this action and the action of S on U by right multiplication, where  $\ker(\lambda)=\mathscr{L}^U$ . Furthermore, for every  $x\in U$  we assume that we have a faithful representation  $\zeta$  of the stabiliser  $U_{L_x^U}$  and a function  $\mu_x:\operatorname{Stab}_U(L_x^U)\longrightarrow (U_{L_x^U})\zeta$  defined by

$$(u)\mu_x = (u|_{L^U})\zeta$$
 for all  $u \in U^1$ ;

see (3.5). We also assume that we have the left handed analogues  $\rho: U \longrightarrow (U)\rho$  and  $\nu_x: \operatorname{Stab}_U(R_x^U) \longrightarrow (R_x^U U)\zeta'$  (where  $\zeta'$  is any faithful representation of  $R_x^U U$ ) of  $\lambda$  and  $\mu_x$ , respectively. Recall that we write

$$S_x = (\operatorname{Stab}_S(L_x^U))\mu_x$$
 and  $_xS = (\operatorname{Stab}_S(R_x^U))\nu_x$ .

In order to apply the algorithms described in this section, it is necessary that certain fundamental computations can be performed.

**Assumptions.** We assume that we can compute the following:

- (I) the product xy;
- (II) the value  $(x)\lambda$ ;
- (III) an element  $\bar{s} \in U^1$  such that  $xs\bar{s} = x$  whenever  $(x)\lambda \sim (x)\lambda \cdot s = (xs)\lambda$ ;
- (IV) the value  $(x's)\mu_x$ , for some choice of  $x' \in U$  such that xx'x = x, whenever  $(x)\lambda = (s)\lambda$  and  $(x)\rho = (s)\rho$ ;

for all  $x, y, s \in U$ . We also require the facility to perform the analogous computations involving the functions  $\rho$  and the  $\nu_x$  for all  $x \in S$ .

We will prove that  $\overline{s} \in S^1$  in Assumption (III) exists. By the definition of  $\lambda$ , since  $(x)\lambda \sim (xs)\lambda$ , it follows that  $L_x^U \sim L_{xs}^U$  under the action of S on  $U/\mathscr{L}^U$  defined in (3.2). Hence, by Proposition 2.3(a), there exists  $\overline{s} \in S^1$  such that

$$L_{xs}^{U} \cdot \overline{s} = L_{x}^{U}$$
 and  $(s\overline{s})|_{L_{x}^{U}} = \mathrm{id}_{L_{x}^{U}}$ 

and so  $xs\overline{s} = x$ . Given the orbit graph of  $(x)\lambda \cdot S^1$ , it is possible to compute  $\overline{s}$  by finding a path from  $(xs)\lambda$  to  $(x)\lambda$  and using Algorithm 2. However, this is often more expensive than computing  $\overline{s} \in U$  directly from s. Some details of how to compute Assumptions (I) to (IV) in the special cases given in Section 4 can be found in the next section.

If  $\overline{s} \in U^1$  satisfies Assumption (III) for some  $x \in S$ , then we note that  $xs\overline{s} = x$  implies that  $xs\overline{s}s = xs$  and so by Lemma 3.6:

$$(\overline{s}s)|_{L_{xs}^U} = \mathrm{id}_{L_{xs}^U}$$
.

Note that if  $s \in \operatorname{Stab}_S(R_x^U)$ , then  $(sx)\rho = (x)\rho$  and, by Proposition 3.10(b),  $sx \in L_x^S \cap R_x^U$ . In particular,  $(sx)\lambda = (x)\lambda$ , and so, by Assumption (IV), we can compute:

(V) the value  $((s)\nu_x)\Psi = (x'sx)\mu_x$  whenever  $s \in \operatorname{Stab}_S(R_x^U)$ .

We will refer to this as Assumption (V), even though it is not an assumption.

It will follow from the comments in Subsection 5.2 that the algorithms described in this paper can be applied to any subsemigroup of the full transformation monoid, the symmetric inverse monoid, the partition monoid, the general linear monoid, or any subsemigroup of a regular Rees 0-matrix semigroup over a group. However, we would like to stress that the algorithms in this section apply to any subsemigroup of a finite regular semigroup. In the worst case, the functions  $\lambda: U \longrightarrow U/\mathscr{L}^U$  and  $\rho: U \longrightarrow U/\mathscr{R}^U$  defined by  $(x)\lambda = L_x^U$  and  $(x)\rho = R_x^U$ , and the natural mappings  $\mu_x: \operatorname{Stab}_S(L_x^U) \longrightarrow S_{L_x^U}$  and  $\nu_x: \operatorname{Stab}_S(R_x^U) \longrightarrow R_x^U S$  for every  $x \in U$ , fulfil the required conditions, although it might be that our algorithms are not very efficient in this case.

## 5.2 Computational prerequisites

In this section, we describe how to perform the computations required in Assumptions (I) to (IV) for semigroups of transformations, partial permutations, partitions, and matrices, and for subsemigroups of a Rees 0-matrix semigroup.

#### **Transformations**

A transformation x can be represented as a tuple  $((1)x, \ldots, (n)x)$  where n is the degree of x.

- (I) The composition xy of transformations x and y is represented by  $((1)xy, \ldots, (n)xy)$ , which can be computed by simple substitution in linear time O(n);
- (II) The value  $(x)\lambda = \operatorname{im}(x)$  can be found by sorting and removing duplicates from  $((1)x, \ldots, (n)x)$  with complexity  $O(n\log(n))$ ;
- (III) If x, s are such that im(x) and im(xs) have equal cardinality, then the transformation  $\overline{s}$  defined by

$$(i)\overline{s} = \begin{cases} (j)x & \text{if } i = (j)xs \in \text{im}(xs) \\ i & \text{if } i \notin \text{im}(xs) \end{cases}$$

has the property that  $xs\overline{s} = x$  (finding  $\overline{s}$  has complexity O(n));

(IV) If x and s are transformations such that  $\ker(x) = \ker(s)$  and  $\operatorname{im}(x) = \operatorname{im}(s)$  and x' is any transformation such that xx'x = x, then, from Proposition 4.2(b),  $(x's)\mu_x$  is just the restriction  $(x's)|_{\operatorname{im}(x)}$  of x's to  $\operatorname{im}(x)$ , which can be determined in  $|\operatorname{im}(x)|$  steps from x and s.

The analogous calculations can be made in terms of the kernel of a transformation, and the left actions of S; the details are omitted.

#### Partial permutations

A partial permutation x can be represented as a tuple  $((1)x, \ldots, (m)x)$  where m is the largest value where x is defined, and (i)x = 0 if x is not defined at i. The values required under Assumptions (I), (II), and (IV) can be computed for partial permutations in the same way they were computed for transformations. Assumption (III) is described below:

(III) if x, s are such that  $\operatorname{im}(x)$  and  $\operatorname{im}(xs)$  have equal cardinality, then the partial permutation  $\overline{s} = s^{-1}$  has the property  $xs\overline{s} = x$  (finding  $s^{-1}$  has complexity O(n)).

#### Matrices over finite fields

The values required in Assumptions (I) to (IV) can be found for matrix semigroups in a similar way as they are found for transformation and partial permutations, using elementary linear algebra; the details will appear in [30].

#### Rees 0-matrix semigroups

It should be clear from the definition of a Rees 0-matrix semigroup, and by Proposition 4.9, how to compute the values in Assumptions (I) and (II). Assumptions (III) and (IV) are described below:

- (III) if  $x = (a, b, i), s = (j, g, k) \in S$  are such that  $(x)\lambda = i \sim (s)\lambda = k$ , then  $p_{i,j} \neq 0$ . So, if  $l \in I$  is such that  $p_{k,l} \neq 0$ , then  $\overline{s} = (l, p_{k,l}^{-1} g^{-1} p_{i,j}^{-1}, i)$  has the property that  $xs\overline{s} = x$ ;
- (IV) if  $x = (i, g, j) \in S$ , then there exist  $k \in I$  and  $l \in J$  such that  $p_{j,k}, p_{l,i} \neq 0$ . One choice for  $x' \in U$  is  $(k, p_{j,k}^{-1} g^{-1} p_{l,i}^{-1}, l)$ . So, if  $h \in G$  is arbitrary and s = (i, h, j), then  $(x)\lambda = (s)\lambda$  and  $(x)\rho = (s)\rho$ . Hence the action of  $(x's)\mu_x$  on  $m \in \mathbf{n}$  (defined in Proposition 4.9(b)) is given by

$$m \cdot (x's)\mu_x = (m)g^{-1}h.$$

#### **Partitions**

A partition  $x \in P_n$  can be represented as a 2n-tuple where the first n entries correspond to the indices of the blocks containing  $\{1, \ldots, n\}$  and entries n+1 to 2n correspond to the indices of the blocks containing  $\{-1, \ldots, -n\}$ . For example, the partition  $x \in P_6$  shown in Figure 1 is represented by (1, 2, 3, 4, 5, 5, 1, 6, 4, 6, 5, 5).

Given  $x \in P_n$  represented as above, it is possible to compute  $x^* \in P_n$  in 2n steps (linear complexity). This will be used in several of the assumptions.

- (I) The composition xy of partitions x and y can be found using a variant of the classical Union-Find Algorithm;
- (II) The value  $x^*x$  can be found using (I);
- (III) If  $x, s \in P_n$  are such that  $(x)\lambda = x^*x \sim (xs)\lambda = s^*x^*xs$ , then  $x^*x$  and  $s^*x^*xs$  have equal number of transverse blocks. A partition  $\bar{s}$  with the property that  $xs\bar{s} = x$  can then be found using a variant of the Union-Find Algorithm;
- (IV) If  $x, s \in P_n$  are such that  $(x)\lambda = (s)\lambda$  and  $(x)\rho = (s)\rho$ , then, by Proposition 4.11(b),  $(x^*s)\mu_x$  is a permutation of the transverse blocks of x, which can be found in linear time (complexity O(n)).

In practice, it is possible to compute the values required by these assumptions with somewhat better complexity than that given above. However, this is also more complicated to describe and so we opted to describe the simpler methods.

## 5.3 Components of the action

We start by describing a procedure for calculating  $\alpha \cdot S^1 = \{\alpha \cdot s : s \in S^1\}$  or  $\Omega \cdot S^1 = \{\alpha \cdot s : \alpha \in \Omega, s \in S^1\}$ . These are essentially the same as the standard orbit algorithm for a group acting on a set (see, for example, [46, Section 4.1]), but without the assumption that S is a group. An analogous algorithm can be used for left actions. We will refer to  $\alpha \cdot S^1$  as the component of the action of S containing  $\alpha$ .

In this subsection we present algorithms for computing: the components of an action of a semigroup S; elements of S that act on points in the component in a specified way; generators for the stabiliser of a set. Examples of how these algorithms can applied can be found in Section 6. The algorithms in this subsection and the next are somewhat similar to those described in [24].

Suppose that  $X = \{x_1, \ldots, x_m\}$  is a generating set for a monoid S acting on the right on a set  $\Omega$ . If  $\alpha \in \Omega$  and  $\alpha \cdot S = \{\beta_1 = \alpha, \beta_2, \ldots, \beta_n\}$ , then the *orbit graph* of  $\alpha \cdot S$  is just the directed graph with vertices  $\{1, \ldots, n\}$  and an edge from i to  $g_{i,j}$  labelled with j if  $\beta_i \cdot x_j = \beta_{g_{i,j}}$ . The orbit graph of  $\Omega \cdot S$  is defined analogously. A *Schreier tree* for  $\alpha \cdot S$  is just a spanning tree for the orbit graph with root at  $\beta_1$ . Schreier trees for  $\alpha \cdot S$  will be represented using a 2-dimensional array

$$v_2 \quad \dots \quad v_n \\ w_2 \quad \dots \quad w_n$$

such that  $\beta_{v_i} \cdot x_{w_i} = \beta_j$  and  $v_j < j$  for all j > 1.

The orbit graph of  $\Omega \cdot S^1$  may not be connected and so it has a forest of (not necessarily disjoint) Schreier trees rooted at some elements of  $\Omega$ . To simplify things, we may suppose without loss of generality that there is an  $\alpha \in \Omega$  such that  $\alpha \cdot S = \Omega \cdot S$ . This can be achieved by adding an artificial  $\alpha$  to  $\Omega$  (and perhaps some further points) and defining the action of S on these values so that  $\alpha \cdot S$  contains all of the roots of the Schreier forest for  $\Omega \cdot S$ .

Note that unlike the orbit graph of a component of a group acting on a set, the orbit graph of a component of a semigroup acting on a set is, in general, not strongly connected. This makes several of the steps required below more complicated than in the group case.

#### Algorithm 1 Compute a component of an action

```
Input: S := \langle X \rangle where X := \{x_1, \dots, x_m\}, S acts on a set \Omega on the right, and \alpha \in \Omega
Output: \alpha \cdot S^1, a Schreier tree for \alpha \cdot S^1, and the orbit graph of \alpha \cdot S^1
                                                                                [C will equal \alpha \cdot S^1 by the end of the algorithm]
 1: C := \{\beta_1 := \alpha\}, n := 1
 2: for \beta_i \in C, j \in \{1, ..., m\} do
                                                                                            [loop over: values in C, the generators]
         if \beta_i \cdot x_j \notin C then
              n := n + 1, \, \beta_n := \beta_i \cdot x_j, \, C \leftarrow C \cup \{\beta_n\};
                                                                                                                          [add \beta_i \cdot x_i to C]
 4:
              v_n := i, w_n := j, g_{i,j} := n
                                                                                         [update the Schreier tree and orbit graph]
 5:
          else if \beta_i \cdot x_j = \beta_r \in C then
 6:
                                                                                                                  [update the orbit graph]
              g_{i,j} := r;
 7:
 8:
          end if
 9: end for
10: return C, (v_2, \ldots, v_n, w_2, \ldots, w_n), \{g_{i,j} : i \in \{1, \ldots, n\}, j \in \{1, \ldots, m\}\}
```

The Schreier tree for  $\alpha \cdot S^1$  produced by Algorithm 1 can be used to obtain elements  $u_i \in S^1$  such that  $\beta_1 \cdot u_i = \beta_i$  for all *i* using Algorithm 2. The standard method for a group acting on a set, such as the procedure U-Beta in [46, p80], cannot be used here, due to the non-existence of inverses in semigroups.

## Algorithm 2 Trace a Schreier tree

```
Input: a Schreier tree (v_2, \ldots, v_n, w_2, \ldots, w_n) for \alpha \cdot S^1, and \beta_i \in \alpha \cdot S^1

Output: u \in S^1 such that \alpha \cdot u = \beta_i

1: u := 1_S, j := i

2: while j > 1 do

3: u := x_{w_j} u and j := v_j

4: end while

5: return u
```

The right action of S on  $\Omega$  induces an action of S on  $\mathcal{P}(\Omega)$ . In Algorithm 3, Algorithms 1 and 2 are used to obtain the Schreier generators from Proposition 2.3(c) for the stabiliser  $S_{\Sigma}$  of a subset  $\Sigma$  of  $\Omega$  under this induced action.

### Algorithm 3 Compute Schreier generators for a stabiliser

```
Input: the component \Sigma \cdot S^1 of \Sigma \subseteq \Omega under the action of S on \mathcal{P}(\Omega), a Schreier tree and orbit graph
      for \Sigma \cdot S^1
Output: Schreier generators Y for the stabiliser S_{\Sigma}
 1: Y := \{ id_{\Sigma} \}
 2: find the s.c.c. of \Sigma in \Sigma \cdot S^1 := \{\Sigma_1 := \Sigma, \dots, \Sigma_r\}
 3: for i \in \{1, ..., r\}, j \in \{1, ..., m\} do
                                                                                                       [loop over: \Sigma \cdot S^1, the generators of S^1]
                                                                                                           [g_{i,j} \text{ is from the orbit graph of } \Sigma \cdot S^1]
           set k := g_{i,j}
 4:
           if \Sigma_k \sim \Sigma_1 then
                                                                                                                              [\Sigma_k \text{ is in the s.c.c. of } \Sigma_1]
 5:
                find u_i, u_k \in S^1 such that \Sigma_1 \cdot u_i = \Sigma_i and \Sigma_1 \cdot u_k = \Sigma_k
                                                                                                                                                 [Algorithm 2]
 6:
                find \overline{u_k} \in U such that \Sigma_k \cdot \overline{u_k} = \Sigma_1 and (u_k \overline{u_k})|_{\Sigma_1} = \mathrm{id}_{\Sigma_1}
 7:
                                                                                                                                       [Proposition 2.3(a)]
                 Y \leftarrow Y \cup \{(u_i x_i \overline{u_k})|_{\Sigma_1}\}
 8:
           end if
 9:
10: end for
11: return Y
```

In Algorithm 4 we give a more specialised version of Algorithm 3, which we require to find  $(\operatorname{Stab}_S(L_x^U))\mu_x = S_x$  where  $\mu_x$  is the function defined at the start of this section.

## **Algorithm 4** Compute Schreier generators for $S_x$

```
Input: the component (x)\lambda \cdot S^1 of (x)\lambda, a Schreier tree and orbit graph for (x)\lambda \cdot S^1
Output: Schreier generators Y for the stabiliser S_x
 1: set Y := \emptyset, x_1 := x
 2: find the s.c.c. \{(x_1)\lambda, \dots, (x_r)\lambda\} of (x)\lambda in (x)\lambda \cdot S^1
                                                                               [loop over: the s.c.c. of (x)\lambda, the generators of S]
 3: for i \in \{1, ..., r\}, j \in \{1, ..., m\} do
          set k := g_{i,j}
                                                                                             [g_{i,j} \text{ is from the orbit graph of } (x)\lambda \cdot S^1]
 4:
          if (x_k)\lambda \sim (x_1)\lambda then
                                                                                                          [(x_k)\lambda \text{ is in the s.c.c. of } (x_1)\lambda]
 5:
               find u_i, u_k \in S^1 such that (x_1)\lambda \cdot u_i = (x_i)\lambda and (x_1)\lambda \cdot u_k = (x_k)\lambda
                                                                                                                                    [Algorithm 2]
 6:
               find \overline{u_k} \in U such that xu_k\overline{u_k} = x
                                                                                                                            [Assumption (III)]
 7:
               Y \leftarrow Y \cup \{(x'u_ix_i\overline{u_k})\mu_x\}
                                                                                                                            [Assumption (IV)]
 8:
 9:
          end if
10: end for
11: return Y
```

Algorithms 3 and 4 can also be used to find generators for the stabiliser of any value in the component  $\Sigma \cdot S^1$  or  $(x)\lambda \cdot S^1$ , if we have a Schreier tree for the s.c.c. rooted at that value. Algorithm 1 returns a Schreier tree for the entire component (possibly including several strongly connected components) rooted at the first point in the component. It is possible to find a Schreier tree for any s.c.c. rooted at any value in the component by finding a spanning tree for the subgraph of the orbit graph the s.c.c. induces. Such a spanning tree can be found in linear time using a depth first search algorithm, for example.

## 5.4 Individual Green's classes

#### Data structures

By Corollary 3.13, we can represent the  $\mathscr{R}$ -class  $R_x^S$  of any element  $x \in S$  as a quadruple consisting of:

- the representative x;
- the s.c.c.  $\{(x)\lambda = \alpha_1, \dots, \alpha_n\}$  of  $(x)\lambda$  under the action of S (this can be found using Algorithms 1 and any algorithm to find the strongly connected components of a digraph);
- a Schreier tree for  $\{\alpha_1, \ldots, \alpha_n\}$ ;
- the stabiliser group  $S_x$  found using Algorithm 4.

The  $\mathscr{L}^S$ -class of x in S can be represented using the analogous quadruple using the s.c.c. of  $(x)\rho$ , and the group  $_xS$ . The Green's  $\mathscr{H}$ - and  $\mathscr{D}$ -classes of x in S are represented using the quadruple for  $R_x^S$  and the quadruple for  $L_x^S$ .

## Size of a Green's class

Having the above data structures, it follows from Corollaries 3.13(b) and 3.14(b) that  $|R_x^S| = n \cdot |S_x|$  where n is the length of the s.c.c. of  $(x)\lambda$  in  $(x)\lambda \cdot S^1$ . Similarly,  $|L_x^S|$  is the length of the s.c.c. of  $(x)\rho$  multiplied by |xS|. Suppose that  $x' \in U$  is such that xx'x = x. Then, by Proposition 3.11(b),  $|H_x^S| = |S_x \cap (xS)\Psi|$ , where  $\Psi: xS \longrightarrow U_x$  is defined by  $((s)\nu_x)\Psi = (x'sx)\mu_x$  for all  $s \in xS$ . The group xS can be found using the analogue of Algorithm 4, and we can compute the values  $(x'sx)\mu_x$  for all  $s \in xS$  by Assumption (V). The size of the  $\mathscr{D}$ -class  $D_x^S$  is just

$$|D_x^S| = \frac{|L_x^S| \cdot |R_x^S|}{|H_x^S|},$$

and so  $|D_x^S|$  can be found using the values of  $|L_x^S|$ ,  $|R_x^S|$ , and  $|H_x^S|$ .

#### Elements of a Green's class

Corollary 3.18(a) states that

$$R_x^S = \{xsu : s \in \operatorname{Stab}_S(L_x^U), u \in S^1, (x)\lambda \cdot u \sim (x)\lambda\}.$$

If  $s, t \in \operatorname{Stab}_S(L_x^U)$  are such that  $(s)\mu_x = (t)\mu_x$ , then, by Lemma 3.6, xs = xt. It follows that if M is any subset of  $\operatorname{Stab}_S(L_x^U)$  such that  $(M)\mu_x = S_x$  and  $(s)\mu_x \neq (t)\mu_x$  for all  $s, t \in M$  such that  $s \neq t$ , then

$$R_x^S = \{xsu : s \in M, u \in S^1, (x)\lambda \cdot u \sim (x)\lambda\}.$$

If  $Y = \{(y_1)\mu_x, \ldots, (y_k)\mu_x\}$  is a set of generators for  $S_x$  (from Algorithm 4), then every element of  $S_x$  is of the form  $(s)\mu_x$  where  $s \in \langle y_1, \ldots, y_k \rangle$ . Thus the set M can be found by computing the elements of  $S_x$ , and expressing each element as a product of the generators Y. An algorithm for finding the elements of  $R_x^S$  is given in Algorithm 5.

## Algorithm 5 Elements of an $\mathcal{R}$ -class

```
Input: x \in S

Output: the elements Y of the \mathscr{R}-class R_x^S

1: Y := \varnothing

2: find the s.c.c. \{(x)\lambda = \alpha_1, \dots, \alpha_n\} of (x)\lambda, and a Schreier tree for this s.c.c. [Algorithm 1]

3: find the group S_x [Algorithm 4]

4: for i \in \{1, \dots, n\}, (s)\mu_x \in S_x do [Loop over: the s.c.c., elements of the group]

5: find u_i \in S^1 such that \alpha_1 \cdot u_i = \alpha_i [Algorithm 2]

6: Y \leftarrow Y \cup \{xsu_i\}

7: end for

8: return Y
```

The elements of an  $\mathscr{L}$ -class can be found using an analogous algorithm. By the proof of Proposition 3.11(b),  $\phi_1: S_x \cap (_xS)\Psi \longrightarrow H_x^S$  defined by  $((s)\mu_x)\phi_1 = xs$  is a bijection. It follows that if we can compute the intersection of groups  $S_x \cap (_xS)\Psi$ , then we can obtain the elements of  $H_x^S$ . As mentioned above,  $(_xS)\Psi$  can be determined using Algorithm 4 and by Assumption (V).

The elements of a  $\mathscr{D}$ -class are slightly more complicated to compute. In Algorithm 6 we show how to find the  $\mathscr{R}$ -classes in a given  $\mathscr{D}$ -class of S and this combined with Algorithm 5 gives a method for finding the elements of a  $\mathscr{D}$ -class.

## Classes within classes

By Proposition 3.21, if  $x \in S$ ,  $x' \in U$  is such that xx'x = x, and:

- $\mathcal{C}$  is a subset of  $\operatorname{Stab}_U(L_x^U)$  such that  $\{(c)\mu_x:c\in\mathcal{C}\}$  is a left transversal of  $S_x\cap(_xS)\Psi$  in  $(_xS)\Psi$  where  $\Psi:_xS\longrightarrow U_x$ , defined by  $((s)\nu_x)\Psi=(x'sx)\mu_x$ , is the embedding from Proposition 3.11(a);
- $\{u_i \cdot (x)\rho : 1 \le i \le m\}$  is the s.c.c. of  $(x)\rho$  under the left action of S, where  $u_i \in S^1$  for all i,

then  $\{u_ixc:c\in\mathcal{C},\ 1\leq i\leq m\}$  is a set of  $\mathscr{H}^S$ -class representatives for  $L_x^S$ , and hence a set of  $\mathscr{R}^S$ -class representatives for  $D_x^S$ . Using this result in Algorithm 6 we show how to find the  $\mathscr{R}^S$ -classes of a  $\mathscr{D}^S$ -class. Since  $(u_ixc)\lambda=(x)\lambda$ , it follows that  $S_{u_ixc}=S_x$  for all i and all  $c\in\mathcal{C}$ . Therefore, the data structures for  $R_x^S$  and  $R_{u_ixc}^S$  are identical except for the representatives.

From Proposition 3.22, algorithms analogous to Algorithm 6, can be used to find the  $\mathcal{L}^S$ -classes in a  $\mathcal{D}^S$ -class, the  $\mathcal{H}^S$ -classes in an  $\mathcal{R}^S$ -classes in an  $\mathcal{L}^S$ -class.

#### Testing membership

Using Corollary 3.16 and 3.20, in Algorithms 7 and 8 we show how the data structures described at the start of the section can be used to test membership in an  $\mathcal{R}$ - or  $\mathcal{D}$ -class.

Using Proposition 3.17, an algorithm analogous to Algorithm 7 (for  $\mathscr{R}$ -classes) can be used to test membership in an  $\mathscr{L}$ -class. Testing membership in an  $\mathscr{H}$ -class can then be accomplished by testing membership in the corresponding  $\mathscr{L}$ - and  $\mathscr{R}$ -classes.

#### Regularity and idempotents

An  $\mathscr{R}$ -class R of S is regular if and only if there is  $x \in R$  such that  $H_x^U$  is a group. Since  $y \in H_x^U$  if and only if  $(x)\lambda = (y)\lambda$  and  $(x)\rho = (y)\rho$ , it is possible to verify that an  $\mathscr{R}$ -class is regular only by considering

## **Algorithm 6** $\mathcal{R}$ -classes in a $\mathcal{D}$ -class

```
Input: x \in S
Output: \mathscr{R}-class representatives \mathfrak{R} of the \mathscr{D}-class D_x^S
 1: \mathfrak{R} := \varnothing
 2: find the s.c.c.s of (x)\lambda and (x)\rho and their Schreier trees
                                                                                                                                  [Algorithm 1]
 3: find S_x \cap (_xS)\Psi
                                                                                                    [Algorithm 4 and Assumption (V)]
 4: find \mathcal{C} \subseteq \operatorname{Stab}_U(L_x^U) such that \{(c)\mu_x : c \in \mathcal{C}\} is a left transversal of S_x \cap ({}_xS)\Psi in ({}_xS)\Psi
                                                                                                                       [Proposition 3.21(a)]
 5: for (y)\rho in the s.c.c. of (x)\rho do
 6:
          find u_i \in S such that u_i \cdot (x)\rho = (y)\rho
                                                                                                                                  [Algorithm 2]
          for c \in \mathcal{C} do
 7:
               \Re \leftarrow \Re \cup \{u_ixc\}
 8:
          end for
 9:
10: end for
11: return \Re
```

## **Algorithm 7** Test membership in an $\mathcal{R}$ -class

```
Input: y \in U and the data structure of an \mathscr{R}-class R_x^S

Output: true or false

1: if (x)\rho = (y)\rho and (y)\lambda \sim (x)\lambda then

2: find u \in S^1 such that (x)\lambda \cdot u = (y)\lambda [Algorithm 2]

3: find \overline{u} \in U^1 such that (y)\lambda \cdot \overline{u} = (x)\lambda and xu\overline{u} = x [Assumption (III)]

4: return (x'y\overline{u})\mu_x \in S_x [Assumption (IV)]

5: else

6: return false

7: end if
```

## Algorithm 8 Test membership in a $\mathcal{D}$ -class

```
Input: y \in U and the data structure for the \mathscr{D}-class of x \in S
Output: true or false
 1: if (x)\lambda \sim (y)\lambda and (x)\rho \sim (y)\rho then
          find u_1, u_2 \in S^1 such that (x)\lambda \cdot u_1 = (y)\lambda and u_2 \cdot (x)\rho = (y)\rho
                                                                                                                                      [Algorithm 2]
 2:
          find \overline{u_1} \in U^1 such that (y)\lambda \cdot \overline{u_1} = (x)\lambda and xu_1\overline{u_1} = x
                                                                                                                              [Assumption (III)]
 3:
          find \overline{u_2} \in U^1 such that \overline{u_2} \cdot (y)\rho = (x)\rho, and \overline{u_2}u_2x = x
                                                                                                       [the analogue of Assumption (III)]
 4:
          find S_x \cap (_xS)\Psi
 5:
                                                                                                       [Algorithm 4 and Assumption (V)]
          find \mathcal{C} \subseteq \operatorname{Stab}_U(L_x^U) such that \{(c)\mu_x : c \in \mathcal{C}\} is a left transversal of S_x \cap ({}_xS)\Psi in ({}_xS)\Psi
 6:
                                                                                                                           [Proposition 3.21(a)]
 7:
          for c \in \mathcal{C} do
 8:
               if (x'\overline{u_2}y\overline{u_1}c)\mu_x in (_xS)\Psi then
 9:
                    return true
10:
               end if
11:
          end for
12:
13: end if
14: return false
```

the value  $(x)\lambda$  and the s.c.c. of  $(x)\rho$ . A similar approach can be used to compute the idempotents in an  $\mathcal{R}$ -class.

How we test if the  $\mathcal{H}$ -class in U corresponding to  $(x)\lambda$  and  $(y)\rho$  is a group, depends on the context. For example:

**Transformation semigroups:** in the full transformation monoid,  $(x)\lambda$  and  $(y)\rho$  are an image set and kernel of a transformation, respectively. In this case, the  $\mathscr{H}$ -class corresponding to  $(x)\lambda$  and  $(y)\rho$  is a group if and only if  $(x)\lambda$  contains precisely one element in every kernel class of  $(x)\rho$ . If  $(x)\lambda$  and  $(y)\rho$  satisfy this property, then it is relatively straightforward to compute an idempotent with kernel  $(y)\rho$  and image  $(x)\lambda$ ;

**Partial permutations:** in the symmetric inverse monoid,  $(x)\lambda$  and  $(y)\rho$  are the image set and domain of a partial permutation, respectively. In this case, the  $\mathscr{H}$ -class corresponding to  $(x)\lambda$  and  $(y)\rho$  is a group if and only if  $(x)\lambda = (y)\rho$ . Given such  $(x)\lambda = (y)\rho$ , the partial identity function with domain  $(x)\lambda$  is the idempotent in the  $\mathscr{H}$ -class;

**Matrix semigroups:** In this case, there is no simple criteria for the  $\mathcal{H}$ -class of  $(x)\lambda$  and  $(y)\rho$  to be a group.

Rees matrix semigroups: in a Rees 0-matrix semigroup  $\mathcal{M}^0[G; I, J; P]$ ,  $(x)\lambda = j \in J$  and  $(y)\rho = k \in I$ . In this case, the  $\mathscr{H}$ -class corresponding to  $(x)\lambda$  and  $(y)\rho$  is a group if and only if  $p_{j,k} \neq 0$ . The idempotent in the  $\mathscr{H}$ -class is then  $(k, p_{j,k}^{-1}, j)$ ;

**Partitions:** in the partition monoid,  $(x)\lambda = x^*x$  and  $(y)\rho = yy^*$ . In this case, the  $\mathscr{H}$ -class  $L_{x^*x} \cap R_{yy^*}$  contains an idempotent if and only if the number of transverse blocks of  $yy^*x^*x$  equals the number of transverse blocks of  $x^*x$ . If the previous condition is satisfied, then the idempotent contained in  $L_{x^*x} \cap R_{yy^*}$  is  $yy^*x^*x$ .

## Algorithm 9 Regularity of an $\mathcal{R}$ -class

Algorithms analogous to Algorithms 9 and 10 can be used to test regularity and find the idempotents in an  $\mathscr{L}$ -class. A  $\mathscr{D}$ -class D in S is regular if and only if any (equivalently, every)  $\mathscr{R}$ -class in D is regular. Hence Algorithms 6 and 9 and can be used to verify if a  $\mathscr{D}$ -class is regular or not. It is possible to calculate the idempotents in a  $\mathscr{D}$ -class D by creating the  $\mathscr{R}$ -class representatives using Algorithm 6 and finding the idempotents in every  $\mathscr{R}$ -class of D using Algorithm 10.

## **Algorithm 10** Idempotents in an $\mathcal{R}$ -class

```
Input: x \in S
Output: the idempotents E of R_x^S
 1: E := \emptyset
 2: find the s.c.c. of (x)\rho in S^1 \cdot (x)\rho
                                                                                                [analogue of Algorithm 1]
 3: for (y)\rho in the s.c.c. of (x)\rho do
        if the \mathcal{H}-class H in U corresponding to (x)\lambda and (y)\rho is a group then
 4:
             find the identity e of H
 5:
             E \leftarrow E \cup \{e\}
 6:
        end if
 7:
 8: end for
 9: return E
```

## 5.5 The global structure of a semigroup

In this section, we provide algorithms for determining the structure of an entire semigroup, rather than just its individual Green's classes as in the previous sections. Unlike the previous two subsections, the algorithms described in this section differ significantly from the analogous procedures described in [24].

In order to streamline the presentation of some algorithms in this section, we will perform them on  $U^1$  and  $S^1$ , instead of U and S. Of course, if we want to perform a calculation on U or S instead of  $U^1$  or  $S^1$ , then we can simply perform whatever calculation we require in  $U^1$  or  $S^1$ , then return the answer for U or S.

#### The main algorithm

Algorithm 11 is the main algorithm for computing the size, the Green's structure, testing membership, and so on in S. This algorithm could be replaced by an analogous algorithm which enumerates  $\mathscr{L}$ -classes, rather than  $\mathscr{R}$ -classes. In the case of transformation semigroups, in many test cases, the algorithm for  $\mathscr{R}$ -classes has better performance than the analogous algorithm for  $\mathscr{L}$ -classes. This is one reason for presenting this algorithm and not the other.

Since Green's  $\mathscr{R}$ -relation is a left congruence, it follows that representatives of the  $\mathscr{R}$ -classes of S can be obtained from the identity by left multiplying by the generators. The principal purpose of Algorithm 11 is to determine the action of S on its  $\mathscr{R}$ -class representatives by left multiplication. In particular, we enumerate  $\mathscr{R}$ -class representatives of S, which we will denote by  $\mathfrak{R}$ . Since we are calculating an action of S, we may also discuss the Schreier tree and orbit graph of this action, as we did in Algorithm 1. At the same time as finding the action of S on its  $\mathscr{R}$ -class representatives, we also calculate  $(S)\rho$ . Since  $(x)\rho = (y)\rho$  if and only if  $x\mathscr{R}^U y$ , it follows that

$$(S)\rho = (\mathfrak{R})\rho = \{(x)\rho : x \in \mathfrak{R}\}.$$

In Algorithm 11, we find an addition parameter, which is denoted by  $K_i$ . This parameter is used later in Algorithm 13, which allows us to factorise elements of a semigroup over its generators.

If the subsemigroup S we are trying to compute is  $\mathscr{R}$ -trivial, then Algorithm 11 simply exhaustively enumerates the elements of S. In such a case, unfortunately, Algorithm 11 has poorer performance than a well-implemented exhaustive algorithm, since it contains some superfluous steps. For example, the calculations of  $(S)\lambda$  and the groups  $S_x$  are unnecessary steps if S is  $\mathscr{R}$ -trivial. On a more positive note, in some cases, it is possible to detect if a semigroup is  $\mathscr{R}$ -trivial with relatively little effort. For example, if S is a transformation semigroup of degree n, then it is shown in [41] that S is  $\mathscr{R}$ -trivial if and only if its action on the points in  $\{1,\ldots,n\}$  is acyclic. In other words, it is possible to check whether a transformation semigroup is  $\mathscr{R}$ -trivial in polynomial time  $O(n^3)$ . At least in this case, it would then be possible to use the  $\mathscr{L}$ -class version of Algorithm 11 instead of the  $\mathscr{R}$ -class version, or indeed, an exhaustive algorithm such as that in [35].

From Algorithm 11, it is routine to calculate the size of S as:

$$|S| = \sum_{i=1}^{r} |S_{z_i}| \cdot |\text{s.c.c. of } (z_i)\lambda| \cdot |\{y \in \Re : (y)\lambda = (z_i)\lambda\}| - 1.$$
 (5.1)

The elements of S are just the union of the sets of elements of the  $\mathcal{R}$ -classes of S, which can be found using Algorithm 5.

The idempotents of S can be found by determining the idempotents in the  $\mathscr{R}$ -classes of S using Algorithm 10. Similarly, it is possible to test if S is regular, by checking that every  $\mathscr{R}$ -class or  $\mathscr{D}$ -class is regular using Algorithm 9. Note that the existence of an idempotent in an  $\mathscr{R}^S$ -class depends only on the values of  $\lambda$  and  $\rho$  of the elements of that class. Hence if there are at least two distinct  $\mathscr{R}$ -class representatives x and y in S such that  $(x)\lambda = (y)\lambda$  and  $(x)\rho = (y)\rho$ , then S is not regular, since the disjoint  $\mathscr{R}$ -classes  $R_x^S$  and  $R_y^S$  cannot contain the same idempotents.

The  $\mathscr{D}$ -classes of  $S^1$  are in 1-1 correspondence with the strongly connected components of the orbit

The  $\mathscr{D}$ -classes of  $S^1$  are in 1-1 correspondence with the strongly connected components of the orbit graph of  $\mathfrak{R}$ , which is obtained in Algorithm 11. Since we also find  $(S^1)\rho$  in Algorithm 11, it is possible to find the  $\mathscr{D}$ -classes of  $S^1$ , and their data structures, by finding the strongly connected components of the orbit graph of  $\mathfrak{R}$ . The  $\mathscr{L}$ - and  $\mathscr{H}$ -classes of S and  $S^1$  can then be found from the  $\mathscr{D}$ -classes using the analogues of Algorithm 6 (for finding the  $\mathscr{R}$ -classes in a  $\mathscr{D}$ -class).

In Algorithms 12, 13, 14, and 15, we will assume that Algorithm 11 has been performed and that the result has been stored somehow. So, for example, if we wanted to check membership of several elements in the semigroup U in its subsemigroup S, we perform Algorithm 11 only once.

#### Algorithm 11 Enumerate the $\mathcal{R}$ -classes of a semigroup

```
Input: S^1 := \langle X \rangle where X := \{x_1, ..., x_m\}
Output: data structures for the \mathscr{R}-classes with representatives \mathfrak{R} in S^1, the set (S^1)\rho, and Schreier trees
     and orbit graphs for \mathfrak{R} and (S^1)\rho
 1: set \mathfrak{R} := \{y_1 := 1_S\}, M := 1
                                                                                                        [initialise the list of \mathcal{R}-class reps]
 2: set (S^1)\rho := \{\beta_1 := (1_S)\rho\}, N := 1
                                                                                                                                   [initialise (S)\rho]
 3: find (S^1)\lambda = (1_S)\lambda \cdot S
                                                                                                                                      [Algorithm 1]
 4: find representatives (z_1)\lambda, \ldots, (z_r)\lambda of the s.c.c.s of (S^1)\lambda
                                                                                                        [standard graph theory algorithm]
 5: find the groups S_{z_i} for i \in \{1, \ldots, r\}
                                                                                                                                      [Algorithm 4]
                                                                       [loop over: existing \mathscr{R}-representatives, generators of S^1]
 6: for y_i \in \Re, j \in \{1, ..., m\} do
          find n \in \{1, ..., r\} such that (z_n)\lambda \sim (x_j y_i)\lambda
 7:
          find u \in S^1 such that (z_n)\lambda \cdot u = (x_iy_i)\lambda
                                                                                                                                      [Algorithm 2]
 8:
          find \overline{u} \in U^1 such that (x_i y_i) \lambda \cdot \overline{u} = (z_n) \lambda and z_n u \overline{u} = z_n
                                                                                                                              [Assumption (III)]
 9:
          if (x_iy_i)\rho \not\in (S^1)\rho then
                                                                                                        [x_iy_i\overline{u} \text{ is a new } \mathcal{R}\text{-representative}]
10:
               N := N + 1, \ \beta_N := (x_j y_i \overline{u}) \rho = x_j \cdot (y_i) \rho, \ (S^1) \rho \leftarrow (S^1) \rho \cup \{\beta_N\}
                                                                                                                         [add (x_i y_i) \rho \ to (S^1) \rho]
11:
               v_N := i, w_N := j, g_{i,j} := N
                                                                                [update the Schreier tree and orbit graph of (S^1)\rho]
12:
          else if (x_j y_i) \rho = \beta_k \in (S^1) \rho then
13:
               g_{l,i} := k where l is such that (y_i)\rho = \beta_l;
                                                                                                          [update the orbit graph of (S^1)\rho]
14:
               for y_l \in \mathfrak{R} with (y_l)\rho = (x_jy_i)\rho and (y_l)\lambda = (z_n)\lambda do
15:
                                                                                                                                         [x_i y_i \mathcal{R}^S y_l]
                    if (y_l'x_jy_i\overline{u})\mu_{z_n} \in S_{z_n} = S_{y_l} then
16:
                         G_{i,j} := l
                                                                                                               [update the orbit graph of \mathfrak{R}]
17:
                         go to line 6
18:
                    end if
19:
               end for
20:
          end if
21:
          M := M + 1, y_M := x_j y_i \overline{u}, \mathfrak{R} \leftarrow \mathfrak{R} \cup \{y_M\}
                                                                                                                                 [add \ x_i y_i \overline{u} \ to \ \mathfrak{R}]
22:
          V_M := i, W_M := j, K_M := \text{the index of } (x_j y_i) \lambda \text{ in } (S^1) \lambda
                                                                                                             [update the Schreier tree of \mathfrak{R}]
23:
24:
          G_{i,j} := M
                                                                                                               [update the orbit graph of \mathfrak{R}]
25: end for
26: return the following:
         • the \mathscr{R}-representatives: \mathfrak{R}
         • the Schreier tree for \mathfrak{R}: (V_2, \ldots, V_M, W_2, \ldots, W_M)
         • the orbit graph of \Re: \{G_{i,j} : i = 1, ..., M, j = 1, ..., m\}
         • the set (S^1)\rho
         • the Schreier tree for (S^1)\rho: (v_2,\ldots,v_N,w_2,\ldots,w_N)
         • the orbit graph of (S^1)\rho: \{g_{i,j}: i=1,\ldots,N, j=1,\ldots,m\}
         • the parameters: (K_1, \ldots, K_M)
```

#### Testing membership in a semigroup

Algorithm 12 Test membership in a semigroup

In Algorithm 12, we give a procedure for testing if an element of U belongs to S. This can be easily modified to return the  $\mathcal{R}$ -class representative in S of an arbitrary element of U, if it exists (simply return the element x in line 12). We require such an algorithm when it comes to factorising elements of S over the generators in Algorithm 13.

```
Input: S^1 := \langle X \rangle where X := \{x_1, \dots, x_m\} and \overline{y \in U^1}
Output: true or false
 1: let (S^1)\lambda = (1_S)\lambda \cdot S and (S^1)\rho = S \cdot (1_S)\rho
                                                                                                                                         [Algorithm 1]
 2: if (y)\lambda \notin (S^1)\lambda or (y)\rho \notin (S^1)\rho then
                                                                                                                                                 [y \not\in S^1]
 3:
           return false
 4: end if
 5: let (z_1)\lambda, \ldots, (z_r)\lambda be representatives of the s.c.c.s of (S^1)\lambda
 6: let \mathfrak{R} denote the \mathscr{R}-class representatives of S^1
                                                                                                                                       [Algorithm 11]
 7: find n \in \{1, \ldots, r\} such that (z_n)\lambda \sim (y)\lambda
 8: find u \in S^1 such that (z_n)\lambda \cdot u = (y)\lambda
                                                                                                                                         [Algorithm 2]
 9: find \overline{u} \in U^1 such that (y)\lambda \cdot \overline{u} = (z_n)\lambda and yu\overline{u} = y
                                                                                                                                 [Assumption (III)]
```

 $[x\mathscr{R}^{S^1}y]$ 

14: end for 15: return false

if  $(x'y\overline{u})\mu_x \in S_{z_n} = S_x$  then

return true

end if

11:

12:

13:

10: for  $x \in \mathfrak{R}$  such that  $(x)\lambda = (z_n)\lambda$  and  $(x)\rho = (y)\rho$  do

#### Factorising elements over the generators

In this part of the paper, we describe how to factorise an element of S as a product of the generators of S using the output of Algorithm 11.

Corollary 3.18(a) says that

$$R_x^S = \{xsu : s \in \operatorname{Stab}_S(L_x^U), u \in S^1, (xu)\lambda \sim \lambda(x)\}.$$

Suppose that  $y \in S$  is arbitrary and that  $\mathfrak{R}$  denotes a set of  $\mathscr{R}$ -class representatives of S. If we can write y = xsu, where  $x \in \mathfrak{R}$  such that  $x\mathscr{R}^S y$ ,  $s \in \operatorname{Stab}_S(L_x^U)$ , and  $u \in S^1$  such that  $(xu)\lambda = (y)\lambda$ , then it suffices to factorise each of x, s, and u individually.

The element  $x \in \Re$  can be found using the alternate version of Algorithm 12 mentioned above. Algorithm 2, applied to  $(S)\lambda$ , can be used to find u such that  $(xu)\lambda = (y)\lambda$ .

Suppose that  $\overline{u} \in U^1$  is such that  $(y)\lambda \cdot \overline{u} = (x)\lambda$  and  $xu\overline{u} = x$ . Since  $(xs)\lambda = (x)\lambda$ , it follows from Lemma 3.6 that  $xsu\overline{u} = xs$ . If  $x' \in U$  is any element such that xx'x = x, then from Proposition 3.9(a), x is the identity of the group  $L_x^U \cap R_x^S$  under multiplication \* defined by a\*b = ax'b. In particular,  $xx'y\overline{u} = y\overline{u}$  since  $y\overline{u} \in L_x^U \cap R_x^S$ . This implies that

$$xx'y\overline{u}=y\overline{u}=xsu\overline{u}=xs$$

and so, by Lemma 3.6,  $(x'y\overline{u})\mu_x = (s)\mu_x$ . Since  $(y\overline{u})\lambda = (x)\lambda$  and  $(y\overline{u})\rho = (x)\rho$ , by Assumption (IV), we can compute  $(x'y\overline{u})\mu_x$ . Thus for any  $y \in S$  we can determine  $x, s, u \in S$  (as above) such that y = xsu.

We still require a factorisation of  $x, s, u \in S$  over the generators of S (here we suppose that  $u \neq 1_S$ ). Tracing the Schreier tree of  $\mathfrak{R}$  returned by Algorithm 11 and using the parameter  $K_i$ , we can factorise  $x \in \mathfrak{R}$ ; more details are in Algorithm 13. We may factorise u over the generators of S using Algorithm 2 applied to the s.c.c. of  $(x)\lambda$  in  $(S)\lambda$ .

Any algorithm for factorising elements of a group can be used to factorise  $(s)\mu_x$  as a product of the generators of  $S_x$  given by Algorithm 4. For example, in a group with a faithful action on some set, such a factorisation can be obtained from a stabiliser chain, produced using the Schreier-Sims algorithm. The generators of  $S_x$  are of the form  $(u_i x_k \overline{u_j})\mu_x$ , where  $u_i$  is obtained by tracing the Schreier tree of  $(S)\lambda$ ,  $x_k$  is one of the generators of S, and  $\overline{u_j}$  is obtained using Assumption (III). Therefore to factorise s over the generators of S, it suffices to factorise those  $\overline{u_j}$  where  $\overline{u_j} \neq 1_S$  over the generators of S.

Suppose that  $\{\alpha_1, \ldots, \alpha_K\}$  is a s.c.c. of  $(S)\lambda$  for some  $K \in \mathbb{N}$ . Then from the orbit graph of  $(S)\lambda$  we can find

$$a_2,\ldots,a_K,b_2,\ldots,b_K$$

such that

$$\alpha_j \cdot x_{a_i} = \alpha_{b_i}$$

and  $b_j < j$  for all j > 1. In other words,  $(a_2, \ldots, a_K, b_2, \ldots, b_K)$  describes a spanning tree, rooted at  $\alpha_1$ , for the component of the orbit graph of  $(S)\lambda$ , whose edges have the opposite orientation to those in the usual Schreier tree. We refer to  $(a_2, \ldots, a_K, b_2, \ldots, b_K)$  as a reverse Schreier tree.

It follows that for any  $i \in \{1, ..., K\}$  we can use an analogue of Algorithm 2 to obtain  $v \in S^1$  such that  $\alpha_j \cdot v = \alpha_1$ . However, if  $u \in S^1$  is obtained using Algorithm 2 such that  $\alpha_1 \cdot u = \alpha_j$  and  $z \in S$  is such that  $(z)\lambda = \alpha_1$ , then it is possible that  $(uv)\mu_z \neq (1_U)\mu_z$ . So, if  $w \in \operatorname{Stab}_S(L_z^U)$  is such that  $(w)\mu_z = ((uv)\mu_z)^{-1}$ , then  $\alpha_j \cdot vw = \alpha_1 (uvw)\mu_z = (1_U)\mu_z$ . We have factorisations of v, since it was obtained by tracing the reverse Schreier tree, and w, since it can be given as a power of uv (u and v are factorised), over the generators of u. Hence u is factorized over the generators of u, and it has the properties required of u from above.

We note that all the information required to factorise any element of S is returned by Algorithm 11 except the factorisation of  $(s)\mu_x$  in  $S_x$ . So, Algorithm 13 is just concerned with putting this information together. There is no guarantee that the word produced by Algorithm 13 is of minimal length.

```
Algorithm 13 Factorise an element over the generators
```

```
Input: S := \langle X \rangle where X := \{x_1, \dots, x_m\} and s \in S
Output: a word in the generators X equal to s
 1: let \theta: X^+ \longrightarrow S be the unique homomorphism extending the inclusion of X in S
 2: suppose that (S)\lambda := \{(z_1)\lambda, \dots, (z_K)\lambda\} for some z_1, \dots, z_K \in S
                                                                                                                                 [Algorithm 1]
 3: let \mathfrak{R} := \{y_1, \dots, y_r\} be the \mathscr{R}-representatives of S
                                                                                                                                [Algorithm 11]
 4: let (V_2, \ldots, V_r, W_2, \ldots, W_r) be the Schreier tree for \mathfrak R
                                                                                                                                [Algorithm 11]
 5: let (K_1, \ldots, K_r) denote the additional parameter return from Algorithm 11
 6: find y_i \in \Re such that y_i \mathcal{R}^S s
                                                                                                [the modified version of Algorithm 7]
 7: find a word \omega_1 \in X^+ such that (\omega_1)\theta = u \in S where (y_i)\lambda \cdot u = (s)\lambda [factorise u using Algorithm 2]
 8: find \overline{u} \in U^1 such that (s)\lambda \cdot \overline{u} = (y_i)\lambda, and y_i u\overline{u} = y_i
                                                                                                                          [Assumption (III)]
 9: compute (y_i's\overline{u})\mu_{y_i} \in S_{y_i}
                                                                                                                          [Assumption (IV)]
10: find a word \omega_2 \in X^+ such that (\omega_2)\theta = y_i' s \overline{u}
                                                                                                                                   [factorise \ s]
                     [factorise (y_i's\overline{u})\mu_{y_i} over the generators of S_{y_i} and then factorise these generators over X
11: \omega_3 := \varepsilon (the empty word), j = i
                                                                                         [trace the Schreier tree of \Re, factorise y_i]
12: while j > 1 do
          find \beta \in X^+ such that (z_{K_i})\lambda \cdot (\beta)\theta = (y_j)\lambda
13:
          set \omega_3 := x_{W_i} \omega_3 \beta and j := V_i
                                                                                                              [(\omega_3)\theta = y_i][(\omega_3\omega_2\omega_1)\theta = y_iy_i's\overline{u}u = s]
15: end while
16: return \omega_3\omega_2\omega_1
```

#### The partial order of the *I*-classes

Recall that there is a partial order  $\leq_{\mathscr{J}}$  on the  $\mathscr{J}$ -classes of a semigroup S, which is induced by containment of principal two-sided ideals. If S is finite, then  $\mathscr{J} = \mathscr{D}$ , and so  $\leq_{\mathscr{J}}$  is also a partial order on the  $\mathscr{D}$ -classes of S. More precisely, if A and B are  $\mathscr{J}$ -classes of S, then we write  $A \leq_{\mathscr{J}} B$  if  $S^1 a S^1 \subseteq S^1 b S^1$  for any (and every)  $a \in A$  and  $b \in B$ .

The penultimate algorithm in this paper allows us to calculate the partial order of the  $\mathscr{D}$ -classes of S. This algorithm is based on [24, Algorithm Z] and the following proposition which appears as Proposition 5.1 in [20]. The principal differences between our algorithm and Algorithm Z in [24] are that our algorithm applies to classes of semigroups other than transformation semigroups, and it takes advantage of information already determined in Algorithm 11.

**Proposition 5.2** (cf. Proposition 5.1 in [20]). Let S be a finite semigroup generated by a subset X, and let D be a  $\mathscr{D}$ -class of S. If R and L are representatives of the  $\mathscr{R}$ - and  $\mathscr{L}$ -classes in D, then the set  $XR \cup LX$  contains representatives for the  $\mathscr{D}$ -classes immediately below D under  $\leq_{\mathscr{J}}$ .

We described above that we find the  $\mathscr{D}$ -classes (or representatives for the  $\mathscr{D}$ -classes) of S by finding the s.c.c.s of the orbit graph of the  $\mathscr{R}$ -class representatives  $\mathfrak{R}$  of S. Thus, after performing Algorithm 11 and finding the s.c.c.s of the orbit graph of  $\mathfrak{R}$ , we know both the  $\mathscr{D}$ -classes of S and the  $\mathscr{R}$ -classes contained in the  $\mathscr{D}$ -classes. In Algorithm 11, we obtain the  $\mathscr{R}$ -class representatives of S by left multiplying existing representatives. Therefore we have already found the information required to determine the set XR in Proposition 5.2. In particular, we do not have to multiply the  $\mathscr{R}$ -class representatives of a  $\mathscr{D}$ -class by each of the generators in Algorithm 14, or determine which  $\mathscr{D}$ -classes correspond to the elements of XR.

In Algorithm 14 we represent the partial order of the  $\mathscr{D}$ -classes  $D_1, \ldots, D_n$  of S as  $P_1, \ldots, P_n$  where  $P_i$  contains the indices of the  $\mathscr{D}$ -classes immediately below  $D_i$  (and maybe some more). The elements of  $P_i$  are obtained by applying Proposition 5.2 to  $D_i$ .

```
Algorithm 14 The partial order of the D-classes of a semigroup
```

```
Input: S := \langle X \rangle where X := \{x_1, \dots, x_m\}
Output: the partial order of the \mathscr{D}-classes of S
 1: let \Re denote the \mathscr{R}-representatives of S
                                                                                                                              [Algorithm 11]
 2: let \Gamma := \{G_{i,j} : 1 \le i \le |\mathfrak{R}|, 1 \le j \le m\} be the orbit graph of \mathfrak{R}
                                                                                                                              [Algorithm 11]
 3: find the \mathscr{D}-classes D_1, \ldots, D_n of S and D_i \cap \mathfrak{R} for all i
                                                                                                                     [find the s.c.c.s of \Gamma]
 4: let P_i := \emptyset for all i
                                                                                                            [initialise the partial order]
     for i \in \{1, ..., n\} do
                                                                                                                [loop over the \mathscr{D}-classes]
          for y_j \in \mathfrak{R} \cap D_i, x_k \in X do
                                                                        [loop over: \mathcal{R}-classes of the \mathcal{D}-class, generators of S
 6:
              find l \in \{1, \ldots, n\} such that y_{G_{i,k}} \in D_l
                                                                                                               [use the orbit graph of \mathfrak{R}]
 7:
               P_i \leftarrow P_i \cup \{l\}
 8:
          end for
 9:
          find the \mathcal{L}-class representatives \mathfrak{L} in D_i
                                                                                 the analogue of Algorithm 6, Proposition 3.22
10:
          for z_i \in \mathfrak{L}, x_k \in X do
                                                                       [loop over: \mathcal{L}-classes of the \mathcal{D}-class, generators of S]
11:
              find l such that z_i x_k \in D_l
                                                               [use the analogue of Algorithm 7 to find the \mathscr{R}-rep. of z_i x_k]
12:
               P_i \leftarrow P_i \cup \{l\}
13:
          end for
14:
15: end for
16: return P_1, \ldots, P_n.
```

### The closure of a semigroup and some elements

The final algorithm (Algorithm 15) we present describes a method for taking the closure of a subsemigroup S of U with a set V of elements of U. The purpose of this algorithm is to reuse whatever information is known about S to make subsequent computations involving  $\langle S, V \rangle$  more efficient.

At the beginning of this procedure we form  $(\langle S, V \rangle)\lambda$  by adding the new generators V to  $(S)\lambda$ . This can be achieved in GAP using the AddGeneratorsToOrbit function from the ORB package [31], which has the advantage that the existing information in  $(S)\lambda$  is not recomputed. The orbit  $(S)\lambda$  is extended by a breadth-first enumeration with the new generators without reapplying the old generators to existing values in  $(S)\lambda$ .

## 5.6 Optimizations for regular and inverse semigroups

Several of the algorithms presented in this section become more straightforward if it is known a priori that the subsemigroup S of U is regular. For example, the  $\mathscr{R}$ -classes of S are just the  $\mathscr{R}$ -classes of U intersected with S, and so are in 1-1 correspondence with  $(S)\rho$ . The algorithms become simpler still under the assumption that U is an inverse semigroup since in this case we may define  $(x)\rho = (x^{-1})\lambda$  for all  $x \in U$  and so it is unnecessary to calculate  $(S)\rho$  and  $(S)\lambda$  separately.

The first algorithm where an advantage can be seen is Algorithm 6. Suppose that S is a regular subsemigroup of U, and  $x \in S$ . Then, by Corollary 3.12,  $S_x = ({}_xS)\Psi$  and so it is no longer necessary to compute  $({}_xS)\Psi$  or the cosets of  $S_x \cap ({}_xS)\Psi$  in  $({}_xS)\Psi$  in Algorithm 6. If U is an inverse semigroup with unary operation  $^{-1}: x \mapsto x^{-1}$ , then Algorithm 6 becomes simpler still. In this case, it is unnecessary to find the s.c.c. of  $(x)\rho$ . This follows from the observation that we may take  $(x)\rho = (x^{-1})\lambda$ , which implies that  $(S)\rho = (S)\lambda$  and

$$u^{-1} \cdot (x^{-1})\rho = (u^{-1}x^{-1})\rho = (xu)\lambda = (x)\lambda \cdot u.$$

### Algorithm 15 The closure of a semigroup and some elements

```
Input: S := \langle X \rangle where X := \{x_1, \dots, x_m\} and any existing data structures for S, and V \subseteq U
Output: data structures for the \mathscr{R}-classes with representatives \Re in \langle S, V \rangle, the set (\langle S, V \rangle)\rho, and Schreier
     trees and orbit graphs for \Re and (\langle S, V \rangle)\rho
 1: set \widehat{\mathfrak{R}} := \{\widehat{y_1}, \dots, \widehat{y_k}\} to be the \mathscr{R}-class representatives of S
                                                                                                                                              [Algorithm 11]
 2: set \Re := \{y_1 := \widehat{y_1}\}, M := 1
                                                                                                                 [initialise the list of new \mathcal{R}-reps]
 3: define (1)\iota = 1
                                                                                              [keep track of indices of old and new \mathcal{R}-reps]
 4: set (\langle S, V \rangle) \rho := (S) \rho, N := |(S) \rho|
                                                                                                                                      [initialise (\langle S, V \rangle)\rho]
 5: set the Schreier tree of (\langle S, V \rangle)\rho to be that of (S)\rho
                                                                                                       [initialise the Schreier tree of (\langle S, V \rangle)\rho]
 6: set the orbit graph of (\langle S, V \rangle)\rho to be that of (S)\rho
                                                                                                               [initialise orbit graph of (\langle S, V \rangle)\rho]
 7: extend (S)\lambda to (\langle S, V \rangle)\lambda
                                                                                                                                      [as described above]
 8: find representatives (z_1)\lambda, \ldots, (z_r)\lambda of the s.c.c.s of (\langle S, V \rangle)\lambda
 9: find the groups S_{z_i} for i \in \{1, \ldots, r\}
                                                                                                                                                [Algorithm 4]
10: for x_j \widehat{y_k} v = \widehat{y_i} \in \mathfrak{R} do
                                                                                 [j, k \text{ are obtained from the Schreier tree for } \widehat{\mathfrak{R}}, k < i]
           find n \in \{1, ..., r\} such that (z_n)\lambda \sim (\widehat{y_i})\lambda (in (\langle S, V \rangle)\lambda)
11:
           find u \in S^1 such that (z_n)\lambda \cdot u = (\widehat{y_i})\lambda
                                                                                                                                                [Algorithm 2]
12:
           find \overline{u} \in U^1 such that (\widehat{y}_i)\lambda \cdot \overline{u} = (z_n)\lambda and z_n u\overline{u} = z_n
                                                                                                                                        [Assumption (III)]
13:
           for y_l \in \mathfrak{R} with (y_l)\rho = (\widehat{y_i})\rho and (y_l)\lambda = (z_n)\lambda do
14:
                if (y_l'\widehat{y_l}\overline{u})\mu_{z_n}\in S_{z_n}=S_{y_l} then
                                                                                                                                                  [\widehat{y_i}\mathscr{R}^{\langle S,V\rangle}y_l]
15:
                      G_{(k)\iota,j} := l
                                                                                                                        [update the orbit graph of \mathfrak{R}]
16:
                      define (i)\iota := l
                                                                                                               [the old index i is the new index l]
17:
                      go to line 10
18:
                end if
19:
20:
           end for
           M := M + 1, y_M := \widehat{y_i}\overline{u}, \mathfrak{R} := \mathfrak{R} \cup \{y_M\}
                                                                                                                                              [add \ \widehat{y_i}\overline{u} \ to \ \mathfrak{R}]
21:
           V_M := (k)\iota, W_M := j
                                                                                                                     [update the Schreier tree of \mathfrak{R}]
22:
           K_M := \widehat{K_i}
                                                            [the index of (x_i \hat{y_k}) \lambda in (\langle S, V \rangle) \lambda from Algorithm 11 applied to S]
23:
           G_{(k)\iota,j} := M
                                                                                                                        [update the orbit graph of \mathfrak{R}]
24:
           (i)\iota := M
                                                                                                            [the old index i is the new index M]
25:
26: end for
27: return apply Algorithm 11 to the data structures for \langle S, V \rangle determined so far, and return the output
```

Similar simplifications can be made in Algorithm 8.

The unary operation in the definition of U means that given an inverse subsemigroup S of U and  $x \in S$ , that we can find  $x^{-1}$  without reference to S. Or put differently, the inverse of x is the same in every inverse subsemigroup of U. For example, the symmetric inverse monoid has this property, but the full transformation monoid does not. More precisely, there exist distinct inverse subsemigroups S and T of  $T_n$  and  $x \in S \cap T$  such that the inverse of x in S is distinct from the inverse of x in T.

As noted above, in a regular subsemigroup S of U, the  $\mathscr{R}$ -classes are in 1-1 correspondence with the elements of  $(S)\rho$  and the  $\mathscr{L}$ -classes are in 1-1 correspondence with  $(S)\lambda$ . It follows that the search for  $\mathscr{R}$ -class representatives in Algorithm 11 is redundant in this case. Hence the  $\mathscr{R}$ -classes,  $\mathscr{L}$ -classes, size, and elements, of a regular subsemigroup can be determined from  $(S)\lambda$  and  $(S)\rho$  using Algorithms 2 and 4 alone. The  $\mathscr{D}$ -classes of a regular subsemigroup S are then in 1-1 correspondence with the s.c.c.s of  $(S)\lambda$  (or  $(S)\rho$ ). In the case that U is an inverse semigroup, it suffices to calculate either  $(S)\lambda$  or  $(S)\rho$ , making these computations simpler still. The remaining algorithms in Subsection 5.5 can also be modified to take advantage of these observations, but due to considerations of space, we do not go into the details here.

The simplified algorithms alluded to in this section have been fully implemented in the SEMIGROUPS package for GAP; see [29].

# 6 Examples

In this section we present some examples to illustrate the algorithms from the previous section.

One of the examples is that of a semigroup of partial permutations. Similar to permutations, a partial permutation can be expressed as a union of the components of its action. Any component of the action of a partial permutation f is either a permutation with a single cycle, or a chain  $[i\ (i)f\ (i)f^2\ ...\ (i)f^r]$  where  $i\in \text{dom}(f)\setminus \text{im}(f)$  and  $(i)f^r\in \text{im}(f)\setminus \text{dom}(f)$ , for some r>1. For the sake of brevity, we will use disjoint component notation when writing a specific partial permutation f, i.e. we write f as a juxtaposition of disjoint cycles and chains. For example,

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 4 & - & 2 & 6 & - \end{pmatrix} = \begin{bmatrix} 1 & 5 & 6 \end{bmatrix} (2 & 4).$$

We include fixed points in the disjoint component notation for a partial permutation f so that it is possible to deduce the domain and image of f from the notation, and so that the notation for f is unique (up to the order of the components, and the order of elements in a cycle); see [25] for further details.

Throughout this section, we will denote by S the submonoid of the symmetric inverse monoid on  $\{1, \ldots, 9\}$  generated by

$$x_1 = (1\ 4)(2\ 6)(3\ 8)(5)(7)(9), \quad x_2 = (1\ 5\ 4\ 2\ 7\ 6)(3\ 9\ 8) \quad x_3 = (2\ 5\ 6), \quad x_4 = (1\ 3\ 2),$$
 (6.1)

by T the submonoid of the full transformation monoid on  $\{1,\ldots,5\}$  generated by

$$x_1 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 2 & 4 & 5 \end{pmatrix}, \quad x_2 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix}, \quad x_3 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 3 & 2 & 2 \end{pmatrix}, \tag{6.2}$$

and we will use the notation of Sections 4.1 and 4.2.

Since both S and T are submonoids, adjoining an identity to either would be redundant, and so in the interest of saving space, we will not do this.

## Components of the action

Applying Algorithm 1 to S and  $\alpha_1 = (x_1)\lambda = \{1, \dots, 9\}$ , we obtain:

$$(S)\lambda = \{ \begin{array}{cccc} \alpha_1 = \{1,\ldots,9\}, & \alpha_2 = \{2,5,6\}, & \alpha_3 = \{1,2,3\}, & \alpha_4 = \{1,4,7\}, \\ \alpha_5 = \{1\}, & \alpha_6 = \{4,6,8\}, & \alpha_7 = \{5,7,9\}, & \alpha_8 = \{5\}, \\ \alpha_9 = \varnothing, & \alpha_{10} = \{3\}, & \alpha_{11} = \{4\}, & \alpha_{12} = \{2\}, \\ \alpha_{13} = \{6\}, & \alpha_{14} = \{8\}, & \alpha_{15} = \{9\}, & \alpha_{16} = \{7\} \end{array} \}.$$




Figure 3: The orbit graph of  $(S)\lambda$  with loops and (all but one of the) edges to  $\alpha_9$  omitted, and the Schreier tree indicated by solid edges.

The Schreier tree is:

and the orbit graph is:

i	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$g_{i,1}$	1	2	6	4	11	3	7	8	9	14	5	13	12	10	15	16 13 . 9 9
$g_{i,2}$	1	4	7	2	8	3	6	11	9	15	12	16	5	10	14	13 .
$g_{i,3}$	2	2	8	9	9	12	13	13	9	9	9	8	12	9	9	9
$g_{i,4}$	3	5	3	10	10	9	9	9	9	12	9	5	9	9	9	9

Diagrams of the Schreier tree and orbit graphs of  $(S)\lambda$  can be found in Figure 3. The strongly connected components of  $(S)\lambda$  are:

$$\{\{\alpha_1\}, \{\alpha_2, \alpha_4\}, \{\alpha_3, \alpha_6, \alpha_7\}, \{\alpha_5, \alpha_8, \alpha_{10}, \alpha_{11}, \alpha_{12}, \alpha_{13}, \alpha_{14}, \alpha_{15}, \alpha_{16}\}, \{\alpha_9\}\}.$$

From the Schreier tree, we deduce that

$$\alpha_1 = (x_1)\lambda, \quad \alpha_2 = (x_1x_3)\lambda, \quad \alpha_3 = (x_1x_4)\lambda, \quad \alpha_5 = (x_1x_3x_4)\lambda, \quad \text{and} \quad \alpha_9 = (x_1x_3x_2x_3)\lambda.$$

In this case, we set  $\bar{s} = s^{-1}$  in Assumption (III). Using Algorithms 2 and 4, after removing redundant generators, we obtain Schreier generators for the stabilisers of  $x_1, x_1x_3, x_1x_4, x_1x_3x_4$ , and  $x_1x_3x_2x_3$ :

$$\begin{array}{lclcl} S_{x_1} & = & \langle x_1, x_2 \rangle \cong D_{12}, & S_{x_1x_3} & = & \langle (2\ 6), (2\ 6\ 5) \rangle \cong \operatorname{Sym}(\{2,5,6\}), \\ S_{x_1x_4} & = & \langle (1\ 3\ 2), (1\ 2) \rangle \cong \operatorname{Sym}(\{1,2,3\}), & S_{x_1x_3x_4} & = & \operatorname{Sym}(\{1\}) \cong \mathbf{1}, \\ S_{x_1x_3x_2x_3} & = & \operatorname{Sym}(\varnothing) \cong \mathbf{1}, \end{array}$$




Figure 4: The orbit graph of  $(T)\lambda$  with loops omitted, and the Schreier tree indicated by solid edges.

where 1 denotes the trivial group.

Applying Algorithm 1 to T (defined in (6.2)) and  $(x_1)\lambda = \{1, 2, 3, 4, 5\}$ , we obtain:

the Schreier tree is:

and the orbit graph is:

Diagrams of the Schreier tree and orbit graphs of  $(T)\lambda$  can be found in Figure 4.

The strongly connected components of  $(T)\lambda$  are:

$$\{\{\alpha_1\}, \{\alpha_2\}, \{\alpha_3, \alpha_4, \alpha_5\}, \{\alpha_6, \alpha_7, \alpha_8\}\}.$$

From the Schreier tree for  $(T)\lambda$  and Algorithms 2,

$$\alpha_1 = (x_1)\lambda, \quad \alpha_2 = (x_1x_3)\lambda, \quad \alpha_3 = (x_1x_3^2)\lambda, \quad \text{and} \quad \alpha_6 = (x_1x_3^2x_1x_2x_3)\lambda.$$

Using Algorithm 4, after removing redundant generators, we obtain Schreier generators for the stabilisers  $T_y$  where  $y = x_1, x_1x_3, x_1x_3^2, x_1x_3^2x_1x_2x_3$ :

$$\begin{array}{lcl} T_{x_1} & = & \langle x_1, x_2 \rangle \cong D_{12}, & T_{x_1 x_3} & = & \langle (2\ 3), (1\ 2\ 3) \rangle \cong \operatorname{Sym}(\{1, 2, 3\}), \\ T_{x_1 x_3^2} & = & \langle (1\ 3) \rangle \cong \operatorname{Sym}(\{1, 3\}), & T_{x_1 x_3^2 x_1 x_2 x_3} & = & \operatorname{Sym}(\{3\}) \cong \mathbf{1}, \end{array}$$

where 1 denotes the trivial group.

## Individual Green's classes

Let S be partial permutation semigroup defined in (6.1). The s.c.c. of  $(x_1x_3x_4)\lambda$  contains 9 values:

$$\{\alpha_5 = \{1\}, \alpha_8 = \{5\}, \alpha_{10} = \{3\}, \alpha_{11} = \{4\}, \alpha_{12} = \{2\}, \alpha_{13} = \{6\}, \alpha_{14} = \{8\}, \alpha_{15} = \{9\}, \alpha_{16} = \{7\}\},$$

and  $|S_{x_1x_3x_4}| = 1$ . It follows, by Corollary 3.13(b), that the size of the  $\mathscr{R}$ -class of  $x_1x_3x_4$  in S is  $9 \cdot 1 = 9$ . One choice for the Schreier tree (rooted at  $\alpha_5 = (x_1x_3x_4)\lambda$ ) of the s.c.c. of  $(x_1x_3x_4)\lambda$  is:

A diagram of the Schreier tree can be found in Figure 5.

Since  $x_1x_3x_4 = [2 \ 1]$ , using the Schreier tree and Algorithm 5, the elements of  $R_{x_1x_3x_4}^S$  are:




Figure 5: Schreier trees for the strongly connected component of  $(x_1x_3x_4)\lambda$  in  $(S)\lambda$ , and for  $(x_1x_3x_4)\rho$  in

Since  $x_1x_3x_4^3$  is an idempotent, it also follows that  $R_{x_1x_3x_3}^S$  is a regular  $\mathscr{R}$ -class. We will calculate the  $\mathscr{R}$ -classes in  $D_{x_1x_3x_4}^S$  using Algorithm 6. Since  $(x_1x_3x_4)\rho=\{2\}$ , it follows immediately that  $x_1x_3x_4S=\{\mathrm{id}_{\{2\}}\}$  is trivial. Set  $x=x_1x_3x_4$  and  $x'=x^{-1}=[1\ 2]$ . The embedding  $\Psi: {}_{x}S \longrightarrow U_{x}$  (from Proposition 3.11(a)) defined by

$$((s)\nu_x)\Psi = (x'sx)\mu_x$$

maps  $\operatorname{id}_{\{2\}}$  to  $\operatorname{id}_{\{1\}}$ . Hence  $S_x \cap (_xS)\Psi = S_x = \{\operatorname{id}_{\{1\}}\}$ . It also follows from Proposition 3.21 that the  $\mathscr{R}^S$ -class representatives of  $D_x^S$  are in 1-1 correspondence with the s.c.c. of  $(x)\rho = \{2\}$ . Using the left analogue of Algorithm 1, we obtain

$$S \cdot (x)\rho = \{ \beta_1 = \{2\}, \beta_2 = \{6\}, \beta_3 = \{4\}, \beta_4 = \{3\}, \beta_5 = \{7\}, \beta_6 = \{5\}, \beta_7 = \varnothing, \beta_8 = \{1\}, \beta_9 = \{8\}, \beta_{10} = \{9\} \}$$

with Schreier tree:

A diagram of the Schreier tree can be found in Figure 5. Hence the  $\mathscr{R}^S$ -class representatives of  $D_x^S$  are:

Since the number of  $\mathscr{R}^S$ -classes in  $D_x^S$  is 9 and each  $\mathscr{R}^S$ -class has size 9, it follows that  $|D_x^S| = 81$ .

We will demonstrate how to use Algorithm 7 to check if the partial permutation  $y = \begin{bmatrix} 1 & 5 \end{bmatrix} \begin{bmatrix} 2 & 7 \end{bmatrix} \begin{bmatrix} 3 & 9 \end{bmatrix}$ is  $\mathscr{R}^S$ -related to either of the generators  $x_3$  and  $x_4$  of S. Since  $(y)\rho = \{1,2,3\}$  and  $(x_3)\rho = \{2,5,6\}$ , it follows that  $(y, x_3) \notin \mathcal{R}^S$ . However,  $(y)\rho = (x_4)\rho$  and

$$(y)\lambda = \{5, 7, 9\} = \alpha_7 \sim \alpha_3 = \{1, 2, 3\} = (x_4)\lambda.$$

Tracing the Schreier tree of  $(S)\lambda$  from  $\alpha_3$  to  $\alpha_7$ , we obtain  $u=x_2$  such that  $(x_4)\lambda \cdot u=(y)\lambda$ . It follows that  $\overline{u}=x_2^{-1}$  has the property that  $(y)\lambda \cdot \overline{u}=(x)\lambda$ . Also setting  $x_4'=x_4^{-1}$ , it follows that  $y\in R_x^S$  since

$$(x_4'y\overline{u})\mu_x = (x_4^{-1}yx_2^{-1})\mu_x = (1\ 2\ 3) \in S_{x_4} = S_{x_1x_4} = Sym(\{1, 2, 3\}).$$

#### The main algorithm

We now determine the global structure of the transformation semigroup T defined in (6.2). We will do the same thing for the partial permutation semigroup S defined in (6.1) in the next subsection.




Figure 6: The orbit graph of  $(T)\rho$  and  $\Re$  with loops omitted, and the Schreier tree indicated by solid edges.

If x is a transformation of degree  $n \in \mathbb{N}$ , then the kernel  $\ker(x)$  of x is a partition of  $\{1, \ldots, n\}$ . If the classes of  $\ker(x)$  are  $A_1, A_2, \ldots, A_r$ , for some r, then to avoid writing too many brackets we write  $\ker(x) = \{A_1 | \cdots | A_r\}$ .

Applying Algorithm 11 to T defined in (6.2), we find that the  $\mathcal{R}$ -class representatives of S are:

$$y_1 = 1_T,$$
  $y_2 = x_3,$   $y_3 = x_2x_3,$   $y_4 = x_3^2,$   $y_5 = x_1x_2x_3,$   $y_6 = x_3x_2x_3,$   $y_7 = x_2x_3^2,$   $y_8 = x_3x_1x_2x_3,$   $y_9 = (x_2x_3)^2,$   $y_{10} = x_1x_2x_3^2,$   $y_{11} = x_3^2x_1x_2x_3,$   $y_{12} = x_1(x_2x_3)^2,$ 

and that

The Schreier tree of the orbit graphs of  $\mathfrak{R}$  and  $(T)\rho$  are both equal:

and the orbit graph is:

A diagrams of the Schreier tree and orbit graph can be found in Figure 6.

It is coincidentally the case that the  $\mathscr{R}$ -class representatives of T are obtained by left multiplying previous  $\mathscr{R}$ -class representatives by a generator. In other words, the u and  $\overline{u}$  in line 8 and 9 of Algorithm 11 are just the identity of T in this example. Hence the additional parameters returned by Algorithm 11 in this case are (1, 2, 2, 3, 2, 3, 3, 3, 3, 3, 3, 6, 3).

Recall that the strongly connected components of  $(T)\lambda$  have representatives:  $\alpha_1 = (x_1)\lambda = \{1, \dots, 5\}$ ,  $\alpha_2 = (x_1x_3)\lambda = \{1, 2, 3\}$ ,  $\alpha_3 = (x_1x_3^2)\lambda = \{1, 3\}$ ,  $\alpha_6 = (x_1x_3^2x_1x_2x_3)\lambda = \{3\}$ , and sizes: 1, 1, 3, and 3,

respectively. We saw above that  $|T_{x_1}| = 12$ ,  $|T_{x_1x_3}| = 6$ ,  $|T_{x_1x_3^2}| = 2$ , and  $|T_{x_1x_3^2x_1x_2x_3}| = 1$ . It follows from Corollary 3.13(b) and (c) that

$$|T| = |R_{x_1}^T| + 3|R_{x_1x_3}^T| + 7|R_{x_1x_3^2}^T| + |R_{x_1x_3^2x_1x_2x_3}^T|$$
  
=  $(1 \cdot 1 \cdot 12) + (3 \cdot 1 \cdot 6) + (7 \cdot 3 \cdot 2) + (1 \cdot 3 \cdot 1) = 75.$ 

The orbit graph of  $\Re$  has 5 strongly connected components:

$$\{y_1 = x_1\}, \quad \{y_2 = x_3, \quad y_3 = x_2x_3, y_5 = x_1x_2x_3\}, \\ \{y_4 = x_3^2, y_6 = x_3x_2x_3, y_7 = x_2x_3^2, y_9 = (x_2x_3)^2, y_{10} = x_1x_2x_3^2, y_{12} = x_1(x_2x_3)^2\}, \\ \{y_8 = x_3x_1x_2x_3\}, \quad \{y_{11} = x^3x_1x_2x_3\}$$

and so there are five  $\mathscr{D}$ -classes in T.

## Optimizations for inverse semigroups

The semigroup S defined in (6.1) is an inverse semigroup, since the inverses of the generators can be obtained by taking powers. From Section 5.6, it follows that the  $\mathscr{R}$ -class representatives of S are in 1-1 correspondence with the values in  $(S)\lambda$  and that  $(S)\rho = (S)\lambda$ . Hence the number of  $\mathscr{R}$ -classes in S is 16, and by tracing the Schreier tree of  $(S)\lambda$ , the  $\mathscr{R}$ -class representatives are:

The strongly connected components of  $(S)\lambda$  are in 1-1 correspondence with the  $\mathscr{D}$ -classes of S, and so S has five  $\mathscr{D}$ -classes. Representatives of  $\mathscr{L}$ -classes can be obtained by taking the inverses of the  $\mathscr{R}$ -class representatives  $\mathfrak{R}$ .

It follows from Corollary 3.13(c) that

$$|S| = (1^2 \cdot 12) + (2^2 \cdot 6) + (3^2 \cdot 6) + (9^2 \cdot 1) + 1 = 172.$$

## Testing membership

In this subsection, we will use Algorithm 12 to test if the following transformations belong to the semigroup T defined in (6.2):

$$x = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 3 & 1 \end{pmatrix}$$
 and  $y = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 3 & 2 & 2 \end{pmatrix}$ .

Although  $(x)\lambda = \{1,2,3\} = \alpha_2 \in (T)\lambda$ ,  $(x)\rho = \{1,5|2|3,4\} \notin (T)\rho$ , and so  $x \notin T$ . Firstly,

$$(y)\lambda = \{2,3\} = \alpha_5 \in (T)\lambda$$

and so the representative of the s.c.c. of  $(y)\lambda$ , which we chose above, is  $\alpha_3$ . Tracing the Schreier tree for  $(T)\lambda$  from  $\alpha_5$  back to  $\alpha_3$ , using Algorithm 2, we find  $u=x_1x_2$  such that  $\alpha_3 \cdot u=\alpha_5$ . Since  $x_1, x_2$  are permutations,  $\overline{u}=x_2^{-1}x_1^{-1}$  has the property that  $\alpha_5 \cdot \overline{u}=\alpha_3$  and  $yu\overline{u}=y$ . The  $\mathscr{R}$ -class representative  $y_6 \in \mathfrak{R}$  is the only one such that  $(y_6)\lambda = \alpha_3 = (y)\lambda \cdot \overline{u}$  and  $(y_6)\rho = \{1,4,5|2,3\} = (y)\rho$ . Thus to check that  $y \in T$ , it suffices to show that the permutation  $(y_6'y\overline{u})\mu_x$  belongs to the group  $T_{y_6}=T_{x_1x_3^2}=\operatorname{Sym}(\{1,3\})$ . We know that  $(y_6'y\overline{u})\mu_x$  is a permutation on  $\{1,3\}$ , and so it must belong to  $T_{y_6}$ , and so  $y \in S$ .

### **Factorization**

In the previous subsection we showed that

$$y = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 3 & 2 & 2 \end{pmatrix}$$

is an element of T defined in (6.2). In this subsection, we will show how to use Algorithm 13 to factorize y as a product of the generators of T. Recall that we will write y = xsu, where  $x \in \Re$ ,  $u \in S$  is such

that  $(x)\lambda \cdot u = (y)\lambda$ , and  $s = x'y\overline{u}$ , and that we factorise each of x, s, u separately. From the previous subsection, the chosen  $\mathcal{R}$ -class representative for y is:

$$x = y_6 = x_3 x_2 x_3 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 1 & 3 & 3 \end{pmatrix},$$

and one choice for  $x' \in T_5$  is:

$$x' = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 2 & 1 & 4 & 5 \end{pmatrix}.$$

From the previous subsection,  $u = x_1 x_2$  and  $\overline{u} = x_2^{-1} x_1^{-1}$ . It follows that

$$s = x'y\overline{u} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 2 & 1 & 4 & 5 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 3 & 2 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 3 & 4 & 5 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 3 & 1 & 1 & 1 \end{pmatrix}$$

and so  $(s)\mu_x = s|_{im(x)} = s|_{\{1,3\}} = (1\ 3)$ , which is the only generator of  $T_x$ . From Algorithm 4, one choice for s such that  $s|_{im(x)} = (1\ 3)$  is  $x_3^2x_1x_2^2$ . Hence

$$y = xsu = x_3x_2x_3 \cdot x_3^2x_1x_2^2 \cdot x_1x_2 = x_3x_2x_3^3x_1x_2^2x_1x_2.$$

Note that  $x_3x_2x_3x_2^2$  is a minimal length word in the generators that is equal to y.

## The $\mathcal{D}$ -class structure

We showed above that the partial permutation semigroup S defined in (6.1) has five  $\mathscr{D}$ -classes  $D_1$ ,  $D_2$ ,  $D_3$ ,  $D_4$ , and  $D_5$  with representatives  $x_1$ ,  $x_1x_3$ ,  $x_1x_4$ ,  $x_1x_3x_4$ , and  $x_1x_3x_2x_3$ , respectively. The  $\mathscr{D}$ -class  $D_1$  has only one  $\mathscr{R}$ -class and one  $\mathscr{L}$ -class. If we left multiply the unique  $\mathscr{R}$ -class representative  $x_1$  of  $D_1$  by the generators of S, then we obtain the  $\mathscr{R}$ -class representatives:

$$x_1^2 \mathcal{D}^S x_1$$
,  $x_2 x_1 \mathcal{D}^S x_1$ ,  $x_3 x_1 = (2\ 5)(6) \mathcal{D}^S x_1 x_3$ ,  $x_4 x_1 = [1\ 8][2\ 4][3\ 6] \mathcal{D}^S x_1 x_4$ 

and so  $D_2, D_3 \leq_{\mathscr{J}} D_1$ . Note that, since S is inverse, we have not performed Algorithm 11, and so we have not previously left multiplied the  $\mathscr{R}$ -representatives of S by its generators.

Right multiplying the unique  $\mathscr{L}$ -class representative  $x_1$  of  $D_1$  by the generators of S we obtain:

$$x_1^2 \mathscr{D}^S x_1$$
,  $x_1 x_2 \mathscr{D}^S x_1$ ,  $x_1 x_3 = (2)(5 \ 6) \mathscr{D}^S x_1 x_3$ ,  $x_1 x_4 = [4 \ 3][6 \ 1][8 \ 2] \mathscr{D}^S x_1 x_4$ ,

which yields no additional information.

Continuing in this way, we obtain the partial order of the  $\mathscr{D}$ -classes of S. A picture of the egg-box diagrams of the  $\mathscr{D}$ -classes of S and the partial order of  $\mathscr{D}$ -classes of S can be seen in Figure 7. An analogous computation can be used to find the partial order of the  $\mathscr{D}$ -classes of the transformation semigroup T and this is included in Figure 7.

# 7 Experimental results

In this section we compare Algorithm 11, the optimized variants of Algorithm 11 for regular and inverse semigroups (described in Section 5.6) when they apply, and the Froidure-Pin algorithm [13] when computing the size of various standard examples; see Figures 8, 9, 10, 11, and 12. The three variants of Algorithm 11 used for this comparison are implemented in the GAP [15] package SEMIGROUPS [29], and the Froidure-Pin algorithm is implemented in LIBSEMIGROUPS [28]. The latter is a highly optimized C++ library, whereas the former is written largely in the high-level GAP programming language. If the three variants of Algorithm 11 were also implemented in C/C++, then a further improvement in performance could be expected.

For the partition monoid, full transformation monoid, and symmetric inverse monoid, each of which contain large subgroups, the performance of Algorithm 11 is considerably better than that of the Froidure-Pin algorithm. The monoid of order-preserving transformations on  $\{1,\ldots,n\}$  is  $\mathscr{H}$ -trivial, but far from  $\mathscr{J}$ -trivial, and Algorithm 11 is still faster than the Froidure-Pin algorithm in this case, albeit by a lesser margin. The monoid of unitriangular boolean matrices is  $\mathscr{J}$ -trivial, and in Figure 12, we see that, as expected, the Froidure-Pin algorithm outperforms our algorithms.

All of the computations in this section were performed using a 3.1 GHz Intel Core i7 processor with 16GB of RAM.




Figure 7: The partial order of the  $\mathscr{D}$ -classes of S (left) and T (right), group  $\mathscr{H}$ -classes indicated by shaded boxes.

n	$ P_n $	Algorithm 11	Algorithm 11 (regular)	Froidure-Pin
4	4 140	11	7	6
5	115 975	30	21	174
6	4 213 597	135	97	7 621
7	190 899 322	817	576	-
8	10 480 142 147	5 985	4 102	-
9	682 076 806 159	73 390	29.852	_

Figure 8: Time in milliseconds to find the size of the partition monoid  $P_n$ .

n	$ T_n $	Algorithm 11	Algorithm 11 (regular)	Froidure-Pin
5	3 125	6	5	2
6	46 656	13	9	62
7	823 543	35	20	989
8	16 777 216	130	79	21 430
9	387 420 489	691	374	-
10	10 000 000 000	4 135	$2\ 359$	-
11	285 311 670 611	31 699	$15\ 227$	-

Figure 9: Time in milliseconds to find the size of the full transformation monoid  $T_n$ .

n	$ I_n $	Algorithm 11	Algorithm 11 (inverse)	Froidure-Pin
6	13 327	10	7	10
7	130 922	14	10	151
8	$1\ 441\ 729$	24	14	2 137
9	$17\ 572\ 114$	33	20	28 151
10	234 662 231	57	33	_
11	$3\ 405\ 357\ 682$	110	54	-
12	53 334 454 417	210	97	_
13	896 324 308 634	418	180	_
14	$16\ 083\ 557\ 845\ 279$	858	351	_
15	306 827 170 866 106	1 748	710	_

Figure 10: Time in milliseconds to find the size of the symmetric inverse monoid  $I_n$ .

n	$ O_n $	Algorithm 11	Algorithm 11 (regular)	Froidure-Pin
6	462	8	7	1
7	1 716	14	14	2
8	6 435	29	26	10
9	24 310	61	59	28
10	92 378	141	109	145
11	352 716	291	242	611
12	1 352 078	631	523	2 639
13	5 200 300	1 380	1 116	11 074
14	20 058 300	3 120	2 542	$45\ 532$
15	77 558 760	7 246	5 649	-

Figure 11: Time in milliseconds to find the size of the monoid  $O_n$  of order-preserving transformations on  $\{1, \ldots, n\}$ .

n	$ U_n $	Algorithm 11	Froidure-Pin
3	8	32	0
4	64	593	1
5	1 024	-	215
6	32 768	-	7 820

Figure 12: Time in milliseconds to find the size of a transformation semigroup  $U_n$  isomorphic to the monoid of  $n \times n$  unitriangular boolean matrices.

## Acknowledgements

The authors would like to thank Wilf Wilson for his careful reading of several early drafts of this paper, and for the many improvements he suggested. We would also like to thank the two anonymous referees for their comments, in particular, for pointing out the "identity crisis" in the submitted version of the paper.

## References

- [1] Matrix group recognition project, https://www.math.auckland.ac.nz/~henrik/mgrp/, June 2008.
- [2] J. Araújo, P. V. Bünau, J. D. Mitchell, and M. Neunhöffer. Computing automorphisms of semigroups. J. Symbolic Comput., 45(3):373–392, 2010.
- [3] Martin Beaudry. Membership testing in commutative transformation semigroups. *Inform. and Comput.*, 79(1):84–93, 1988.
- [4] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [5] John J. Cannon. Computation in finite algebraic structures. Ph.D. thesis, University of Sydney, 1969.
- [6] John J. Cannon. Computing the ideal structure of finite semigroups. Numer. Math., 18:254–266, 1971/72.
- [7] J.-M. Champarnaud and G. Hansel. AUTOMATE, a computing package for automata and finite semigroups. J. Symbolic Comput., 12(2):197–220, 1991.
- [8] F. G. Cousineau, J. F. Perrot, and J.-M. Rifflet. Apl programs for direct computation of a finite semigroup. In *APL Congress*, volume 73, pages 67–74, Amsterdam, 1973. North-Holland Publishing Co.
- [9] J. East and R.D. Gray. Diagram monoids and Graham-Houghton graphs: idempotents and generating sets of ideals. *Preprint*, 2015, arXiv:1404.2359.
- [10] D. G. Fitz-Gerald. Computing the maximum generalized inverse of a Boolean matrix. *Linear Algebra and Appl.*, 16(3):203–207, 1977.
- [11] D.G. FitzGerald and K.W. Lau. On the Partition Monoid and some Related Semigroups. Bull. Aust. Math. Soc., 83(2):273–288, 2011.
- [12] George E. Forsythe. SWAC computes 126 distinct semigroups of order 4. *Proc. Amer. Math. Soc.*, 6:443–447, 1955.
- [13] Véronique Froidure and Jean-Eric Pin. Algorithms for computing finite semigroups. In *Foundations* of computational mathematics (Rio de Janeiro, 1997), pages 112–126. Springer, Berlin, 1997.
- [14] Harold N. Gabow. Path-based depth-first search for strong and biconnected components. *Information Processing Letters*, 74(34):107 114, 2000.
- [15] The GAP Group. GAP Groups, Algorithms, and Programming, Version 4.7.8, June 2015.
- [16] R. Gray and N. Ruškuc. Generating sets of completely 0-simple semigroups. *Comm. Algebra*, 33(12):4657–4678, 2005.
- [17] John M. Howie. Fundamentals of semigroup theory, volume 12 of London Mathematical Society Monographs. New Series. The Clarendon Press Oxford University Press, New York, 1995. Oxford Science Publications.
- [18] H. Jürgensen. Computers in semigroups. Semigroup Forum, 15(1):1–20, 1977/78.
- [19] Janusz Konieczny. Green's equivalences in finite semigroups of binary relations. Semigroup Forum, 48(2):235–252, 1994.

- [20] Gerard Lallement and Robert McFadden. On the determination of Green's relations in finite transformation semigroups. J. Symbolic Comput., 10(5):481–498, 1990.
- [21] Charles R. Leedham-Green. The computational matrix group project. In *Groups and computation*, *III (Columbus, OH, 1999)*, volume 8 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 229–247. de Gruyter, Berlin, 2001.
- [22] S. A. Linton, G. Pfeiffer, E. F. Robertson, and N. Ruškuc. *Monoid GAP 3 package, Version 2.4.* University of St Andrews, September 1997.
- [23] S. A. Linton, G. Pfeiffer, E. F. Robertson, and N. Ruškuc. Groups and actions in transformation semigroups. *Math. Z.*, 228(3):435–450, 1998.
- [24] S. A. Linton, G. Pfeiffer, E. F. Robertson, and N. Ruškuc. Computing transformation semigroups. J. Symbolic Comput., 33(2):145–162, 2002.
- [25] Stephen Lipscomb. Symmetric inverse semigroups, volume 46 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1996.
- [26] A. Markov. The impossibility of certain algorithms in the theory of associative systems. II. *Doklady Akad. Nauk SSSR (N.S.)*, 58:353–356, 1947.
- [27] D. B. McAlister. Semigroup for Windows. University of Northern Illinois, 2006.
- [28] J. D. Mitchell et al. libsemigroups C++ library version 0.0.3, December 2016.
- [29] J. D. Mitchell et al. Semigroups GAP package, Version 2.8.0, May 2016.
- [30] J. D. Mitchell and Markus Pfeiffer. Computing with semigroups of matrices over finite fields. 2015.
- [31] J Müller, F. Noeske, and M. Neunhöffer. Orb GAP package, Version 4.6, May 2013.
- [32] T. E. Nordahl and H. E. Scheiblich. Regular \*-semigroups. Semigroup Forum, 16(3):369-377, 1978.
- [33] Jan Okniński. Semigroups of matrices, volume 6 of Series in Algebra. World Scientific Publishing Co. Inc., River Edge, NJ, 1998.
- [34] Jean-François Perrot. Contribution à l'étude des monoïdes syntactiques et de certains groupes associés aux automates finis. Université de Paris VI, Paris, 1972. Thèse présentée à l'Université Paris VI, Paris, pour l'obtention du Doctorat d'État ès Sciences, Spécialité: Mathématiques.
- [35] Jean-Eric Pin. Semigroupe 2.01: a software for computing finite semigroups. Laboratoire d'Informatique Algorithmique: Fondements et Applications (LIAFA), CNRS et Université Paris 7, 2.01 edition, 2009.
- [36] Emil L. Post. Recursive unsolvability of a problem of Thue. J. Symbolic Logic, 12:1–11, 1947.
- [37] John Rhodes and Benjamin Steinberg. The q-theory of finite semigroups. Springer Monographs in Mathematics. Springer, New York, 2009.
- [38] Boris M. Schein. Regular elements of the semigroup of all binary relations. *Semigroup Forum*, 13(2):95–102, 1976/77.
- [39] Robert Sedgewick. *Algorithms*. Addison-Wesley Series in Computer Science. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, 1983.
- [40] Ákos Seress. Permutation group algorithms, volume 152 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2003.
- [41] Imre Simon. Piecewise testable events. In Automata theory and formal languages (Second GI Conf., Kaiserslautern, 1975), pages 214–222. Lecture Notes in Comput. Sci., Vol. 33. Springer, Berlin, 1975.
- [42] Charles C. Sims. Computational methods in the study of permutation groups. In *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, pages 169–183. Pergamon, Oxford, 1970.

- [43] W. A. Stein et al. Sage Mathematics Software (Version 5.12). The Sage Development Team, 2013. http://www.sagemath.org.
- [44] Robert Tarjan. Depth-first search and linear graph algorithms. SIAM J. Comput., 1(2):146–160, 1972.
- [45] S. Wilcox. Cellularity of Diagram Algebras as Twisted Semigroup Algebras. J. Algebra, 309(1):10–31, 2007.
- [46] D. Holt with B. Eick and E. O'Brien. *Handbook of computational group theory*. CRC Press, Boca Raton, Ann Arbor, London, Tokyo, 2004.